

REFERENCES

Afif, C.R. (2001):

Intravascular catheter-related infections.
In: D S, ed. Current therapy of infectious diseases.
St.Louis: Mosby, 416-418.

Anwar, H.; Strap, J.L. and Chen, K.(1992):

Dynamic interactions of biofilms of mucoid *Pseudomonas aeruginosa* with tobramycin and piperacillin.
Antimicrob. Agents Chemother. 36:1208-14.

Appelgren, P.; Hellstroem, I.; Weitzberg, E.; Soderlund, V. and Bindeslev, L. (2001):

Risk factors for nosocomial intensive care infection: a longterm prospective analysis.
Acta. Anaesthesiol. Scand. 45:710-719.

Arciola, C.R. (2006):

Details about the preparation of congo red agar media. E-mail sent by professor Carla Reneta Arciola. Personal communication.

Arciola, C.R.; Baldassarri, L. and Montanaro, L. (2001):

Presence of *icaA* and *icaD* genes and slime production in a collection of staphylococcal strains from catheter associated infections.
J. Clin. Microbio. 39:2151-2156.

Aygun, G.; Yasar, H.; Yilmaz, M.; Karasahin, K.; Dikmen, Y.; Polat, E.; Sidan, A. and Altas, K. (2006)

The value of Gram staining of catheter segments for rapid detection of peripheral venous catheter infections.
Diagn. Microbiol. Infect. Dis. 54: 165-167.

Assadian, O. (2004):

Skin antiseptic in reducing the risk of central venous catheter-related infections.
Crit. Care Med. 32:887–888.

Baird, D. (1996):

Staphylococcus: cluster-forming gram-positive cocci
In: Collee J. G.; Fraser A. G. ; Marmion B. P. and Simmons, A. (eds.)
Mackie & McCartney Practical Medical Microbiology. Churchill
Livingstone. P: 245-274.

Baron, E. J.; Peterson, L. R.; and Finegold, S. M. (1994):

Microorganisms encountered in the blood.
In: Bailey & Scott's Diagnostic Microbiology, ed 9. Mobsy, Missouri.
P: 193-208.

Bloemberg, G.V.; O'Toole, G.A.; Lugtenberg, B.J. and Kolter, R. (1997):

Green fluorescent protein as a marker for *Pseudomonas* spp.
Appl. Environ. Microbiol. 63:4543-4551.

Blot, F.; Schmidt, E.; Nitenberg, G.; Tancrede, G.; Leclercq, B.;**Laplanche, A. and Andreumont, A. (1998):**

Earlier positivity of central-venous- versus peripheral-blood cultures is
highly predictive of catheter-related sepsis.
J. Clin. Microbiol . 36:105-109.

Bouza, E., Burillo, A. and Munoz, P. (2002):

Catheter-related infections: diagnosis and intravascular treatment.
Clin. Microbiol. Infect. 8: 265–274

Brun-Buisson, C.; Abrar, F.; Legrand, P.; Huet, Y.; Larabi, S. and Rapin, M. (1987):

Diagnosis of central venous catheter related sepsis. Critical level of
quantitative tip cultures.
Arch. Intern. Med. 147: 873-877.

Burke, (1922):

Notes on Gram stain with description of a new method.
J. of Bacteriology 7:159.

Cappelli, G.; Tetta, C. and Canaud, B. (2005):

Is biofilm a cause of silent chronic inflammation in haemodialysis
patients? A fascinating working hypothesis.
Nephrol. Dial. Transplant. 19:1-4.

Carratala, J. (2002):

The antibiotic-lock technique for therapy of 'highly needed' infected catheters.

Clin. Microbiol. Infect. 8:282-289.

Casado-Flores, J.; Valdivielso-Serna, A.; Perez-Jurado, L.; Pozo-Román, J.; Monleón-Luque, M.; García-Pérez, J.; Ruiz-Beltran, A. and García-Teresa, M. (1991) :

Subclavian vein catheterization in critically ill children: Analysis of 322 cannulations.

Intensive Care Med. 17:350-354.

Chatzinikolaou, L. and Raad, I.(2006):

Catheter- related infections

Iranian Journal of Clinical Infectious Disease 1:5-10

Cheesbrough, M. (2000):

Culturing blood. In: District Laboratory Practice in Tropical Countries. Part 2. Cambridge University Press. P: 124-130.

Christensen, G.; Simpson, W.; Bisno, A. and Beachey, E. (1982):

Adherence of slime-producing strains of Staphylococcus epidermidis to smooth surfaces.

Infect. Immun. 37:318-326.

Civetta, J.; Civetta, J. and Bail, S. (1996):

Decreasing catheter-related infections and hospital costs by continuous quality improvement.

Crit. Care Med. 24: 1660-1665.

Collee, J. and Marr, W. (1996):

Specimen collection, culture containers and media

In: Collee J. G.; Fraser A. G.; Marmion, B.P.; et al. (eds.) Mackie & McCartney Practical Medical Microbiology ,14th edition. Churchill Livingstone 95-111.

Collee, J.; Miles, R. and Watt, B. (1996):

Tests for identification of bacteria.

In: Collee J. G.; Fraser, A. G.; Marmion B. P.; et al. (eds.). Mackie & McCartney Practical Medical Microbiology. 14th ed. Churchill Livingstone 131-150.

Collignon, P. (1994):

Intravascular catheter associated sepsis: a common problem. The Australian study on intravascular catheter associated sepsis.
Med. J. Aust. 161:374-378.

Collignon, P.; Chan, R. and Munro, R. (1987):

Rapid diagnosis of intravascular catheter-related sepsis.
Arch. Intern. Med. 147:1906

Costerton, B. and Lang, E.(1998):

Center for Biofilm Engineering (CBE), Montana State University,
www.erc.montana.edu.

Costerton, J. (1999):

Introduction to biofilm.
Int. J. Antimicrob. Agents; 11:217-221.

Costerton, J.; Stewart, P. and Greenberg, E. (1999):

Bacterial Biofilms: A Common Cause of Persistent Infections.
Science; 284: 1318 -1322

Costerton, W.; Veeh, R.; Shirtliff, M.; Pasmore, M.; Post, C. and Ehrlich, G. (2003):

The application of biofilm science to the study and control of chronic bacterial infections.
J. Clin. Invest. 112:1466-1477.

Coutlee, F.; Lemieux, C. and Paradis, F. (1988):

Value of direct catheter staining in the diagnosis of intravascular-catheter-related infection.
J. Clin. Microbiol. 26:1088-1090.

Cramton, S.; Ulrich, M.; Götz, F. and Doring, G. (2001) :

Anaerobic conditions induce expression of polysaccharide intercellular adhesin in *Staphylococcus aureus* and *Staphylococcus epidermidis*.
Infect. Immun. 43: 4079–4085.

Crnich, C. and Maki, D. (2002):

The promise of novel technology for the prevention of intravascular device-related bloodstream infection. Pathogenesis and short-term devices.
Clin. Infect. Dis.; 34:1232–1242.

Crump, J. and Collignon, P. (2000):

Intravascular catheter-associated infections.
Eur. J. Clin. Microbiol. Infect. Dis. 19:1-8.

Dahlberg, P.J.; Agger, W.A. and Singer, J.R. (1995):

Subclavian hemodialysis catheter infections: a prospective, randomized trial of an attachable silver impregnated cuff for prevention of catheter-related infections.
Infect. Control. Hosp. Epidemiol. 16:506–511.

Darouiche, R.O.; Raad, I.I. and Heard, S.O. (1999):

A comparison of two antimicrobial- impregnated central venous catheters. Catheter Study Group.
N. Engl. J. Med. 340:1–8.

DesJardin, J.; Falagas, M.; Ruthazer, R.; Griffith, J.; Wawrose, D.; Schenkein, D.; Miller, K. and Snyderman, D. (1999):

Clinical utility of blood cultures drawn from indwelling central venous catheters in hospitalized patients with cancer.
Ann. Intern. Med. 131:641-7.

Dimick, J. B.; Swoboda, S.; Talamini, M.A.; Pelz, R.K.; Hendrix, C.W. and Lipsett, P.A. (2003):

Risk of colonization of central venous catheters: catheters for total parenteral nutrition vs other catheters.
American Journal of Critical Care, 12: 328-335.

DK, H.(2000):

Infections due to percutaneous intravascular devices.
In: Mandell, G. B.J. and Dolin, R., (eds.) Principles and Practice of Infectious Diseases. Vol. 2. Philadelphia: Churchill Livingstone, P: 3005-3028.

Donlan, R.M. (2000):

Role of biofilms in antimicrobial resistance.
ASAIO J.; 46:547-552

Donlan, R.M. (2001) a :

Biofilms and device-associated infections.
Emer. Infect. Dis. 7:277-281.

Donlan, R.M. (2001) b:

Biofilm formation: a clinically relevant microbiological process.
Clin. Infect. Dis.; 33:1387-1392.

Donlan, R.M. (2002):

Biofilms: microbial life on surfaces.
Emerg. Infect. Dis.; 8:881-890.

Donlan, R.M. and Costerton, J.W. (2002):

Biofilms: survival mechanisms of clinically relevant microorganisms.
Clin. Microbiol. Rev.; 15:167-193.

Edmond, M.B.; Wallace, S.E.; McClish, D.K.; Pfaller M.A.; Jones, R.N. and Wenzel, R.P.(1999):

Nosocomial bloodstream infections in United States hospitals: a 3-year analysis.
Clin. Infect. Dis.; 29:239-244.

Estrada, B. (2002):

Catheter-related infections in children: an update.
Infect. Med.; 19:541.

Fan, S.T.; Teoh-Chan, C.H. and Lau, K.F. (1989):

Evaluation of central venous catheter sepsis by differential quantitative blood culture.
Eur. J. Clin. Microbiol. Infect. Dis.; 8:142-4.

Farmer, J.J. (1999):

Enterobacteriaceae: Introduction and Identification.
In: Patrick, R.M.; Boron, E.J.; Pfaller, M.A.; Tenover, F.C. and Tenover, R.H. (eds.) Manual of Clinical Microbiology. 7th ed. American Society for Microbiology. P: 442-446.

Fatkenheuer, G.; Cornely, O. and Seifert, H. (2002):

Clinical management of catheter-related infections.
Clin. Microbiol. Infect. 8:545-50.

Fey, P.D.; Ulphani, J.S.; Götz, F.; Heilmann, C.; Mack, D. and Rupp, M.E. (1999):

Characterization of the relationship between polysaccharide intercellular adhesin and hemagglutination in *Staphylococcus epidermidis*.
J. Infect. Dis .43: 1561–1564.

Fidalgo, S.; Vázquez, F.; Mendoza, M.C.; Perez, F. And Méndez, F.J. (1990):

Bacteremia due to *Staphylococcus epidermidis*: microbiological, epidemiological, clinical, and prognostic features.
Rev. Infect. Dis. 12:520-8.

Flemming, H.C.; Wingender, J. and Mayer, C. (2000):

Physico-chemical properties of biofilms.
In: Evans, L.V. (ed.) Biofilms: recent advances in their study and control.
Amsterdam: Harwood Academic Publishers; P: 19–34.

Forbes, B.A.; Sahm, D.F. and Weissfeld, J. R. (1998):

Overview of bacterial identification, methods and strategies.
In: Belly, A. and Forbes, B.A. (eds.). Baily and Scott's Diagnostic Microbiology. 10th ed. New York: Mosby. P: 427-442.

Freeman, D.J.; Falkiner, F.R. and Keane, C.T. (1989):

New method for detecting slime production by coagulase negative staphylococci.
J. Clin. Pathol. 42:872-874.

Friedman, N.B.; Kaye, K.S. and Stout, J.E. (2002):

Health care–associated bloodstream infections in adults.
Ann. Intern. Med. 137:791-797.

Furfaro, S.; Gauthier, M.; Lacroix, J.; Nadeau, D.; Lafleur, L. and Mathews, S. (1991):

Arterial catheter-related infections in children: a 1-year cohort analysis.
Am. J. Dis. Child. 145:1037–1043.

Fux, C.A.; Stoodley, P.; Hall-Stoodley, L. and Costerton, J.W. (2003):

Bacterial biofilms: a diagnostic and therapeutic challenge.
Expert Rev. Anti. Infect. Ther. 1:667-683.

Fux, C.A.; Wilson, S. and Stoodley, P. (2004):

Detachment characteristics and oxacillin resistance of *Staphylococcus aureus* biofilm emboli in an in vitro catheter infection model.
J. Bacteriol. 186:4486-4491.

Garland, J.S.; Dunne, W.M.; Havens, P.; Hintermeyer, M.; Bozzette, M.A.; Wincek, J.; Bromberger, T. and Seavers, M. (1992):

Peripheral intravenous catheter complications in critically ill children: a prospective study.
Pediatrics 89:1145–1150.

Garner, J.S.; Jarvis, W.R.; Emori, T.G.; Horan T.C. and Huges, T.M. (1988):

CDC definition for nosocomial infections.
Am. J. Infect. Control. 16:128-140.

Garner, J.S.; Jarvis, W.R.; Emori, T.G.; Horan T.C. and Huges, T.M. (1996):

CDC definition for nosocomial infections.
In: Olmsted, R.N.(ed.) Infection Control and Applied Epidemiology: Principle and Practice. St Louis, MO: Mosby, P: 1–20.

Gaynes, R. (2002):

Health care–associated bloodstream infections: a change in thinking
Ann. Intern. Med. 137:850-851.

Gilbert, P.; Maira-Litran, T.; McBain, A.J.; Rickard, A.H. and Whyte, F.W. (2002):

The physiology and collective recalcitrance of microbial biofilm communities.
Adv. Microbial. Physiol. 46:203-256.

Goldmann, D.A. and Pier, G.B.(1993):

Pathogenesis of infections related to intravascular catheterization.
Clin. Microbiol. Rev. 6: 176-192.

Götz, F. (2002):

Staphylococcus and biofilms
Mol. Microbiol. 43:1367-1376.

Grupo de Hospitales Castrillo (2000):

Prospective evaluation of umbilical catheters in newborn infants.
An. Esp. Pediatr. 53: 470 – 478.

Hoiby, N.; Johansen, H.K.; Moser, C.; Song, Z.; Oana, C. and Arsalan, K. (2001):

Pseudomonas aeruginosa and the in vitro and in vivo biofilm mode of growth.
Microbes Infect. 3:23-35.

Hussain, M.; Herrmann, M.; Peters, G. Perdreau-Remington, F. and Peters, G. (1997):

A 140-kilodalton extracellular protein is essential for the accumulation of *Staphylococcus epidermidis* strains on surfaces.

Infect. Immun. 65:519-24.

Jeremy, M.Y.; Douglas, J.B.; Esther, M.V. and Greenberg, E.P. (2004):

Quorum sensing in *staphylococcus aureus* biofilms.

J.Bacteriol. 186:1838-1850.

Kacica, M.A.; Horgan; M.J.; Ochoa, L.; Sandler, R.; Lepow, M.L. and Venezia, R.A. (1993):

Prevention of gram-positive sepsis in neonates weighing less than 1500 g.

J. Pediatr. 125:253–258.

Karchmer, A.W.(2000):

Nosocomial bloodstream infections: organisms, risk Factors, and implications.

Clin. Infect. Dis. 31:139- 143.

Keren, I.; Shah, D.; Spoering, A.; Kaldalu N. and Lewis, K. (2004):

Specialized persister cells and the mechanism of multidrug tolerance in *Escherichia coli*.

J. Bacteriol. 186:8172-8180.

Kiem, S.; Oh, S.W.; Peck, K.R.; Yong, N.L.; Young, J.L.; Hoon, J.S. and Soo, E.H.:(2004):

Phase variation of biofilm formation in *staphylococcus aureus* by IS256 Insertion and its impact on the capacity adhering to polyurethane surface.

J. Korean Med. Sci. 19: 779-782.

Kite, P.; Eastwood, K.; Sugden, S. and Percival, S. L. (2004):

Use of in vivo-generated biofilms from hemodialysis catheters to test the efficacy of a novel antimicrobial catheter lock for biofilm eradication in vitro.

J.Clin. Microbiol. 42:3073-3076.

Kline, A.M. (2005):

Pediatric catheter-related bloodstream infections, latest strategies to decrease risk.

AACN Clinical Issues 16: 185–198.

Knobloch, J.K.; Bartscht, K.; Sabottke, A.; Rohde, H.; Feucht, H. and Mack, D. (2001):

Biofilm formation by *Staphylococcus epidermidis* depends on functional RsbU, an activator of the *sigB* operon: differential activation mechanisms due to ethanol and salt stress.
J. Bacteriol. 43: 2624–2633.

La Quaglia, M.; Lucas, A.; Thaler, H.; Friedlander-Klar, H. and Exelby, P.R.; Groeger, J.S. (1992):

A prospective analysis of vascular access device-related infections in children.
J. Pediatr. Surg. 27:840-842.

Lakshmi, K.S.; Jayashree, M.; Singhi, S. and Ray, P. (2007):

Study of nosocomial primary bloodstream infections in a pediatric intensive care unit.
Journal of Tropical Pediatrics 53:87-92.

Landers, S.; Moise, A.A.; Fraley, J.K.; Smith, E.O. and Baker, C.J.(1991):

Factors associated with umbilical catheter related sepsis in neonates.
Am. J. Dis. Child. 145:675–80.

Lauderdale, T. L.; Kimberle, C. C. and Murray P. R. (1999):

Reagents, stains and media.
In: Patrick, R.M.; Boron, E.J.; Pfaller, M.A.; Tenover, F.C. and Tenover, R.H. (eds.) Manual of Clinical Microbiology. 7th ed. American Society for Microbiology. P: 1663-1707.

Lauer, B. A.; Reller, L. B. and Mirrett, S. (1981):

Comparison of acridine orange and Gram stains for detection of microorganisms in cerebrospinal fluid and other clinical specimens.
J. Clin. Microbiol. 14:201-205.

Leid, J.G.; Shirtliff, M.E.; Costerton, J.W. and Stoodley, A.P.(2002):

Human leukocytes adhere to, penetrate, and respond to *Staphylococcus aureus* biofilms.
Infect. Immun. 70:6339-6345.

Lewis, K.(2000):

Programmed death in bacteria.
Microbiol. Mol. Biol. Rev. 64:503-514.

Lewis, K.(2001):

Riddle of biofilm resistance
Antimicrob. Agents and Chemother. ; 45: 999-1007.

Ludo, M.M.; Jozef, D.J. and Aimé, D.O. (2001):

Microbiology and risk factors for catheter exit-site and -Hub colonization in neonatal intensive care unit patients.
Infect. control. hosp. epidemiol. 22: 357-362.

Maki, D.G. and Mermel, L.A. (1991):

Infections due to infusion therapy. In: Bennett, J.V.and Brachman, P.S. (eds). Hospital Infections. 4th ed. Philadelphia: Lippencott-Raven, P: 689–724.

Maki, D.G. and Ringer, M. (1987):

Evaluation of dressing regimens for prevention of infection with peripheral intravenous catheters.
J. Am. Med. Assoc. 258:2396-2403.

Maki, D.G.; Ringer, M. and Alvarado, C.J. (1991):

Prospective randomized trial of povidone-iodine, alcohol, and chlorhexidine for prevention of infection associated with central venous and arterial catheters.
Lancet 338:339–43.

Maki, D.G.; Weise, C. and Sarafin, H. (1977):

A semiquantitative culture method for identifying intravenous-catheter-related infection.
N. Engl. J. Med. 296: 1305–1309.

Manzoni, P.; Farina, D.; Leonessa, M.; Antoniellid'Oulx, E.; Galletto, P.; Mostert,M.; Miniero, R. and Gomirato, G.(2006):

Risk factors for progression to invasive fungal infection in preterm neonates with fungal colonization.
Pediatrics. 118:2359-2364.

Mathur, T.; Singhal, S.; Khan, S.; Upadhyay, D.J.; Fatma, T. and Rattan, A. (2006):

Detection of biofilm formation among the clinical isolates of staphylococci: An evaluation of three different screening methods.
Indian Journal of Medical Microbiology 24: 25-29.

McKenney, D.; Pouliot, K.L.; Wang, Y.; Murthy, V.; Ulrich, M.; Döring, G.; Lee, J.C.; Goldmann, D.A. and Pier, G.B. (1999):

Broadly protective vaccine for *Staphylococcus aureus* based on an *in vivo*-expressed antigen.
Science 43: 1523–1527.

McKinley, S.; Mackenzie, A.; Finfer, S.; Ward, R. and Penford, J. (1999):

Incidence and predictors of central venous catheter related infection in intensive care patients.
Anaesth. Intens. Care 27:164-169.

Mermel, L.A.; Farr, B.M.; Sherertz, R.J.; Raad, I.I; O’Grady, N.; Harris, J.S. and Craven, D.E. (2001):

Guidelines for the management of intravascular catheter-related infections.
Clin. Infect. Dis.; 32:1249-1272.

Miles, R. and Amyes, S. (1996):

Laboratory control of antimicrobial therapy. In: Collee, J. G.; Fraser, A. G.; Marmion, B.P. and Simmons, A. (eds.) Mackie & McCartney Practical Medical Microbiology. Churchill Livingstone. P: 151-178.

Møretrø, T.; Hermansen, L.; Holck, A.L.; Sidhu, M.S.; Rudi, K. and Langsrud, S. (2003):

Biofilm formation and the presence of the intercellular adhesion locus *ica* among staphylococci from food and food processing environments.
Appl. Environ. Microbiol. 69: 5648-5655.

Moro, M.L.; De Toni, A.; Stolfi, I.; Carrieri, M.P.; Braga, M. and Zunin, C. (1996):

Risk factors for nosocomial sepsis in newborn intensive and intermediate care units.
Eur. J. Pediatr. 155:315-322.

Mueller-Premru, M.; Gubina, M.; Kaufmann, M.E.; Primožic, J. and Cookson, B.D. (1999):

Use of semi-quantitative and quantitative culture methods and typing for studying the epidemiology of central venous catheter-related infections in neonates on parenteral nutrition.
J. Med. Microbiol. 48: 451-460.

Mulloy, R.H.; Jadavji, T. and Russell, M.L. (1991):

Tunneled central venous catheter sepsis: Risk factors in a pediatric hospital.

J. Parenter. Enteral Nutr. 15:460-463.

Nasia, S.; Jason, P.F. and Dennis, G.M. (2005):

Meta-Analysis: Methods for Diagnosing Intravascular Device–Related Bloodstream Infection

Ann. Intern. Med.142:451-466.

Odetola, F.O.; Moler, F.W.; Dechert, R.E.; VanDerElzen, K. and Chenoweth, C. (2003):

Nosocomial catheter-related bloodstream infections in a pediatric intensive care unit: Risk and rates associated with various intravascular technologies.

Pediatr. Crit. Care Med. 4:432-436.

O'Grady, N.P.; Alexander, M.; Dellinger, E.P.; Gerberding, J.L.; Heard, S.O.; Maki, D.G.; Masur, H.; McCormick, R.D.; Mermel, L.A.; Pearson, M.L.; Raad, I.I.; Randolph, A. and Weinstein, R.A. (2002):

Guidelines for the prevention of intravascular catheter-related infections. Centers for Disease Control and Prevention.

MMWR Recomm. Rep. 51:1-29

Öncü, S.; Özsüt, H.; Yildirim, A.; Ay, P.; Cakar, N.; Eraksoy, H. and Calangu, S. (2003):

Central venous catheter related infections: Risk factors and the effect of glycopeptide antibiotics.

Ann. Clin. Microbiol. Antimicrob. 2: 3-9.

Oncü, S. and Sakarya, S.(2003):

Central venous catheter - related infections: an overview with special emphasis on diagnosis, prevention and management.

The Internet Journal of Anesthesiology.

O'Toole, G. (2003):

To build a biofilm.

J. Bacteriol. 185:2687-2689.

Paragioudaki, M.; Stamouli, V.; Kolonitsiou, F.; Anastassiou, E.D.; Dimitracopoulos, G. and Spiliopoulou, I. (2004):

Intravenous catheter infections associated with bacteraemia: a 2-year study in a University Hospital.

Clin. Microbiol. Infect. 10: 431–435.

Parsek, M.R. and Singh, P.K. (2003):

Bacterial biofilms: an emerging link to disease pathogenesis.

Annu. Rev. Microbiol. 57:677-701.

Pascual, A. (2002):

Pathogenesis of catheter-related infections: lessons for new designs.

Clin. Microbiol. Infect. 8:256-64.

Pelletier, S.J.; Crabtree, T.D. and Gleason, T.G. (2000):

Bacteremia associated with central venous catheter infection is not an independent predictor of outcomes.

J. Am. Coll. Surg. 190:671-680.

Pittet, D.; Li, N.; Woolson, R.F. and Wenzel, R.P. (1997):

Microbiological factors influencing the outcome of nosocomial bloodstream infections: a 6-year validated, population-based model.

Clin. Infect. Dis. 24:1068-78.

Pittet, D.; Tarara, D. and Wenzel, R.P. (1994):

Nosocomial bloodstream infection in critically ill patients: Excess length of stay, extra costs, and attributable mortality.

JAMA 271: 1598-1601.

Pratten, J.; Foster, S.J.; Chan, P.F.; Wilson, M. and Nair, S.P. (2001):

Staphylococcus aureus accessory regulators: expression within biofilms and effect on adhesion.

Microbes Infect. 43: 633–637.

Quirynen, M.; Brex, M. and Steenberghe, D. (2000):

Biofilms in the oral cavity: impact of surface characteristics.

In: Evans, L.V. (ed.) Biofilms: recent advances in their study and control. Amsterdam: Harwood Academic Publishers. P: 167-187.

Raad, I.I. and Bodey, G.P. (1992):

Infectious complications of indwelling vascular catheters.

Clin. Infect. Dis. 15:197-208.

Raad, I.I. and Darouiche, R.O. (1996):

Catheter-related septicemia: Risk reduction.

Infect. Med. 13: 807–823.

Raad, I.I.; Hohn, D.C. and Gilbreath, B.J. (1994):

Prevention of central venous catheter-related infections by using maximal sterile barrier precautions during insertion.

Infect. Control. Hosp. Epidemiol. 15:231–238.

Rachid, S.; Ohlsen, K.; Witte, W.; Hacker, J. and Ziebuhr, W. (2000):

Effect of subinhibitory antibiotic concentrations on polysaccharide intercellular adhesin expression in biofilm-forming *Staphylococcus epidermidis*.

Antimicrob. Agents Chemother. 43: 3357–3363.

Rani SA, Pitts B and Stewart PS, (2005):

Rapid diffusion of fluorescent tracers into *Staphylococcus epidermidis* biofilms visualized by time lapse microscopy.

Antimicrob. Agents Chemother., 49:728-732 .

Rao, S.D.; Preetha, M.J.; Lavi, R. and Macaden, R. (2005):

Infections related to vascular catheters in a pediatric intensive care unit

Indian Pediatrics. 42:667-672 .

Raul, R.M. and Melvin, P.W. (2001):

Laboratory diagnosis of bacteremia and fungemia.

Infect. Dis. Clin. North Am. 15: 1-25.

Richet, H.; Hubert, B. and Nitemberg, G. (1990):

Prospective multicenter study of vascular-catheter-related complications and risk factors for positive central- catheter cultures in intensive care unit patients.

J. Clin. Microbiol. 28: 2520-2525.

Roberts, M.E. and Stewart, P.S.(2005):

Modeling protection from antimicrobial agents in biofilms through the formation of persister cells.

Microbiology 151:75-80.

Rowin, M.E.; Patel, V.V. and Christenson, J.C. (2003):

Pediatric intensive care unit nosocomial infections.

Crit. Care Clin. 19:473–487.

Ryder, M.A. (2001):

The role of biofilm in vascular catheter-related infections.
N. Dev. Vasc. Dis. 2:15-25.

Ryder, M.A. (2005):

Catheter-related infections: it's all about biofilm.
Topics in advanced practice nursing e Journal, 5.

Sachdev, A.; Gupta, D. and Soni, A. (2002):

Central venous catheter colonization and related bacteremia in pediatric intensive care unit.
Indian Pediatrics 39: 752-760.

Sadoyama, G. and Pinto, P.G. (2003):

Comparison between the Jugular and Subclavian Vein as Insertion Site for Central Venous Catheters: Microbiological Aspects and Risk Factors for Colonization and Infection.
BJID 7:142-148.

Safdar, N.; Jason, P. and Dennis, G.(2005):

Meta-analysis: methods for diagnosing intravascular device-related bloodstream infection.
Ann. Intern. Med. 142:451-466.

Sahm, D.F.; Marsilio, M.K. and Piazza, G.(1999):

Antimicrobial resistance in key bloodstream bacterial isolates: electronic surveillance with the Surveillance Network Database USA.
Clin. Infect. Dis. 29:259-263.

Saiman, L.; Ludington, E. and Dawson, J.D. (2001):

Risk factors for Candida species colonization of neonatal intensive care unit patients.
Pediatr .Infect .Dis. 20:1119–1124.

Shama, A.; Patole, S.K. and Whitehall, J.S.(2002):

The dilemma of removing umbilical venous catheters in high-risk neonates with nosocomial sepsis.
Ind. Ped. 39:209.

Shanks, R.M.; Donegan, N.P. and Graber, M.L. (2005):

Heparin stimulates staphylococcus aureus biofilm formation infection and Immunity.

Infect. Immun. 73: 4596-4606.

Sherertz, R.J. Heard, S.O. and Raad, I.I., (1997):

Diagnosis of triple-lumen catheter infection: comparison of roll plate, sonication, and flushing methodologies.

J. Clin. Microbiol. , 35:641-646.

Shimandle, R. B.; Johnson, D. and Baker M. (1999):

Safety of peripheral intravenous catheters in children.

Infect. Control Hosp. Epidemiol. 20:736-740.

Singh-Naz, N.; Sprague, B.M.; Patel, K.M. and Pollack, M.M. (1996):

Risk factors for nosocomial infection in critically ill children: a prospective cohort study.

Crit. Care Med. 24:875–878.

Slaughter, S.E. (2004):

Intravascular catheter-related infections: strategies for combating this common foe.

Postgrad. Med. 116:59-66.

Slonim, A.D.; Kurtines, H.C. and Sprague, B.M. (2001):

The costs associated with nosocomial bloodstream infections in the pediatric intensive care unit.

Pediatr. Crit. Care Med.; 2: 170–174.

Smith, T.L. ;Pearson,M.L. and Wilcox, K.R. (1999):

Emergence of vancomycin resistance in *Staphylococcus aureus*.

N. Engl. J. Med.; 340:493-501.

Soifer, N.E.; Borzak, S.; Edlin, B.R. and Weinstein, R.A. (1998):

Prevention of peripheral venous catheter complications with an intravenous therapy team: a randomized controlled trial.

Arch. Intern. Med. 158:473–477.

Spafford, P.S.; Sinkin, R.A.; Cox, C.; Reubens, L. and Powell, K.R. (1994):

Prevention of central venous catheter-related coagulase-negative staphylococcal sepsis in neonates.

J. Pediatr.125:259–263.

Stenzel, J.P.; Green, T.P. and Fuhrman, B.P.(1989):

Percutaneous central venous catheterization in a pediatric intensive care unit: a survival analysis of complications.
Crit. Care Med.; 17:984–988.

Stewart, P.S. (2002):

Mechanisms of antibiotic resistance in bacterial biofilms.
Int. J. Med. Microbiol.; 292:107-113.

Stewart, P.S. and Costerton, J.W. (2001):

Antibiotic resistance of bacteria in biofilms.
Lancet. , 358:135-138.

Stewart, P.S. and Dirckx, P. (2001):

Center for Biofilm Engineering (CBE), Montana State University.
www.erc.montana.edu.

Stickler, D.J. (1996):

Bacterial biofilms and the encrustation of urethral catheters.
Biofouling 94:293-305.

Stickler, D.J.; Morris, N. and Moreno, M.C.(1998):

Studies on the formation of crystalline bacterial biofilms on urethral catheters.
Eur. J. Clin. Microbiol. Infect. Dis. 17:649-652.

Stoodley, P. and Dirckx, P. (2003)

Center for Biofilm Engineering (CBE), Montana State University.
www.erc.montana.edu.

Swinscow, T.D.V. (1997)

Statistics at Square One.
BMJ Publishing Group

Tunney, M.M.; Jones, D.S. and Gorman, S.P.(1999):

Biofilm and biofilm-related encrustations of urinary tract devices.
In: Doyle RJ, ed. Methods in enzymology, vol. 310. Biofilms. San Diego: Academic Press, 558–66.

Vandecasteele, S.J.; Peetermans, W.E. and Merckx, R. (2003):

Expression of Biofilm-Associated Genes in *Staphylococcus epidermidis* during In Vitro and In Vivo Foreign Body Infections
J. Infect. Dis. 188:730-737.

Vuong, C.; Saenz, H.L.; Götz, F. and Otto, M. (2000):

Impact of the *agr* quorum-sensing system on adherence to polystyrene in *Staphylococcus aureus*.

J. Infect. Dis 43: 1688–1693.

Wilson, J.N.; Grow, J.B.; Demong, C.V.; Prevedel, A.E. and Owens, J.C. (1962):

Central venous pressure in optimal blood volume maintenance.

Arch. Surg. 85:563–578.

Woznicova, V.; Votava, M. and Skalka, B. (1993):

Comparison of 2 methods of detecting slime production by coagulase-negative staphylococci.

Cesk. Epidemiol. Mikrobiol. Imunol. , 42:51-53.

Yogaraj, J.S.; Elward, A.M. and Fraser, V.J. (2002):

Rate, risk factors, and outcomes of nosocomial primary bloodstream infection in pediatric intensive care unit patients.

Pediatrics.110:481–485.

Zufferey, J.; Rime, B. and Francioli P. (1988):

Simple method for rapid diagnosis of catheter-associated infection by direct acridine orange staining of catheter tips.

J. Clin. Microbiol. 26:175-177.