

REFERENCES

REFERENCES

- Abbott M.W. and Williams M.M. (2006):**
Postnatal depressive symptoms among Pacific mothers in Auckland: prevalence and risk factors. *Aust N Z J Psychiatry*. Mar;40(3):230-8.
- Abiodun O.A.(2006):**
Postnatal depression in primary care populations in Nigeria. *Gen Hosp Psychiatry*. 2006 Mar-Apr;28(2):133-6.
- Adewuya A.O. and Afolabi O.T.(2005):**
The course of anxiety and depressive symptoms in Nigerian postpartum women. *Arch Womens Ment Health*. 2005 Nov;8(4):257-9. Epub 2005 Jun 17.
- Adewuya A.O., (2002):**
Prevalence and correlates of depression in late pregnancy among nigerian women. *Depress Anxiety*. Jul 14; [Epub ahead of print] *Depression and Anxiety* 0:1-7.
- Adewuya A.O.(2005):**
The maternity blues in Western Nigerian women: prevalence and risk factors. *Am J Obstet Gynecol*. 2005 Oct;193(4):1522-5.
- Adewuya A.O., Fatoye F.O., Ola B.A., et al., (2005):**
Sociodemographic and obstetric risk factors for postpartum depressive symptoms in Nigerian women. *J Psychiatr Pract*. 2005 Sep;11(5):353-8.
- Adouard F., Glangeaud-Freudenthal N.M., Golse B.(2005):**
Validation of the Edinburgh postnatal depression scale (EPDS) in a sample of women with high-risk pregnancies in France. *Arch Womens Ment Health*. Jun;8(2):89-95. Epub 2005 May 11.
- Agoub M., Moussaoui D., Battas O.(2005):**
Prevalence of postpartum depression in a Moroccan sample. *Arch Womens Ment Health*. 2005 May;8(1):37-43. Epub May 4.
- Alder E.M.,and Cox J.L.(1983):**
Breast feeding and post-natal depression. *J Psychosom Res*. 1983;27(2):139-44.
- Al-Nuaim L.A.(2004):**
Views of women towards cesarean section. *Saudi Med J*. 2004 Jun;25(6):707-10. Comment in: *Saudi Med J*. 2004 Nov;25(11):1774; author reply 1774-5.
- Alvarado R., Perucca E., Neves E.,et al.,(1993):**
Depressive disorders during pregnancy and associated factors. *Rev Chil Obstet Ginecol*. 1993;58(2):135-41.

Andrews-Fike C.(1999):

A review of Postpartum Depression. Prim Care Companion J Clin Psychiatry. 1999 Feb;1(1):9-14.

Anoop S., Saravanan B., Joseph A., et al., (2004):

Maternal depression and low maternal intelligence as risk factors for malnutrition in children: a community based case-control study from South India. Arch Dis Child. 2004 Apr;89(4):325-9.

Appleby L., Gregoire A., Platz C.,et al.,(2004):

Screening women for high risk of postnatal depression. Rinsho. Aug;59(8):1555-9.

Asher I., Kaplan B., Modai I., et al., (1995):

Mood and hormonal changes during late pregnancy and puerperium.Clin Exp Obstet Gynecol.;22(4):321-5.

Astbury J., Brown S., Lumley J., et al., (1994):

Birth events, birth experiences and social differences in postnatal depression. Aust J Public Health. 1994 Jun;18(2):176-84. Erratum in: Aust J Public Health Oct;19(5):524.

Austin M.P.(2004):

Antenatal screening and early intervention for "perinatal" distress, depression and anxiety: where to from here? Arch Womens Ment Health. 2004 Feb;7(1):1-6. Epub 2003 Dec 15.

Austin M.P., Hadzi-Pavlovic D., Saint K., et al., (2005):

Antenatal screening for the prediction of postnatal depression: validation of a psychosocial Pregnancy Risk Questionnaire. Acta Psychiatr Scand.Oct;112(4):310-7.

Austin M.P., Hadzi-Pavlovic D., Leader L., et al., (2005):

Maternal trait anxiety, depression and life event stress in pregnancy: relationships with infant temperament. Early Hum Dev. Feb;81(2):183-90.

Austin M.P.and Lumley J.(2003):

Antenatal screening for postnatal depression: a systematic review. Acta Psychiatr Scand. 2003 Jan;107(1):10-7. Comment in: Evid Based Ment Health. 2003 Aug;6(3):77.

Aydin N., Inandi T., Karabulut N.(2005):

Depression and associated factors among women within their first postnatal year in Erzurum province in eastern Turkey. Women Health ;41(2):1-12.

Azidah A.K, Shaiful B.I, Rusli N., et al., (2006):

Postnatal depression and socio-cultural practices among postnatal mothers in Kota Bahru,Malaysia.Med J Malaysia Mar;61(1):76-83.

Bacchus L., Mezey G., Bewley S.(2004):

Domestic violence: prevalence in pregnant women and associations with physical and psychological health. *Eur J Obstet Gynecol Reprod Biol*, Mar 15;113(1):6-11.

Baker L., Cross S., Greaver L., et al., (2005):

Prevalence of postpartum depression in a native American population. *Matern Child Health J*. 2005 Mar;9(1):21-5. Comment in: *Matern Child Health J*. Mar;9(1):1-2.

Beck C.T.(2001):

Predictors of postpartum depression: an update. *Nurs Res*. 2001 Sep-Oct;50(5):275-85.

Beck C.T.(2002):

Revision of the postpartum depression predictors inventory. *J Obstet Gynecol Neonatal Nurs*. 2002 Jul-Aug;31(4):394-402.

Beck C.T.(2006):

Postpartum depression: it isn't just the blues. *Am J Nurs*. 2006 May;106(5):40-50; quiz 50-1.

Beck C.T.and Indman P.(2005):

The many faces of postpartum depression. *J Obstet Gynecol Neonatal Nurs*. 2005 Sep-Oct;34(5):569-76.

Beeghly M., Olson K.L., Weinberg M.K., et al., (2003):

Prevalence, stability, and socio-demographic correlates of depressive symptoms in Black mothers during the first 18 months postpartum. *Matern Child Health J*. Sep;7(3):157-68.

Beeghly M., Weinberg M.K., Olson K.L., et al., (2002):

Stability and change in level of maternal depressive symptomatology during the first postpartum year. *J Affect Disord*. Sep;71(1-3):169-80.

Bellantuono C., Migliarese G., Imperadore G.(2006):

Pharmacologic therapy of depression during pregnancy. *Recenti Prog Med*. Feb;97(2):94-107.

Berle J.O. and Spigset O.(2003):

Psychiatric disorders during pregnancy and lactation. *Tidsskr Nor Laegeforen*. Aug 14;123(15):2037-40.

Berle J.O., Aarre T.F., Mykletun A., et al., (2003):

Screening for postnatal depression. Validation of the Norwegian version of the Edinburgh Postnatal Depression Scale, and assessment of risk factors for postnatal depression. *J Affect Disord*. Sep;76(1-3):151-6.

Bernazzani O., Marks M.N., Bifulco A., et al., (2005):

Assessing psychosocial risk in pregnant/postpartum women using the Contextual Assessment of Maternity Experience (CAME)--recent life adversity, social support and maternal feelings. *Soc Psychiatry Psychiatr Epidemiol*. Jun;40(6):497-508.

Binetti P.(2006):

Therapeutic education and the increased risk of postpartum depression. Clin Ter. Mar-Apr;157(2):111-6.

Blackmore E.R., Jones I, Doshi M., et al., (2006):

Obstetric variables associated with bipolar affective puerperal psychosis. Br J Psychiatry. Jan;188:32-6.

Blenning C.E and Paladine H.(2005):

An approach to the postpartum office visit. Am Fam Physician. 2005 Dec 15;72(12):2491-6. Comment in: Am Fam Physician. Dec 15;72(12):2443-4.

Bloch M., Daly R.C., Rubinow D.R.(2003):

Endocrine factors in the etiology of postpartum depression. Compr Psychiatry. May-Jun;44(3):234-46.

Bloch M., Rotenberg N., Koren D, et al., (2005):

Risk factors associated with the development of postpartum mood disorders. J Affect Disord. Sep;88(1):9-18.

Bloch M., Rotenberg N., Koren D.,et al., (2006):

Risk factors for early postpartum depressive symptoms. Gen Hosp Psychiatry. Jan-Feb;28(1):3-8.

Bodnar L.M. and Wisner K.L.(2005):

Nutrition and depression: implications for improving mental health among childbearing-aged women. Biol Psychiatry. Nov 1;58(9):679-85.

Bowen A. and Muhajarine N.(2006):

Prevalence of antenatal depression in women enrolled in an outreach program in Canada. J Obstet Gynecol Neonatal Nurs. Jul-Aug;35(4):491-8.

Boyce P. and Hickey A.(2005):

Psychosocial risk factors to major depression after childbirth. Soc Psychiatry Psychiatr Epidemiol. Aug;40(8):605-12. Epub 2005 Aug 17.

Boyce P.M.(2003):

Risk factors for postnatal depression: a review and risk factors in Australian populations. Arch Womens Ment Health. Aug;6 Suppl 2:S43-50.

Bozoky I., Corwin E.J.(2002):

Fatigue as a predictor of postpartum depression. J Obstet Gynecol Neonatal Nurs. 2002 Jul-Aug;31(4):436-43.

Brinsmead M., Smith R., Singh B.,et al.,(1985):

Peripartum concentrations of beta endorphin and cortisol and maternal mood states. Aust N Z J Obstet Gynaecol. Aug;25(3):194-7.

Brown S., Bruinsma F., Darcy M.A., et al., (2004):

Early discharge: no evidence of adverse outcomes in three consecutive population-based Australian surveys of recent mothers,

conducted in 1989, 1994 and 2000. *Paediatr Perinat Epidemiol.* May;18(3):202-13.

Brown J.C., Scott K.M., Silvers K.M.(2006):

Fish consumption in pregnancy and omega-3 status after birth are not associated with postnatal depression. *J Affect Disord.* 2006 Feb;90(2-3):131-9. Epub 2005 Dec 1.

Buist A.(2006):

Perinatal depression--assessment and management. *Aust Fam Physician.* Sep;35(9):670.

Burt V.K. and Stein K.(2002):

Epidemiology of depression throughout the female life cycle. *J Clin Psychiatry*;63 Suppl 7:9-15.

Chaaya M., Campbell O.M., El Kak F., et al., (2002):

Postpartum depression: prevalence and determinants in Lebanon. *Arch Womens Ment Health.* Oct;5(2):65-72.

Chandran M., Tharyan P., Muliylil J., et al., (2002):

Post-partum depression in a cohort of women from a rural area of Tamil Nadu, India. Incidence and risk factors. *Br J Psychiatry.* Dec;181:499-504.

Chang J., Zhang Y., Cui L.(1995):

Impact of placental hormone withdrawal on postpartum depression. *Zhonghua Fu Chan Ke Za Zhi.* Jun;30(6):342-4.

Chaudron L.H., Kitzman H.J., Peifer K.L., et al., (2005):

Prevalence of maternal depressive symptoms in low-income Hispanic women. *J Clin Psychiatry.* 2005 Apr;66(4):418-23.

Chaudron L.H., Klein M.H., Remington P., et al.,(2002):

Predictors, prodromes and incidence of postpartum depression. *J Psychosom Obstet Gynaecol.* Jun;22(2):103-12.

Chee C.Y., Lee D.T., Chong Y.S., et al., (2005):

Confinement and other psychosocial factors in perinatal depression: a transcultural study in Singapore. *J Affect Disord.* 2005 Dec;89(1-3):157-66. Epub 2005 Oct 28.

Chen C.H.(1995):

Etiology of postpartum depression--a review. *Gaoxiong Yi Xue Ke Xue Za Zhi.* Jan;11(1):1-7.

Chen T.H., Lan T.H., Yang C.Y.,et al., (2006):

Postpartum mood disorders may be related to a decreased insulin level after delivery. *Med Hypotheses.* 2006;66(4):820-3.

Chrousos G.P., Torpy D.J., Gold P.W.(1998):

Interactions between the hypothalamic-pituitary-adrenal axis and the female reproductive system: clinical implications. *Ann Intern Med.* Aug 1;129(3):229-40.

Clement S.(2001):

Psychological aspects of caesarean section. *Best Pract Res Clin Obstet Gynaecol.* Feb;15(1):109-26.

Clemmens D., Driscoll J.W., Beck C.T.(2004):

Postpartum depression as profiled through the depression screening scale. *MCN Am J Matern Child Nurs.* 2004 May-Jun;29(3):180-5.

Corwin E.J., Brownstead J., Barton N., et al., (2005):

The impact of fatigue on the development of postpartum depression. *J Obstet Gynecol Neonatal Nurs.* Sep-Oct;34(5):577-86.

Craig E.A.(2003):

Mental illness in women who have young children: current literature. *Aust J Midwifery.* Mar;16(1):5-9.

Da Costa D., Dritsa M., Rippen N.,et al., (2006):

Health-related quality of life in postpartum depressed women. *Arch Womens Ment Health.* Mar;9(2):95-102. Epub 2005 Oct 18.

Danaci A.E., Dinc G., Deveci A., et al., (2002):

Postnatal depression in turkey: epidemiological and cultural aspects. *Soc Psychiatry Psychiatr Epidemiol.* 2002 Mar;37(3):125-9.

Dearing E., Taylor B.A., McCartney K.(2004):

Implications of family income dynamics for women's depressive symptoms during the first 3 years after childbirth. *Am J Public Health.* Aug;94(8):1372-7.

Demyttenaere K., Lenaerts H., Nijs P.,et al., (1995):

Individual coping style and psychological attitudes during pregnancy and predict depression levels during pregnancy and during postpartum. *Acta Psychiatr Scand.* Feb;91(2):95-102.

Denenberg R.(1995):

Women, immunity and sexual hormones.Sidahora. 1995 Apr-May;:27-30.

Dennis C.L. and Creedy D.(2003):

Psychosocial and psychological interventions for preventing postpartum depression. *Cochrane Database Syst Rev.* Jul;8(3):76.

Dennis C.L. and Ross L.(2005):

Relationships among infant sleep patterns, maternal fatigue, and development of depressive symptomatology. *Birth.* Sep;32(3):187-93.

Dennis C.L.(2004):

Can we identify mothers at risk for postpartum depression in the immediate postpartum period using the Edinburgh Postnatal Depression Scale? *J Affect Disord.* Feb;78(2):163-9.

Dennis C.L., Janssen P.A., Singer J.(2004):

Identifying women at-risk for postpartum depression in the immediate postpartum period. *Acta Psychiatr Scand.* Nov;110(5):338-46.

Dobie S.A. and Walker E.A.(1992):

Depression after childbirth. J Am Board Fam Pract. Sep-Oct;5(5):544-5.

Dorn L.D., Susman E.J., Petersen A.C.(1993):

Cortisol reactivity and anxiety and depression in pregnant adolescents: a longitudinal perspective. Psychoneuroendocrinology. ;18(3):219-39.

Eberhard-Gran M., Eskild A., Tambs K., et al., (2002):

Depression in postpartum and non-postpartum women: prevalence and risk factors. Acta Psychiatr Scand. Dec;106(6):426-33.

Eberhard-Gran M., Tambs K., Opjordsmoen S., et al., (2005):

A comparison of anxiety and depressive symptomatology in postpartum and non-postpartum mothers. Soc Psychiatry Psychiatr Epidemiol. 2003 Oct;38(10):551-6.

Eberhard-Gran M., Tambs K., Opjordsmoen S., et al., (2004):

Depression during pregnancy and after delivery: a repeated measurement study. J Psychosom Obstet Gynaecol. Mar;25(1):15-21.

Edwards D.R., Porter S.A., Stein G.S.(1994):

A pilot study of postnatal depression following caesarean section using two retrospective self-rating instruments. J Psychosom Res. 1994 Feb;38(2):111-7.

Eilat-Tsanani S., Merom A., Romano S., et al., (2006):

The effect of postpartum depression on women's consultations with physicians. Isr Med Assoc J. 2006 Jun;8(6):406-10.

Ekuklu G., Tokuc B., Eskiocak M., et al., (2004):

Prevalence of postpartum depression in Edirne, Turkey, and related factors. J Reprod Med. 2004 Nov;49(11):908-14.

Faisal-Cury A., Tedesco JJ, Kahhale S., Menezes P.R., et al., (2004):

Postpartum depression: in relation to life events and patterns of coping. Arch Womens Ment Health. Apr;7(2):123-31. Epub Dec 15.

Fatoye F.O., Adeyemi A.B., Oladimeji B.Y.(2004):

Postpartum depression following normal vaginal delivery among Nigerian women. Psychol Rep. Jun;94(3 Pt 2):1276-8.

Feksi A., Harris B., Walker R.F., et al.,(1984):

Maternity blues' and hormone levels in saliva. J Affect Disord. Jun;6(3-4):351-5.

Felice E., Saliba J., Grech V., et al., (2004):

Prevalence rates and psychosocial characteristics associated with depression in pregnancy and postpartum in Maltese women. J Affect Disord. Oct 15;82(2):297-301.

Felice E., Saliba J., Grech V., et al., (2006):

Validation of the Maltese version of the Edinburgh Postnatal Depression Scale. Arch Womens Ment Health. Mar;9(2):75-80.

Ferguson S.S., Jamieson D.J., Lindsay M.(2002):

Diagnosing postpartum depression: can we do better? *Am J Obstet Gynecol.* May;186(5):899-902.

Figueiredo B., Bifulco A., Pacheco A., et al., (2006):

Teenage pregnancy, attachment style, and depression: a comparison of teenage and adult pregnant women in a Portuguese series. *Attach Hum Dev.* Jun;8(2):123-38.

Fisher J.R., Hammarberg K., Baker H.W.(2005):

Assisted conception is a risk factor for postnatal mood disturbance and early parenting difficulties. *Fertil Steril.* Aug;84(2):426-30.

Flaherty R.J.(1984):

Postpartum thyroiditis. *Am Fam Physician.* 1984 Feb;29(2):195-7.

Flores D.L and Hendrick V.C.(2002):

Etiology and treatment of postpartum depression. *Curr Psychiatry Rep.* Dec;4(6):461-6.

Forman D. N., Videbech P., Hedegaard M. et al.,(2000):

Postpartum depression: identification of women at risk. *Br J Obstet Gynaecol.* 2000 Oct ;107:1210-17.

Freeman M.P.(2006):

Omega-3 fatty acids and perinatal depression: A review of the literature and recommendations for future research. *Prostaglandins Leukot Essent Fatty Acids.* Aug 21.

Freeman M.P., Wright R., Watchman M., et al., (2005):

Postpartum depression assessments at well-baby visits: screening feasibility, prevalence, and risk factors. *J Womens Health (Larchmt).* Dec;14(10):929-35.

Gale S. and Harlow B.L.(2003):

Postpartum mood disorders: a review of clinical and epidemiological factors. *J Psychosom Obstet Gynecol.* Dec;24(4):257-66.

Garnier J.P., Bousquet B., Dreux C., et al.,(1985):

Changes in urinary levels of MHPG, VMA and 5-HIAA and plasma tryptophan in post-partum blues. Initial results. *Encephale.* Jan-Feb;11(1):35-7.

Garvey M.J., Tollefson G.D.(1984):

Postpartum depression. *J Reprod Med.* Feb;29(2):113-6. **Gavin N.I., Gaynes B.N., Lohr K.N., et al., (2005):**

Perinatal depression: a systematic review of prevalence and incidence. *Obstet Gynecol.* Nov;106(5 Pt 1):1071-83.

George G.C., Milani T.J., Hanss-Nuss H., et al., (2005):

Compliance with dietary guidelines and relationship to psychosocial factors in low-income women in late postpartum. *J Am Diet Assoc.* Jun;105(6):916-26.

Gjerdingen D.K. and Chaloner K.M.(1994):

The relationship of women's postpartum mental health to employment and social support. *J Fam Pract.* May;38(5):465-72.

Godfroid I.O. and Charlot A.(1996):

Postpartum psychiatry. *Rev Med Brux.* Feb;17(1):22-3, 6. Universite Libre de Bruxelles.

Goldbort J.(2006):

Transcultural analysis of postpartum depression. *MCN Am J Matern Child Nurs.* Mar-Apr;31(2):121-6.

Goodman J.H.(2004):

Postpartum depression beyond the early postpartum period. *J Obstet Gynecol Neonatal Nurs.* Jul-Aug;33(4):410-20.

Gotlib I.H., Whiffen V.E., Wallace P.M., et al.,(1991):

Prospective investigation of postpartum depression: factors involved in onset and recovery. *J Abnorm Psychol.* May;100(2):122-32.

Goyal D., Murphy S.O., Cohen J.(2006):

Immigrant Asian Indian women and postpartum depression. *J Obstet Gynecol Neonatal Nurs.* Jan-Feb;35(1):98-104.

Groer M., Davis M., Casey K., et al., (2005):

Neuroendocrine and immune relationships in postpartum fatigue. *MCN Am J Matern Child Nurs.* 2005 Mar-Apr;30(2):133-8.

Gross K.H., Wells C.S., Radigan-Garcia A., et al., (2002):

Correlates of self-reports of being very depressed in the months after delivery: results from the Pregnancy Risk Assessment Monitoring System. *Matern Child Health J.* Dec;6(4):247-53.

Guo S.F.(1993):

Postpartum depression. *Zhonghua Fu Chan Ke Za Zhi.* 1993 Sep;28(9):532-3, 569.

Halbreich U. and Karkun S.(2006):

Cross-cultural and social diversity of prevalence of postpartum depression and depressive symptoms. *J Affect Disord.* Apr;91(2-3):97-111.

Halbreich U.(2004):

Prevalence of mood symptoms and depressions during pregnancy: implications for clinical practice and research. *CNS Spectr.* 2004 Mar;9(3):177-84.

Hall L.A., Kotch J.B., Browne D., et al., (1996):

Self-esteem as a mediator of the effects of stressors and social resources on depressive symptoms in postpartum mothers. *Nurs Res.* Jul-Aug;45(4):231-8.

Hanley J.(2006):

The assessment and treatment of postnatal depression. *Nurs Times.* Jan 3-9;102(1):24-6.

Hanna B., Jarman H., Savage S., et al., (2004):

The early detection of postpartum depression: midwives and nurses trial a checklist. *J Obstet Gynecol.* Mar-Apr;33(2):191-7.

Harris B., Lovett L., Newcombe RG., et al., (1994):

Maternity blues and major endocrine changes: Cardiff puerperal mood and hormone study II. *BMJ.* Apr 9;308(6934):949-53.

Harris B., Lovett L., Smith J., et al., (1996):

Cardiff puerperal mood and hormone study. III. Postnatal depression at 5 to 6 weeks postpartum, and its hormonal correlates across the peripartum period. *Br J Psychiatry.* Jun;168(6):739-44.

Harris B., Othman S., Davies J.A., et al.,(1992):

Association between postpartum thyroid dysfunction and thyroid antibodies and depression. *BMJ.* Jul 18;305(6846):152-6.

Hatton D.C., Harrison-Hohner J., Coste S., et al., (2005):

Symptoms of postpartum depression and breastfeeding. *J Hum Lact.* Nov;21(4):444-9; quiz 450-4.

Heh S.S. and Fu Y.Y.(2003):

Effectiveness of informational support in reducing the severity of postnatal depression in Taiwan. *J Adv Nurs.* Apr;42(1):30-6.

Heh S.S., Coombes L., Bartlett H.(2004):

The association between depressive symptoms and social support in Taiwanese women during the month. *Int J Nurs Stud.* Jul;41(5):573-9.

Heidrich A., Schleyer M., Spingler H.,et al., (2005):

Postpartum blues: relationship between not-protein bound steroid hormones in plasma and postpartum mood changes. *J Affect Disord.* 1994 Feb;30(2):93-8.

Henderson J.J., Evans S.F., Straton J.A., et al., (2003):

Impact of postnatal depression on breastfeeding duration. *Birth.* Sep;30(3):175-80.

Hendrick V., Altshuler L.L., Suri R.(2005):

Hormonal changes in the postpartum and implications for postpartum depression.

Henshaw C.(2003):

Mood disturbance in the early puerperium: a review. *Arch Womens Ment Health.* 2003 Aug;6 Suppl 2:S33-42. Epub 2003 Aug 9.

Henshaw C., Boath L., Bradley E.(2006):

Antidepressant prevention of postnatal depression. *Cochrane Database Syst Rev.* Apr 18;(2):CD004363.

Henshaw C., Foreman D., Cox J.(2004):

Postnatal blues: a risk factor for postnatal depression. *J Psychosom Obstet Gynaecol.* 2004 Sep-Dec;25(3-4):267-72.

Heron J., Thomas G., Connor O., et al., (2004):

The course of anxiety and depression through pregnancy and the postpartum in a community sample. *J Affective Disorders* 80:65-73.

Hight N. and Drummond P.(2004):

A comparative evaluation of community treatments for postpartum depression: implications for treatment and management practices. *Aust N Z J Psychiatry*. Apr;38(4):212-8.

Hiltunen P., Raudaskoski T., Ebeling H., et al., (2004):

Does pain relief during delivery decrease the risk of postnatal depression? *Acta Obstet Gynecol Scand*. Mar;83(3):257-61.

Hobfoll S.E., Ritter C., Lavin J., et al., (1995):

Depression prevalence and incidence among inner-city pregnant and postpartum women *J Consult Clin Psychol*. Jun;63(3):445-53.

Hoffbrand S., Howard L., Crawley H.(2001):

Antidepressant drug treatment for postnatal depression. *Cochrane Database Syst Rev*. ;(2):CD002018.

Honey K.L., Bennett P., Morgan M.(2003):

Predicting postnatal depression. *J Affect Disord*. Sep;76(1-3):201-10.

Horowitz J.,and Goodman J.H.(2005):

Identifying and treating postpartum depression. *J Obstet Gynecol Neonatal Nurs*. 2005 Mar-Apr;34(2):264-73.

Horowitz J.A.(2006):

Community-based postpartum depression screening within the first month after delivery. *Contemp Nurse*. 2006 Feb-Mar;21(1):85-93.

Howell E.A., Mora P.A., Horowitz C.R., et al., (2005):

Racial and ethnic differences in factors associated with early postpartum depressive symptoms. *Obstet Gynecol*. Jun;105(6):1442-50.

Huang C.M., Carter P.A., Guo J.L.(2004):

A comparison of sleep and daytime sleepiness in depressed and non-depressed mothers during the early postpartum period. *J Nurs Res*. Dec;12(4):287-96.

Husain N., Bevc I., Husain M., et al., (2006):

Prevalence and social correlates of postnatal depression in a low income country. *Arch Womens Ment Health*. Jul;9(4):197-202.

Inandi T., Bugdayci R., Dundar P., et al., (2005):

Risk factors for depression in the first postnatal year: a Turkish study. *Soc Psychiatry Psychiatr Epidemiol*. Sep;40(9):725-30.

Inandi T., Elci O.C., Ozturk A., (2006):

Risk factors for depression in postnatal first year, in eastern Turkey. *Int J Epidemiol*. Dec;31(6):1201-7.

Iranfar S., Shakeri ., Ranjbar M., et al., (2005):

Is unintended pregnancy a risk factor for depression in Iranian women? *East Mediterr Health J.* 2005 Jul;11(4):618-24.

Irfan N. and Badar A.(2003):

Determinants and pattern of postpartum psychological disorders in Hazara division of Pakistan. *J Ayub Med Coll Abbottabad.* Jul-Sep;15(3):19-23.

Jadresic E. and Araya R.(1995):

Prevalence of postpartum depression and associated factors in Santiago, Chile. *Rev Med Chil.* Jun;123(6):694-9.

Jadresic E., Jara C., Araya R.(1993):

Depression in pregnancy and puerperium: study of risk factors. *Acta Psiquiatr Psicol Am Lat.* Mar;39(1):63-74.

Jardri R., Pelta J., Maron M., et al., (2006):

Predictive validation study of the Edinburgh Postnatal Depression Scale in the first week after delivery and risk analysis for postnatal depression. *J Affect Disord.* Jul;93(1-3):169-76.

Joffe H.and Cohen L.S.(1998):

Estrogen, serotonin, and mood disturbance: where is the therapeutic bridge? *Biol Psychiatry.* Nov 1;44(9):798-811.

Jones H.W.and Venis J.A.(2001):

Identification and classification of postpartum psychiatric disorders. *J Psychosoc Ment Health Serv.* Dec;39(12):2330.

Jung V., Short R., Letourneau N., et al., (2006):

Interventions with depressed mothers and their infants: Modifying interactive behaviours. *J Affect Disord.* 2006 Sep 7(9):221-33

Jurczynska J. and Zieleniewski W.(2004):

Clinical implications of occurrence of antithyroid antibodies in pregnant women and in the postpartum period. *Przegl Lek.*;61(8):864-7.

Kalantaridou S.N., Makrigiannakis A., Zoumakis E., et al., (2004):

Stress and the female reproductive system. *J Reprod Immunol.* Jun;62(1-2):61-8.

Kaplan B.J.(1983):

Causes and attributions of depression during pregnancy. *Women Health.* Winter;8(4):23-32.

Kaplinsky N., Pines A, Olchovsky D., et al.,(1983):

Transient post-partum hypothyroidism. *Acta Obstet Gynecol Scand.* ;62(1):91-2.

Kendler K.S., Kessler RC., Neale M.C.,et al., (1993):

The prediction of major depression in women: toward an integrated etiologic model. *Am J Psychiatry.* Aug;150(8):1139-48.

Kennedy H.P., Beck C.T., Driscoll J.W.(2002):

A light in the fog: caring for women with postpartum depression. *J Midwifery Women's Health*. 2002 Sep-Oct;47(5):318-30.

Kiernan K. and Pickett K.E.(2006):

Marital status disparities in maternal smoking during pregnancy, breastfeeding and maternal depression. *Soc Sci Med*. Jul;63(2):335-46.

Kim J. and Buist A.(2005):

Postnatal depression: a Korean perspective. *Australas Psychiatry*. Mar;13(1):68-71.

Kitamura T., Shima S., Sugawara M., et al.,(1996):

Clinical and psychosocial correlates of antenatal depression: a review. *Psychother Psychosom*;65(3):117-23.

Kitamura T., Sugawara M., Sugawara K., et al., (1996):

Psychosocial study of depression in early pregnancy. *Br J Psychiatry*. Jun;168(6):732-8.

Kitamura T., Yoshida K., Okano T., et al., (2006):

Multicentre prospective study of perinatal depression in Japan: incidence and correlates of antenatal and postnatal depression. *Arch Womens Ment Health*. May;9(3):121-30.

Kleeb B. and Ragoth C.J.(2005):

Influence of prophylactic information on the frequency of baby blues. *Z Geburtshilfe Neonatol*. Feb;209(1):22-8.

Kohl C., Walch T., Huber R., et al., (2005):

Measurement of tryptophan, kynurenine and neopterin in women with and without postpartum blues. *J Affect Disord* Jun;86(2-3):135-42.

Koo V., Lynch J., Cooper S.(2003):

Risk of postnatal depression after emergency delivery. *J Obstet Gynaecol Res*. Aug;29(4):246-50.

Kuijpers J.L., Vader H.L, Drexhage H.A., et al.,(2001):

Thyroid peroxidase antibodies during gestation are a marker for subsequent depression postpartum. *Eur J Endocrinol*. Nov;145(5):579-84.

La Coursiere D.Y., Baksh L., Bloebaum L., et al., (2006):

Maternal body mass index and self-reported postpartum depressive symptoms. *Matern Child Health J*. Jul;10(4):385-90.

Lanczik M. and Brockington I.F.(1999):

Postpartum dysphoric syndrome. Psychopathology, diagnosis and etiology. *Fortschr Neurol Psychiatr*. Feb;67(2):60-7.

Latorre-Latorre J.F., Contreras-Pezzotti L.M., Herran-Falla O.F.(2006):

Postnatal depression in a Colombian city. Risk factors. *Aten Primaria*. Apr 15;37(6):332-8.

Learoyd D.L, Fung H.Y., McGregor A.M.(1992):

Postpartum thyroid dysfunction. *Thyroid*. Spring;2(1):73-80.

Lee D.T., Yip A.S., Leung T.Y., et al., (2004):

Epidemiology of postnatal depression. Prospective multivariate study of sociocultural risk factors in a Chinese population in Hong Kong. *Br J Psychiatry*. Jan;184:34-40.

Leiferman J.(2002):

The effect of maternal depressive symptomatology on maternal behaviours associated with child health. *Health Educ Behav*. Oct;29(5):596-607.

Limlomwongse N. and Liabsuetrakul T.(2006):

Cohort study of depressive moods in Thai women during late pregnancy and 6-8 weeks of postpartum using the Edinburgh Postnatal Depression Scale (EPDS). *Arch Womens Ment Health*. May;9(3):131-8.

Lindahl V., Pearson J.L., Colpe L.(2005):

Prevalence of suicidality during pregnancy and the postpartum. *Arch Womens Ment Health*. Jun;8(2):77-87.

Logsdon M.C., Wisner K., Billings D.M., et al., (2006):

Raising the awareness of primary care providers about postpartum depression. *Issues Ment Health*. Jan;27(1):59-73.

Lumley J.and Austin M.P.(2001):

What interventions may reduce postpartum depression. *Curr Opin Obstet Gynecol*. Dec;13(6):605-11.

Maes M., Ombelet W., Verkerk R., et al.,(2001):

Effects of pregnancy and delivery on the availability of plasma tryptophan to the brain: relationships to delivery-induced immune activation and early post-partum anxiety and depression. *Psychol Med*. Jul;31(5):847-58.

Magiakou M.A., Mastorakos G., Rabin Det al., (1996):

Hypothalamic corticotropin-releasing hormone suppression during the postpartum period: implications for the increase in psychiatric manifestations at this time. *J Clin Endocrinol Metab*. May;81(5):1912-7.

Mahomed K., Gulmezoglu A.M., Nikodem VC., et al.,(1995):

Labor experience, maternal mood and cortisol and catecholamine levels in low-risk primiparous women. *J Psychosom Obstet Gynaecol*. Dec;16(4):181-6.

Mallikarjun P.K, Oyebode F.(2005):

Prevention of postnatal depression. *J R Soc Health*. Sep;125(5):221-6.

Martinez-Schallmoser L., Telleen S., MacMullen N.J.(2003):

The effect of social support and acculturation on postpartum depression in Mexican American women. *J Transcult*. Oct;14(4):329-38.

Matthey S., Barnett B., Howie P., et al., (2003):

Diagnosing postpartum depression in mothers and fathers: whatever happened to anxiety? *J Affect Disord*. Apr;74(2):139-47.

- McConnell J., Baker M., Marks L.(2005):**
Experiencing a dilemma--accounts of working with PND. *Community Pract.* Jul;78(7):251-4.
- McCoy S.J., Beal J.M., Payton M.E., et al., (2005):**
Correlations of visual analog scales with Edinburgh Postnatal Depression Scale. *J Affect Disord.* Jun;86(2-3):295-7.
- McCoy S.J., Beal J.M., Shipman S.B., et al., (2006):**
Risk factors for postpartum depression: a retrospective investigation at 4-weekspostnatal and a review of the literature. *J Am Osteopath Assoc.* Apr;106(4):193-8.
- McCoy S.J., Beal J.M., Watson G.H.(2003):**
Endocrine factors and postpartum depression. *J Reprod Med.* Jun;48(6):402-8.
- McGill H., Burrows V.L., Holland L.A., et al., (1995):**
Postnatal depression: a Christchurch study. *N Z Med J.* May 10;108(999):162-5.
- McLearn K.T., Minkovitz C.S., Strobino D.M., et al., (2006):**
Maternal depressive symptoms at 2 to 4 months post partum and early parenting practices. *Arch Pediatr Adolesc Med.* Mar;160(3):279-84.
- McMahon C., Barnett B., Kowalenko N., et al., (2005):**
Psychological factors associated with persistent postnatal depression: past and current relationships, defence styles and the mediating role of insecure attachment style. *J Affect Disord.* Jan;84(1):15-24.
- Metastasio P., Verdecchia P., D'Urzo E., et al., (1995):**
Postpartum depression. *Minerva Ginecol.* Jan-Feb;47(1-2):43-5.
- Mezey G., Bacchus L., Bewley S., et al., (2004):**
Domestic violence, lifetime trauma and psychological health of childbearing women. *BJOG.* Feb;112(2):197-204.
- Mian A.I.(2005):**
Depression in pregnancy and the postpartum period: balancing adverse effects of untreated illness with treatment risks. *J Psychiatr Pract.* Nov;11(6):389-96.
- Milgrom J., Ericksen J., Negri L., et al., (2005):**
Screening for postnatal depression in routine primary care: properties of the Edinburgh Postnatal Depression Scale in an Australian sample. *Aust N Z J Psychiatry.* Sep;39(9):833-9.
- Miller E.H.(2004):**
Women and insomnia. *Clin Cornerstone.* 2004;6 Suppl 1B:S8-18.
- Miller R.L., Pallant J.F., Negri L.M.(2006):**
Anxiety and stress in the postpartum: is there more to postnatal distress than depression? *BMC Psychiatry.* Mar 24;6:12.

- Moraes IG, Pinheiro RT, Silva RA, et al., (2006):**
Prevalence of postpartum depression and associated factors. *Rev Saude Publica*. Feb;40(1):65-70.
- Mosack V. and Shore E.R.(2006):**
Screening for depression among pregnant and postpartum women. *J Community Health* . Spring;23(1):37-47.
- Moses-Kolko E.L. and Roth E.K.(2004):**
Antepartum and postpartum depression: healthy mom, healthy baby. *J Am Med Womens Assoc*. Summer;59(3):181-91.
- Murphy-Eberenz K., Zandi P.P., March D., et al., (2006):**
Is perinatal depression familial? *J Affect Disord*. Jan;90(1):49-55.
- Nobbs E., Collins N.L., Lobel M., et al., (1999):**
Psychosocial predictors of postpartum depressed mood in socioeconomically disadvantaged women. *Womens Health*. Spring;1(1):51-75.
- Newhan D.J., Hostetter A., Arnold A., et al., (1983):**
The treatment of postpartum depression: minimizing infant exposures. *J Clin Psychiatry* ;63 Suppl 7:31-44.
- O'Boyle A.L., Magann E.F., Ricks R.E. et al., (2005):**
Depression screening in the pregnant soldier wellness program. *South Med J*. Apr;98(4):416-8.
- O'Hara M.W., Schlechte J.A., Lewis D.A., et al., (1984):**
Prospective study of postpartum blues. Biologic and psychosocial factors. *Arch Gen Psychiatry*. Sep;48(9):801-6.
- O'Hara M.W., Schlechte J.A., Lewis D.A., et al., (1991):**
Controlled prospective study of postpartum mood disorders: psychological, environmental, and hormonal variables. *J Abnorm Psychol*. Feb;100(1):63-73.
- Okano T. and Nomura J.(1992):**
Endocrine study of the maternity blues. *Prog Neuropsychopharmacol Biol Psychiatry*. ;16(6):921-32.
- Olshansky E.(2003):**
A theoretical explanation for previously infertile mothers' vulnerability to depression. *J Obst Scholarsh*. ;35(3):263-8.
- Onozawa K., Kumar R.C., Adams D., et al., (2003):**
High EPDS scores in women from ethnic minorities living in London. *Arch Womens Ment Health*. Aug;6 Suppl 2:S51-5.
- Owoeye A.O., Aina O.F., Morakinyo O.(2006):**
Risk factors of postpartum depression and EPDS scores in a group of Nigerian women. *Trop Doct*. Apr;36(2):100-3.
- Park Y.J., Shin H.J., Ryu H., et al., (2004):**
The predictors of postpartum depression. *Taehan Kanho Hakhoe Chi*. Aug;34(5):722-8.

Parker J.(2004):

Does traumatic birth increase the risk of postnatal depression? Br J Community Nurs. Feb;9(2):74-9.

Paterson J.A., Davis J., Gregory M.,et al., (1994):

A study on the effects of low haemoglobin on postnatal women. Midwifery. Jun;10(2):77-8.

Parskad and Kulkurai (2003):

Thyroid and adrenal measures during late pregnancy and the puerperium in women who have been major depressed or who become dysphoric postpartum. J Affect Disord. Oct-Nov;29(2-3):201-11.

Paulson J.F., Dauber S., Leiferman J.A.(2006):

Individual and combined effects of postpartum depression in mothers and fathers on parenting behavior. Pediatrics. Aug;118(2):659-68.

Pearson Murphy B.E., Steinberg S.I., Hu F.Y., et al.,(2001):

Neuroactive ring A-reduced metabolites of progesterone in human plasma during pregnancy: elevated levels of 5 alpha-dihydroprogesterone in depressed patients during the latter half of pregnancy. J Clin Endocrinol Metab. Dec;86(12):5981-7.

Pedersen C.A.(1999):

Postpartum mood and anxiety disorders: a guide for the nonpsychiatric clinician with an aside on thyroid associations with postpartum mood. Thyroid. Jul;9(7):691-7.

Perfetti J., Clark R., Fillmore C.M.(2005):

Postpartum depression: identification, screening, and treatment. WMJ. 2004;103(6):56-63.

Pesavento F., Marconcini E., Drago D.(2005):

Quality of life and depression in normal and in high-risk pregnancy. Analysis of a sample of 100 women. Minerva Ginecol. Aug;57(4):451-60.

Pop V.J., Essed G.G., de Geus CA.,et al., (1993):

Prevalence of post partum depression--or is it post-puerperium depression? Acta Obstet Gynecol Scand. Jul;72(5):354-8.

Pop V.J., Wijnen H.A., van Montfort M., t al., (1995):

Blues and depression during early puerperium: home versus hospital deliveries. Br J Obstet Gynaecol. Sep;102(9):701-6.

Rahman A., Bunn J.E., Lovel H., et al., (2006):

Maternal depression increases infant risk of diarrhoeal illness - a cohortstudy Arch Dis Child.Sep 11; 322- 25.

Rahman A., Iqbal Z., Bunn J., et al., (2004):

Impact of maternal depression on infant nutritional status and illness: a cohort study. Arch Gen Psychiatry. Sep;61(9):946-52.

- Reck C., Weiss R., Fuchs T., et al., (2004):**
 Psychotherapy for postpartum depression with a focus on mother-infant Interaction. *Nervenarzt*. Nov;75(11):1068-73.
- Reynolds O.P., Records K. I., Rice M.J., et al.,(2005):**
 A comparative study of postpartum depression in abused and non-abused women. *Arch Psychiatr Nurs*. Dec;19(6):281-90.
- Redmond G.P.(2004):**
 Thyroid dysfunction and women's reproductive health. *Thyroid*. ;14 Suppl 1:S5-15.
- Rees A.M., Austin M.P., Parker G.(2005):**
 Role of omega-3 fatty acids as a treatment for depression in the perinatal period. *Aust N Z J Psychiatry*. Apr;39(4):274-80.
- Rhonda C. B.and Jay D.A. (2004):**
 Mood disorder in women from adolescence to late life: An overview.*Clin Obstet Gynecol*. 47(3):515-26.
- Rich-Edwards J.W., Kleinman K., Abrams A., et al., (2006):**
 Sociodemographic predictors of antenatal and postpartum depressive symptoms among women in a medical group practice. *J Epidemiol Community Health*.Mar;60(3):2217.
- Rossler R. A.,and Fallahpour M. H. (2003):**
 Postpartum depression: do we still need this diagnostic term? *Acta Psychiatr Scand Suppl*. ;(418):51-6.
- Robertson E., Grace S., Wallington T., et al.,(2004):**
 Antenatal risk factors for postpartum depression: a synthesis of recent literature. *Gen Hosp Psychiatry*. Jul-Aug;26(4):289-95.
- Robertson E., Grace S., Wallington T., et al.,(2006):**
 Women and omega-3 Fatty acids.*Obstet Gynecol Surv*. Oct;59(10):722-30; qu6.
- Ross L.E., Murray B.J., Steiner M.(2005):**
 Sleep and perinatal mood disorders: a critical review. *J Psychiatry Neurosci*. 2005 Jul;30(4):247-56.
- Rouillon F., Thalassinos M., Miller H.D., et al., (1992):**
 Folates and post partum depression. *J Affect Disord*. Aug;25(4):235-41.
- Russell J.A., Douglas A.J., Ingram C.D.(2001):**
 Brain preparations for maternity--adaptive changes in behavioral and neuroendocrine systems during pregnancy and lactation. An overview. *Prog Brain Res*. ;133:1-38.
- Ryan D., Millis L., Misri N.(2005):**
 Depression during pregnancy. *Can Fam Physician*. Aug;51:1087-93.

- Saisto T. and Halmesmaki E.(2003):**
Fear of childbirth: a neglected dilemma. *Acta Obstet Gynecol Scand.* Mar;82(3):201-8.
- Sanderson C.A., Cowden B., Hall D.M., et al., (2002):**
Is postnatal depression a risk factor for sudden infant death? *Br J Gen Pract.* Aug;52(481):636-40.
- Sandler R.M., Wiemann C.M., Rickert V.I., et al., (1980):**
Moderate to severe depressive symptoms among mothers followed four years postpartum. *J Adolesc Health.* Jun;38(6):712-8.
- Shneeberg N.G. and Perloff W.H. (1980):**
In the shadow of maternal depressed mood: experiences of parenthood during the first year after childbirth. *J Psychosom Obstet Gynaecol.* Mar;25(1):23-34.
- Sichel D.A., Cohen L.S, Robertson L.M., et al., (1995):**
Prophylactic estrogen in recurrent postpartum affective disorder. *Biol Psychiatry.* Dec 15;38(12):814-8.
- Sierra Manzano J.M., Carro Garcia T., Ladron Moreno E.(2002):**
Variables associated with the risk of postpartum depression. *Edinburgh Postnatal Depression Scale.* *Br J Gen Pract.* Oct;52(483):854.
- Small R., Astbury J., Brown S., et al., (1994):**
Depression after childbirth. Does social context matter? *Med J Aust.* Oct 17;161(8):473-7.
- Smith R. and Thomson M.(1991):**
Neuroendocrinology of the hypothalamo-pituitary-adrenal axis in pregnancy and the puerperium. *Baillieres Clin Endocrinol Metab.* Mar;5(1):167-86.
- Sobrinho L.G and Almeida-Costa J.M.(1992):**
Hyperprolactinaemia as a result of immaturity or regression: the concept of maternal subrountine. A new model of psychoendocrine interactions. *Psychother Psychosom.* ;57(3):128-32.
- Steiner M., Dunn E., Born L.(2003):**
Hormones and mood: from menarche to menopause and beyond. *J Affect Disord.* Mar;74(1):67-83.
- Stern G., Kruckman L.(1983):**
Multi-disciplinary perspectives on post-partum depression: an anthropological critique. *Soc Sci Med.*;17(15):1027-41.
- Stuart S., O'Hara M.W., Gorman L.L.(2003):**
The prevention and psychotherapeutic treatment of postpartum depression. *Arch Womens Ment Health.* Aug;6 Suppl 2:S57-69.
- Sutter-Dallay A.L., Marcesche V.M., Glatigny-Dallay E., et al., (2004):**
Women with anxiety disorders during pregnancy are at increased risk of intense postnatal depressive symptoms: a prospective survey of the MATQUID cohort. *Eur Psychiatry.* Dec;19(8):459-63.

Taveras E.M., Capra A.M., Braveman P.A., et al., (2003):

Clinician support and psychosocial risk factors associated with breastfeeding discontinuation. *Pediatrics*. Jul;112(1 Pt 1):108-15.

Tcheremissine O.V. and Lieving L.M.(2005):

Pharmacotherapy of postpartum depression: current practice and future directions. *Expert Opin Pharmacother*. Oct;6(12):1999-2005.

Templeton L., Velleman R., Persaud A., et al., (2003):

The experiences of postnatal depression in women from black and minority ethnic communities in Wiltshire, UK. *Ethn Health*. Aug;8(3):207-21.

Thome M., Alder E.M., Ramel A.(2004):

A population-based study of exclusive breastfeeding in Icelandic women: is there a relationship with depressive symptoms and parenting stress? *Int. J. Nurs. Stud*. Jan;43(1):11-20.

Ugarriza D.N. and Robinson M.K.(1997):

Assessment of postpartum depression: a review of the research. *Online J Knowl Synth*. Nov 17;4:6.

Unterman R.R., Posner N.A., Williams K.N.(1990):

Postpartum depressive disorders: changing trends. *Birth*. Sep;17(3):131-7; discussion 138.

Verkerk G.J., Denollet J., Van Heck G.L., et al., (2005):

Personality factors as determinants of depression in postpartum women: a prospective 1-year follow-up study. *Psychosom Med*. Jul-Aug;67(4):632-7.

Verkerk G.J., Pop V.J., Van Son M.J., et al., (2003):

Prediction of depression in the postpartum period: a longitudinal follow-up study in high-risk and low-risk women. *J Affect Disord*. Nov;77(2):159-66.

Walker L., Timmerman G.M., Kim M., et al., (2002):

Relationships between body image and depressive symptoms during postpartum in ethnically diverse, low income women. *Women Health* ;36(3):101-21.

Wang S.Y. and Chen C.H.(2006):

Psychosocial health of Taiwanese postnatal husbands and wives. *J Psychosom Res*. Mar;60(3):303-7.

Wang S.Y., Jiang X.Y., Jan W.C., et al., (2003):

A comparative study of postnatal depression and its predictors in Taiwan and mainland China. *Am J Obstet Gynecol*. Nov;189(5):1407-12.

Warner R., Appleby L., Whitton A., et al., (1996):

Demographic and obstetric risk factors for postnatal psychiatric morbidity. *Br J Psychiatry*. May;168(5):607-11.

- Webster J., Linnane J., Roberts J., et al., (2003):**
Identify, Educate and Alert (IDEA) trial: an intervention to reduce postnatal depression. *BJOG*. Sep;110(9):842-6.
- Wewerinke A., Honig A., Heres M.H., et al.,(2006):**
Psychiatric disorders in pregnant and puerperal women. *Ned Tijdschr Geneesk*. Feb 11;150(6):294-8.
- Wickberg B. and Hwang C.P.(1997):**
Screening for postnatal depression in a population-based Swedish sample. *Acta Psychiatr Scand*. Jan;95(1):62-6.
- Widner B., Laich A., Sperner-Unterweger B., et al.,(2002):**
Neopterin production, tryptophan degradation, and mental depression--what is the link? *Brain Behav Immun*. Oct;16(5):590-5.
- Wilson L.M., Reid A.J., Midmer D.K., et al., (1996):**
Antenatal psychosocial risk factors associated with adverse postpartum family outcomes. *CMAJ*. Mar 15;154(6):785-99.
- Wilson R.G. and Davis R.E.(1983):**
Clinical chemistry of vitamin B6. *Adv Clin Chem*.3;23:1-68.
- Wisner K.L., Perel J.M., Peindl K.S., et al., (2004):**
Timing of depression recurrence in the first year after birth. *J Affect Disord*. Mar;78(3):249-52.
- Wolf A.W., De Andraca I., Lozoff B.(2002):**
Maternal depression in three Latin American samples. *Soc Psychiatry Psychiatr Epidemiol*. Apr;37(4):169-76.
- Yamashita H. and Yoshida K.(2003):**
Screening and intervention for depressive mothers of new-born infants. *Seishin Shinkeigaku Zasshi* ;105(9):1129-35.
- Yoshida K. and Yamashita H.(2003):**
Treatment strategy for women with puerperal psychiatric disorders--psychopharmaco-therapy and its impact on fetus and breast-fed infants. *Seishin Shinkeigaku Zasshi*. ;105(9):1136-44.
- Zinga D., Phillips S.D., Born L.(2005):**
Postpartum depression: we know the risks, can it be prevented? *Rev Bras Psiquiatr*. Oct;27 Suppl 2:S56-64. Epub 2005 Nov 10.
- Zlotnick C., Miller I.W., Pearlstein T., et al., (2006):**
A preventive intervention for pregnant women on public assistance at risk for postpartum depression. *Am J Psychiatry*. Aug;163(8):1443-5.
- Zonana J. and Gorman J.M.(2005):** The neurobiology of postpartum depression. *CNS Spectr*. Oct;10(10):792-9, 805.