

References

Baigre RS, Hag A, Morgan CD, et al : The spectrum of right ventricular involvement in inferior wall myocardial infarction: a clinical, hemodynamic and noninvasive study. *J Am Coll Cardiol* 1983; 1: 396-404.

Bairey CN, Shah PM, Lew AS et al.:

Electrocardiographic differentiation of occlusion of the left circumflex versus the right coronary artery as a cause of inferior acute myocardial infarction. *Am. J. Cardiol.*, 1987, 60:456-9.

Bakker AJ, Gorgels JPMC, Van Vlies B, et al.: Contribution of creatine kinase MB mass concentration at admission to early diagnosis of myocardial infarction. *Br. Heart J* 1994; 72:122.

Balcells E, Powers ER, Lepper W, et al: Detection of myocardial viability by contrast echocardiography in acute infarction predicts recovery of resting function and contractile reserve. *J Am Coll Cardiol* 2003; 41:827.

Baroldi G: Diseases of the coronary arteries. In Silver MD (ed); cardiovascular pathology, Vol. 1, New York, Churchill Livingstone, 1982, p. 317-391.

Barren HV, Cannon CP, Murphy SA, et al.: Association between white blood cell count, epicardial blood flow, myocardial perfusion, and clinical outcomes in the setting of acute myocardial infarction. *Circulation* 2000; 102:2329.

Barry I., Huey George A, Beller Kaiser, et al.: A comprehensive analysis of myocardial infarction due to left circumflex artery occlusion: Comparison with infarction due to right coronary artery and left descending artery occlusion. *J Am Coll Cardiol.* 1988; 12:1156.

Basha BJ, Sowers JR, Kossouw JE, et al.:

Atherosclerosis: an update. *Am. Heart J.* 1996; 131:1192-1202.

Bayes de Luna A: Clinical Electrocardiology: A Textbook 2nd updated edition. Armonk, N Y: Futura Publ, 1998.

Bellamy RF and Lowensohn HS: Effect of systole on coronary pressure- flow relation in the right ventricle of the dog. Am J Physiol 1980; 238:H4S1- 6.

Berger PB and Ryan TJ: Inferior myocardial infarction. High-Risk subgroups. Circulation 1990; 81 (2): 401.

Berger PB, Ruocco Jr NA, Ryan TJ, et al.: Incidence and prognostic implications of heart block complicating inferior myocardial infarction treated with thrombolytic therapy: results from TIMIII. J Am Coll Cardiol 1992; 20:533 - 40.

Berger PB, Ryan TJ: Inferior myocardial infarction: high risk subgroups. Circulation 1990;

Bernard JG and Shahbudin HR: Acute myocardial infarction. 2nd edition. International Thomson publishing. New York, Boston, London, Paris'. P. 173: 1997.

Berry C, Zalewski A, Kovach R, et al.: Surface electrocardiogram in the detection of transmural myocardial ischemia during coronary occlusion. Am J Cardiol. 1989,63: 21 -26 .

Billadello JJ, Smith JL, Ludbrook PA, et al.: Implications of "reciprocal" ST segment depression associated with acute myocardial infarction identified by positron tomography. J AM Coll Cardiol. 1983, May; 20(5):477-81.

Birnbaun Y, Sclarovsky S, Herz I, et al.: Admission clinical and electrocardiographic characteristics predicting in-hospital development of high-degree atrioventricular block in inferior wall acute myocardial infarction. Am. J. Cardiol 1997; 80: 1134-8. .

Bosimini E, Giannuzzi P, Temporelli PL, et al.: Klectrocardiographic evolutionary changes and left ventricular remodeling after acute myocardial infarction: results of the G1SS1-3 echo substudy. J Am Coll Cardiol 2000; 35:127-35. .

Bounhoure JP, Galinier M, Assoun B, et al.: Inferior wall myocardial infarction and atrioventricular block; angiography and prognosis. Arch Mal Coeur Vaiss. 1994 Apr; 87(4):445-50. . "

Bowers TR, O'Neill WW, Pica M, et al.: Patterns of Coronary Compromise Resulting in Acute Right Ventricular Ischemic Dysfunction. Circulation. 2002; 106; 1104-9..

Braat SH, Brugada P, den Dulk K, et al.: Value of lead V4R for recognition of the infarct coronary artery in acute inferior myocardial infarction. Am J Cardiol 1984; 53:1538-1541. .

Braat, SH, Gorgels, APM, Bar, FW, et al.: Value of the ST-T segment in lead V4R in inferior wall acute myocardial infarction to predict the site of coronary arterial occlusion. A m J Car dial 1988, 62,140-142.

Braunwald E, Antman EM, Beasley JW, et al.: ACC/AHA guidelines for the management of patients with unstable angina and non ST- segment elevation myocardial infarction. A report of the American College of Cardiology/American Heart Association Task Force on Guidelines (Committee on the Management of Patients With Unstable Angina). J Am Coll Cardiol 2002; 36:970-1062.

Braunwald E, Zipes DP, and Libby P.: Coronary angiography and intravascular ultrasonography. In Heart Disease 6th edition, Philadelphia: W.B. Saunders. 2001, P.397-400. .

BRAUNWALD'S HEART DISEASE: A Textbook of Cardiovascular Medicine.? 2005. .

Burns RJ, Gibbons RJ, Yi Q, et al.: The relationships of left ventricular ejection fraction, end-systolic volume index and infarct size to six-month mortality after hospital discharge following myocardial infarction treated by thrombolysis. J Am Coll Cardiol 2002; 39:30-6.

Carlos V, Serrano JR, Luiz A, et al: Sinus bradycardia as a predictor of right coronary artery occlusion in patients with inferior myocardial infarction. International J. of Cardio 1999; 68: 75-82.

Cheitlin MD., Armstrong \VF, Alpert JS, et al.: ACC/AHA/ASF 2003 Guideline Update for the Clinical Application of Echocardiography A Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (ACC/AHA/ASE Committee to Update the 1997 Guidelines for the Clinical Application of Echocardiography).

Chia BL, Yip JW, Tan HC, et al.: Usefulness of ST elevation 11/ III ratio and ST deviation in lead I for identifying the culprit artery in inferior wall acute myocardial infarction. Am J Cardiol 2000, Augl; 86:341-343.

Clemmensen P, Bates ER, Califf RM, et al.: Complete atrioventricular block complicating inferior wall acute myocardial infarction treated with reperfusion therapy. TAMI Study Group. Am J Cardiol 1991; 67:225-30.

Cohen JN, Guha NH, Broder MI, et al.: Right ventricular infarction, clinical and hemodynamic features. Am J Cardiol 1974; 33; 209.
Collinson PO: Troponin T or troponin I or CK-MB (or none) Eur Heart J 1998; 19: 16-24.

Corr PB and Gillis RA.: Effect of autonomic neural influence on the cardiovascular changes induced by coronary occlusion. Am Heart J 1975; 89:767-74.

Corr PB, Pearle DL, Hinton JR, et al.: Site of myocardial infarction. A determination of cardiovascular changes induced in the cat by coronary occlusion. Circ. Res. 1976; 39:840.

Dell'Italia LJ, Starling MR, Boros BL, et al.: First transit and equilibrium radionuclide angiography in patients with inferior transmural myocardial infarction: criteria for the diagnosis of associated hemodynamically significant right ventricular infarction Am Coll Cardiol 1984; 4:923-30.

Dell'Italia LJ, Starling MR and O'Rourke RA: Physical examination for exclusion of hemodynamically important right ventricular infarction. Ann Intern Med 1983; 95:608-611.

Dell'Italia LJ: Right ventricular infarction. J Intensive Care Med 1986; 1:246-56.

Derumeaux G, Loufoua J, Pontier G, et al.: Tissue Doppler imaging differentiates transmural from nontransmural acute myocardial infarction after reperfusion therapy. Circulation 2001; 103: 589.

Dwyer EM Jr: The predictive accuracy of the electrocardiogram identifying the presence and location of myocardial infarction and coronary artery disease. Ann N Y Acad Sci. 1990;601:67-76.

Dwyer EM, Cofluis S, Greinberg H, et al.: Inferior myocardial infarction and right coronary artery occlusive disease. A correlative study, Br Heart J. 1975, May; 37(5):464-70.

EL-Allaf M, Chapelle J, EL-AHaf D, et al.: Differentiating muscle damage from myocardial injury by means of the serum Creatine Kinase (CK) Isoenzyme MB mass measure/total CK activity ratio. Clin. Chem. 1986; 32:291.

Eskola M, Mikkelsen J, Nikus K, et al.: Fatality of myocardial infarction in relation to the coronary anatomy: role of culprit lesion location. Ann Med 2004; 36:474-479.

Feigl D, Ashkenasy J, and Kishon Y: Early and late atrioventricular block in acute inferior myocardial infarction. J. Am. Coll. Cardiol., 4:35 38, 1984.

Finn AV and Antnian EM: Isolated right ventricular infarction. N Engl J Med 2003, 349: p 17.

Fiol M, Carrillo A, Cygankiewicz I, et al.: New criteria based on ST changes in 12-leads surface ECG to detect proximal versus distal right coronary artery occlusion in the case of acute infroposterior myocardial infarction. A.N.E. 2004; 9(4):383-388.

Derumeaux G, Loufoya J, pontier G, et al.: Tissue Doppler imaging differentiates transmural from nontransmural acute myocardial infarction after rcperfusion therapy. Circulation 2001; 103: 589.

Dwyer EM Jr: The predictive accuracy of the electrocardiogram identifying the presence and location of myocardial infarction and coronary artery disease. Ann N Y Acad Sci. 1990:601:67-76.

Dwyer EM, Cofluia S, Greinberg H, et al.: Inferior myocardial infarction and right coronary artery occlusive disease. A correlative study, Br Heart J. 1975, May; 37(5):464-70.

EL-Allaf M, Chapelle J, EL-AHaf D, et al.: Differentiating muscle damage from myocardial injury by means of the serum Creatine Kinase (CK) Isoenzyme MB mass measure/total CK activity ratio. Clin. Chem. 1986; 32:291.

Eskola M, Mikkelsen J, Nikus K, et al.: Fatality of myocardial infarction in relation to the coronary anatomy: role of culprit lesion location. *Ann Med* 2004; 36:474-479.

Feigl D, Ashkenasy J, and Kishon Y: Early and late atrioventricular block in acute inferior myocardial infarction. *J. Am. Coll. Cardiol.*, 4:35-38, 1984.

Finn AV and Antnian EM: Isolated right ventricular infarction. *N Engl J Med* 2003, 349: p 17.

Fiol M, Carrillo A, Cygankiewicz I, et al.: New criteria based on ST changes in 12-lead surface ECG to detect proximal versus distal right coronary artery occlusion in the case of acute inferior myocardial infarction. *A.N.E.* 2004; 9(4):383-388.

Fiol M, Cygankiewicz I, Guindo J, et al.: Evolving myocardial infarction with ST elevation: ups and downs of ST in different leads identifies the culprit artery and location of the occlusion. *A.N.E.* 2004; 9: 180-186.

Fleet RP, Dupuis G, Marchand A, et al.: ACC/AHA guideline update for the management of patients with unstable angina and non-ST-segment elevation myocardial infarction. *Circulation* 2002, 106:1893.

Force T, Kemper A, Perkins L, et al.: Overestimation of the infarct size by quantitative two-dimensional echocardiography: The role of tethering and analytic procedures. *Circulation* 1986; 73: 1360.

Franch RH, King SB, Douglas JS, et al.: Techniques of cardiac catheterization including coronary arteriography. In: *The Heart, a textbook of cardiovascular medicine* by Hurst JW et al. 7th Ed. New York, McGraw Hill, 1990; Chapter 3: 1858.

Fuchs RM, Achuff SC, Brunwald L et al.: Electrocardiographic localization of coronary artery narrowing: Studies during myocardial ischemia and infarction in patients with one-vessel disease, *Circulation* 1982;66:1168.

Gibbons GH, and Dzau VJ: The emerging concept of vascular remodeling. *New England Journal of Medicine*, 1994;330, 1431-38

Gibson RS, Crampton RS, Watson DD, et al.: Precordial ST-segment depression during acute inferior myocardial

infarction: clinical, scintigraphic and angiographic correlations. *Circulation* 1982; 66:732-41.

Fiol M, Cygankiewkz I, Guindo J, et al.: Evolving myocardial infarction with ST elevation: ups and downs of ST in different leads identifies the culprit artery and location of the occlusion. *A.N.E.* 2004; 9: 180-186.

Fleet RP, Dupuis G, Marchand A, et al.: ACC/AHA guideline update for the management of patients with unstable angina and non-ST-segment elevation myocardial infarction. *Circulation* 2002, 106:1893.

Force T, Kemper A, Perkins L, et al.: Overestimation of the infarct size by quantitative two-dimensional echocardiography: The role of tethering and analytic procedures. *Circulation* 1986; 73: 1360.

Franch RH, King SB, Douglas JS, et al.: Techniques of cardiac catheterization including coronary arteriography. In: *The Heart, a textbook of cardiovascular medicine* by Hurst JW et al. 7th Ed. New York, McGraw Hill, 1990; Chapter 3: 1858.

Fuchs RM, Achuff SC, Brunwald L et al.: Electrocardiographic localization of coronary artery narrowing: Studies during myocardial ischemia and infarction in patients with one-vessel disease, *Circulation* 1982;66:1168.

Gibbons GH, and Dzau VJ: The emerging concept of vascular remodeling. *New England Journal of Medicine*, 1994;330, 1431-38

Gibson RS, Crampton RS, Watson DD, et al.: Precordial ST-segment depression during acute inferior myocardial infarction: clinical, scintigraphic and angiographic correlations. *Circulation* 1982; 66:732-41.

Goldstein JA: Pathophysiology and clinical management of right heart ischemia. *Curr Opin Cardiol* 1999; 14: 329-39.

Goldstein JA, Harjai KJ, Boura J, et al.: Comparison of effectiveness of primary angioplasty for proximal versus distal RCA culprit lesion during acute myocardial infarction. *Am J Cardiol* 2002; 90: 1193-97.

Goldstein JA, Lee DT, Pica MC, et al.: Patterns of coronary compromise leading to bradyarrhythmias and hypotension in inferior

myocardial infarction. *Coronary Artery Disease* 2005 August 16(5):265-274.

Graner LE, Greshen, BJ and Orlando, MM:

Bradycardia and its complication in the prehospital phase of AMI *AM. J. Cardiol*, 1973, 32:607.

Gupta A, Lokhandwala YY, Kerkar PG, et al.:

Electrocardiograph[^] differentiation between right coronary and left circumflex coronary arterial occlusion in isolated inferior wall myocardial infarction. *Indian Heart J.* 1999

May-Jun; 51(3):281-4.

Hamm CW, Goldman BU, Heeschen C, et al.:

Emergency room triage of patients with acute chest pain by means of rapid testing for cardiac troponin T or troponin I: *NEnglJMed* 1997; 337: 1684-53.

Harjai KJ, Boura J, Grines L, et al.: Comparison of

L'flicthencsb of priniar) angioplasty for proximal versus distal right coronary artery culprit lesion during acute myocardial infarction. *Am J Cardiol.* 2002 Dec I; 90(11) 193-7.

Hasdai D, Birubbaum Y, Herz I, et al.: SI segment depression in lateral limb leads in inferior wall acute myocardial infarction. Implication regarding the culprit artery and the site of obstruction. *Eur Heart J.* 1995, 16:1549.

Hasdai D, Birnbaur Y, Porter A et al.: Maximal precordial ST-segment depression in leads V4-V6 in patients with inferior wall acute myocardial infarction indicates coronary artery disease involving the left anterior descending coronary artery system. *Int J Cardiol.* 1997, Feb; 58(3):273-8.

Haupt HM, Hutchins GM, and Moore GW: Right ventricular infarction: role of the moderator band artery in determining infarct size. *Circulation* 1983; 67: 1268-1272.

Healy BP: Pathology coronary atherosclerosis. In; Hurst J.W., (ed in chief): *The Heart, Arteries and Veins.* Seventh edition. New York, McGraw-Hill Inc., PP. 924, 1990.

Herz I, Assali AR, Adler Y, et al.: New electrocardiographic criteria for predicting either the right or left circumflex artery as the culprit coronary artery in inferior wall acute myocardial infarction. *Am J Cardiol.* 1997. 80: 1343-5.

Huey BL, Seller GA, Kaiser DL, et al.: A comprehensive analysis of myocardial infarction due to left Circumflex artery occlusion: Comparison with infarction due to right coronary artery and left anterior descending artery occlusion. *J. Am. Coll. Cardiol.* 1988; 12; 1156.

Iseri LT, Humphrey SB, Siner EJ, et al.: Prehospital brady-asystolic cardiac arrest. *Ann Inter Med* 1978; 88: 741-5.

Jeroudi MO, Cheirif J, Habib G, et al.: Prolonged wall motion abnormality after chest pain in patients with unstable angina. *Am Heart J* 1994; 124: 1241- 50.

Kinch JV and Ryan T.I: Right Ventricular infarction. *N EngJ Med.* 1994; 330:1211.

Kinn JW, Ajhmi SC, Samyn JG, et al.: Rapid hemodynamic improvement after reperfusion during right ventricular infarction. *J Am Coll Cardiol* 1995; 26: 1230.

Kleber AG: ST-segment elevation in the electrocardiogram: A sign of myocardial ischemia. *Cardiovasc Res* 2000; 45:111. Klock FJ, Baird MG, Bateman TM, et al.: ACC/AHA/ASNC guideline for the clinical use of cardiac radionuclide imaging. American collage of cardiology .2003.

Kontos MC, Desai PV, Jesse RL, et al.: Usefulness of the admission electrocardiogram for identifying the infarct-related artery in inferior wall acute myocardial infarction. *Am J Cardiol* 1997, 79:182-184.

Kontos MC, Jesse RL, Anderson F, et al.: Comparison of myocardial perfusion imaging and cardiac troponin I in patients admitted to the emergency department with chest pain. *Circulation* 1999; 99:2073.

Kontos MC, Kurdziel K, McQueen R, et al.: Comparison of 2D echocardiology and myocardial perfusion imaging for diagnosing myocardial infarction in emergency department patients. *Am Heart J* 2002; 143: 659.

Kosuge M, Kimura K, Ishikawa T, et al.: Implication of the absence of ST segment elevation in lead V4R in patient who have inferior wall acute myocardial infarction with right ventricular involvement. *Clin Cardiol* 2001; 24: 255-230.

Kosuge M, Kimura K, Ishikawa T, et al.: New electrocardiographic criteria for predicting the site of coronary artery occlusion in inferior wall acute myocardial infarction. *Am J Cardiol* 1998; 82:1318-22.

Kourtesis F, Lichstein E, Cliaocla KD, et al.: Incidence and significance of left anterior hemi block complicating acute inferior myocardial infarction. *Circulation* 1976; 53:784.

Kyriakidis MK, Kouraouklis CB, Papaioannou JT, et al.: Sinus node coronary arteries studied with angiography. *Am. J. Cardiol.* 1983. 31: 749-750.

Lee TH and Goldman L: Evaluation of the patient with acute chest pain. *N Engl J Med* 2000, 342: 1187.

Levin DC, Beckmann CF, Garnic JD, et al.: Frequency and clinical significance of failure to visualize the conus artery during coronary arteriography. *Circulation.* 1981, Apr; 63(4):833-7.

Lew AS, Weiss AT, Shah PK, et al.: Precordial ST-segment depression during acute inferior myocardial infarction: Early thallium-201 scintigraphic evidence of adjacent posterolateral or inferoseptal involvement. *J. Am. Cardiol.*, 1985; 5: 203.

Licka m, Zimmermann R, Zehelein J, et al.: Troponin T concentration 72 hours after myocardial infarction as a serological estimate of infarct size. *Heart* 2002; 87:520-524.

Lin TC, Lee WS, Kong CW et al.: Congenital absence of the left circumflex coronary artery. *Jpn Heart J.* 2003, Nov; 44(6): 1015-20.
Lindh B, Diderholm E, Lagerqvist B, et al.:

Mechanisms behind the prognostic value of troponin 'T' in -unstable coronary artery disease: a FRISC II substudy. J Am Coll Cardiol 2001; 38:979-86.

Little WC, Rogers E\V, and Sodums MT: Mechanisms of anterior ST-segment depression during acute inferior myocardial infarction. Ann. Intern. Med., 1984; 100:226.

Lopez-Sendon J, Coma-Canella I? Alcasena S et al.:

Electrocardiographic findings in acute right ventricular infarction: sensitivity and specificity of electrocardiographic alterations in right precordial leads V4R, V3R, V2 and V3. J Am Coll Cardiol 1985; .6:1273-79.

Mak KH, Chia BL, Tan AT, et al.: Simultaneous ST-segment elevation in lead VI and depression in lead V2. A discordant ECG pattern indicating right ventricular infarction. J Electrocardiol. 1994, Jul; 27(3):203-7.

Mak KH, Chia BL, Tan AT, et al.: Simultaneous ST-segment elevation in lead VI and depression in lead V2. A discordant ECG pattern indicating right ventricular infarction. J Electrocardiol 1994 Jul; 27(3):203-7

Mark AL: The Bezold -Jarish reflex revisited: clinical implications of inhibitory reflex originating in the heart. J Am Coll Cardiol 1983; 1:90-102.

Matetzky S, Novikov M, Gruberg L, et al.: The significance of persistent ST elevation versus early resolution of ST segment elevation after primary PTCA. J Am Coll Cardiol 1999; 34:1932-8.

Matetzky S, Pierr C, Dohn F, et al.: Significance of ST segment elevations in posterior chest leads VI to V9 in patients with acute inferior myocardial infarction: Application for thrombolytic therapy. J. Am. coll. Cardiol. 1998;31:506.

McAllister RG, Frisking GC and Sinclair- Smith BC:

Tricuspid regurgitation following inferior myocardial infarction. Arch Intern MED 1976; 136:95-9.

Meine TJ, Al-Khatib SM, Alexander JH, et al.: Incidence, Predictors, and Outcomes of High-Degree Atrioventricular Block Complicating Acute Myocardial Infarction Treated With Thrombolytic Therapy. *Am Heart J*. 2005; 149(4): 670-674.

Meyers GB, Klein HA and Kiratzki T: Correlation of electrocardiographic and pathologic findings in large anterolateral infarcts. *Am J Cardiol* 1948; 36: 838.

Mikkelsen J, Eskola M, Nikus K, et al.: Fatality of myocardial infarction in relation to the coronary anatomy: role of culprit lesion location. *Ann Med* 2004, Oct; 36(6):474-9.

Mill MR, Wileox BR and Anderson RH: Surgical Anatomy of the Heart. In: Cohn LIT, Edmunds LH Jr, eds. *Cardiac Surgery in the Adult*. New York: McGraw-Hill, 2003:3152.

Mirvis DM: Physiologic basis for anterior ST-depression in patients with acute inferior wall myocardial infarction. *Am Heart J*, 1988; 116:1308.

Moller JE, Egstrup K, Kobler L, et al.: Prognostic importance of systolic and diastolic function after acute myocardial infarction. *Am Heart J* 2003; 145: 147.

Morris F and Brady WJ: ABC of clinical electrocardiography Acute myocardial infarction—Part I. *BMJ* 2002 April; 324:831 -834.

Mukharji J, Murray S, Lewis SE, et al.: Is anterior ST depression with acute transmural inferior infarction due to posterior infarction. A vector cardiograph and scintigraphic study. *J. Am. Coll. Cardiol.* 1984; 4: 28.

Mulvagh SL, DeMaria AN, Feinstein SB, et al.: Contrast echocardiography: current and future applications. *J Am Soc Echocardiography* 2000; 13:331.

Muscholl MW, Oswald M, Mayer C, et al.: Prognostic value of 2D echocardiography in patients presenting with acute chest pain and non diagnostic ECG for ST-segment elevation myocardial infarction. *Inter J of Cardiol* 2002; 84:217-225.

Nair R and Glancy DL: ECG discrimination between right and left circumflex coronary arterial occlusion in patients with acute inferior

myocardial infarction: value of old criteria and use of lead aVR. Chest. 2002, Jul; 122(1): 134-9.

Norell MS, Lyons JP, Gershlick AH, et al.: Assessment of left ventricular performance during percutaneous transluminal coronary angioplasty: a study by intravenous digital subtraction ventriculography. Br Heart J. 1988, Apr; 59(4): 419-28.

Orford JL, Selwyn AP, Ganz P, et al.: The comparative pathobiology of atherosclerosis and restenosis. Am J Cardiol 2003; 86(4B):6H-11H.

O'Rourke RA and Dell' Italia LJ: Diagnosis and management of right ventricular myocardial infarction. Curr Probl Cardio. 2004 Jan; 29(1):6-47.

Pasternak RC, Braunwald E, and sobel BE: Acute myocardial infarction. In: Braunwald E. (ed.): Heart disease. A Textbook of Cardiovascular Medicine.4th edition, 1992.

Phibbs B, Marcus F, Marriott HJC, et al.: Q-wave versus non Q wave myocardial infarction: A meaningless distinction. J Am Coll Cardiol 1999; 33:576.

Pohost GM, Hung L and Doyle M: Clinical use of cardiovascular MRI. Circulation 2003; 108:647.

Polak JF, Holman BL, Wynne J, et al.: Right ventricular ejection fraction: an indicator of increased mortality in patients with congestive heart failure associated with coronary artery disease. J Am Coll Cardiol 1983; 2:217-24.

Ramires JAF, Solimene MC, Savioli RM, et al.: Mortality is not increased with inferior myocardial infarction associated with right ventricular infarction and atrioventricular block. Coron Artery Dis 1993; 4: 965-70.

Serrano CV Jr, Bortolotto LA, Cesar LA, et al.: Sinus bradycardia as a predictor of right coronary artery occlusion in patients with inferior myocardial infarction. Int J Cardiol. 1999 Jan; 68(0):75-82.

Sctaro JF and Cabin HS: Right ventricular infarction. Cardiol Clin. 1992, Feb; 10 (1):69-90.

Sgarbossa EB, Pinski SL, BarbagelataA, et al.: Llectrocardiographic diagnosis of evolving acute myocardial infarction in the presence of left bundle-branch block. *N lingl J Mcd* 1996; 334:431-7.

Shah PK, Maddahi J, Staniloff H ct al.: Variable spectrum and prognostic implications of left and right ventricular ejection fractions in patients with and without clinical heart failure after acute myocardial infarction. *Am J Cardiol* 1986; 58:387-393.

Shah PK, Pickler M, and Berraan DS: Left ventricular ejection fraction determined by radionuclide venuiculography in early stages of first transmural myocardial infarction. *Am. J. Cardiol.*, 1980; 45: 542.

Shiraki H, Yoshikawa T, Anzai T, et al.: Association between Preinfarction Angina and a Lower Risk of Right Ventricular Infarction N. *Engl. J. Med.*, April 2,1998; 338(14): 941 -947.

Sobl BE, and Shell WE: Serum enzyme determinations in the diagnosis and assessment of myocardial infarction. *Circulation*1979; 67: 233.

Serrano CV Jr, Bortolotto LA, Cesar LA, et al.: Sinus bradycardia as a predictor of right coronary artery occlusion in patients with inferior myocardial infarction. *Int .1 Cardiol.* 1999 Jan; 68(0:75-82.

Sctaro JF and Cabin HS: Right ventricular infarction. *Cardiol Clin.* 1992, Feb; 10 (1):69-90.

Sgarbossa EB, Pinski SL, BarbagelataA, et al.: Llectrocardiographic diagnosis of evolving acute myocardial infarction in the presence of left bundle-branch block. *N lingl J Mcd* 1996; 334:431-7.

Shah PK, Maddahi J, Staniloff H ct al.: Variable spectrum and prognostic implications of left and right ventricular ejection fractions in patients with and without clinical heart failure after acute myocardial infarction. *Am J Cardiol* 1986; 58:387-393.

Shah PK, Pickler M, and Berraan DS: Left ventricular ejection fraction determined by radionuclide venuiculography in early stages of first transmural myocardial infarction. *Am. J. Cardiol.*, 1980; 45: 542.

Shiraki H, Yoshikawa T, Anzai T, et al.: Association between Preinfarction Angina and a Lower Risk of Right Ventricular Infarction N. *Engl. J. Med.*, April 2,1998; 338(14): 941 -947.

Soblc BE, and Shell WE: Serum enzyme determinations in the diagnosis and assessment of myocardial infarction. *Circulation* 1979; 67: 233.

Sonndergaard E, Moller JE, Poulsen SH, et al.:

Pseudonormal and restrictive filling patterns predict left ventricular dilatation and cardiac death after a first myocardial infarction. *J Am Coll Cardiol* 2000; 36: 1841.

Srinivas VS, Cannon CP, Gibson CM, et al.: Myoglobin level at 12 hours identify patients at low risk for 30- days mortality after thrombolysis of acute myocardial infarction: Thrombolysis of acute myocardial infarction 10 B substudy. *Am heart J* 2001;142:29-36.

Stewart S and Haste M: Prediction of right ventricular and posterior wall ST elevation by coronary care nurse: The 12-lead electrocardiograph versus the 18-lead ECG. *Heart & Lung* 1996; 25:14-23.

Strasberg B, Pinchas A, Barbash GI, et al.: Importance of reciprocal ST segment depression in leads V5 and V6 as an indicator of disease of the left anterior descending coronary artery in acute inferior wall myocardial infarction. *Br Heart J*. 1990, Jun; 63(6):339-41.

Takahashi T, Sakunia M, Komaki K, et al.: Greater impairment of the right ventricular systolic function in patients with anterior myocardial infarction because of the extent of proximal lesion *Circ J* 2003; 67:396-400.

Tatum JL, Jesse RL, Kontos MC, et al.: Comprehensive strategy for the evaluation and triage of the chest pain patient. *Ann Emerg Med*. 1997 Jan; 29(1): 116-125.

Tsuka Y, Sugiura T, Hatada K, et al.: Clinical significant of ST segment elevation in lead VI in patients with acute inferior wall Q-wave myocardial infarction. *Am Heart J*. 2001, Apr; 141 (4):615-20.

Turhan H, Yilmas MB, Yetkin E, et al.: Diagnostic value of aVL ST deviation for right ventricular involvement in patient with acute MI. *Ann Noninvasive electro-cardiol* 2003;8:185-189.

Uretsky BF, Farguhar DS, and Berez in AF:

Symptomatic myocardial infarction without chest pain: Prevalence and clinical course. *Am. J. Cardiol*. 1977, 40:498.

Wackers FJ, Brown KA, Heller GV, et al: American society of nuclear cardiology positron statement on radionuclide imaging in patient with suspected acute ischemic syndromes in the emergency department or chest pain center. J Nucl Cardiol 2002; 9:246.

Warner RA, Nil HE, Mookherjee S, et al.: Improved electrocardiographic criteria for the diagnosis of combined inferior myocardial infarction and left anterior hemiblock. Am J Cardiol. 1983. Mar 1; 51(5):718-22.

Wellens HJ et al.: The ECG in the Acute Myocardial Infarction and Unstable Angina. Boston, Mass: Kluwer Academic Publishers; 2003.
Wellens HJ: Acute myocardial infarction and left bundle branch block N Engl J Med. 1996; 334:528.

Wellens HJ: The value of the right precordial leads of the electrocardiogram. N Engl J Med 1999; 340:381-383.

Wikins ML, Pryor AD, Maynard C, et al.: An electrocardiographic acuteness score for quantifying the timing of myocardial infarction to guide decision regarding reperfusion therapy. Am J Cardiol 1995; 75:617-20.

Wong CK and Freeilman SB: Usefulness of continuous ST monitoring in inferior wall acute myocardial infarction for describing the relation between precordial ST depression and inferior ST elevation. Am J Cardiol. 1993, Sep 1; 72(7):532-7.

Wong CK and Freedman: Electrocardiographic identification of the infarct-related artery in acute inferior myocardial infarction, Int J Cardiol 1996, 54:5-11.

Wong CK, Freedman SB, Bautovitch GJ et al.: Mechanism and significance of precordial ST depression during inferior wall acute myocardial infarction associated with severe narrowing of the right coronary artery. Am J Cardiol. 1993;71: 1025-1030.

Wong TW, Huang XH, Liu W, et al.: New electrocardiographic criteria for identifying the culprit artery in inferior wall acute myocardial infarction-usefulness of T-wave amplitude ratio in leads II/III and T-wave polarity in the right V5 lead. Am J Cardiol. 2004 Nov 1;94(9):1168-71.

Zalenski RJ, Cooke D, Rydman R, et al.: Assessing the diagnostic value of an ECG constraining leads V4R, V8 and V9: The 15 lead ECG. *Ann. Emerg. Med.*, 1993; 22: 786.

Zehender M, Rasper W, Kaunder E, et al.: Comparison of diagnostic accuracy, time dependency and prognostic impact of abnormal Q Waves, Combined electrocardiographic criteria, and SI abnormalities in right ventricular infarction. *Br. Heart S.* 1994; 72:119.

Zehender M, Kasper W, Kaunder E, et al.: Right ventricular infarction as an independent predictor of prognosis after acute inferior myocardial infarction. *N Engl J Med* 1993; 328:981-8.

Zeiber AM, Drexler H, Saurbier B, et al.: Endothelium-mediated coronary blood flow modulation in humans: effects of age, atherosclerosis, hypercholesterolemia, and hypertension. *J Clin Invest.* 1993; 92:652-662

Zimetbaum PJ and Josephson ME: Use of the Electrocardiogram in Acute Myocardial Infarction *N Engl J Med* 2003; 348:933-40.

Zipes DP: The clinical significance of bradycardiac rhythm in acute myocardial infarction. *Am J cardiol* 1969; 24:814-25