

REFERENCES

Arey, L.B. (1974):

Human Histology, A. textbook in outline form 4th Ed.,
pp. 252-260.

W.B. Saunders Co., Philadelphia, London and Toronto.

Baert, L.L. (1979):

Microdissection in the study of renal structure. The
normal nephron.

Acta Anat. 104:52-60.

Bowman, W. (1940):

On the structure and use of the Malpighian bodies of
the kidney with observations on the circulation through
the gland.

Reprinted in Med. Classics, 5. 249-291.

Boycott, A.B. (1911):

A case of unilateral aplasia of the kidney in a rabbit.

J. Anat. Physiol., London, Vol. 4

Cited by Vimtrup (1926).

Brenda D.P. and Rack, J.H. (1970):

Histological laboratory methods. E.S. Livingstone:

Edinburgh and London.

Culling, C.F.A. (1974):

*Hand-book of Histopathological and Histochemical
Techniques. 3rd ed. PP., 211-425.
Butterworths, London.*

Darmady, E.M. and Stranack F. , (1957):

*Microdissection of the nephrons in disease.
Br. Med. Bull., 13-21.*

Darmady, E.M., Offer, J. and Woodhouse, M.A. (1973):

*The parameters of the aging kidney.
J. Path., 109:195-207.*

Drasch, O. (1877):

*Über das vorkommen Zweierlei verschiedner Gefassknaul
in der Niere. Sitaungsb, D.K. Akad. D. Wissensch., 761
79-102.
Cited by Mueller(1955).*

Emery, J.L. and MacDonald M.S. (1960):

*Involuting and scarred glomeruli in the kidneys of
infants.
Am. J. Pathol. 36; 713-732,*

Faarup, B. (1965):

On the morphology of the juxtaglomerular apparatus.

Acta Anat., 60:20-38.

Fetterman, G.H.; Shuplock, N.A., Philipp, B.S. and Gregg, H.S.

(1965):

The growth and maturation of human glomeruli and proximal convolutions from term to adulthood.

Pediatrics, 35. 601-619.

Freund, J.E. and Williams, F.J. (1975):

Elementary Business Statistics. The modern approach,

2nd edition, PP. 267-283 and 475-477.

Prentice-Hall International Inc., London.

Friedman, H.H., Grayzel, D.M. and Ledrer, M. (1948):

Kidney lesions in stillborn and newborn infants, congenital glomerulosclerosis.

Am. J. Pathol. 18. 699-706.

Friis, C. (1980):

Postnatal development of the pig kidney:

Ultrastructure of the glomerulus and the proximal tubule.

J. Ant. 130; 3 pp. 513-526.

Horster M., Kemler B.H. and Valtin H. (1971):

Intracortical distribution of number and volume of glomeruli during postnatal maturation in the dog.
J. Clin. Invest., 50. Hanover, New Hampshire 769-800.

Horster M. and Valtin, H. (1971):

Post natal development of renal function: Micropuncture and clearance studies in the dog.
J. Clin. Invest., 50:779.

Kampmeier, O.F. (1926):

The metanephros or so socalled permanent kidney in part provisional and vestigial.
Anatomical Record 33, 115-120.

Key, E.A. (1865):

On circulations for hallendena I. Niurarne.
Farh unde skandin. Naturf., stockholm, PP. 685-728.
Cited by Mueller (1955).

Kulz, L. (1899):

Untersuchungen uber das postfotale wachstum der menschen.
Niere Diss for the M.D., Kiel, 20 PP. Namburg, a.s.
Cited by Maluf (1978).

Leeson, T.S. and Leeson C.R. (1981):

Histology, 4th edition, pp. 432-442.

W.B. Saunders company/Philadelphia/London/Toronto
Holt-Saunders Japan, LTD/Tokyo.

Loggie, I.M.H., Kleinman, L.I. and Van Maanen, E.F. (1975):

Renal function and diuretic therapy in infants and children.

J. Ped., 86: 485-496.

MacDonald, M.S. and Emery, J.L. (1959):

The late intra-uterine and postnatal development of human renal glomeruli.

J. Anat., 93: 331-341.

Maluf, N.S.R. (1978):

Anatomy of the kidneys of newly born pigmy hippopotamus (Chaeropsis liberiensis morton).

Vet. Med. Anat. Histol. 7:28.

Noberg, E. (1929):

Anzahl und Grobe der Glomeruli renales beim Menschen nebst Methoden diese Zahlenmässig Festzustellen. Z. Mikr-Anat. Foursch., 18, 271-311.

Cited by Maluf (1978).

Muller C. B. (1955):

Anatomy of the glomerulus.

Am. J. Med. Stout D.G. and 18:267-276.

Nash, M.A. and Edelmann, C.M. (1973):

The developing kidney.

Nephron 11,71-90

Oliver, J.R. (1945):

*New directions in renal morphology: A method, its results
and its future.*

Harvey lecture, series XL., 102.

Paatela, M. (1963):

*Renal microdissection in infants with special
reference to the congenital nephrotic syndrome.*

Annals Paediat. 9,, 21.

Peter, K. (1907):

Über die Nierenkanäle des Menschen und einiger sauge

thiere. Verh. Anat. Ges., 21 ti Versammlung. Würzburg.

1909 untersuehungen iiber Bau und Entwi ekelung der Niere.

Jena.

Cited by Vimtrup , (1926).

Potter, E.L. (1972):

Normal and abnormal development of the kidney. Chicago year book Publ., XI-305pp.

Cited by Maluf (1978).

Roosen-Runge, E.C. (1949):

Retardation of postnatal development of the kidney in persons with early cerebral lesions.

Am. J. of diseases of children, 77, 185-200.

Sorensen, H.F. and Ledet T. (1972):

Quantitative studies of the renal corpuscles. Intraglomerular, Interglomerular and interfocal variation in the normal kidney.

Acta Pathologica et microbiologica Scandinavica 80. 721-728.

Souster L.P. and Emery J.L. (1980):

The sizes of glomeruli in fetuses and infants,

J. Anat. 130, 595-602.

Tisher, C.C., Bulger, R.E. and Trump, B.F. (1968):

Human renal ultrastructure. III. The distal tubule in healthy individuals.

Lab. Invest. 181:655-668.

Trautmann, A. and Fiebiger, J. (1957):

Fundamentals of the histology of domestic animals.
translated and revised from the 8th and 9th German
Eds., pp. 245-254,
Constock Publishing Associates Ithaca, New York.

Vimtrup Bj. (1926):

*On the number, shape, Structure, and Surface are of
the glomerulus in the kidneys of man and mammals.*
Am. J. Anat. 41:123-151.

Yadava, R.R. and Calhoun, M.L. (1958):

Comparative histology of the kidney of domestic animals.
Am. J. Vet. Res., 19:958-968.

Zolnai B. and Palkovits, M. (1965):

Glomerulonetric III.
Acta Biologica Scientiarian Hungaricae, 15 490-423.