

References

Abbate, M.; Brown, D. and Bonventre, J.V. (1999):

Expression of NCAM recapitulates tubulogenic development in kidneys.

Physiol., 277: 454-463.

Abuljadayel, I.S. (2003) :

Induction of stem cell-like plasticity in mononuclear cells derived from unmobilised adult human peripheral blood.

Current Medical Research and Opinion, 19;355–375.

Aiuti, A.; Slavin, S.; Aker, M.; Ficara, F.; Deola, S.; Mortellaro, A.; Morecki, S.; Andolfi, G.; Tabucchi, A.; Carlucci, F.; Marinello, E.; Cattaneo, F.; Vai, S.; Servida, P.; Miniero, R.; Roncarolo, M.G. and Bordignon, C. (2002):

Correction of ADA-SCID by stem cell gene therapy combined with non myeloablative conditioning.

Science, 296:2410-2413.

Åkerud, P.; Holm, P.C.; Castelo-Branco, G.; Sousa, K.; Rodriguez, F.J. and Arenas, E. (2002):

Persephin-overexpressing neural stem cells regulate the function of nigral dopaminergic neurons and prevent their degeneration in a model of Parkinson's disease.

Molecular and Cellular Neuroscience, 21: 205-222.

Akiyama, Y.; Radtke, C. and Kocsis, J. (2002):

Remyelination of the rat spinal cord by transplantation of identified bone marrow stromal cells.

J Neurosci;22:6623–6630.

Alexander, D.B. and Goldberg, G.S. (2003) :

Transfer of biologically important molecules between cells through gap junction channels.

Curr. Med. Chem.,10:2045–2058.

Alison, M.R.; Poulson, R.; Jeffery, R.; Dhillon, A.P.; Quaglia, A.; Jacob, J.; Novelli, M.; Prentice, G.; Williamson, J. and Wright, N.A., (2000):

Hepatocytes from non-hepatic adult stem cells.
Nature, 406:257-263.

Anderson, D.J. (2001):

Stem cells and pattern formation in the nervous system: the possible vs the actual.

Neuron., 30:19–35.

Andrews, R.G.; Briddell, R.A.; Knitter, G.H.; Rowley, S.D.; Appelbaum, F.R. and McNiece, I.K. (1995):

Rapid engraftment by peripheral blood progenitor cells mobilized by recombinant human stem cell factor and recombinant human granulocyte colony-stimulating factor in nonhuman primates.

Blood., 85 (1):15-20.

Anglani, F.; Forino, M.; Toso, E.; and Torregrossa, R., (2004):

Histomorphology and molecular biology of kidney.

Curr. Biol., 8:145–155.

Anversa, P.; Olivetti, G. and Meggs, L.G. (1993):

Cardiac anatomy and ventricular loading after myocardial infarction.
Circulation, 87: 22–27.

Arndt-Jovin, D.J. and Jovin, T.M. (1977):

Analysis and sorting of living cells according to deoxyribonucleic acid content.

J. Histochem. Cytochem., 25:585–589.

Asakura, A. and Rudnicki, M.A. (2002):

Side population cells from diverse adult tissues are capable of in vitro hematopoietic differentiation.

Exp. Hematol., 30:1339–1345.

Asakura, A.; Komaki, M. and Rudnicki, M. (2001):

Muscle satellite cells are multipotential stem cells that exhibit myogenic, osteogenic, and adipogenic differentiation.

Differentiation, 68: 245–53.

Bannerman, P.; Nichols, W.; Puhalla, S.; Oliver, T.; Berman, M. and Pleasure, D. (2000):

Early migratory rat neural crest cells express functional gap junctions: evidence that neural crest cell survival requires gap junction function.

J. Neurosci. Res., 61: 605–615.

Baserga, R. and Rubin, R. (1993):

Cell cycle and growth control.

Crit. Rev. Eukaryot. Gene Exp., 3:47-61.

Baserga, R.; Estensen, R.D. and Petersen, R.O. (1965):

Inhibition of DNA synthesis in Ehrlich ascites cells by actinomycin D. II. The presynthetic block in the cell cycle.

Proc. Natl. Acad. Sci. USA, 54:1141-1148.

Beltrami, A.P.; Barlucchi, L.; Torella, D.; Baker, M.; Limana, F.; Chimenti, S.; Kasahara, H.; Rota, J.; Musso, E.; Urbanek, K.; Leri, A.; Kajstura, J.; Nadal-Ginard, B.; and Anversa, P. (2003):

Adult cardiac stem cells are multipotent and support myocardial regeneration.

Cell, 114: 763–776.

Benedetti, S.; Pirola, B.; Pollo, B.; Magrassi, L.; Bruzzone, M.G.; Rigamonti, D.; Galli, R.; Selleri, S. D.; Fraja, C.; Vescovi, A.; Cattaneo, E. and Finocchiaro, G. (2000):

Gene therapy of experimental brain tumors using neural progenitor cells.

Nature Medicine, 6: 447-450.

Bernard-Kargar, C. and Ktorza, A. (2001):

Endocrine pancreas plasticity under physiological and pathological conditions.

Diabetes, 50 (Suppl. 1): 30-35.

Bhattacharya, S.; Jackson, J.D.; Das, A.V.; Thoreson, W.B.; Kuszynski, C.; James, J.; Joshi, S. and Ahmad, I. (2003):

Direct identification and enrichment of retinal stem cells/progenitors by Hoechst dye efflux assay.

Invest. Ophthalmol. Vis. Sci., 44 :2764–2773.

Bianchi, G.; Muraglia, A.; Daga, A. and Corte G. (2001):

Microenvironment and stem properties of bone-marrow derived mesenchymal cells.

Wound Repair Regen., 9: 460-466.

Bianco, P.; Riminucci, M.; Grothos, S.; and Gehron, P. (2001):

Bone marrow stromal cells :nature, biology, and potential applications. Stem cells , 19(3):180-192.

Bizzozzero, G. (1894):

The growth and regeneration of organism

Br. Med. J., 1:728-782.

Bjerknes, M. and Cheng, H. (1999):

Clonal analysis of mouse intestinal epithelial progenitors. Gastroenterology, 116:7–14.

Bjerknes, M.; Cheng, H.; Hay, K.; and Gallinger, S. (1997):

APC mutation and the crypt cycle in murine and human intestine.

Am. J. Pathol., 150: 833–839 .

Bjornson, C.R.; Rietze, R.L. and Reynolds, B.A. (2000):

Turning brain into blood: a hematopoietic fate adopted by adult neural stem cells *in vivo*.

J. Clin. Invest., 105: 1663-1668.

Borlongan, C.V. and Hess, D.C. (2003):

G-CSF-mobilized human peripheral blood for transplantation therapy in stroke.

Cell Transplantation, 12: 447-448.

Bouwens, L. (1998):

Transdifferentiation versus stem cell hypothesis for the regeneration of islet beta-cells in the pancreas.

Microsc. Res. Tech., 5:332-336.

Bree, R.T.; Stenson-Cox, C.; Greal, M.; Byrnes, L.; Gorman, A.M. and Samali, A. (2002):

Cellular longevity: role of apoptosis and replicative senescence. Biogerontology, 3:195–206.

Broxmeyer, H.E.; Srour, E.F.; Hangoc, G.; Cooper, S.; Anderson, S.A. and Bodine, D.M. (2003):

High-efficiency recovery of functional hematopoietic progenitor and stem cells from human cord blood cryopreserved for 15 years. Proceedings of the National Academy of Sciences USA, 100: 645-650.

Cai, J.; Wu, Y. and Mirua, T. (2002):

Properties of a fetal multipotent neural stem cells, identifying them as nonependymal.

Neuron, 35: 865–75.

Camargo, F.D.; Green, R.; Capetenaki, Y.; Jackson, K.A. and Goodell, M.A. (2004):

Single hematopoietic stem cells generate skeletal muscle through myeloid intermediates.

Nat. Med., 9:1520–1527.

Cao, Y.; Zhao, Z.; Gruszczynska-Biegala, J. and Zolkiewska, A. (2003):

Role of metalloprotease disintegrin ADAM12 in determination of quiescent reserve cells during myogenic differentiation *in vitro*.

Mol. Cell Biol., 23: 6725–38.

Catherine, M.; Verfillie, E.; Martin, F.; Pera, R. and Peterm, L. (2002):

Stem cell :hype and reality

American society of hematology;(Annual proceedings,12-17).

Chacko, D.M.; Das, A.V.; Zhao, X.; James, J.; Bhattacharya, S. and Ahmad, I. (2003):

Transplantation of ocular stem cells: the role of injury in incorporation and differentiation of grafted cells in the retina.

Vision Research, 43: 937-946.

Charrasse, S.; Meriane, M.; Comunale, F.; Blangy, A. and Gauthier-Rouviere, G. (2002):

N- cadherin-dependent cell contact.

Science, 158: 953–965.

Chenn, A. and Walsh, C.A. (2002):

Regulation of cerebral cortical size by control C. N-cadherin-dependent cell-cell contact regulates Rho GTPases and beta-catenin localization in mouse C2C12 myoblasts.

J. Cell Biol., 105: 1669–74.

Clyne, R.K. and Kelly, T.J. (1995):

Genetic analysis of an ARS element from the fission yeast *Schizosaccharomyces pombe*.

EMBO. J., 14 : 6348-6357.

Cossu, G. and Mavilio, F. (2000):

Myogenic stem cells for the therapy of primary myopathies: wishful thinking or therapeutic perspective?

J. Clin. Invest., Oncogene, 18:1537–1544.

Cottler-Fox, M.; Lapidot, T.; Petit, I.; Kollet, O.; Dippersio, J.; Link. D. and Devine, S. (2003):

Stem cells mobilization .

American society of hematology ;(Annual proceedings,22-33).

Davidson, E. H. (1986):

Gene activity in early development

Academic press, Orlando, Florida, 3rd edition.

De -Bari, C.; Dell 'Accio, F.; Tylzanowski, P. and Luyten, F.P. (2001):

Multipotent mesenchymal stem cells from adult human synovial membrane.

Arthritis Rheumatism, 44: 1928-1942.

Deng, W.; Obrocka, M.; Fischer, I. and Prockop, D.J. (2001):

In vitro differentiation of human marrow stromal cells into early progenitors of neural cells by conditions that increase intracellular cyclic AMP.

Biochem. Biophys. Res. Commun., 282:148–152.

Deryugina, E.I. and Muller-Sieburg, C.E. (1993):

Stromal cells in long-term cultures keys to the elucidation of hematopoietic development?.

Crit. Rev. Immunol., 13:115–119.

Doetsch, F.; Caille, I.; Lim, D.A.; Garcia-Verdugo, J.M. and Alvarez-Buylla, A. (1999):

Subventricular zone astrocytes are neural stem cells in the adult mammalian brain.

Cell, 97: 703–16.

Doetsch, F.; Garcia-Verdugo, J.M. and Alvarez-Buylla, A. (1997):

Cellular composition and three-dimensional organization of the subventricular germinal zone in the adult mammalian brain.

J. Neurosci., 17: 5046–61.

Dowdy, S.F.; Hinds, P.W.; Louie, K.; Reed, S.I.; Arnold, A. and Weinberg, R.A. (1993):

Physical interaction of the retinoblastoma protein with human D cyclins.

Cell, 73:499-511.

Eglitis, M. and Mezey, E. (1997):

Hematopoietic cells differentiate into both microglia and macroglia in the brains of adult mice.

Proc. Natl. Acad. Sci. USA, 94:4080–4085.

Eiges, R.; Schuldiner, M.; Drukker, M.; Yanuka, O.; Itskovitz-Eldor, J. and Benvenisty, N. (2001):

Establishment of human embryonic stem cell-transduced clones carrying a marker of undifferentiated cells.

Curr. Biol., 11: 514-518.

El Ansary, M., (2006)

Introduction to stem cells.

Clinical Pathology Department, Faculty of Medicine, Cairo University, Personal communication.

Falciatori, I.; Borsellino, G.; Haliassos, N.; Battistini, L.; Bernardi, G.; Stefanini, M. and Vicini, E. (2004):

Identification and enrichment of spermatogonial stem cells displaying side-population phenotype in immature mouse testis.

FASEB. J., 18:376–388.

Fallon, J.; Reid, S.; Kinyamu, R.; Opole, I.; Opole, R.; Baratta, J.; Korc, M.; Endo, T.L.; Duong, A.; Nguyen, G.; Karkehabadhi, M.; Twardz, D.; Patel, S. and Loughlin, S. (2000):

In vivo induction of massive proliferation, directed migration, and differentiation of neural cells in the adult mammalian brain.

Proceedings of the Nat. Acad. of Sci. USA, 97:14686-14691.

Fausto, N.; Lemire, J.M. and Shojiri, N. (1993):

Cell lineages in hepatic development and the identification of progenitor cells in normal and injured liver.

Proc. Soc. Exp. Biol. Med., 204: 237–41.

Ferrari, G. and Mavilio, F. (2002):

Myogenic stem cells from the bone marrow: a therapeutic alternative for muscular dystrophy?.

Neuromuscul. Disord, 12 (suppl 1): 7–10.

Ferrari, G.; Cusella, D.; Coletta, M. and Angelis, G. (1998):

Muscle regeneration by bone marrow-derived myogenic progenitors.
Science, 279: 1528-1530.

Ferrari, G.; Stornaiuolo, A. and Mavillo, F. (2001):

Failure to correct murine muscular dystrophy.
Nature, 411:1014–1015.

Fortunel, N.O.; Out, H.H.; Ng, H.H.; Chen, J.; Mu, X.; Chevassut T.; Li, X.; Joseph, M.; Bailey, C.; Hatzfeld, J.A.; Hatzfeld, A.; Usta F.; Vega, V.B.; Long, P.M.; Libermann, T.A. and Lim, B. (2003).

Comment on Stemness: transcriptional profiling of embryonic and adult stem cells and (a stem cell molecular signature).
Science, 12 : 302:393.

Friedrich, U.; Griesse, E.; Schwab, M.; Fritz ,P.; Thon, K. and Klotz U. (2000):

Telomere length in different tissues of elderly patients.
Mech. Ageing Dev., 119:89–99.

Gage, F.H. (2000):

Mammalian neural stem cells
Science, 287:1433-1438.

Garbuzova-Davis, S.; Willing, A.E.; Zigova, T.; Saporta, S.; Justen, E.B.; Lane, J.C.; Hudson, J.E.; Chen, N.; Davis, C.D. and Sanberg P.R. (2003):

Intravenous administration of human umbilical cord blood cells in a mouse model of amyotrophic lateral sclerosis: distribution, migration, and differentiation.

Journal of Hematotherapy & Stem Cell Research, 12: 255–270.

Gerald, D.; Fischbach, A. and Ruth, L. (2004):

Neural stem cells.
Circ. Res., 92:598-608.

Gibson, M.C. and Schultz, E. (1982):

The distribution of satellite cells and their relationship to specific fiber types in soleus and extensor digitorum longus muscles.

Anat. Rec., 202: 329–37.

Gibson, M.C. and Schultz, E. (1983):

Age-related differences in absolute numbers of skeletal muscle satellite. Cells, Muscle, Nerve, 6: 574–80.

Gill, S.S.; Patel, N.K.; Hotton, G.R.; O'Sullivan, K.; McCarter, R.; Bunnage, M.; Brooks, D.J.; Svendsen, C.N. and Heywood P. (2003):

Direct brain infusion of glial cell line-derived neurotrophic factor in Parkinson disease.

Nature Medicine, 9: 589-595.

Ginis, I.; Luo, Y. and Miura, T. (2003):

Differences between human and mouse embryonic stem cells.

Dev. Biol., 12:45-56.

Goodell, M.A. (2002):

Stem cell identification and sorting using the Hoechst 33342 side population (SP). In: Current Protocols in Cytometry. eds. Robinson JP, Darzynkiewicz Z, Dean PN, PP 1-11, Willy-Liss Inc., New York.

Goodell, M.A.; Rosenzweig, M. and Kim, H. (1997):

Dye efflux studies suggest that hematopoietic stem cells expressing low or undetectable levels of CD34 antigen exist in multiple species.

Nat. Med., 3:1337–1345.

Gronthos, S.; Mankani, M.; Brahimi, J.; Robey, P.G. and Shi, S. (2000) :

Postnatal human dental pulp stem cells (DPSCs) in vitro and in vivo. Nati. Acad. of Sci. USA, 97:13625-13630.

Grove, J.E.; Lutzko, C.; Priller, J.; Henegariu, O.; Theise, N.D.; Kohn, D.B. and Krause, D.S. (2002):

Marrow-Derived Cells as Vehicles for Delivery of Gene Therapy to Pulmonary Epithelium.

Stem Cells, 12: 645-651.

Gu, D.; Lee, M. S.; Krah, T. and Sarvetnick, N. (1994):

Transitional cells in the regenerating pancreas.

Development, 120: 1873-1881.

Gupta, P.; Oegema, T.R.; Brazil, J.J.; Dudek, A.Z.; Slungaard, A. and Verfaillie, C.M. (1998):

Structurally specific heparan sulfates support primitive human hematopoiesis by formation of a multimolecular stem cell niche.

Blood, 92 :4641–4651.

Gussoni, E.; Soneoka, Y. and Strickland, C.D. (1999):

Dystrophin expression in the mdx mouse restored by stem cell transplantation.

Nature, 401:36-45.

Hao, H.N.; Zhao, J. and Thamos, R.L. (2003):

Fetal human hematopoietic stem cells can differentiate sequentially into neural stem cells and then astrocytes in vitro.

J. Hematother Stem Cell Res., 12:23-32.

Hamlin, J.L.; Mosca, P.J. and Levenson, V.V. (1994):

Defining origins of replication in mammalian cells.

Biochim. Biophys. Acta., 1198 : 85-111.

Heichman, K.A. and Roberts, J.M., (1994):

Rules to replicate.

Cell, 79: 557-562.

Hematti, P.; Sloand, E.M.; Carvallo, C.A.; Albert, M.R.; Yee, C.L.; Fuehrer, M.M.; Blancato, J.K.; Kearns, W.G.; Barrett, J.A.; Childs, R.W.; Vog, J.C. and Dunbar, C.E. (2002):

Absence of donor-derived keratinocyte stem cells in skin tissues cultured from patients after mobilized peripheral blood hematopoietic stem cell transplantation.

Exp. Hematol., 30: 943–949.

Herrera, P.L. (2000):

Adult insulin- and glucagon-producing cells differentiate from two independent cell lineages.

Development, 127:2317-2322.

Herzenberg, L.A. and De Rosa, S.C. (2000):

Monoclonal antibodies and the FACS: complementary tools for immunobiology and medicine.

Immunol. Today, 21: 383-390.

Hess, D.; Li, L.; Martin, M.; Sakano, S.; Hill, D.; Strutt, B.; Thyssen, S.; Gray, D.A. and Bhatia, M. (2003):

Bone marrow-derived stem cells initiate pancreatic regeneration.

Nat. Biotechnol., 21:763–770.

Hierlihy, A.M.; Seale, P.; Lobe, C.G.; Rudnicki, M.A. and Megeney, L.A. (2002):

The post-natal heart contains a myocardial stem cell population.

F.E.B.S. Lett., 530: 239–243.

Horwitz, E.M.; Gordon, P.L.; Koo, W.K.; Marx, J.C.; Neel, M.D.; Mc Nall, R.Y.; Muul, L. and Hofmann, T. (2002):

Isolated allogeneic bone marrow-derived mesenchymal cells engraft and stimulate growth in children with osteogenesis imperfecta: Implications for cell therapy of bone.

Nati. Acad. Sci. USA, 99: 8932-8937.

Horwitz, E.; Prockop, D.; and Fitzpatrick, L. (1999):

Osteogenesis imperfecta calls for caution.

Nat. Med., 5:466–7.

Hou, Z.; Nguyen, Q. and Frenkel, B. (1998):

Osteoblast-specific gene expression after transplantation of marrow cells: implications for skeletal gene therapy.

Proc. Natl. Acad. Sci. U S A., 96:7294–9.

Howard, A. and Pelt, S.R. (1951):

Nuclear incorporation of ^{32}P as demonstrated by autoradiography.

Exp. Cell Res., 2:178-187.

Quoted from Stem cell biology and gene therapy. Edited by Peter J. Quesenberry, Gary S. Stein, Bernard G. Forget, Sherman M. Weissman, chapter (3), PP 41-66.

Copyright , 1998 Wiley-Liss, Inc.

ISBNs: 0-471-14656-0 (Paper); 0-471-22395-6 (Electronic).

Hunter, T. and Pines, J. (1994):

Cyclins and cancer II: Cyclin D and CDK inhibitors come of age.

Cell, 79:573-582.

Jackson, K.A.; Majka, S.M.; Wang, H.; Pocius, J.; Hartley, C.J.; Majesky, M.W.; Entman, M.L.; Michael, L.H.; Hirschi, K.K. and Goodell, M.A. (2001):

Regeneration of ischemic cardiac muscle and vascular endothelium by adult stem cells.

J. Clin. Invest., 107:1395–1402.

Jensen, G.S. and Drapeau, C. (2002):

The use of in situ bone marrow stem cells for the treatment of various degenerative diseases.

Medical Hypotheses, 59:422-428.

Jiang, Y.; Jahagirdar, B.N.; Reinhardt, R.L.; Schwartz, R.E.; Keene, C.D.; Ortiz-Gonzalez, X.R.; Reyes, M.; Lenvik, T.; Lund, T.; Blackstad, M.; Du, J.; Aldrich, S.; Lisberg, A.; Low, W.C.; Largaespada, D.A. and Verfaillie, C.M. (2002):

Pluripotency of mesenchymal stem cells derived from adult marrow. Nature, 418 : 41–49.

Jin, H.K.; Carter, J.E.; Huntley, G.W. and Schuchman, E.H. (2002):

Intracerebral transplantation of mesenchymal stem cells into acid sphingomyelinase-deficient mice delays the onset of neurological abnormalities and extends their life span.

J. Clin. Invest., 109: 1183-1191.

Jordan, C.T.; Mckearn, J.P. and Lemischka, I.R. (1990):

Cellular and developmental properties of fetal hematopoietic stem cells.
Cell, 61:953–963.

Kakinuma, S.; Tanaka, Y.; Chinzei, R.; Watanabe, M.; Shimizu-Saito, K.; Hara, Y.; Teramoto, K.; Arii, S.; Sato, C.; Takase, K.; Yasumizu, T. and Teraoka, H. (2003):

Human Umbilical Cord Blood as a Source of Transplantable Hepatic Progenitor Cells.

Stem Cells, 21: 217-227.

Kale, S.; Karihaloo, A.; Clark, P.R.; Kashgarian, M.; Krause, D.S. and Cantley, L.G. (2003):

Bone marrow stem cells contribute to repair of the ischemically injured renal tubule.

J. Clin. Invest. , 112 : 42–49.

Kanazawa, Y. and Verma, I.M. (2003):

Little evidence of bone marrow-derived hepatocytes in the replacement of injured liver.

Proc. Natl. Acad. Sci. U S A, 100:11850–11853.

Kang, X.Q.; Zang, W.J.; Bao, L.J.; Li, D.L.; Song ,T.S.; Xu, X.L. and Yu, X.J. (2005):

Fibroblast growth factor-4 and hepatocyte growth factor induce differentiation of human umbilical cord blood derived mesenchymal stem cells into hepatocytes.

World J. Gastroentero., 111: 7461–65.

Kawada, H. and Ogawa, M. (2001):

Bone marrow origin of hematopoietic progenitors and stem cells in murine muscle.

Blood, 98:2008–2013.

Kikyo, N. and Wolffe, A.P. (2000):

Reprogramming nuclei: insights from cloning, nuclear transfer and heterokaryons.

J. Cell Sci., 113:11–20.

Kim, B.; Seo, J.H.; Bubien, J.K. and Oh, Y.S. (2002 a):

Differentiation of adult bone marrow stem cells into neuroprogenitor cells in vitro.

Neuroreport, 13:1185–1188.

Kim, M.; Turnquist, H.; Jackson, J.; Sgagias, M.; Yan, Y.; Gong, M.; Dean, M.; Sharp, J.G. and Cowan, K. (2002 b):

The multidrug resistance trans- porter ABCG2 (breast cancer resistance protein 1) effluxes Hoechst 33342 and is overexpressed in hematopoietic stem cells.

Clin. Cancer Res., 8:22–28.

King, R.W.; Deshaies, R.J.; Peters, J.M. and Kirschner, M.W. (1997):

How proteolysis drives the cell cycle.

Science, 274:1652-1659.

King, R.W.; Jackson, P.K. and Kirschner, M.W. (1994):

Mitosis in transition.

Cell, 79: 563-571.

Kollet, O.; Shivtiel, S.; Chen, Y.Q. ; Suriawinata, J.; Thung, S.N.; Dabeva, M.D.; Kahn, J.; Spiegel, A.; Dar, A.; Samira, S.; Goichberg, P.; Kalinkovich, A.; Arenzana-Seisdedos, F.; Nagler, A.; Hardan, I.; Revel, M.; Shafritz, D.A. and Lapidot, T. (2003):

HGF, SDF-1, and MMP-9 are involved in stress-induced human CD34+ stem cell recruitment to the liver.

J. Clin. Invest, 112: 160–69.

Korbling, M.; Katz, R.; Khanna, A. ; Ruifrok, A.C.; Rondon, G.; Albitar, M.; Champlin, R.E. and Estrov, Z. (2002):

Hepatocytes and epithelial cells of donor origin in recipients of peripheral-blood stem cells.

N. Engl. J. Med., 346:738–746.

Kotton, N.; Cardoso, W.V; Sanderson, E.A.; Summer, R.S.; Williams, M.C. and Fine, A. (2001):

Bone marrow-derived cells as progenitors of lung alveolar epithelium. *Development*, 128 :5181–5188.

Krause, D.S.; Ito, T. and Fackler, M.J. (1994):

Characterization of murine CD34, a marker for hematopoietic stem and progenitor cells.

Blood, 84:691–701.

Krause, D.S.; Theise, N.D. and Collector, M.I. (2001):

Multi-organ, multi-lineage engraftment by a single bone marrow-derived stem cell.

Cell, 105:369–377.

Lagasse, E.; Connors, H.; Al-Dhalimy, M.; Reitsma, M.; Dohse, M.; Osborne, L.; Wang ,X.; Finegold, M.; Weissman, I.L. and Grompe, M. (2000):

Purified hematopoietic stem cells can differentiate into hepatocytes in vivo.

Nat. Med., 6:1229–1234.

Lalande, M.E. and Miller, R.G. (1979):

Fluorescence flow analysis of lymphocyte activation using Hoechst 33342 dye.

J. Histochem. Cytochem., 27: 394–397.

Lee, J.Y.; Musgrave, D.; Pelinkovic, D.; Fukushima, K.; Cummins, J.; Usas, A.; Robbins, P.; Fu, F. H. and Huard J. (2001):

Effect of bone morphogenetic protein-2-expressing muscle-derived cells on healing of critical-sized bone defects in mice.

The Journal of Bone and Joint Surgery, 83: 1032-1039.

Levin, M. (2002):

Isolation and community: a review of the role of gap junctional communication in embryonic patterning.

J. Membr. Biol., 185:177–192.

Ley, K., (2001):

Physiology of inflammation, P 546.
Oxford University Press, New York .

Liechty, K.; MacKenzie, T. and Shaaban, A. (2000):

Human mesenchymal stem cells engraft and demonstrate site-specific differentiation after in utero transplantation in sheep.
Nat. Med., 6:1282–6.

Lipsett, M. and Finegood, D.T. (2002):

β -cell neogenesis during prolonged hyperglycemia in rats.
Diabetes, 51: 1834-1841.

Lobo, M.V.; Alonso, F.J.; Redondo, C.; Lopez-Toledano, M.A.; Caso E.; Herranz, A.S.;Paino, C.L.; Reimers, D. and Bazan, E. (2003):

Cellular characterization of epidermal growth factor–expanded free-floating neurospheres.
J. Histochem. Cytochem., 51:89–103.

Lu, B.; Jan, L. and Jan, Y.N. (2000):

Control of cell divisions in the nervous system: symmetry and asymmetry.
Annu. Rev. Neurosci., 23:531–556.

Lu ,M.; Grove, E.A. and Miller, R.J. (2002).

Abnormal development of the hippocampal dentate gyrus in mice lacking the CXCR4 chemokine receptor.
Proc. Natl. Acad. Sci. USA, 99:7090–7095.

Luskin, M.B. (1993):

Restricted proliferation and migration of postnatally generated neurons derived from the forebrain subventricular zone.
Neuron, 11: 173–89.

MacLachlan, T.K.; Sang, N. and Giordano, A. (1995):

Cyclins, cyclin-dependent kinases and Cdk inhibitors: Implications in cell cycle control and cancer.

Crit. Rev. Eukaryot. Gene Exp., 5 : 127-156.

Mahmood, A.; Lu, D.; Wang, L.; Li, Y.; Lu, M. and Chopp, M. (2001):

Treatment of traumatic brain injury in female rats with intravenous administration of bone marrow stromal cells.

Neuro-surgery, 49:1196–1203.

Macy- Tarnouski, T. and Aleksander, L.S. (2006):

Adult stem cells and their ability to differentiate.

Med. Sci. Monit., 12 (8):154-163.

Makino, S.; Fukuda, K.; Miyoshi, S.; Konishi, F.; Kodama, H.; Pan, J.; Sano, M.; Takahashi, T.; Hori, S.; Abe, H.; Hata, J.; Umezawa, A. and Ogawa, S. (1999):

Cardiomyocytes can be generated from marrow stromal cells in vitro.

J. Clin. Invest., 103:697–705 .

Martin, C.M.; Meeson, A.P.; Robertson, S.M.; Hawke, T.J.; Richardson, J.A.; Bates, S.; Goetsch, S.C.; Gallardo, T.D. and Garry, D.J. (2004):

Persistent expression of the ATP-binding cassette transporter, Abcg2, identifies cardiac SP cells in the developing and adult heart.

Dev. Biol., 265: 262–275.

Mason, R.J.; Williams, M.C. and Moses, H.L. (1997):

Stem cells in lung development, disease, and therapy.

Cell Mol. Biol., 16 : 355–363.

Matsumoto, K. and Nakamura, T. (2001):

Hepatocyte growth factor renotropic role and potential therapeutics for renal diseases.

Kidney Int., 59: 2023-2038.

McKinney-Freeman, S.L.; Jackson, K.A.; Camargo, F.D.; Ferrari, G.; Mavilio, F. and Goodell, M.A. (2002):

Muscle-derived hematopoietic stem cells are hematopoietic in origin.

Proc. Natl. Acad. Sci. USA, 99 :1341–1346.

Meller, D.; Pires, R.T. and Tseng, S.C. (2002):

Ex vivo preservation and expansion of human limbal epithelial stem cells on amniotic membrane cultures.

British Journal of Ophthalmology, 86: 463-471.

Mezey, E.; Chandross, K.J.; Harta, G.; Maki, R.A.; and McKercher, S.R. (2000):

Turning blood into brain: cells bearing neuronal antigens generated in vivo from bone marrow.

Science, 290:1779–1782.

Migishima, F.; Oikawa, A.; Kondo, S.; Ema, H.; Morita, Y.; Nakauchi, H.; Yokoyama, M.; Song, S.Y.; Nishijima, M.; Okabe, M. and Shinohara, N. (2003):

Full reconstitution of hematopoietic system by murine umbilical cord blood.

Transplantation, 75:1820–1826.

Minasi, M.G.; Riminucci, M.; De Angelis, L.; Borello, U.; Berarducci, B.; Innocenzi, A.; Caprioli, A.; Sirabella, D.; Baiocchi, M.; De Maria, R.; Boratto, R.; Jaffredo, T.; Broccoli, V.; Bianco, P. and Cossu, G. (2002):

The mesoangioblast: a multipotent, self-renewing cell that originates from the dorsal aorta and differentiates into most mesodermal tissues.

Development, 129: 2773-2783.

Mitchell, K.; Weiss, M.L.; Mitchell, B.M.; Martin, P.; Davis, D.; Morales, L.; Helwig, B.; Beerenstrauch, M.; Abou-Easa, K.; Hildreth, T.; Troyer, D. and Medicetty, S. (2003):

Matrix Cells from Wharton's Jelly Form Neurons and Glia.

Stem Cells, 21:51-60.

Miura, M.; Gronthos, S.; Zhao, M.; Lu, B.; Fisher, L.W.; Robey, P.G. and Shi, S. (2003):

SHED: Stem cells from human exfoliated deciduous teeth.

Nat. Acad. Sci. USA, 100: 5807-5812.

Montanaro, F.; Liadaki, K. and Schienda, J. (2003):

Demystifying SP cell purification: viability, yield distinct progenitor populations in skeletal muscle are bone marrow derived and exhibit different cell fates during vascular regeneration.

J. Clin. Invest., 111:71–79.

Moore, K.A.; Sethi, R. and Doanes, A.M. (1997):

Rac1 is required for cell proliferation and G2/M progression.

Biochem. J., 326: 17–20.

Morgan, D.O. (1995):

Principles of CDK regulation.

Nature, 374:131-134.

Morrison, S.J.; Wright, D.E. and Wessman, I.L. (1997):

Cyclophosphamide/granulocyte colony stimulating factor induces haematopoietic stem cells to proliferate prior to mobilization

Proc. Nat. Acad. Sci. USA, 94:1908-1913.

Morshead, C.M.; Reynolds, B.A. and Craig, C.G. (1994):

Neural stem cells in the mammalian adult forebrain: a relatively quiescent subpopulation of subependymal cells.

Neuron, 13: 1071–82.

Moss, F.P. and Lebond, C.P. (1971):

Satellite cells as the source of nuclei in muscles of growing rats.

Anat. Rec., 170:421–435.

Nakamura, K.; Izumiyama-Shimomura, N.; Sawabe, M.; Arai, T.; Aoyagi, Y.; Fujiwara, M.; Tsuchiya, E.; Kobayashi, Y.; Kato, M.; Oshimura, M.; Sasajima, K.; Nakachi, K. and Takubo, K. (2002):

Comparative analysis of telomere lengths and erosion with age in human epidermis and lingual epithelium.

J. Invest. Dermatol., 119:1014– 1019.

National institute of health(2004):

The stem cell.

National institute of health(2005):

resources for stem cell research.

Nilsson, S.; Dooner, M.and Weier, H. (1999):

Cells capable of bone production engraft from whole bone marrow transplants in nonablated mice.

J. Exp. Med., 189:729–34.

Oh, S.H.; Miyazaki, M.; Kouchi, H.; Inoue, Y.; Sakaguchi, M.; Tsuji T.; Shima, N.; Higashio, K. and Namba, M. (2000):

Hepatocyte growth factor induces differentiation of adult rat bone marrow cells into a hepatocyte lineage in vitro.

Biochem. Biophys. Res. Commun., 279: 500–504.

Okamoto, R.; Yajimi, T.; Yamazaki, M.; Kanai, T.; Mukai, M.;***Okamoto, S.; Ikeda, Y.; Hibi, T.; Inazawa, J. and Watanabe, M., (2002):***

Damaged epithelia regenerated by bone marrow-derived cells in the human gastrointestinal tract.

Nat. Med., 8:1011–1017.

Okumura, K.; Nakamura, K.; Hisatomi, Y.; Nagano, K.; Tanaka, Y.;***Terada, K.; Sugiyama, T.; Umeyama, K.; Matsumoto, K.; Yamamoto, T. and Endo, F. (2003):***

Salivary gland progenitor cells induced by duct ligation differentiate into hepatic and pancreatic lineages.

Hepatology, 38: 104-113.

Oliver, J.A.; Maarouf, O.; Cheema, F.H.; Martens, T.P. and Al-Awqati, Q. (2004):

The renal papilla is a niche for adult kidney stem cells.

J. Clin. Invest., 114: 795-804.

Olmsted-Davis, E.A.; Gugala, Z.; Camargo, F.; Gannon, F.H.; Jackson, K.; Kienstra, K.A.; Shine, H.D.; Lindsey, R.W.; Hirschi, K.K.; Goodell, M.A.; Brenner, M.K. and Davis, A.R. (2003):

Primitive adult hematopoietic stem cells can function as osteoblast precursors.

Proc. Natl. Acad. Sci. USA, 100:15877–15882.

Orlic, D.; Kajstura, J.; Chimenti, S.; Limana, F.; Jakoniuk, I.; Quaini, F.; Nadal-Ginard, B.; Bodine, D.M.; Leri, A. and Anversa, P. (2001, a):

Mobilized bone marrow cells repair the infarcted heart, improving function and survival

Nat. Acad. Sci. USA , 98: 10344-10349.

Orlic, D.; Kajstura, J.; Chimenti, S.; Jakoniuk, I.; Anderson, S.M.; Li, B.; Pickel, J.; McKay, R.; Nadal-Ginard, B.; Bodine, D.M.; Leri, A. and Anversa, P. (2001, b):

Bone marrow cells regenerate infarcted myocardium.

Nature, 410 : 701–705.

Osawa, M.; Hanada, K. and Hamada, H. (1996):

Long-term lymphohematopoietic reconstitution by a single CD34-low/negative hematopoietic stem cell.

Science, 273:242–245.

Oshima, H.; Rochat, A.; Kedzia, C.; Kobayashi, K. and Barrandon, Y. (2001):

Morphogenesis and renewal of hair follicles from adult multipotent stem cells.

Cell, 104: 233–245.

Otmar, E.; Pfister, R.; Frédéric, T.; Mouquet, A.; Mohit, H.; Jain, J.; Ross, G.; Summer, K.; Michiel, L.; Helmes, I.; Alan, A.; Fine, F.; Wilson, F.; Colucci, M.; Ronglih, J. and Liao, K. (2005):

CD31⁻ but Not CD31⁺ Cardiac Side Population Cells Exhibit Functional Cardiomyogenic Differentiation .

Whitaker Cardiovascular Institute, Boston University School of Medicine.

Oyamada, Y.; Komatsu, K.; Kimura, H.; Mori, M. and Oyamada, M. (1996):

Differential regulation of gap junction protein (connexin) genes during cardiomyocytic differentiation of mouse embryonic stem cells in vitro. *Exp. Cell Res.*, 229:318–326.

Pardee, A.B. (1989):

G1 events and regulation of cell proliferation. *Science*, 246:603-608.

Perin, E.C.; Dohmann, H.F.; Borojevic, R.; Silva, S.A.; Sousa, A.L.; Mesquita, C.T.; Rossi, M.I.; Carvalho, A.C.; Dutra, H.S.; Dohmann, H.J.; Silva, G.V.; Belem, L.; Vivacqua, R.; Rangel, F.O.; Esporcatte, R.; Geng, Y.J.; Vaughn, W.K.; Assad, J.A.; Mesquita, E.T. and Willerson, J.T. (2003):

Transendocardial, autologous bone marrow cell transplantation for severe, chronic ischemic heart failure. *Circulation*, 107:2294–2302.

Petersen, B.E. (2001):

Hepatic stem cells: coming full circle. *Blood Cells, Mol. Diss.*, 27: 590-598.

Petersen, B.E.; Bowen, W.C. and Patrene, K.D. (1999):

Bone marrow as a potential source of hepatic oval cells. *Science*, 284:1168–1170.

Petersen, B.E.; Goff, J.P.; Greenberger, J.S. and Michalopoulos, G.K. (1998):

Hepatic oval cells express the hematopoietic stem cell marker Thy-1 in the rat. *Hepatology*, 27: 433–455.

Pfeffer, M.A. and Braunwald, E. (1990):

Ventricular remodeling after myocardial infarction. *Circulation*, 81:1161–1172.

Poulsom, R.; Forbes, S.J. and Hodivala-Dilke, K. (2001):

Bone marrow contributes to renal parenchymal turnover and regeneration.

J. Pathol., 195:229–235.

President's Council on Bioethics (2003 a):

Human cloning and human dignity: an ethical inquiry. The President's Council on Bioethics. Washington, DC, USA.

President's Council on Bioethics (2003 b):

Beyond therapy: biotechnology and the pursuit of happiness. A report of the President's Council on Bioethics. 1st edition. P 328. ReganBooks. New York, USA.

President's Council on Bioethics (2004):

Monitoring stem cell research, Washington.

Prost. S.; Bellamy, C.O.; Clarke, A.R.; Wyllie, A.H. and Harrison, D.J. (1998):

p53- independent DNA repair and cell cycle arrest in embryonic stem cells

FEBS. Lett., 425:499–504.

Prusa, A.R.; Marton, E.; Rosner, M.; Bernaschek, G.; and Hengstschlager, M. (2003):

Oct-4-expressing cells in human amniotic fluid: a new source for stem cell research.

Human Reproduction, 18:1489-1493.

Quaini, F.; Urbanek, K.; Beltrami, A.P.; Finato, N.; Beltrami, C.A.; Nadal-Ginard, B.; Kajstura, J.; Leri, A. and Anversa, P. (2002):

Chimerism of the transplanted heart.

N. Engl. J. Med., 346:5–15.

Rainsford, E. and Reen, D. J. (2002):

Interleukin 10, produced in abundance by human newborn T cells, may be the regulator of increased tolerance associated with cord blood stem cells transplantation .

B. J. haemato., 116:702-709.

Ramer, M.S.; Bishop, T.; Dockery, P.; Mobarak, M.S.; O'Leary, D.; Fraher, J.P.; Priestley, J.V. and McMahon, S.B. (2000):

Functional regeneration of sensory axons into the adult spinal cord
Nature, 403:312-316.

Reynolds, B.A. and Weiss, S. (1992):

Generation of neurons and astrocytes from isolated cells of the adult mammalian central nervous system.

Science, 255: 1707–10.

Rietze, R.L.; Valcanis, H.; Brooker, G.F.; Thomas, T.; Voss, A.K. and Bartlett, P.F. (2001):

Purification of a pluripotent neural stem cell from the adult mouse brain.
Nature, 412:736–739.

Robinson, D.; Ash, H.; Yayon, A.; Nevo, Z. and Aviezer, D. (2001):

Characteristics of cartilage biopsies used for autologous chondrocytes transplantation.

Cell Transplant, 10: 203-208.

Ros, J.E.; Roskams, T.A.; Geuken, M.; Havinga, R.; Splinter, P.L.; Petersen, B.E.; LaRusso, N.F.; Vander, D.S.; Kolk, D.M.; Kuipers, F.; Faber, H.J; Muller, M. and Jansen, P.L. (2003):

ATP binding cassette transporter gene expression in rat liver progenitor cells.

Gut., 52: 1060–67.

Rozental, R.; Morales, M.; Mehler, M.F.; Urban, M.; Kremer, M.; Dermietzel, R.; Kessler, J.A. and Spray, D.C. (1998):

Changes in the properties of gap junctions during neuronal differentiation of hippocampal progenitor cells.

J. Neurosci., 18:1753–1762.

Salingcarnboriboon, R.; Yoshitake, H.; Tsuji, K; Obinata, M.; Amagasa, T.; Nifuji, A. and Noda, M. (2003):

Establishment of tendon-derived cell lines exhibiting pluripotent mesenchymal stem cell-like property.

Experimental Cell Research, 287: 289-300.

Sanchez-Ramos, J.; Song, S.; Cardozo-Pelaez, F.; Hazzi, C.; Stedeford, T.; Willing, A.; Freeman, T.B.; Saporta, S.; Jansse, W.; Patel, N.; Cooper, D.R. and Sanberg, P.R. (2000):

Adult bone marrow stromal cells differentiate into neural cells in vitro. Exp. Neurol., 164:247–256.

Sanchez-Ramos, J.; Song, S.; Kamath, S.G.; Zigova, T.; Willing, A.; Cardozo-Pelaez, F.; Stedeford, T.; Chopp, M. and Sanberg, P.R. (2001):

Expression of neural markers in human umbilical cord blood. Experimental Neurology, 171:109-115.

Saporta, S.; Kim, J.J.; Willing, A.E.; Fu, E.S.; Davis, C.D, and Sanberg, P.R. (2003):

Human umbilical cord blood stem cells infusion in spinal cord injury: engraftment and beneficial influence on behavior.

Journal of Hematotherapy & Stem Cell Research, 12:271-278.

Schuelke, M.; Wagner, K.R.; Stolz, L.E.; Hubner, C.; Riebel, T.; Komen, W.; Braun, T.; Tobin, J.F. and Lee, S.J. (2004):

Myostatin mutation associated with gross muscle hypertrophy in a child. N. Engl. J. Med., 350: 2682–88.

Schwartz, R.E.; Reyes, M.; Koodie, L.; Jiang, Y.; Blackstad, M.; Lund, T.; Lenvik, T.; Johnson, S.; Hu, W.S. and Verfaillie, C.M. (2002):

Multipotent adult progenitor cells from bone marrow differentiate into functional hepatocyte-like cells.

J. Clin. Invest., 109:1291–1302.

Seaberg, R.M.; Smukler, S.R.; Kieffer, T.J.; Enikolopov, G.; Asghar, Z, Wheeler, M.B.; Korbitt, G. and Kooy, D. (2004):

Clonal identification of multipotent precursors from adult mouse pancreas that generate neural and pancreatic lineages.

Nat. Biotechnol., 22:1115-1124.

Seale, P.; Sabourin, L.; Girgis-Gabardo, A.; Mansouri, A.; Gruss, P. and Rudnicki, M.A. (2006):

Pax7 is required for the specification of myogenic satellite cells.
Cell, (102): 777-86.

Seigel, G.M.; Sun, W.; Salvi, R.; Campbell, L.M.; Sullivan, S. and Reidy, J.J. (2003):

Human corneal stem cells display functional neuronal properties.
Molecular Vision, 9:159-163.

Seri, B.; Garcia-Verdugo, J.M.; McEwen, B.S. and Alvarez-Buylla, A. (2001):

Astrocytes give rise to new neurons in the adult mammalian hippocampus.

J. Neurosci., 21: 7153–60.

Sharma, A.D.; Cantz, T.; Richter, S.; Eckert, K.; Henschler, R.; Wilkens, L.; Jochheim-Richter, A.; Arseniev, L. and Ott, M. (2005):

Human cord blood stem cells generate human cytokeratin 18-negative hepatocyte-like cells in injured mouse liver.

Am. J. Pathol., 167: 555–64.

Shen, C.N.; Slack, J.M. and Tosh, D. (2000):

Molecular basis of transdifferentiation of pancreas to liver.

Nat. Cell Biol., 2:879–887.

Sherr, C.J., (1993):

Mammalian G1 cyclins.

Cell, 73:1059-1065.

Sherr, C.J. (1996):

Cancer cell cycles.
Science, 274:1672-1677.

Shi, S.; Gronthos, S.; Chen, S.; Reddi, A.; Counter, C.M.; Robe, P.G. and Wang, C.Y. (2002):

Bone formation by human postnatal bone marrow stromal cells is enhanced by telomerase expression.
Nature Biotechnology, 20: 587-591.

Shih, C.C.; Weng, Y.; Mamelak, A.; LeBon, T.; Hu, M.C. and Forman, S.J. (2001):

Identification of a candidate human neurohematopoietic stem-cell population.
Blood, 98: 2412–22.

Smith, L.L.; Coller, H.A. and Roberts, J.M. (2003):

Telomerase modulates expression of growth-controlling genes and enhances cell proliferation.
Nat. Cell Bio., 15:474–479.

Song, X. and Xie, T. (2003):

Wingless signaling regulates the maintenance of ovarian somatic stem cells in *Drosophila*.
Development, 130: 3259–3268.

Sordling, A.; Drummond-Barbosa, D.; and Toskiokai, E. (2001):

Adult stem cells and their ability to differentiate.
Med. Sci. Monit., 12 (8):154-163.

Steidl, U.; Kronenwett, R.; Martin, S. and Haas, R. (2003):

Molecular biology of hematopoietic stem cells.
Vitam. Horm., 66:1–28. 1132.

Steptoe, R.J.; Ritchie, J.M. and Harrison, L.C. (2003):

Transfer of hematopoietic stem cells encoding autoantigen prevents autoimmune diabetes.

J. Clin. Invest, 111:1357-1363.

Stillman, B. (1996):

Cell cycle control of DNA replication.
Science, 274:1659-1664.

Suzuki, A.; Nakauchi, H. and Taniguchi, H. (2004):

Prospective isolation of multipotent pancreatic progenitors using flow-cytometric cell sorting.
Diabetes, 53::2143-2152.

Suzuki, A.; Zheng, Y.Y.; Kaneko, S.; Onodera, M.; Fukao, K.; Nakauchi, H. and Taniguchi, H. (2002):

Clonal identification and characterization of self-renewing pluripotent stem cells in the developing liver.
J. Cell Biol., 156:173–184.

Tang, D.G.; Tokumoto, Y.M.; Apperly, J.A.; Lloyd, A.C. and Raff, M.C. (2001):

Lack of replicative senescence in cultured rat oligodendrocyte precursor cells.
Science., 291:868–871.

Taylor, G.; Lehrer, M. and Sensen, P.J. (2000):

Involvement of follicular stem cells in forming not only the follicle but also the epidermis.
Cell, 102:451–461.

Terada, N.; Hamazaki, T.; Oka, M.; Hoki, M.; Mastalerz, D.M.; Nakano, Y.; Meyer, E.M.; Morel, L.; Petersen, B.E. and Scott, E.W. (2002):

Bone marrow cells adopt the phenotype of other cells by spontaneous cell fusion.
Nature, 416:542–545.

Terasima, L. and Yasukawa, M . (1966):

Synthesis of G1 protein preceding DNA synthesis in cultured mammalian cells.

Exp. Cell Res., 44:669-672.

Quoted by stem cell biology and gene therapy book, chapter (3). Edited by Peter J. Quesenberry, Gary S. Stein, Bernard G. Forget, Sherman M. Weissman

Copyright □ 1998 Wiley-Liss, Inc.

ISBNs: 0-471-14656-0 (Paper); 0-471-22395-6 (Electronic)

Theise, N.D.; Henegariu, O.; Grove, J .; Jagirdar, J.; Kao, P.N.; Crawford, J.M.; Badve, S.; Saxena, R. and Krause, D.S . (2002):

Radiation pneumonitis in mice: a severe injury model for pneumocyte engraftment from bone marrow.

Exp. Hematol., 30:1333–1338.

Theise, N.D.; Nimmakayalu, M. and Gardner, R. (1999):

Liver from bone marrow in humans.

Hepatology, 12: 34-39.

Thomas, E.D. (1999):

Bone marrow transplantation .

Semin. Hematol., 36:95-103.

Thorgeirsson, S.S. (1996):

Hepatic stem cells in liver regeneration.

FASEB. J., 10: 1249–56.

Thorgeirsson, S.S. and Grisham, J.W. (2006):

Hematopoietic cells as hepatocyte stem cells: a critical review of the evidence.

Hepatology, 43: 2–8.

Tisdale ,J.F. and Dunbar, C.E. (2002):

Plasticity and hematopoiesis.

Curr. Opin. Hematol., 9:268–273.

Tomita, S.; Li, R.K.I.; Weisel, R.D.; Mickle, D.A.; Kim, E.J.; Sakai, T. and Jia, Z.Q. (1999):

Autologous transplantation of bone marrow cells improves damaged heart function.

Circulation, 100: 247–256.

Tsai, R.V.L.; Kittappa, R.; and McKay, R.D., (2002):

Plasticity, niches, and the use of stem cells.

Develop. Cell., 2:707–712 .

Tucker, R.P. (2004):

Neural crest cells: a model for invasive behavior.

Int. J. Biochem .Cell. Bio., 136:173–177.

Tzukerman, M.; Rosenberg, T.; Ravel, Y.; Reiter, I.; Coleman, R. and Skorecki, K. (2003):

An experimental platform for studying growth and invasiveness of tumor cells within teratomas derived from human embryonic stem cells.

Proc. Natl. Acad. Sci. USA., 100 (23):13507-12.

Wagner, K.R. (2005):

Muscle regeneration through myostatin inhibition.

Curr. Opin. Rheumatol., 17: 720–24.

Weimann, J.M.; Johansson, C.B.; Trejo, A. and Blau, H.M. (2003):

Stable reprogrammed heterokaryons form spontaneously in Purkinje neurons after bone marrow transplantation.

Nat. Cell Biol., 5:959–966.

Weir, G.C. and Bonner-Weir, S. (2004):

Beta-cell precursors : a work in progress.

Nat. Biotechnol., 22:1-2.

Weiss, M.L.; Mitchell, K.E.; Hix, J.E.; Medicetty, S.; El-Zarkouny, S.Z.; Grieger, D. and Troyer, D.L. (2003):

Transplantation of porcine umbilical cord matrix cells into the rat brain.

Experimental Neurology, 54:87-88.

Wencker, D.; Chandra, M.; Nguyen, K.; Miao, W.; Garantziotis, S.; Factor, S.M.; Shirani, J.; Armstrong, R.C. and Kitsis, R.N. (2003):

A mechanistic role for cardiac myocyte apoptosis in heart failure.

J. Clin. Invest., 111: 1497–1504.

Whelock, M.J.; and Johnson, K.R. (2003):

Cadherin-mediated cellular signaling.

Curr. Opin. Cell Biol., 15:509–514.

White, E. (1994):

p53, guardian of Rb.

Nature , 371:21-22.

Wilmut, I.; Schnieke, A.E.; McWhir, J.; Kind, A.J. and Campbell, K.H. (1997):

Viable offspring derived from fetal and adult mammalian cells.

Nature, 385:810–813.

Wolf, N.S.; Kone, A.; Priestley, G.V. and Bartelmez, S.H. (1993):

In vivo and in vitro characterization of long-term repopulating primitive hematopoietic cells isolated by sequential hoechst-33342-rhodamine-123 FACS selection.

Exp. Hematol., 21:614–622.

Wright, D.E.; Wagers, A. J.; Gulati, A.P.; Johnson, F.L. and Weissman, I.L. (2001) :

Physiological migration of haematopoietic stem cells and progenitor cells.

Science, 294:1933-1936.

Wright, N. A. (2000):

Epithelial stem cell repertoire in the gut: clues to the origin of cell lineages, proliferative units and cancer.

Int. J. Exp. Pathol., 117:143-147 .

Wulf, G.G.; Jackson, K.A. and Goodell, M.A. (2001):

Somatic stem cell plasticity: current evidence and emerging concepts.

Exp. Hematol., 29:1361–1370.

Wulf, G.G.; Luo ,K.L.; Jackson, K.A .; Brenner, M.K. and Goodell, M.A. (2003):

Cells of the hepatic side population contribute to liver regeneration and can be replenished by bone marrow stem cells.

Haematologica, 88:368–378.

Ying, Q.Y.; Nichols, J.; Evans, E.P. and Smith, A.G. (2002):

Changing potency by spontaneous fusion.

Nature., 416:545–548.

Zhang, J.; Niu, C.; Ye, L.; Huang, H.; He, X.; Tong, W.G.; Ross, J.; Haug, J.; Johnson, T.; Feng, J.Q.; Harris, S.; Wiedemann, L.M.; Mishina, Y. and Li, L. (2002):

Identification of the haematopoietic stem cell niche and control of the niche size.

Nature, 425:836–841.

Zhao, Y.; Glesne, D. and Huberman, E. (2003):

A human peripheral blood monocyte-derived subset acts as pluripotent stem cells.

Nat. Acad. Sci. USA, 100: 2426-2431.

Zuk, P.A.; Zhu, M.; Ashjian, P.; De Ugarte, D.A.; Huang, J.I.; Mizuno, H.; Alfonso, Z.C.; Fraser, J.K.; Benhaim, P. and Hedrick, M.H. (2002):

Human adipose tissue is a source of multipotent stem cells.

Mol. Biol.of the Cell, 13: 4279-4295.