
References

Abernethy VE & Lieberthal W.(2002): acute renal failure in critically ill patient. *Critical Care Clin*;18:203-222.

Abramson S & Niles JL.(1999): Anticoagulation in continuous renal replacement therapy. *Curr Opin Nephrol Hypertens* ; 8:701-706.

Alex S, Evers & Mervyn Maze.(2004): Anesthetic pharmacology : Physiologic principles and clinical practice; Ch 21.

Allgren RL, Marbury TC & Rahman SN.(1997): Anaritide in acute tubular necrosis. Auriculin Anaritide Acute Renal Failure Study Group. *The New England Journal of Medicine* 20; 336(12): 828–834.

Aronson S & Blumenthal R.(2008): Perioperative renal dysfunction and cardiovascular anesthesia concerns and controversies. *J Cardiothoracic vasc Anesth* 12:567-586.

Arroyo V, Guevara M & Ginès P. (2002): Hepatorenal syndrome in cirrhosis: Pathophysiology and treatment. *Gastroenterology* 122 (6): 1658–76 .

Balakrishnan PV & Dubose TD.(2001): Osmotic nephropathy in Militoris BA, Finn WV(edt). *Acute renal failure* 1st ed.391-396.

Ballerman BJ, Zeidel ML & Gunning ME. (1991): Vasoactive peptides and the kidney. In Brenner BM, Rector FCJ(eds). *The kidney* 4th ed .510-583.

Balogh Z, McKinley BA & Cocanour CS. (2003): Supranormal trauma resuscitation causes more cases of abdominal compartment syndrome. *Archives of Surgery*; 138(6): 637–642.

Bellomo R, Kellum JA & Ronco C.(2007): Defining and classifying acute renal failure: from advocacy to consensus and validation of the RIFLE criteria. *Intensive Care Medicine*; 33(3): 409–413.

Better OS, Rubinstein I, Winaver JM & Knochel JP.(1997): Mannitol therapy revisited (1940–1997). *Kidney International*; 52: 886–894.

Birck R, Krzossok S & Markowetz F. (2003): Acetylcysteine for prevention of contrast nephropathy: meta-analysis. *Lancet* ;362:598–603.

Blantz RC.(1998): Pathophysiology of pre-renal azotemia. *Kidney International*; 53: 512–523.

Bonventre JV.(2002): Kidney ischemic preconditioning. *Current Opinion in Nephrology and Hypertension*; 11:43-48.

Bonventre JV.(2003): Dedifferentiation and proliferation of surviving epithelial cells in acute renal failure. *Journal of the American Society of Nephrology*; 14(supplement 1): S55–S61.

Bourgoin A, Leone M & Delmas A.(2005): Increasing mean arterial pressure in patients with septic shock: effects on oxygen variables and renal function. *Critical Care Medicine*; 33(4): 780–786.

Bove T, Calabro MG & Landoni G. (2004): The incidence and risk of acute renal failure after cardiac surgery. *Journal of Cardiothoracic and Vascular Anesthesia*; 18(4): 442–445.

Brezis M & Epstein FH. (1993): Pathophysiology of acute renal failure. In: *Toxicology of the kidney*. 2nd ed. Hook JB, Goldstein RS (eds). New York, Raven,; 129-152.

Caimmi PP, Pagani L & Micalizzi E. (2003): Fenoldopam for renal protection in patients undergoing cardiopulmonary bypass. *Journal of Cardiothoracic and Vascular Anesthesia* ; 17(4): 491–494.

Carmichael P & Carmichael AR.(2003): Acute renal failure in the surgical setting. *ANZ Journal of Surgery* ; 73(3): 144–153.

Clark LC, Farghaly H, Saba SR & Vesely DL.(2000): Amelioration with vessel dilator of acute tubular necrosis and renal failure established for 2 days. *American Journal of Physiology. Heart and Circulation Physiology*; 278: H1555–H1564.

Cohen MV, Baines CP & Downey JM.(2000): Ischemic preconditioning: from adenosine receptor of K ATP channel. *Annual Review of Physiology*; 62: 79–109.

Colson p, Rychwaert F & Coriat P. (1999): Renin angiotensin system and anesthesia. *Anesth analg*; 89 : 1145.

Corwin HL, Gettinger A & Rodriguez RM.(1999): Efficacy of recombinant human erythropoietin in the critically ill patient: a randomized, double-blind, placebo-controlled trial. *Crit Care Med*, 27:2346-2350.

Cuhaci B, Lee J & Ahmed Z.(2000): Sodium bicarbonate controversy in lactic acidosis. *Chest* ; 118: 882-883.

David C. Mendelssohn, Salim K. Mujais & Steven D. Soroka3. (2009): A prospective evaluation of renal replacement therapy modality eligibility. *Nephrology Dialysis Transplant* 24(2):555-561.

David Humes.(2007): Principles of tissue engineering .3th ed. Elsevier. PP.801-810.

Davison AM, Cumming AD, Swainson CP & Turner N. (1999): Diseases of the kidney and urinary system. *Investigations of renal disease*; 18: 418-470.

Desjars Ph, Stafford M & Smith M.(1999): Acute renal failure following cardiac surgery .Nephrol Dial Transplant.14: 1158-1162.

Dharnidharka VR, kwon C & Stevens G. (2002): Serum cystatin c is superior to serum creatinine as a marker of kidney function: a meta-analysis. Am J Kidney Dis;40:221-226.

Di Giantomasso D, May CN & Bellomo R.(2002): Norepinephrine and vital organ blood flow. Intensive Care Medicine ; 28: 1804–1809.

Dishart MK & Kellum JA.(2000): An evaluation of pharmacological strategies for the prevention and treatment of acute renal failure. Drugs; 59: 79–91.

Druml W, Schwarzenyhofer M & Apsner R.(1998): Fat soluble vitamins in patients with acute renal failure. Miner electrolyte Metab;22-24

E.F. van Bommel. (2003): Renal replacement therapy for acute renal failure on the intensive care unit: coming of age?, *Neth J Med* **61** ,pp. 239–248.

Eliot Sykes & Joseph F Cosgrove.(2007): Acute renal failure and critically ill surgical patient.Ann R Coll Surg Engl;89:22-29.

Frankenfield DC & Reynolds HN.(1995): Nutritional effect of continuous haemofiltration.Nutrition;11:388-393.

Gabutti L, Marone C & Colucci G. (2002): Citrate anticoagulation in continuous venovenous hemodiafiltration: a metabolic challenge. Intensive Care Med ; 28:1419-1420.

Ganong WF.(2001): Regulation of extracellular fluid composition and volume. Review of medical physiology 20th ed .Boston.633:670

Garwood S.(2000): New pharmacologic options for renal preservation. *Anesthesiology Clinic of North America*; 18: 753–771.

Gill N, Nally Jr. JV & Fatica RA.(2005): Renal failure secondary to acute tubular necrosis: epidemiology, diagnosis and management. *Chest*; 128(4): 2847–2863.

Goligorsky MS & Noiri E.(1999): Duality of nitric oxide in acute renal injury. *Semin Nephrol* ;19:263-271.

Gray H. (2000): *Anatomy of the human body. The urinary organs* by Henry Gay. 20th ed. Thoroughly rev and re-edited by Warren H. Lewis. Philadelphia: Lea and Febger, 1981, Bartleby, (2000).

Greenberg A.(1998): Hyperkalemia: treatment options. *Semin Nephrol* ; 18:46-57.

Gülberg V, Bilzer M & Gerbes AL (1999): "Long-term therapy and retreatment of hepatorenal syndrome type 1 with ornipressin and dopamine". *Hepatology* 30 (4): 870–881.

Gunning ME. (1991): Vasoactive peptides and the kidney. In Brenner BM, Rector FCJ (eds) *the kidney* 4th ed; 510:583.

Guyton AC & Hall JE. (2000): *Regulation of acid-base balance*, Textbook of medical physiology, Elsevier Saunders, 10th ed ;26:279 –294.

Hawaleshka A & Jacobsohn E. (1998): Ischaemic preconditioning: mechanisms and potential clinical applications. *Canadian Journal of Anaesthesiology*; 45: 670–682.

Hediger MA, Boron WF & Fong P. (1997): The electrogenic Na/Hco₃ cotransporter. *J Am Soc Nephrol*; *Wien Klin Wochenschr*; 109:445.

Hoek FJ, kemperman FA & Krediet RT.(2003): A comparison between cystatin c, plasma creatinine and Cockcroft and Gault formula for the estimation of GFR. *Nephrol Dial transplant*;18:2024-2031.

Hoffmann U, Fischeder M, Kruger B, Drobnik W & Kramer BK. (2004): The value of N-acetylcysteine in the prevention of radiocontrast agent-induced nephropathy seems questionable. *J Am Soc Nephrol*; 15: 407–10.

Hoomberg K, Smir AJ & Girbes AR.(1998): Effects of low dose dopamine on renal and systemic hemodynamics during incremental norepinephrine infusion in healthy volunteers. *Critical Care Medicine*; 26:260-265.

Hotchkiss RS & Karl IE. (2003): The pathophysiology and treatment of sepsis. *N Engl J Med*; 348: 138–50. .

Jacob R.(2005): Acute renal failure. *Indian J Anesth.* :74(5)p367-372.

Jaffe R, Ariel I & Beerl R.(1997): Frequent apoptosis in human kidneys after acute renal hypoperfusion. *Experimental Nephrology*; 5: 399–403.

James M.(2006): Volume expanders: crystalloid vs. plasma colloids vs. synthetic colloids. *ISBT Science Series*; 1(1): 52–58.

Janberg Po.(2004): Renal protection strategies in the perioperative period. *Best practice&Research clinical Anaesthesiology*;18:p 645-660.

Jones D & Bellomo R.(2005): Renal-dose dopamine: from hypothesis to paradigm to dogma to myth and, finally, superstition?
Journal of Intensive Care Medicine ,20(4): 199–211.

Jones DR & Thomas Lee HT.(2008): Perioperative renal protection. *Best Practice & Research Clinical Anaesthesiology*. Vol. 22, No. 1, pp. 193–208.

Jones DR & Lee HT.(2007): Protecting the kidney during critical illness. *Current Spin anesthiol*: 20(2): pp106-112.

Julier K, Da Silva R & Garcia C.(2003): Preconditioning by sevoflurane decreases biochemical markers for myocardial and renal dysfunction in coronary artery bypass graft surgery: a double-blinded, placebocontrolled,multicenter study. *Anesthesiology*; 98: 1315–1327.

Karlesson S, Varpula M, Ruokonen E, Pettila" V, Parviainen I & Ala-Kokko TI .(2007): Incidence, treatment, and outcome of severe sepsis in ICU-Intensive Care Med;33:435—43.

Kashyap VS, Cambria RP, Davison JK & L'Italien GJ.(1997): Renal failure after thoracoabdominal aortic surgery.*Journal of Vascular Surgery*; 26(6): 949–955.

Klaver MJ, Manning L, Palmer LA & Rich GF. (2002): Isoflurane pretreatment inhibits cytokine-induced cell death in cultured rat smooth muscle cells and human endothelial cells. *Anesthesiology*; 97: 24–32.

Knoe BC. (1996): Renal H, K-AT pase-structure, function and regulation. *Miner Electrolyte Metab*; 22:349.

Kribben A, Edelstein CL & Schrier RW.(1999): Pathophysiology of acute renal failure. *Journal of Nephrology*;12(supplement 2): S142–S151.

Lake EW & Humes HD.(1994): Acute renal failure: directed therapy to enhance renal tubular regeneration. *Seminars in Nephrology*; 14: 83–97.

Lameire N, Van BW & Vanholder R.(2005): Acute renal failure. *Lancet Jan 29*; 365(9457): 417–430.

Lameire NH, De Vriese AS & Vanholder R.(2003): Prevention and nondialytic treatment of acute renal failure. *Curr Opin Crit Care*; 9: 481–90.

Landoni G, Biondi-Zoccai GG & Tumlin JA. (2007): Beneficial impact of fenoldopam in critically ill patients with or at risk for acute renal failure: a meta-analysis of randomized clinical trials.

American Journal of Kidney Diseases; 49(1): 56–63.

Landry DW, Levin HR & Gallant EM. (1997): Vasopressin pressor hypersensitivity in vasodilatory septic shock. Critical Care Medicine; 25: 1279–1282.

Lane N & Alien K.(1999): hyponatremia after orthopedic surgery. BM;318: 1363-1364.

Lee HT & Sladen RN .(2004): Perioperative renal protection. In Murray MJ, Coursin DB,et al(eds). Critical care medicine. perioperative management,2"ed.;42:503-520.

Leibowitz AB. (2003): Approach to renal failure. In Apostolakis MJ and Papadakis pj(eds). The intensive care manual; Tata McGraw Hill edition .p103-118.

Levens NR, peach MJ & Carey RM. (1981):Role of the intrarenal rennin-angiotensin system in the control of renal function. Circ Res; 48:157 – 167.

Leverve X & Barnoud D.(1998): Stress metabolism and nutritional support in acute renal failure. Kidney int;53:562-566.

Maqsood Elahi, Sanjay Asopa & Axel Pflueger Nady.(2008): Acute kidney injury following cardiac surgery: impact of early versus late haemofiltration on morbidity and mortality. European Journal of cardiothoracic surgery.P1-36.

McCullough PA, Adam A & Becker CR.(2006): Risk prediction of contrast-induced nephropathy. Am J Cardiol;98:27K–36K.

McCullough PA & Soman SS.(2005): Contrast-induced nephropathy. Crit Care Clin;21:261– 80.

Megyesi J, Andrade L & Vieira Jr JM.(2001): Positive effect of the induction of p21WAF1/CIP1 on the course of ischemic acute renal failure. *Kidney International*; 60: 2164–2172.

Mehta R, McDonald B & Gabbai F.(2001): A randomized, clinical trial of continuous versus intermittent dialysis for acute renal failure. *Kidney International* ; 60:1154-1163.

Mehta RL & Chertow GM.(2000): Selection of dialysis modality In Owen WF, Pereira BJ, Sayegh MH, editors. *Dialysis and transplantation; a companion to Brenner & Rectors' the kidney*. Philadelphia: WB Sanders; p. 403-417.

Mehta RL, Pascual MT, Soroko S & Chertow GM.(2002):Diuretics, mortality and non recovery of renal function in ARF.*JAMA* ;288: 2547-2553.

Miller D.(2000): Renal function monitoring, anesthetic management, In Roland D Miller, *anesthesia text book*,5th ed ; Churchill livingstone,18:1123- 1175.

Miller D.(2005): Anesthesia and the renal and genitor-urinary system, In Ronald D Miller, *Anesthesia text book*, 6th ed, Churchill Livingstone; 54

Molitoris BA.(2003): Transitioning to therapy in ischemic acute renal failure. *Journal of American society of nephrology* ;14:265-267.

Moran SM & Meyers BD.(1985): Course of acute renal failure studied by a model of creatinine kinetics. *Kidney International*; 27: 928–937.

Morelli A, Ricci Z & Bellomo R. (2005): Prophylactic fenoldopam for renal protection in sepsis: a randomized, double-blind, placebo-controlled pilot trial. *Critical Care Medicine*; 33(11): 2451–2456.

Murphy MB, Murray C & Shorten GD.(2001): Fenoldopam: a selective peripheral dopamine-receptor agonist for the treatment of severe hypertension. *New England Journal of Medicine*; 345: 1548–1557.

Murray MJ, Coursin DB & pearl RG.(2004): critical care medicine. Perioperative management. 2nd ed.p 521-532.

Nemoto T, Yokota N & Keane WF. (2001): Recombinant erythropoietin rapidly treats anemia in ischemic acute renal failure. *Kidney Int*; 59:246-251.

Nobert lameire & Win van Biesen.(2005): Acute renal failure *Lancet*; 30 p365-417.

O'Malley CM, Frumento RJ & Hardy MA. (2005): A randomized, double-blind comparison of lactated Ringer's solution and 0.9% NaCl during renal transplantation. *Anesthesia & Analgesia*; 100(5):1518-1524.

Obialo CI, Crowell AK & Okonofua EC.(2002): Acute renal failure mortality in hospitalized African Americans: age and gender considerations. *Journal of the National Medical Association*; 94: 127–134.

O'Reilly P & Tolwani A.(2005): Renal replacement therapy III:IHD,CRRT,SLED. *Critical Care Clin*; 21:367-378.

P. Massoudy, S. Wagner & M. Thielmann.(2008): Acute renal failure in patients with coronary heart disease: coronary artery bypass surgery and acute kidney injury—impact of the off-pump technique, *Nephrol Dial Transplant* **23** (9), pp. 2853–2860.

Palevsky PM.(2005): Renal replacement therapy indications and timing. *Crit Care Clin* ; 21: 347-356.

Pannu N, Klarenbach S & Wiebe N. (2008): Renal replacement therapy in patients with acute renal failure: a systematic review. *Alberta Kidney Disease Network*. (7):793-805.

Patel BM, Chittock DR & Russell JA.(2002): Beneficial effects of short-term vasopressin infusion during severe septic shock. *Anesthesiology*; 96: 576–582.

Paul Hayden.(2007): Intra abdominal hypertension and the abdominal compartment syndrome. *Current Anaesthesia & Critical Care*;18, 311–316.

Per-Olof Janberg.(2007):critical illness and intensive care. *surg (oxford)*;25(9):p391-398.

Peter A & McCullough MD.(2008): Contrast-Induced Acute Kidney Injury. *Journal of the American College of Cardiology*;51:1420-1429.

Pierucci A, Pistolesi V & Morabito S. (2009): Continous renal replacement therapies will remain the most widely adopted dialysis modality in the critically ill, *G Ital Nephrol*;26(1)13-21.

Pieter Evenepoel.(2004): Acute toxic renal failure. *Best Practice & Research Clinical Anaesthesiology* Vol. 18, No. 1, pp. 37–52.

Poullis M.(1999):Mannitol and cardiac surgery. *Thoracic and Cardiovascular Surgeon*;47:58-62.

purday J.(2001): ARF in Allman KG and Wilson IH (eds).*Oxford hand book of anesth.* Oxford university press. p 116-118.

Rahman TM & Treacher DF.(2002): Management of acute renal failure on the intensive care unit. *Clin Med JRCPL*; 2: 108–113.

Reddy VG .(2002): prevention of postoperative ARF. *A review Journal of postgraduate medicine*.48:p 64-70.

Renton MC & Snowden CP.(2005): Dopexamine and its rule in the protection of hepatosplanchnic and renal perfusion in high risk surgical and critically ill patients. *Br J Anesth* ;94:4:459-467.

Roberts I, Alderson P & Bunn F.(2004): Colloids versus crystalloids for fluid resuscitation in critically ill patients. Cochrane Database of Systematic Reviews 2004; 4: CD000567.

Ronco C, Tetta C & Mariano F.(2003): Interpreting the mechanisms of continuous renal replacement therapy in sepsis: the peak concentration hypothesis. Artif Organs; 27:792-801.

Rose BD & Post TW (2001): Clinical physiology of acid-base and electrolyte disorders, 5th ed . New York; McGraw-Hill ; 187-190.

Rosen S & Heyman SN.(2001): Difficulties in understanding human acute tubular necrosis: limited data and flawed animal models. Kidney International; 60: p1220–1224.

Rossert J.(2001): Drug-induced acute interstitial nephritis. Kidney International; 60(2): 804–817.

Sakr Y, Payen D & Reinhart K. (2007): Effects of hydroxyethyl starch administration on renal function in critically ill patients. British Journal of Anaesthesia; 98(2): 216–224.

Schrier RW & Wang W.(2004): Acute renal failure and sepsis. N Engl J Med; 351: 159–69.

Schuller D, Lynch JP & Fine D.(1997): Protocol guided diuretic management: comparison of furosemide by continuous infusion and intermittent bolus. Crit care Med ;25:1669-1675.

Sean M Bagshaw & Rinaldo Bellomo.(2007): Acute renal failure. Surgery(oxford) ; 25:391-398.

Sear JW. (2005): Kidney dysfunction in the post operative period. British Journal of Anesthesia ;95 (1) : 20-32.

- Sear JW.** (2001): Pharmacology of the critically ill. Fundamental of anesthesia and acute medicine series. London:BMJ ;4:50-88.
- Seri I ,Kone BC & Gullans SR.**(1988): Locally formed dopamine inhibits Na-K-ATPase activity in rat renal cortical tubule cells. American Journal of physiology;255:F666-F673.
- Shilliday IR, Quinn KJ & Allison ME.**(1997): Loop diuretics in the management of acute renal failure: a prospective, double-blind, placebo-controlled, randomized study. Nephrology Dialysis Transplantation; 12: 2592–2596.
- Shoemaker.**(2000): Adult acute and chronic renal failure.Todd W and Anton C (eds).In Shoemaker ICU textbook.154:1648-1658.
- Sladen RN.**(2000): Oliguria in the ICU. Systematic approach to diagnosis and treatment. Anesth Clinc North Am;18:p 739-752.
- Stanton BA & Koeppen BM.**(1993): Elements of renal function. In Berne RM, Levy MN(eds):physiology,3rd ed.louis st,Mosby-year book;719-753.
- Star RA.**(1998): Treatment of acute renal failure. Kidney Int;54:1817-1831.
- Sterling.** (1989): The kidney In textbook of physiology patton HD, Fuchs AF, Hille B (eds) 21st ed;524:560.
- Sugrue M, Jones F, Deane SA, Bishop G, Bauman A & Hillman K.**(1999): Intra-abdominal hypertension is an independent cause of postoperative renal impairment. Arch Surg ;134(10): 1082–1085.
- Suwa T, Hogg JC & Klut ME.** (2001): Interleukin-6 changes deformability of neutrophils and induces their sequestration in the lung. American Journal of Respiratory and Critical Care Medicine; 163: 970–976.

Sward K, Valsson F & Odencrants P.(2004): Recombinant human atrial natriuretic peptide in ischemic acute renal failure: a randomized placebo-controlled trial. *Critical Care Medicine*; 32(6): 1310–1315.

Sykes E & Cosgrove JF.(2007): Acute renal failure and the critically ill surgical patient. *Ann R Coll Surg Engl*; 89(1):PP 22-29.

Tang I Y & Murray P T.(2004): Prevention of perioperative acute renal failure: what works? *Best Practice & Research Clinical Anaesthesiology* Vol. 18, No. 1, pp. 91–111

Thadhani R, Pascual M & Bonventre JV.(1996): Acute renal failure. *New England Journal of Medicine*; 334: 1448–1460.

Tisher CC & Madsen KM. (1991): Anatomy of the kidney. In Brenner BM, Rector FCJ (eds). *The Kidney* 4th ed; 3-75.

Tumlin J, Stacul F & Adam A.(2006): Pathophysiology of contrast induced nephropathy. *Am J Cardiol* ;98:14K–20K.

Tumlin JA, Wang A, Murray PT & Mathur VS. (2002): Fenoldopam mesylate blocks reductions in renal plasma flow after radiocontrast dye infusion: a pilot trial in the prevention of contrast nephropathy. *American Heart Journal*; 143: 894–900.

Uchino S, Kellum JA & Bellomo R.(2005): Acute renal failure in critically ill patients: a multinational, multicenter study. *JAMA*; 294: 813–18.

Vanholder E & Sever N.(2000): Rhabdomyolysis. *Journal of the American Society of Nephrology* ; 11: 1553–1561.

Venkataraman R.(2005): Prevention of ARF. *Crit care clin*;21:281-289.

Vincent JL & Gerlach H.(2004): Fluid resuscitation in severe sepsis and septic shock: an evidence-based review.

Critical Care Medicine 2; 32(11 supplementary): S451–S454.

William H, Fissell & H. David Humes.(2008): Principles of regenerative medicine. Elsevier.PP 1114-1125.

Weldon BC & Monk TG .(2000): The patient at risk for acute renal failure. Recognition ,Prevention , and preoperative optimization. Anesthesiology clinic of North America;18:705-717.

Wong F & Blendis L. (2001): "New challenge of hepatorenal syndrome: prevention and treatment". Hepatology 34 (6): 1242–51.

Yanagwa N & Jo OD. (1997): Intracellular acidification inhibits opossum kidney cell phosphate uptake. Am J Physiol; 272 : R 1904.