

References

1- Ackerman AB, Jacobson M and Vitale P. (1991):

Clues to Diagnosis in Dermatopathology. ASCP Press: Chicago, pp 245–248.

2- Adam I. Rubin, Elbert H. Chen, and Désirée Ratner(2005):

Basal-Cell Carcinoma. N Engl J Med;353:2262-9.

3- Alam M and Goldberg LH(2003):

Oblique advancement flap for defects of the lateral nasal supratip; Arch. Dermatol., 139(8):1039-42

4- Ali F, Brown A and Gottwald L(2005):

Basal cell carcinoma with matrical differentiation in a transplant patient: A case report and review of the literature. J Cutan Pathol;32:445–448

5- Aloï FG, Molinero A and Pippione M(1988):

Basal cell carcinoma with matrical differentiation. Am J Dermatopathol;10:509–513.

6- Ambrojo P, Aguilar A and Simon P(1992):

Basal cell carcinoma with matrical differentiation. Am J Dermatopathol;14:293–297.

7- Armstrong BK and Kricke A.(2001):

The epidemiology of UV induced skin cancer. J Photochem Photobiol B;63:8-18.

8- Ascari Raccagni A and Baldari U(2004):

The retroangular flap used in the surgery of nasal tip defects; Dermatol. Surg., 30(8):1131-7.

References

9- Ashbaugh DR (1999):

Quantitative-Qualitative Friction Ridge Analysis: An Introduction to Basic and Advanced Ridgeology. 1st. CRC :8

10- Bale AE, Gailani MR and Leffell DJ (1994):

Nevoid basal cell carcinoma syndrome. J Invest Dermatol;103:126S–130S.

11- Barr RJ, Headley JL and Jensen JL (1986):

Cutaneous keratocysts of nevoid basal cell carcinoma syndrome.
J Am Acad Dermatol;14:572–576.

12- Bartley GB and Messenger MM (2002):

Outcome of tarsoconjunctival flap dehiscence after eyelid reconstruction. Am J Ophthalmol;134:627-30.

13- Batra RS and Kelley LC(2002):

Predictors of extensive subclinical spread in nonmelanoma skin cancer treated with Mohs micrographic surgery. Arch Dermatol;138:1043-51.

14- Basset-Seguin N, Ibbotson S and Emtestam L(2003):

Photodynamic therapy using Metvix™ is as efficacious as cryotherapy in BCC with better cosmetic results. J Eur Acad Dermatol Venereol 15, 226.
Cited by: Foley P. Clinical efficacy of methylaminolevulinate (metvix) photodynamic therapy. J Dermatolog Treat; 14 (Suppl. 3): 15–22.
2001.

15-Baumann L(2007):

Skin ageing and its treatment. J Pathol. Jan ;211(2):241-51.

References

16- Bauman Leslie ,and EdmundWeisberg(2002):

Cosmetic dermayology:principles and practice (1):3-12

17- Beasley NJ., Gilbert RW., Gullane PJ., brown DH., Irish JC., and Neligan PC (2004) :

Scalp and forehead reconstruction using free revascularized tissue transfer : Arch. Facial plast. Surg., 6 (1) : 16-20 .

18- BeckerH.(1983):

rhomboid to –W flap.Ann. plast. Surg.11:115.

19- Benedetto AV, Benedetto EA and Griffin TD(2002):

Basal cell carcinoma presenting as a large pore. J Am Acad Dermatol;47:727-32

20- Bedllow A.j(1999).

Basal cell carcinoma – an in vivo model of human tumor microcirculation Exp Dermatol . ;8(3):222

21- Berman B, Sullivan T and De Araujo T (2003).

Expression of Fas receptor on basal cell carcinomas after treatment with imiquimod 5% cream or vehicle. Br J Dermatol; 149 (Suppl. 66): 59–61.

22-Bowen AR and LeBoit PE(2005).

Fibroepithelioma of Pinkus is a fenestrated trichoblastoma. Am J Dermatopathol;27:149–154.

23- Boyd AS(2004):

Tumors of the epidermis. In: Barnhill R, Crowson AN (eds). Textbook of Dermatopathology, 2nd edn. McGraw-Hill Co: New York,pp 575–634.

24- Boyd AS and Rapini RP(1994):

Cutaneus collision tumors. Am J Dermatopathol;16:253–257.

References

25- Buckingham ED., Quinn FB., and Calhoun KH (2003):

Optimal design of O- to Z- flaps for closure of facial skin defects;
Arch. Facial Plast. Surg., 5 (1) : 92 – 5 .

26- Burget G.,and Menik F.(1985):

The subunit principle in nasal reconstruction. Plast. reconstr. Surg. 76: 236.

27- Burget,G. and Menick, F.(1985):

Aesthetic restoration of one-half of the upper lip. Plast. Reconstr.surg.76:239,

28- Burns DA, Breathnach SM, Cox N and Griffiths CE(2004):

Rook's Textbook of Dermatology. 7th ed. Wiley-Blackwell.

29- Carlos Luiz,Junqueira and Jose Carneiro (2003):

Basic Histology:Text and Atlas,tenth edition,chap.18:369-381.

30- Carlson BM(1994):

Integumentary,skeletal and muscular system.in.Human Embryology and Developmental Biology.1st:153-181.

31- Carroll RP (1983):

Entropion following the Cutler-Beard procedure.Ophthalmology
;90:1052 5.

32- Carroll RP, Ramsay HM and Fryer AA (2003).

Incidence and prediction of nonmelanoma skin cancer post-renal
transplantation: a prospective study in Queensland, Australia. Am J Kidney
Dis; 41: 676–683.

References

33- Carter DM and Lin AN(1993):

Basal cell carcinoma. In: Fitzpatrick TM, Eisen AZ, Wolff K, et al (eds).
Dermatology in General Medicine, 4th edn. McGraw-Hill: New York , pp 840–
847.

34- Chiller K, Passaro D and McCalmont T (2000):

Efficacy of curettage before excision in clearing surgical margins of
nonmelanoma skin cancer. Arch Dermatol; 136: 1327–1332.

**35- Coleman John J., Edwin G. Wilkins, and Victoria M Vanderkam
(2000):**

Plastic surgery: operations, and outcome, vol. (3), chap. 74 : 1167 – 1192 .

36- Cook JL and Perone JB(2003):

A prospective evaluation of the incidence of complications associated with
Mohs micrographic surgery. Arch Dermatol; 139: 143–152.

37- Cook J and Zitelli JA(1998):

Mohs micrographic surgery: a cost analysis. J Am Acad Dermatol ; 39: 698–
703.

38- Crockett D J(1965):

Lymphatic anatomy and lymphoedema. Br J Plast Surg. Jan;18:12-25.

39- Crowson AN(2006):

Basal cell carcinoma: biology, morphology and clinical implications .
Modern Pathology (19); 127–147.

References

40- Crowson AN, Magro CM and Kadin M (1996):

Differential expression of bcl-2 oncogene in human basal cell carcinoma. Hum Pathol;27:355–359.

41- Crowson AN, Magro CM and Pilavdzic D (1997):

Differential stromal p53 expression in human basal cell carcinoma (Abstract). Lab Invest;76:43A

42- Cutler NI and Beard C(1955):

A method for partial and total upper lid reconstruction. Am J Ophthalmol; 39:1-7.

43- Daniel L. Stulberg, Blain Crandell and Robert S. Fawcett(2004):

Diagnosis and Treatment of Basal Cell and Squamous Cell Carcinomas,Journal of American Family physician,;70(8).

44- DE Faria JL(1985):

Basal cell carcinoma of the skin with areas of squamous cell carcinoma: a basosquamous cell carcinoma. J Clin Pathol;38: 1273–1277.

45- DE Vries E, Louwman M, Bastiaens M, de Gruijl F,and Coebergh JW(2004):

Rapid and continuous increases in incidence rates of basal cell carcinoma in the southeast Netherlands since 1973. J Invest Dermatol;123:634-8.

46- Diepgen TL and Mahler V(2002):

The epidemiology of skin cancer. Br J Dermatol;146: Suppl 61:1-6.

References

47- Disa J, Liew S, and Cordeiro P(2001):

Soft tissue reconstruction of the face using the folded / multiple skin island radial forearm free flap ; Ann. Plast. Surg., 47 (6) : 612-9.

48- Ducic Y and Smith JE(2003):

The cervicodeltopectoral flap for single stage resurfacing of the antero lateral defect of the face and neck;Arch. Plast. Reconstr. Surg.,5(2):197-201.

49- El-sahy Nabil (2000):

Plastic and reconstructive surgery of the nose, part II, chap. 16 : 163-169 .

50- Elston DM, Bergfeld WF and Petroff N(1993) :

Basal cell carcinoma with monster cells. J Cutan Pathol;20:70–73.

51- Epstein E(1985):

Fluorouracil paste treatment of thin basal cell carcinomas. Arch Dermatol; 121: 207–213.

52- Ergun SS(2002):

Reconstruction of labiomenal region with local flaps ; Dermatol. Surg., 28 (9) : 863 – 5.

53- Falco O ,Plew G, Wolf HH and Burgdrof WHC (2001):

Dermatology,second edition,chap.56:1475-1489.

54- Farwell D. Gregory, and Neal D. Futran (2003):

Flap choice in head and neck reconstruction; Facial Plast. Surg. Clin. N. Am., 11 : 107 – 119.

55- Fawcet Don,and Ronald p.jenish(2002):

Bloom and fawcet concise histology,second edition, (11):163-176.

56- Feldman JJ(1990):

Facial burn in: macarthy J.G, Ed plastic surgery,Philadelphia,Saunders,P.2153.

References

57- Fink-Puches R, Soyer HP and Hofer A (1998):

Long-term followup and histological changes of superficial nonmelanoma skin cancers treated with topical delta-aminolevulinic acid photodynamic therapy. Arch Dermatol; 134: 821– 826.

58- Fongo A, Ferraris E and Bocca M(1966):

Skin tension lines and wrinkles. Anatomoclinical observations. Minerva Chir. ;21(13):627-30.

59- Frank H.Netter(2006):

Atlas of Human Anatomy 4th Edition:p23.

60- Frederick Menick (2007):

Defects of nose ,lip and cheek:Rebuilding the composite defect.Plast. Reconstr. Surg. 120: 887.

61- Futran N., and Alsarraf R (2000):

Microvascular free flap reconstruction in the head and neck; JAMA, 284 (14) : 1761 – 3.

62-Gahhos F.N. and CuonoC.B.(1990):

Double z rhombic technique for reconstruction of facial wounds.plast.reconstructr.surg.85-869.

63- Gallagher RP, Hill GB and Bajdik CD (1995):

Sunlight exposure, pigmentary factors, and risk of nonmelanocytic skin cancer. Basal cell carcinoma. Arch Dermatol;131: 157-63.

64- Garcia JA, Cohen PR and Herzberg AJ (1995):

Pleomorphic basal cell carcinoma. J Am Acad Dermatol;32: 740–746.

References

65- Gartner leslie p ., and James L. Hiatt (2000):

Color atlas of histology, third edition, chap.8: 113 – 137 .

66- Geisse J, Caro I and Lindholm J(2004):

Imiquimod 5% cream for the treatment of superficial basal cell carcinoma: results from two phase III, randomized, vehicle-controlled studies. J Am Acad Dermatol; 50: 722–733.

67-Geisse JK, Rich P and Pandya A (2002).

Imiquimod 5% cream for the treatment of superficial basal cell carcinoma: a doubleblind, randomized, vehicle-controlled study. J Am Acad Dermatol; 47: 390–398.

68- Goldberg LH(1996):

Basal cell carcinoma. Lancet;347:663–667.

69- Goldman MP. Shiffman MA, Mirrafati SJ, Lam SM and Cueteaux CG(2007):

Simplified Facial Rejuvenation. 2. Springer Berlin Heidelberg; :47-50.

70- Golcman R, Speranzini MB and Golcman B(1998):

The bilobed island flap in nasal ala reconstruction. Br J Plast Surg;51:493-8.

71- Goldstein AM, Bale SJ and Peck GL (1993):

Sun exposure and basal cell carcinoma in the nevoid basal cell carcinoma syndrome. J Am Acad Dermatol;29: 34–41.

72- Gorlin RJ(1995):

Nevoid basal cell carcinoma syndrome. Dermatol Clin N Am;13:113–125.

References

73- Guo HR, Yu HS, Hu H and Monson RR(2001).

Arsenic in drinking water and skin cancers: cell-type specificity (Taiwan, ROC). *Cancer Causes Control*;12:909-16.

74- Griffiths RW (1999):

Audit of histologically incompletely excised basal cell carcinomas: recommendations for management by re-excision. *Br J Plast Surg*; 52: 24–28.

75- Hairston BR., and Nguyen TH (2003):

Innovations in the island pedicle flap for cutaneous facial reconstruction; *Dermatol. Surg.*, 29 (4) : 3 8-85.

76- Haller JC,Cairnduff F and Slack G (2000).

Routine double treatments of superficial basal cell carcinomas using aminolaevulinic acid-based photodynamic therapy.*Br J Dermatol*;143 :1270 – 1275.

77-Hanke CW and Temofeew RK(1986):

Basal cell carcinoma with eccrine differentiation (eccrine epithelioma). *J Dermatol Surg Oncol*;12:820–824.

78- Hartevelt MM, Bavinck JN, Kootte AM,Vermeer BJ and Vandenbroucke JP(1990):

Incidence of skin cancer after renal transplantation in the Netherlands. *Transplantation*;49:506-9.

79- Hayakawa TJ, Pilavdzic D and Magro CM(1998):

The ultrastructure of aggressive—versus indolent—growth basal cell carcinoma: a comparative study (Abstract). *Lab Invest*;78:50A.

References

80- Hendrix JD and Parlette HL(1996).

Micronodular basal cell carcinoma: a deceptive histologic subtype with frequent clinically undetected tumor extension. Arch Dermatol;132:295–298.

81- Hewes EH, Sullivan JH and Beard C(1976):

Lower eyelid reconstruction by tarsal transposition. Am J Ophthalmol; 81:512-4

82- Hoban PR, Ramachandran S and Strange RC(2002):

Environment, phenotype and genetics: risk factors associated with BCC of the skin. Expert Rev Anticancer Ther 2002; 2: 570–579.

83- Howell JB and Mehregan AH(1970):

Pursuit of the pits in nevoid basal cell carcinoma syndrome. Arch Dermatol ;102:586–597.

84- Hsuan J and Selva D(2004).

Early division of a modified Cutler- Beard flap with a free tarsal graft. Eye;18:714-7.

85-Huber A, Huber JD and Skinner RB Jr (2004):

Topical imiquimod treatment for nodular basal cell carcinomas: an open-label series. Dermatol Surg; 30: 429–430.

86- Hudson DA (2004):

Other ways to use tissue expanded flaps; Br. J. Plast. Surg., 57 (4) : 336 – 41

87- Iida N., Ohsumi N., Tonegawa M., and Tsutsumi K (2001) :

Repair of full thickness defect of the nose using an expanded forehead flap and a glabellar flap ; Aesthetic Plast. Surg., 25 (1) : 15-9 .

References

88- Iida N, Ohsumi N, Tonegawa M and Tsutsumi K(1999):

Simple method of designing a bilobed flap. *Plast Reconstr Surg*;104:495-9.

89-Ingham Pw(1998):

The patched gene in development and cancer, *Curr opin Genet Dev*;8(1)88.

90- Inkster Ng SG., and Leatherbarrow B (2001):

The rhomboid flap in medial canthal reconstruction; *Br. J. Ophthalmol.*, 85 (5) : 556 – 9 .

91- Irvine F and McNab AA(2003).

A technique for reconstruction of upper lid marginal defects. *Br J Ophthalmol*;87:279-81.

92- Jakson IT(1997).

Local flaps reconstruction of defects after excision of non melanoma skin cancer. *Clin Plast Surg*;24:747.

93- Jemec GB and Holm EA(2003):

Nonmelanoma skin cancer in organ transplant patients. *Transplantation*; 75:253-7.

94- Jessica L,Fewkes,Mack L,Cheney and Sheldon V Pollack(1992):

Illustrated atlas of cutaneous surgery.chap.14:436-445.

95- Johnson TM, Tschen J and Ho C (1994):

Unusual basal cell carcinomas. *Cutis*;54:85–92.

96- Jones CC,Ansari SJ and Tschen JA(1991):

Cystic fibroepithelioma of pinkus.*J cutaneous pathology*;18:220-222.

97- Kadoi C, Hayasaka S, Kato T, Nagaki Y, Matsumoto M and Hayasaka Y(2000).

The Cutler-Beard bridge flap technique with use of donor sclera for upper eyelid reconstruction. *Ophthalmologica*;214:140-2.

98- Kato N and Ueno H(1993):

Infundibulocystic basal cell carcinoma. Am J Dermatopathol;15:265–267.

99- Khan JA and Garden VS(2002):

Combined Flap Repair of Moderate Lower Eyelid Defects. Ophthal Plast Reconstr Surg;18:202-4.

100- Kim MY, Park HJ and Baek SC (2002):

Mutations of the p53 and PTCH gene in basal cell carcinomas: UV mutation signature and strand bias. J Dermatol Sci;29:1–9.

101- Kersten RC, Anderson RL, Tse DT and Weinstein GL(1986):

Tarsal rotational flap for upper eyelid reconstruction. Arch Ophthalmol; 104:918-22.

102- Kricker A, Armstrong BK, English DR and Heenan PJ(1995):

Does intermittent sun exposure cause basal cell carcinoma? A case-control study in Western Australia. Int J Cancer 1995; 60:489-94.

103- Kuflik EG and Gage AA(1997):

Recurrent basal cell carcinoma treated with cryosurgery. J Am Acad Dermatol;37: 82–84.

104- Kumar P, Orton CI and McWilliam LJ (2000).

Incidence of incomplete excision in surgically treated basal cell carcinoma: a retrospective clinical audit. Br J Plast Surg ; 53: 563–566.

105- Lang PG and Maize JC(1986):

Histologic evolution of recurrent basal cell carcinoma and treatment implications. J Am Acad Dermatol;14:186–196.

106- Larrabee W and Makielski K (1993):

Surgical anatomy of the face.

107- Larrabee Wayne F.and David A(1997):

Principles of facial reconstruction,chap.1:8-26.

108- Lear JT, Tan BB and Smith AG (1997):

Risk factors for basal cell carcinoma in the UK: case-control study in 806 patients. J R Soc Med 1997;90:371-4.

109- Lear JT,and Smith AG(1997:

Basal cell carcinoma , postgrad Med J,73(863):585.

110- Lear JT, Smith AG and Bowers B (1997):

Truncal tumor site is associated with high risk of multiple basal cell carcinoma and is influenced by glutathione S-transferase, GSTT1, and cytochrome P450, CYP1A1 genotypes, and their interaction. J Invest Dermatol ;108:519-22.

111- Leffell DJ and and Brash DE(1996):

sunlight and skin cancer ,SCI Am;,275(1):52.

112- Leibovitch I and Selva D. (2004):

Modified Hughes flap Division at 7 days. Ophthalmology;111:2164-7.

113- Leibovitch I, Selva D, Davis G and Ghabrial R(2004):

Donor site morbidity in free tarsal grafts. Am J Ophthalmol;138:430-3.

114- Lever WF and Shamberg-Lever G(1989)

Histopathology of the Skin, 7th edn. Lippincott: Philadelphia.

References

115- Lichter MD, Karagas MR, Mott LA, Spencer SK, Stukel TA and Greenberg ER(2000):

Therapeutic ionizing radiation and the incidence of basal cell carcinoma and squamous cell carcinoma. Arch Dermatol;136:1007-11.

116- Lopez R., Dekeister C., Sleiman Z., and paoli JR(2003):

The temporal fascioctaneous island flap for oncologic oral and facial reconstruction ; J. Oral and Maxillofac. Surg., 61 (10) : 1150 – 5 .

117- Lovatt TJ, Lear JT and Bastrilles J (2005):

Associations between ultraviolet radiation, basal cell carcinoma site and histology, host characteristics, and rate of development of further tumors. J Am Acad Dermatol;52: 468-73.

118- Maloof AJ and Leatherbarrow B (2000):

The glabellar flap dissected. Eye;14:597-605.

119- Maloof A, Ng S and Leatherbarrow B(2001):

The Maximal Hughes Procedure. Ophthal Plast Reconstr Surg;17:96-102.

120- Mallon E and Dawber R(1996).

Cryosurgery in the treatment of basal cell carcinoma. Assessment of one and two freeze-thaw cycle schedules. Dermatol Surg; 22: 854–858.

121- Marcil I and Stern RS(2000):

Risk of developing a subsequent nonmelanoma skin cancer in patients with a history of nonmelanoma skin cancer: a critical review of the literature and meta-analysis. Arch Dermatol; 136:1524-30.

References

122- Marks R, Gebauer K and Shumack S (2001):

Imiquimod 5% cream in the treatment of superficial basal cell carcinoma: results of a multicenter 6-week dose–response trial. J Am Acad Dermatol; 44: 807–813.

123- Marshall DM., Amjad I., and Wolfe SA (2003):

Use of radial forearm flap deep, central, midfacial defects; Plast. Reconstr. Surg., 111 (1) : 56-64.

124- Mauriello JA Jr and Antonacci R(1994):

Single tarsoconjunctival flap(lower eyelid) for upper eyelid reconstruction (reverse modified Hughes procedure). Ophthalmic Surg;25:374-8.

125- McNab AA, Martin P, Bengner R, O'Donnell B and Kourt G (2001):

A prospective randomized study comparing division of the pedicle of modified Hughes flaps at two or four weeks.Ophthal Plast Reconstr Surg;17:3179.

126- Meadows AE and Manners RM(2003):

A Simple Modification of the Glabellar Flap in Medial Canthal Reconstruction. Ophthal Plast Reconstr Surg;19:313-5.

127- Menick F(1987):

Artistry in aesthetic surgery: Aesthetic perception and the subunit principle in facial aesthetic surgery.clin.plast.surg.14: 726.

128- Menick F(1998):

Facial reconstruction with local and distant tissue:The interface of aesthetic and reconstructive surgery. Plast.reconstr.surg. 102: 1424.

129- Menick F(2002):

A ten-year experience in nasal reconstruction with the three-stage forehead flap. Plast reconstr.surg.

130- Menick K(1994):

Facial reconstruction in regional units. Perspect.Plast. surg. 8: 104.

131- Mikkilineni R and Weinstock MA(2001):

Epidemiology of skin cancer. In: Atlas of Clinical Oncology: Skin Cancer. London, UK: BC Decker, Inc.: 1–15.

132- Millard, D. R. (1986):

Principiplization of Plastic Surgery. Boston: Little,Brown.

133- Miller DL and Weinstock MA(1994):

Nonmelanoma skin cancer in the United States: incidence. J Am Acad Dermatol;30:774–778.

134- Miller SJ(1991):

Biology of basal cell carcinoma (Part I).J Am Acad Dermatol;24:1–13.

135- Miller SJ(1991):

Biology of basal cell carcinoma (Part II). J Am Acad Dermatol 1991;24:161–175.

136- Miller SJ(1995):

Aetiology and pathogenesis of basal cell carcinoma. Clin Dermatol;13: 527-36.

References

137- Misago N, Mihara I and Ansai SI(2002):

Sebaceoma and related neoplasms with sebaceous differentiation. Am J Dermatopathol;24:294–304.

138- Misago N, Satoh T and Narisawa Y(2004):

Basal cell carcinoma with ductal and glandular differentiation: a clinico-pathological and immunohistochemical study of 10 cases. Eur J Dermatol;14:383–387.

139- Moore KL and Persuad TVN(1998):

The integumentary system. In: Before We Are Born: Essentials of Embryology and Birth Defects. 5th ed. :481-96.

140- Morris JL and Gibbins IL(1997):

Autonomic Innervation of the Skin. 1st. Informa Healthcare .

141- Nagao K, Fujii K and Yamada M (2004):

Identification of a novel polymorphism involving a CGG repeat in the PTCH gene and a genome-wide screening of CGG-containing genes. J Hum Genet;49: 97–101.

142- Nagore E, Grau C and Molinero J(2003):

Positive margins in basal cell carcinoma: relationship to clinical features and recurrence risk. A retrospective study of 248 patients. J Eur Acad Dermatol Venereol; 17: 167–170.

143- Nisanci M., Turegun M., Er E., and Segezer M.(2003):

Reconstruction of the middle and lower face with three simultaneous free flaps: combined use of bilateral fibular flaps for maxillomandibular reconstruction; Ann. Plast. Surg., 51 (3) : 301 – 7 .

References

144- Niazi ZBM and Lamberty BGH(1993):

Perineural infiltration in basal cell carcinomas. Br J Plast Surg;6:156–157. 19
Sahl WJ. Basal cell carcinoma: Influence of tumor size on mortality and morbidity. Int J Dermatol 1995;34: 319–321.

145- Nijsten TE and Stern RS(2003):

The increased risk of skin cancer is persistent after discontinuation of psoralen+ultraviolet A: a cohort study. J Invest Dermatol;121:252-8.

146- Ng SG, Inkster CF and Leatherbarrow B(2001):

The rhomboid flap in medial canthal reconstruction. Br JOphthalmol ; 85:556-9.

147- Omid M and Granick MS(2004):

The versatile V-Y flap for facial reconstruction ; Dermatol. Surg., 30 (3) : 415 – 20

148- Oram Y, Orengo I and Griego RD (1995):

Histologic patterns of basal cell carcinoma based upon patient immunostatus. Dermatol Surg;21:611–614.

149- Papel Ira,John Frodel,G.Richard Holt,wayne F.Larrabee,Nathan Nachlas, Steven s.Park,Jonathan.Sykes,and Dean Toriumi(2002):

Facial plastic and reconstructive surgery,second edition, (1):3-14,(45):549-559.

150- Patrinely JR, O'Neal KD, Kersten RC and Soparkar CN(1994):

Total Upper Eyelid Reconstruction With Mucosalized Tarsal Graft and Overlying Bipedicle Flap. Arch Ophthalmol 1999;117:1655-61.

151- Peschen M, Lo JS and Snow SN (1993):

Linear basal cell carcinoma. Cutis;51:287–289.

References

152- Pinho C., Choupina M., Silva P., Ferreira P., Guimaraes I., and Amarante J.: (2003)

A new retroauricular flap for facial reconstruction ; Br. J. Plast. Surg., 56 (6) : 599 – 602.

153- Pinkus H(1953):

Premalignant fibroepithelial tumors of the skin. Arch Dermatol;67:598–603.

154- Poblet E, Jiménez F and Ortega F(2004):

The contribution of the arrector pili muscle and sebaceous glands to the follicular unit structure. J Am Acad Dermatol;51(2):217-22.

155- Porfiris E, Kalokerinos D, Christopoulos A, Damilakos P, Ioannidis A and Georgiou P(2000):

Upper Eyelid Island Orbicularis Oculi Myocutaneous Flap for Periorbital Reconstruction. *Ophthal Plast Reconstr Surg* 2000;16:42-4.

156- Pratt MD and Jackson R(1987):

Nevoid basal cell carcinoma syndrome. J Am Acad Dermatol;16:964–970.

157- Pribitkin ED and Boger G(2001):

Herbal therapy: what every facial plastic surgeon must know. Arch Facial Plast Surg; 3: 127–132.

158- Price MA, Goldberg LH and Levy ML(1994):

Juvenile basal cell carcinoma. Ped Dermatol;11:176–177.

159- Putterman AM(2002):

Bolster for the Tarsconjunctival Flap-Skin Graft. *Ophthal Plast Reconstr Surg*;18:466-7.

References

160- Randle HW(1996):

Basal cell carcinoma: identification and treatment of the high-risk patient.
Dermatol Surg ;22:255–261.

161- Roger I.Ceilley, James Q.Del Rosso and DO FAOCD (2006):

Current modalities and new advances in the treatment of basal cell carcinoma,
International Journal of Dermatology, 45, 489 –498.

162- Rhodes LE, de Rie M and Enstrom Y (2004):

Photodynamic therapy using topical methyl aminolevulinate vs surgery for
nodular basal cell carcinoma: results of a multicenter randomized prospective
trial. Arch Dermatol; 140:17–23.

163- Robinson JK (2004):

Segmental reconstruction of the face; Dermatol. Surg., 30 (1) : 67 – 74 .

164- Rohrich RJ and Zbar RI(1999):

The Evolution of the Hughes Tarsoconjunctival Flap for Lower Eyelid
Reconstruction. Plast Reconstruct Surg;104:518-22.

165- Ross Michael,Gordon I.kaye,and Wojeiech Pawlina(2003):

Histology:Atext and atlas,fourth edition,chap.14:400-421.

166- Russell R.C(1993):

Flaps for closure of facial defects.Instruction,vol.6,St.Louis,Losby Yearbook

167- Russell EB, Carrington PR and Smoller BR(1999):

Basal cell carcinoma: a comparison of shave biopsy versus punch biopsy
techniques in subtype diagnosis. J Am Acad Dermatol;41:69–71.

References

168- Scrivener Y, Grosshans E and Cribier B(2002):

Variations of basal cell carcinomas according to gender, age, location and histopathological subtype. *Br J Dermatol*; 147: 41–47.

169- Scuderi N, Ribuffo D and Chiummariello S(2005):

Total and Subtotal Upper Eyelid Reconstruction with the Nasal Chondro - mucosal Flap: A 10-Year Experience. *Plast Reconstr Surg*;115:1259-65.

170- Sexton M, Jones DB and Maloney ME(1990):

Histologic pattern analysis of basal cell carcinoma: study of a series of 1039 consecutive neoplasms. *J Am Acad Dermatol*;23: 1118-26.

171- Siegle RJ, MacMillan J and Pollack SV (1986):

Infiltrative basal cell carcinoma: a nonsclerosing subtype. *J Dermatol Surg Oncol*;12:830–836.

172- Shrivastava Rajeev, Singh KK and Shrivastava Minakshi(2007):

Soft tissue metastasis in basal cell carcinoma, *Indian Journal of Dermatology*, 52(4); 206-208.

173- Shumack S, Robinson J and Kossard S(2002):

Efficacy of topical 5% imiquimod cream for the treatment of nodular basal cell carcinoma: comparison of dosing regimens. *Arch Dermatol*;138 :1165 –1171

174- Sihota R, Tandon K, Betharia SM and Arora R(1996):

Malignant eyelid tumors in an Indian population. *Arch Ophthalmol*; 114:108-9.

175- Silverman MK, Kopf AW and Grin CM (1991):

Recurrence rates of treated basal cell carcinomas. Part 2: Curettage electrodesiccation. *J Dermatol Surg Oncol*; 17: 720–726.