

REFERENCES:

- Abbas A, Deschamps C and Cassivi SD (2004):** The role of laparoscopic fundoplication in Barrett's esophagus. *Annals of Thoracic Surgery.*, 77(2):393-96.
- Akiyama H (1980):** Surgery for carcinoma of the esophagus. *Curr. Probl. Surg.*, 17: 53.
- Attwood SA, Lewis CJ and Caplin S (2003):** Argon beam plasma coagulation as therapy for high-grade dysplasia in Barrett's esophagus, *Clin Gastroenterol Hepatol.*, 1pp. 258–63.
- Barrett N (1957):** The lower esophagus lined by columnar epithelium. *Surgery* 41 (6):881-94.
- Beilstein M and Silberg D (2002):** Cellular and molecular mechanisms responsible for progression of Barrett's metaplasia to esophageal carcinoma. *Gastroenterol Clin North Am.*, 31:461-79.
- Bergman JJ and Tytgat GN (2005):** New developments in the endoscopic surveillance of Barrett's esophagus. *Gut.*, 54(suppl 1):138–42.
- Blot WJ and McLaughlin JK (1999):** The changing epidemiology of esophageal cancer. *Semin Oncol.*, 26:2–8.
- Boyce H and Boyce G (2003):** Esophagus: Anatomy and structural anomalies. In: Yamada T, Alpers DH, Kaplowitz N, Laine L, Owyang C, Powell DW, eds. *Textbook of Gastroenterology*, 4th ed., vol. 1. Philadelphia: Lippincott Williams & Wilkins., 1148 –65.
- Bruna J (1991):** Types of collateral lymphatic circulation. *Lymphology* 7: 61, 1974. Quoted from: Shackelford's surgery of the alimentary tract, 3rd edition, volume 1, W.B. Saunders company.

- Bulter H (1991):** The veins of the esophagus. *Thorax* 6: 276, 1951.
Quoted from: Shackelford's surgery of the alimentary tract, 3rd edition,
volume 1, W.B. Saunders company.
- Buttar NS, Wang KK and Sebo TJ (2001):** Extent of high-grade
dysplasia in Barrett's esophagus correlates with risk of
adenocarcinoma. *Gastroenterology*., 120: 1630-39.
- Cameron AJ and Lomboy CT (1992):** Barrett's esophagus: age,
prevalence, and extent of columnar epithelium. *Gastroenterology*.,
103(4):1241-45.
- Cameron AJ, Souto EO and Smyrk TC (2002):** Small
adenocarcinomas of the esophagogastric junction: association with
intestinal metaplasia and dysplasia. *Am J Gastroenterol.*, 97:1375-80.
- Campos GM, DeMeester SR and Peters JH (2001):** Predictive
factors of Barrett esophagus: multivariate analysis of 502 patients with
gastroesophageal reflux disease. *Arch Surg.*, 136:1267-73.
- Chak A, Lee T and Kinnard MF (2002):** Familial aggregation of
Barrett's esophagus, esophageal adenocarcinoma, and esophagogastric
junctional adenocarcinoma in Caucasian adults. *Gut.*, 51: 323-28.
- Chandrasoma P, Der R and Dalton P (2001):** Distribution and
significance of epithelial types in columnar-lined esophagus. *Am. J.*
Surg. Pathol., 25, 1188-93.
- Chandrasoma PT, Lokuhetty DM and DeMeester TR (2000):**
Definition of histopathologic changes in gastroesophageal reflux
disease. *Am J Surg Pathol.*, 24:344-51.

Charbel S (2005): The role of esophageal pH monitoring in symptomatic patients on PPI therapy. *Am J Gastroenterol.*, 100: 283-89.

Chaves P, Crespo M and Ribeiro C (2007): Chromosomal analysis of Barrett's cells: demonstration of instability and detection of the metaplastic lineage involved. *Mod Pathol.*, 20:788-96.

Corley DA, Kubo A and Levin TR (2007): Abdominal obesity and body mass index as risk factors for Barrett's esophagus. *Gastroenterology.*, 133: 34-41.

Curvers W, Baak L and Kiesslich R (2008): Chromoendoscopy and narrow-band imaging compared with high-resolution magnification endoscopy in Barrett's esophagus. *Gastroenterology.*, 134: 670-79.

Cuschieri A (2000): Hiatal hernia and reflux esophagitis In: *Minimal invasive surgery*. Hunter J and Sackier M., International edition. McGraw Hill; New York; Ch. 10 p.87-111.

Delattre JF (2000): Functional anatomy of the gastroesophageal junction. *Surg Clin North Am.*, 80(1): 241-60.

Dincer D, Besisik F, Sahin E and Demir K (2002): Intestinal metaplasia of the gastric cardia: a study from Turkey. *Hepatogastroenterol.*, 49: 1153-56.

Dulai GS, Jensen DM and Cortina G (2005): Randomized trial of argon plasma coagulation vs. multipolar electrocoagulation for ablation of Barrett's esophagus. *Gastrointest Endosc.*, 61:232-40.

Dvorak K, Payne CM and Chavarria M (2007): Bile acids in combination with low pH induce oxidative stress and oxidative DNA damage: relevance to the pathogenesis of Barrett's esophagus. *Gut.*, 56: 763-71.

Edelstein ZR, Farrow DC, Bronner MP, Rosen SN and Vaughan TL (2007): Central adiposity and risk of Barrett's esophagus. *Gastroenterology.*, 133: 403-11.

Ell C, May A and Pech O (2007): Curative endoscopic resection of early esophageal adenocarcinomas (Barrett's cancer), *Gastrointest Endosc.*, 65pp. 3-10.

Eloubeidi MA and Provenzale D (1999): Does this patient have Barrett's esophagus? The utility of predicting Barrett's esophagus at the index endoscopy. *Am J Gastroenterol.*, 94: 937-43.

El-Serag HB, Aguirre TV, Davis S, Kuebel M, Bhattacharyya A and Sampliner RE (2004): Proton pump inhibitors are associated with reduced incidence of dysplasia in Barrett's esophagus. *Am J Gastroenterol.*, 99: 1877-83.

Evans JA, Poneros JM and Bouma BE (2006): Optical coherence tomography to identify intramucosal carcinoma and high-grade dysplasia in Barrett's esophagus, *Clin Gastroenterol Hepatol.*, 4 pp. 38-43.

Falk GW (1999): Endoscopic surveillance of Barrett's esophagus: risk stratification and cancer risk. *Gastrointest Endosc.*, 49 (3 pt -2): S 2934.

Fléjou J (2005): "Barrett's esophagus: from metaplasia to dysplasia and cancer". *Gut* 54 Suppl. 1: 6-12.

- Fu M (2004):** Embryonic development of the ganglion plexuses and the concentric layer structure of human gut: a topographical study. *Anat Embryol (Berl)*., 208(1):33–41.
- Galmiche JP, Sacher-Huvelin S and Coron E (2008):** Screening for esophagitis and Barrett's esophagus with wireless esophageal capsule endoscopy: a multicenter prospective trial in patients with reflux symptoms. *Am J Gastroenterol*., 103: 538-45.
- Ganz RA, Overholt BF and Sharma VK (2008):** Circumferential ablation of Barrett's esophagus that contains high-grade dysplasia: a US multicenter registry. *Gastrointest Endosc*., 68: 35-40.
- Gatenby P, Ramus J and Caygill C (2008):** Relevance of the detection of intestinal metaplasia in non-dysplastic columnar-line oesophagus. *Scand. J. Gastroenterol*., 43, 524–30.
- Gerson LB, Shetler K and Triadafilopoulos G (2002):** Prevalence of Barrett's esophagus in asymptomatic individuals. *Gastroenterology*., 123:461-67.
- Ghosh SK (2006):** Physiology of the esophageal pressure transition zone : separate contraction waves above and below . *Am J Physiol Gastrointest Liver Physiol*., 290 (3) : G568 –76.
- Gopal DV, Faigel DO and Magaret N (1999):** Risk factors for dysplasia in patients with Barrett's esophagus (BE): results from a multicenter consortium. (Abstr) *Gastroenterology*., 116(4 Pt 2):A175.
- Gopal DV, Lieberman DA and Margaret N (2003):** Risk factors for dysplasia in patients with Barrett's esophagus (BE): results from a multicenter consortium. *Dig Dis Sci*., 48:1537-41.
- Goyal R and Sivarao D (1999):** Functional anatomy and physiology of swallowing and esophageal motility. In: Catell OD, Richter JE, eds. *The Esophagus*, 3rd ed . Philadelphia: Lippincott Williams& Wilkins., 24 -26.
-

Guelrud M, Herrera I, Essenfled H and Castro J (2001): Enhanced magnification endoscopy: a new technique to identify specialized intestinal metaplasia in Barrett's esophagus. *Gastrointest Endosc.*, 53: 559-65.

Hahn H, Shahsafaei A and Odze RD (2008): Vascular and lymphatic properties of the superficial and deep lamina propria in Barrett's esophagus. *Am. J. Surg. Pathol.*, 32, 1454–61.

Hamamoto Y, Endo T, Nosho K, Arimura Y, Sato M and Imai K (2004): Usefulness of narrow-band imaging endoscopy for diagnosis of Barrett's esophagus. *J Gastroenterol.*, 39: 14-20.

Hampel H, Abraham NS and El-Serag HB (2005): Meta-analysis: obesity and the risk for gastro-esophageal reflux disease and its complications. *Ann Intern Med.*, 143:199.

Hanna S, Rastogi A and Weston AP (2006): Detection of Barrett's esophagus after endoscopic healing of erosive esophagitis. *Am J Gastroenterol.*, 101: 1416-20.

Harrison R, Perry I and Haddadin W (2007): Detection of intestinal metaplasia in Barrett's esophagus: an observational comparator study suggests the need for a minimum of eight biopsies. *Am J Gastroenterol.*, 102:115-61.

Hassall E (1997): Columnar-lined esophagus in children. *Gastroenterol Clin North Am.*, 26:33.

Heath EI, Canto MI and Piantadosi S (2007): Secondary chemoprevention of Barrett's esophagus with celecoxib: results of a randomized trial. *J Natl Cancer Inst.*, 99: 545-57.

Hornby PJ and Abrahams TP (2000): Central control of lower esophageal sphincter relaxation. *Am J Med.*, 108(suppl 4a):90S–98S.

Inadomi JM, Sampliner R, Lagergren J, Lieberman D, Fendrick AM and Vakil N (2003): Screening and surveillance for Barrett esophagus in high-risk groups: a cost–utility analysis. *Ann Intern Med.*, 138: 176–80.

Isenberg G, Sivak MV Jr and Chak A (2005): Accuracy of endoscopic optical coherence tomography in the detection of dysplasia in Barrett's esophagus: a prospective, double-blinded study. *Gastrointest Endosc.*, 62: 825-31.

Jamieson (1997): Hiatal Hernia and gastro-esophageal reflux : Text Book of surgery, The Biological basis of Modern surgical practice: Ed 15, 167-783.

Jobe BA, Hunter JG and Chang EY (2006): Office-based unsedated small-caliber endoscopy is equivalent to conventional sedated endoscopy in screening and surveillance for Barrett's esophagus: a randomized and blinded comparison. *Am J Gastroenterol.*, 101: 2693-703.

Johnston MH, Cash BD and Dykes CA (2006): Cryoablation of dysplasia in Barrett's esophagus (BE) and early stage esophageal cancer [abstract], *Gastrointest Endosc.*, 63 AB223.

Kahrilas PJ, Shaheen NJ and Vaezi M (2008): American Gastroenterological Association Medical Position Statement on the Management of Gastro-esophageal Reflux Disease. *Gastroenterology.*, 135:1383.

Kakushima N, Yahagi N and Fujishiro M (2006): Efficacy and safety of endoscopic submucosal dissection for tumors of the esophagogastric junction, *Endoscopy.*, 38 pp. 170–74.

Kala Z, Dolina J, Marek F and Izakovicova HL (2007): Polymorphisms of glutathione S-transferase M1, T1 and P1 in patients with reflux esophagitis and Barrett's esophagus. *J Hum Genet.*, 52: 527-34.

Kaltenbach T, Crockett S and Gerson LB (2006): Are lifestyle measures effective in patients with gastro-esophageal reflux disease? An evidence-based approach. *Arch Intern Med.*, 166:965.

Kara MA, Ennahachi M, Fockens P, ten Kate FJ and Bergman JJ (2006): Detection and classification of the mucosal and vascular patterns (mucosal morphology) in Barrett's esophagus by using narrow band imaging. *Gastrointest Endosc.*, 64: 155-66.

Kara MA, Peters FP, ten Kate FJ, Van Deventer SJ, Fockens P and Bergman JJ (2005): Endoscopic video autofluorescence imaging may improve the detection of early neoplasia in patients with Barrett's esophagus. *Gastrointest Endosc.*, 61: 679-85.

Kerkhof M, van Dekken H and Steyerberg EW (2007): Grading of dysplasia in Barrett's esophagus: substantial interobserver variation between general and gastrointestinal pathologists. *Histopathology.*, 50:920-27.

Kiesslich R, Gossner L and Goetz M (2006): In vivo histology of Barrett's esophagus and associated neoplasia by confocal laser endomicroscopy. *Clin Gastroenterol Hepatol.*, 4: 979-87.

Konda VJ, Ross AS and Ferguson MK (2008): Is the risk of concomitant invasive esophageal cancer in high-grade dysplasia in Barrett's esophagus overestimated? *Clin Gastroenterol Hepatol.*, 6: 159-64.

- Koppert L, Wijnhoven B, van Dekken H, Tilanus H and Dinjens W (2005):** The molecular biology of esophageal adenocarcinoma. *J Surg Oncol* 92 (3): 169-90.
- Lagergren J, Bergstrom R and Nyren O (1999):** Association between body mass and adenocarcinoma of the esophagus and gastric cardia. *Ann Intern Med.*, 130:883-90.
- Larghi A, Lecca PG and Costamagna G (2008):** High-resolution narrow band imaging endoscopy. *Gut.*, 57: 976-86.
- Larsen W (2001):** Development of the gastrointestinal tract. In: Sherman LS, Potter SS, Scott WJ, eds. *Human Embryology*, 3rd ed. Philadelphia: Churchill Livingstone., 235–64.
- Lauwers GY, Ban S and Mino M (2004):** Endoscopic mucosal resection for gastric epithelial neoplasms: a study of 39 cases with emphasis on the evaluation of specimens and recommendations for optimal pathologic analysis, *Mod Pathol.*, 17pp. 2–8.
- Le Douarin NM (2004):** Neural crest cell plasticity and its limits . *Development.*, 131(19) : 4637-50.
- Libermann-Meffert D and Brauer B(1995):** In surgical anatomy of distal esophagus and cardia in esophagus, stomach and small intestine. Edited by Christopher Wastell, Lioted M. Nyhus and Philip E. Donahue, fifth ed., 31:44.
- Lim CH, Rotimi O, Dexter SP and Axon AT (2006):** Randomized crossover study that used methylene blue or random 4-quadrant biopsy for the diagnosis of dysplasia in Barrett's esophagus. *Gastrointest Endosc.*, 64: 195-99.

- Lin OS, Schembre DB and Mergener K (2007):** Blinded comparison of esophageal capsule endoscopy versus conventional endoscopy for a diagnosis of Barrett's esophagus in patients with chronic gastro-esophageal reflux. *Gastrointest. Endosc.*, 65: 577-83.
- Liu T, Zhang X and So CK (2007):** Regulation of CDX2 expression by promoter methylation, and effects of CDX2 transfection on morphology and gene expression of human esophageal epithelial cells. *Carcinogenesis*, 28: 488-96.
- Liu W (2009):** Metaplastic esophageal columnar epithelium without goblet cells shows DNA content abnormalities similar to goblet cell containing epithelium. *Am. J. Gastroenterol.*, 104, 816–24.
- Locke GR, Zinsmeister AR and Talley NJ (2003):** Can symptoms predict endoscopic findings in GERD? *Gastrointest. Endosc.* , 58: 661-70.
- Lomo L, Blount PL, Sanchez CA (2006):** Crypt dysplasia with surface maturation: a clinical, pathologic and molecular study of a Barrett's esophagus cohort. *Am J Surg Pathol.*, 30:423-35.
- Long JD and Orlando RC (1999):** Esophageal submucosal glands: structure and function . *Am J Gastroenterol.*, 94 (10): 2818 –24.
- Long JD and Orlando RC (2002):** Anatomy, histology, embryology, and developmental abnormalities of the esophagus. In: Feldman M, Fieldman LS, Sleisenger MH, eds. *Gastrointestinal and liver Diseases*. Philadelphia : WB Saunders., 551-60.
- Maley CC, Galipeau PC and Finley JC (2006):** Genetic clonal diversity predicts progression to esophageal adenocarcinoma. *Nat Genet.*, 38:468-73.
- Maley CC and Reid BJ (2005):** Natural selection in neoplastic progression of Barrett's esophagus. *Semin Cancer Biol.*, 15:474-83.
-

Marchand P (1959): The anatomy of the esophageal hiatus of the diaphragm and the pathogenesis of hiatus herniation. *Thorac Surg.*, 37:81–92.

May A, Gossner L and Pech O (2002): Intraepithelial high-grade neoplasia and early adenocarcinoma in short-segment Barrett's esophagus (SSBE): curative treatment using local endoscopic treatment techniques. *Endoscopy.*, 34: 604-10.

McManus DT, Olaru A and Meltzer SJ (2004): Biomarkers of esophageal adenocarcinoma and Barrett's esophagus. *Cancer Res.*, 64:1561–9.

Miros M, Kerlin P and Walker N (1991): Only patients with dysplasia progress to adenocarcinoma in Barrett's esophagus. *Gut.*, 32:1441-46.

Mittal RK and Balaban DH (1997): The esophagogastric junction. *N Engl J Med.*, 336(13): 924-32.

Montgomery E, Bronner MP and Goldblum JR (2001): Reproducibility of the diagnosis if dysplasia in Barrett's esophagus: a reaffirmation. *Hum Pathol.*, 32:368-78.

Morales CP, Souza RF and Spechler SJ (2002): Hallmarks of cancer progression in Barrett's esophagus. *Lancet.*, 360:1587-89.

Oberg S, Johansson J and Wenner J (2001): Endoscopic surveillance of columnar-lined esophagus: frequency of intestinal metaplasia detection and impact of antireflux surgery. *Ann Surg.*, 234:619-26.

O'Connor JB, Falk GW and Richter JE (1999): The incidence of adenocarcinoma and dysplasia in Barrett's esophagus: report on the Cleveland Clinic Barrett's Esophagus Registry. *Am J Gastroenterol.*, 94:2037-42.

Odze RD (2005): Pathology of the gastroesophageal junction. *Semin. Diag. Pathol.*, 22, 256–65.

Odze RD (2005): Unraveling the mystery of the gastroesophageal junction: a pathologist's perspective. *Am J Gastroenterol.*, 100:1853-67.

Odze RD (2006): Diagnosis and grading of dysplasia in Barrett's esophagus. *J Clin Pathol.*, 59:1029-38.

Odze RD (2008): Update on the diagnosis and treatment of Barrett's esophagus and related neoplastic precursor lesions. *Arch. Pathol. Lab. Med.*, 132, 1577–85.

Ormsby AH, Petras RE and Henricks WH (2002): Observer variation in the diagnosis of superficial esophageal adenocarcinoma. *Gut.*, 51: 671-76.

Overholt BF, Lightdale CJ and Wang KK (2005): on behalf of the International Photodynamic Group for High-Grade Dysplasia in Barrett's Esophagus. Photodynamic therapy with porfimer sodium for ablation of high-grade dysplasia in Barrett's esophagus: international, partially blinded, randomized phase III trial. *Gastrointest Endosc.*, 62: 488-98.

Overholt BF, Panjehpour M and Halberg DL (2003): Photodynamic therapy for Barrett's esophagus with dysplasia and/or early stage carcinoma: long term results, *Gastrointest Endosc.*, 58 pp. 183–88.

Pandolfino JE (2003): Ambulatory esophageal pH monitoring using a wireless technique. *Am J Gastroenterol.*, 98: 545–50.

Paraf F, Flejou JF and Pignon JP (1995): Surgical pathology of adenocarcinoma arising in Barrett's esophagus: analysis of 67 cases. *Am J Surg Pathol.*, 19: 183-91.

- Parrilla P, Martinez de Haro LF and Ortiz A (2003):** Long-term results of a randomized prospective study comparing medical and surgical treatment of Barrett's esophagus. *Ann Surg.*, 237: 291-98.
- Paull A (1976):** The histologic spectrum of Barrett's esophagus. *N. Engl. J. Med.*, 295, 476-80.
- Pera M, Pera M and de Bolos C (2007):** Duodenal-content reflux into the esophagus leads to expression of Cdx2 and Muc2 in areas of squamous epithelium in rats. *J Gastrointest Surg.*, 11: 869-74.
- Peter L. Williams, Lawrence H and Bannister (1995):** *Grays anatomy*, 36th ed., 2: 189-92.
- Peters FP, Kara MA and Rosmolen WD (2005):** Endoscopic treatment of high-grade dysplasia and early stage cancer in Barrett's esophagus, *Gastrointest Endosc.*, 61 pp. 506–14.
- Peters FP, Kara MA and Rosmolen WD (2006):** Stepwise radical endoscopic resection is effective for complete removal of Barrett's esophagus with early neoplasia: a prospective study, *Am J Gastroenterol.*, 101pp. 1449–57.
- Peyre CG, DeMeester SR and Rizzetto C (2007):** Vagal-sparing esophagectomy: the ideal operation for intramucosal adenocarcinoma and Barrett with high-grade dysplasia. *Ann Surg.*, 246: 665-71.
- Playford RJ (2006):** New British Society of Gastroenterology (BSG) guidelines for the diagnosis and management of Barrett's esophagus. *Gut.*, 55: 442.
- Poneros JM, Brand S, Bouma BE, Tearney GJ, Compton CC and Nishioka NS (2001):** Diagnosis of specialized intestinal metaplasia by optical coherence tomography. *Gastroenterology.*, 120: 7-12.

- Prasad GA, Wang KK and Buttar NS (2007):** Predictors of stricture formation after photodynamic therapy for high-grade dysplasia in Barrett's esophagus, *Gastrointest Endosc.*, **65** pp. 60–66.
- Ragunath K, Krasner N, Raman VS, Haqqani MT and Cheung WY (2003):** A randomized, prospective cross-over trial comparing methylene blue-directed biopsy and conventional random biopsy for detecting intestinal metaplasia and dysplasia in Barrett's esophagus. *Endoscopy.*, **35**: 998-1003.
- Reid BJ, Haggitt RC and Rubin CE (1988):** Observer variation in the diagnosis of dysplasia in Barrett's esophagus. *Hum Pathol.*, **19**:166-78.
- Reshamwala P and Darwin p (2006):** Endoscopic management of early gastric cancer. *Curr Opin Gastroenterol.*, **22**(5):541-45.
- Rice TW, Blackstone EH and Adelstein DJ (2003):** Role of clinically determined depth of tumor invasion in the treatment of esophageal carcinoma, *J Thorac Cardiovasc Surg.*, **125**pp. 1091–102.
- Ronkainen J, Aro P and Storskrubb T (2005):** Prevalence of Barrett's esophagus in the general population: an endoscopic study. *Gastroenterology.*, **129**: 1825-31.
- Romero Y, Cameron AJ and Schaid DJ (2002):** Barrett's esophagus: prevalence in symptomatic relatives. *Am J Gastroenterol.*, **97**:1127-32.
- Rubenstein JH, Inadomi JM, Brill JV and Eisen GM (2007):** Cost utility of screening for Barrett's esophagus with esophageal capsule endoscopy versus conventional upper endoscopy. *Clin Gastroenterol Hepatol.*, **5**: 312-18.

- Sampliner RE (2002):** For the Practice Parameters Committee of the American College of Gastroenterology. Updated guidelines for the diagnosis, surveillance, and therapy of Barrett's esophagus. *Am J Gastroenterol.*, 97:1888–95.
- Sarosi G, Brown G and Jaiswal K (2008):** Bone marrow progenitor cells contribute to esophageal regeneration and metaplasia in a rat model of Barrett's esophagus. *Dis Esophagus.*, 21: 43-50.
- Schlemper RJ, Riddell RH and Kato Y (2000):** The Vienna classification of gastrointestinal epithelial neoplasia. *Gut* ., 47:251–55.
- Schnell TG, Sontag SJ and Chejfec G (2001):** Long-term nonsurgical management of Barrett's esophagus with high-grade dysplasia. *Gastroenterology.*, 120: 1607-19.
- Shaheen NJ, Green B and Medapalli RK (2005):** The perception of cancer risk in patients with prevalent Barrett's esophagus enrolled in an endoscopic surveillance program. *Gastroenterology.*, 129: 429-36.
- Shaheen N and Ransohoff DF (2002):** Gastro-esophageal reflux, Barrett' esophagus, and esophageal cancer: Scientific review. *JAMA.*, 287: 1972-81.
- Shaheen NJ and Richter JE (2009):** "Barrett's oesophagus". *Lancet* 373 (9666): 850–61.
- Sharma P, Bansal A and Mathur S (2006):** The utility of a novel narrow band imaging endoscopy system in patients with Barrett's esophagus. *Gastrointest Endosc.*, 64: 167-75.
- Sharma P, Dent J and Armstrong D (2006):** The development and validation of an endoscopic grading system for Barrett's esophagus: the Prague C & M criteria. *Gastroenterology.*, 131: 1392-99.

References

- Sharma P, McCallum RW and Lundell L (2000):** The geoprevalence of Barrett's esophagus. (Paper presented at the Sixth OESO World Congress; September 2000, Paris, France).
- Sharma P, McQuaid K and Dent J (2004):** A critical review of the diagnosis and management of Barrett's esophagus: the AGA Chicago Workshop. *Gastroenterology.*, 127:310.
- Sharma VK, Wang KK and Overholt BF (2007):** Balloon-based, circumferential, endoscopic radiofrequency ablation of Barrett's esophagus: 1-year follow-up of 100 patients. *Gastrointest Endosc.*, 65: 185-95.
- Skacel M, Petras RE, Gramlich TL and Goldblum JR (2000):** The diagnosis of low- grade dysplasia in Barrett's esophagus and its implications for disease progression. *Am J Gastroenterol.*, 95:3383-87.
- Skandalakis JE and Ellis H (2000):** Embryologic and anatomic basis of esophageal surgery. *Surg Clin North Am.*, 80 (1):85-155.
- Spechler SJ (2002):** Columnar-lined esophagus. Definitions. *Chest Surg Clin N Am.*, 12 :1-13.
- Spechler SJ (2004):** Intestinal metaplasia at the gastroesophageal junction. *Gastroenterology.*, 126, 567-75.
- Spechler SJ (2005):** Dysplasia in Barrett's esophagus: limitations of current management strategies. *Am J Gastroenterol.*, 100:927-35.
- Srivastava A (2007):** Morphological features are useful in distinguishing Barrett's esophagus from carditis with intestinal metaplasia. *Am. J. Surg. Pathol.*, 31, 1733-41.
- Stein H and Siewert J (1993):** Barrett's esophagus: pathogenesis, epidemiology, functional abnormalities, malignant degeneration and surgical management. *Dysphagia.*, 8 (3):276-88.

References

- Swisher SG, DeFord L and Merriman KW (2000):** Effects of operative volume on morbidity, mortality, and hospital use after esophagectomy for cancer. *J Thorac Cardiovasc Surg.*, 119:1126.
- Theisen J (2006):** Preferred location for the development of esophageal adenocarcinoma within a segment of intestinal metaplasia. *Surg. Endosc.*, 20, 235–38.
- Trudgill NJ, Kapur KC and Riley SA (1999):** Familial clustering of reflux symptoms. *Am J Gastroenterol.*, 94:1172-78.
- Vaezi MF and Richter JE (1997):** Bile reflux in columnar-lined esophagus. *Gastroenterol Clin North Am.*, 26:565–82.
- Vakil N, van Zanten SV, Kahrilas P, Dent J, Jones R, and The Global Consensus Group (2006):** The Montreal definition and classification of gastroesophageal reflux disease: a global evidence-based consensus. *Am J Gastroenterol.*, 101: 1900-20.
- Van Sandick JW, van Lanschot JJ, Kuiken BW, Tytgat GN, Offerhaus GJ and Obertop H (1998):** Impact of endoscopic biopsy surveillance of Barrett's esophagus on pathological stage and clinical outcome of Barrett's carcinoma. *Gut.*, 43:216–22.
- Wallace AS and Burns AJ (2005):** Development of the enteric nervous system, smooth muscle and interstitial cells of Cajal in the human gastrointestinal tract. *Cell Tissue Res.*, 319(3):367–82.
- Wang KK and Sampliner RE (2008):** Practice Parameters Committee of the American College of Gastroenterology. Updated guidelines 2008 for the diagnosis, surveillance, and therapy of Barrett's esophagus. *Am J Gastroenterol.*, 103:788-97.

- Ward EM, Wolfsen HC and Achem SR (2006):** Barrett's esophagus is common in older men and women undergoing screening colonoscopy regardless of reflux symptoms. *Am J Gastroenterol.*, 101: 12-17.
- Westhoff B, Brotze S and Weston A (2005):** The frequency of Barrett's esophagus in high-risk patients with chronic GERD. *Gastrointest Endosc.*, 61: 226-31.
- Wild CP and Hardle LJ (2003):** Reflux, Barrett's esophagus and adenocarcinoma: burning questions. *Cancer.*, 3:676-84.
- Williamson WA, Ellis FH Jr and Gibb SP (1991):** Barrett's esophagus: prevalence and incidence of adenocarcinoma. *Arch Intern Med.*, 151:2212-16.
- Williams VA, Watson TJ and Herbella FA (2007):** Esophagectomy for high grade dysplasia is safe, curative, and results in good alimentary outcome. *J Gastrointest Surg.*, 11: 1589-97.
- Yu C, Zhang X and Huang Q (2007):** High-fidelity DNA histograms in neoplastic progression in Barrett's esophagus. *Lab Invest.*, 87:466-72.
- Yurdaydin C, Krastev Z and Bulajic M (2002):** Gastrointestinal tract cancer in Southeastern Europe: a multinational study with the participation of seven countries. *Gut.*, 51 (Suppl. III): A263.
- Zhang X, Goyal R, Huang Q and Odze Rd (2008):** DNA ploidy abnormalities in basal and superficial regions of the crypts in Barrett's esophagus and associated neoplastic lesions. *Am. J. Surg. Pathol.* 32, 1327-35.