CONTENTS

Subjects	Page			
Introduction	1			
Aim of the work	3			
Review of literature	4			
EPH-gestosis	4			
History-Definitions –				
Classification				
Epidemiology and risk factors.				
Pathogenesis of EPH-gestosis.	18			
Homocysteine	35			
Biochemistry & Metabolism.				
Homocysteine with normal pregnancy.				
Hyperhomocysteinemia.	46			
with general diseases.				
With adverse pregnancy outcome.				

• Hyperhomocysteinemia and EPH-gestosis	51
Homocystein in prediction of EPH-gestosis.	55
••••	
Subjects and methods	57
Results	68
Discussions	84
Summary	90
Conclusions & Recommendations	92
Reference	93
Arabic summary	

LIST OF TABLES

Table No.	Table	Page
Table I	Objective + subjective symptoms of EPH gestosis	10
Table II	Risk factor for EPH gestosis and subjects rating at	12
	the weight of supporting evidence	
Table 1	The maternal age of three groups	69
Table 2	The gestational age of three groups.	70
Table 3	The infant birth weight of the three groups.	71
Table 4	Apgar score at one minute of the three groups.	72
Table 5	Apgar score at five minute of the three groups.	73
Table 6	Diastolic blood pressure of the three groups.	74
Table 7	The correlation coefficient (r) between serum	75
	Homocysteine and different clinical variable in	
	patients with toxemia of pregnancy.	
Table 8	Mean + S.D., t test & p values of maternal age, GA,	76
	IBW, Apgar score at 1, 5 minute & DBP in mild pre-	
T 11 0	eclampsia, eclampsia compared with the control group.	77
Table 9	Serum homocystein level of the three groups.	77
Table 10	Correlation coefficient (r) between serum	78
	homocysteine level and severity of hypertension in	
TD 11 11	patients with mild pre-eclampsia and eclampisa.	70
Table 11	Correlation coefficient (r) between serum	79
	homocysteine level and severity of edema in patients	
Table 12	with mild pre-eclampsia and eclampisa.	80
Table 12	Relation between serum homocysteine level and	80
	severity of proteinuria in patients with pre-eclampsia & eclampsia.	
Table 13	Diagnostic validity test	81
Table 14	The serum homocysteine level and lipid profile in	82
Taule 14	mild pre-eclampsia, eclampsia compared with the	04
	control group	
Table 15	Correlation coefficient (r) between serum	83
14010 13	homocysteine level and different lipid profile in	
	patients with toxacima of pregnancy.	
	patients with toracina of pregnancy.	

LIST OF FIGURES

Fig. No.	Fig.				
Fig. 1	Prostaglandin synthesis.				
Fig. 2	Homocysteine, a demethylated derivative of 3 methionine				
Fig. 3	Pathway of Homocysteine synthesis & metabolism.				
Fig. 4	Coenzymes derived from vitamin B_{12} . 3'				
Fig. 5	Methionine synthesis. The methyl group is derived from the 5-methyltetrahydrofolate.				
Fig. 6	Catabolic pathway of Homocysteine. 40				
Fig. 7	The mean age of patients in study group I & study group II and control group.	69			
Fig. 8	The mean gestational age in study group I & study group II and control group.				
Fig. 9	The mean infant birth weight in study group I & 71 study group II and control group.				
Fig. 10	The mean Apgar score at 1 minute in study group I & study group II and control group.				
Fig. 11	The mean Apgar score at 5 minutes in study group I 73 & study group II and control group.				
Fig. 12	Mean diastolic blood pressure in study group I & 74 study group II and control group.				
Fig. 13	Mean serum homocysteine in study group I & study group II and control group.				
Fig. 14	Homocysteine and severity of hypertension in study group I & study group II.	78			
Fig. 15	Homocystein and severity of edema in study group I & study group II.				
Fig. 16	ROC plot for homocysteine to discriminate between patients and control.				

LIST OF ABBREVIATIONS

ACOG	:	American colleague of obstetric and gynecology
AT III	:	Antithrombin III
BL.Pr.	:	Blood pressure
СЕ	:	Cholesterol esterase
CHD	:	Congenital heart diseases
Co_x	:	Cholesterol oxidase
DBP	:	Diastolic blood pressure
EDD	:	Expected date of delivery
G.A.	:	Gestational age
Нсу	:	Homocystiene
IBW	:	Infant birth weight
IL-6	:	Interleukin-6
LCF	:	Lipid clearing factor
LMP	:	Last menstrual period
MTHFR	:	Methylentetrahydrofolate reductase
MTX	:	Methotrexate
PG	:	Prostaglandin
PG1	:	Prostaglandin I
SAH	:	S. adenosyl homocysteine
SAM	:	S. Adenosyl methionine
STBM	:	Syncytiotrophoblast microvellaus membranes
THcy	:	Total homocystiene
TH	:	Tetrahydrofolat
TNF. α	:	Tumour necrosis factor α
TSSHP	:	International society for the study of hypertension of pregnancy
TXA2	:	Thromoxan A2