

REFERENCES

Allain CC. et al., (1974):

Clinical chemistry and cholesterol. Clin. 20:470

Anderson, A.; Hutberg, B.; Brrattstrom, L. and Lsaksson, A. (1992):

Decreased serum Homocysteine in pregnancy., Eur J. Clin Chem Clin Biochem; 30 : 377-379.

Arngrimsson, R.; Hayward, C.; Nadaud, S.; Roberts, J.M. and Lifton, R. (1997):

Evidence for a familial pregnancy-induced hypertension locus in the eNOS-gene region., Am J. Hum Genet; 61 : 354–362.

Arngrimsson R.; Purandare, S.; Connor, M.; Fisher, S.J. and Ramon, J. (1993) :

Angiotensinogen, a candidate gene involved in preeclampsia., Nat Genet; 4 : 114 – 115.

Aubard, Y.; Darodes, N. and Cantaloube, M. (2000) :

Hyperhomocysteinemia and Pregnancy-review of our present understanding and therapeutic implications.
Eur. J. Obstet. Gynecol. Reprod. Biol. 93 (2) : 157-165.

Bernhart, F. (1939) :

Weinklin Wochenschr 52 : 1009-1013, quated from Obstet, Gynecol; 43 (4) : 599 April 1974.

Bernstein, I.M.; Ziegler, W. and Stirewalt, W.S. et al. (1998) :

Angiotensinogen genotype and plasma volume in nulligravida women., The coll of Obstet. Gynecol; 92 : 171 – 173.

Boston A.G.; and Lathrop L. (1997):

Homocysteinemia in end stage renal disease: Prevalence, etiology, and potential relationship to arteriosclerotic outcomes. Kideny Int. 52: 10-20.

Brattstrom, L.E.; Hultberg, B.L. and Hardebo, J.E. (1985) :

Folic acid responsive postmenopausal Homocysteinemia., Metabolism; 34 : 1073 – 1083.

- Brower, D.A.; Welten, H.T.; Reijngoud, D.I. et al. (1998) :**
Plasma folic acid cutoff value, derived from its relationship with Homocysteine., Clin Chem; 44 : 1545 – 1550.
- Campbell, W.B. and Halushka, P.V. (1996) :**
Lipid-derived autotoxins. In: Goodman & Gilman's pharmacological basis of therapeutics, 4th ed., By Hardman JG and Limbird LG, Pub Mc Grow Hill, New York P. 601-616.
- Catargi, B.; Parrot-roulaud, F.; Cochet C.; Ducassou, D.; Roger, P. and Tabarin, A. (1999) :**
Homocysteine, hypothyroidism, and effect of thyroid hormone replacement., Thyroid; 9 (12) : 1163 – 1166.
- Castro, L.C.; Hobel C.J. Gornbein J. (1994):** Plasma levels of arterial natriuretic peptide in normal and hypertensive pregnancies: Ametaq analysis., Am. J. Obstet Gynecol; 171 (6): 1642-1651.
- Cattaneo, M.; Vecchi, M.; Zighetti, M.L. et al. (1998) :**
High prevalence of hyperhomocysteinemia in patients with inflammatory bowel disease : a pathogenic link with thromboembolic complications., Thromb Haemost, 80 : 542 – 545.
- Chait, A.; Malinow, M.R.; Nevin, D.N.; Morris, C.D.; Eastgard, R.L.; Krisetherton, P.; Pi-Sunyer, F.X.; Oparil, S.; Resnick, L.M. et al. (1999):**
Increased dietary micronutrients decrease serum Homocysteine concentrations in patients at high risk of cardiovascular disease., Am J Clin Nutr; 70 (5) : 881-887.
- Chauveau, P.; Chadeaux, B.; Coude, M.; Aupetit, J.; Hannedouche, T.; Kamoun, P. et al. (1993) :**
Hyperhomocysteinemia, a risk factor for atherosclerosis in chronic uremic patients., Kidney Int Suppl; 41 : S 72-77.
- Chen, H.; Huang, Y.; Liu, G. and Ma, T. (1996) :**
The immuno-logical damage of placenta and its types in pregnancy-induced hypertension., Chung Hua Fu Chan Ko Tsa Chih; 31 : 540 – 548 (English abstract).

Chesley, L.C. (1974) :

A short history of eclampsia., *Obstet Gynecol*; 43 : 599.

Clewell, H.W. (1997) :

Hypertensive emergencies in pregnancy, obstetrics intensive care: A practical Manual 1st edition, edited by Foley, M.R. and Strong, T.H. USA : W.B. Saunders Company.

Cooper, B.W.; Brennecke, S.P. and Wilson, A.N. (1993) :

Fetuses of preeclampsia., *hypertenspregn*; 12 : 1-23.

Cooper, J.C.; Shryke, A.M.; Baker, P.N. and Milton, P.J. (1995) :

The pathogenesis of preeclampsia : Current theories. In : *Progress in reproductive medicine*, volume II by Asch R and Studd J, Pub., The Parthenon Pub Group, New York, London, P. 165 – 174.

Cotter, A.M.; Molloy, A.M.; Scott, J.M. and Daly, S.F. (2001) :

Elevated plasma homocysteine in early pregnancy : a risk factor for the development of severe preeclampsia.
Am. J. Obstet. Gynecol. Oct; 185 (4) : 781-785.

Cotter, A.M.; Molloy, A.M.; Scott, J.M. and Daly, S.F. (2003) :

Elevated plasma homocysteine in early pregnancy: a risk factor for the development of nonsevere pre-eclampsia.
Am. J. Obstet. Gynecol. Oct; 185 (4) : 781-785.

Cstro, L.C.; Hobel, C.J. and Gornbein, J. (1994) :

Plasma levels of arterial natriuretic peptide in normal and hypertensive pregnancies : A meta analysis., *Am J. Obstet Gynecol*; 171 (6) : 1642 – 1651.

Cunningham, F.G.; Mac-Donald, P.C.; Gan, N.F.; Leveno, K.J.; Gilstrap, L.C.; Hankins, G.F. and Clark, S.L. (1997):

Hypertensive disorders of pregnancy. In : *William's Obstetrics*, 20th edition, London : Prentice-Hall Int. Inc., PP. 693-744.

Davey, D.A. (1995):

Hypertensive disorders of pregnancy. In: *Dewhurst's textbook of Obstetrics and Gynecology for post-graduate*, edited by Whitfield CR, 5th edition, Blackwell Science, Oxford, P. 175 – 178.

D'Anna R. Baviera, G., Corrado F. Ientile, R. Granese, D. Stella N.C. (2004):

Plasma homocysteine in early and late pregnancies complicated with pre-eclampsia and isolated intrauterine growth restriction. Acta. Obstet Gynecol. Scan. 83 (2): 155-158.

Dekker, G.A.; De Vries, J.I.P.; Doelitzch, P.M.; Huijgens P.C.; Von Blomerg, B.M.E. Jakobs, c. et al., (1995) :

Underlying disorders associated with sever early onset preeclampsia., Am. J. Obstet Gynecol; 173 : 1042-1048.

Dekker, G.A. and Sibai, B.M. (1991):

Early detection of preeclampsia., Am. J. Obstet Gynecol; 165 : 160 – 172.

Delee, J.B. (1913) :

The principles and practice of Obstet Philadelphia WB Saunders : 342-72., Am. J. Obstet Gynecol; 156 : 1227-1233, 1986.

Delee, J.B. and Greenhill, J.P. (1974) :

Principles and practice of Obstet philadelphia : W.B. Sanders : 328.

Dierkes, J.; Kroesen, M. and Pietrzik (1998) :

Folic acid and vitamin B6 supplementation and plasma homocysteine concentrations in healthy young women., Int. J. Vitam Nutr Res; 68 : 98-103.

Dryer R.L. Teitz N.W. and Saunders W.B. (1970):

In fundaments of clinical chemistry (Phladelphia, P329).

El-Khairi L., Vollset SE., Resfum, H. Ueland, P.M. (2003):

Plasma total homocysteine, pregnancy complications, and pregnancy outcomes: the Hordaland Homocysteine study. Am. J. Clin. Nutr. 77 (2):467-472.

Eskenazi, B.; Fenster, L. and Sidney, S. (1991) :

A multivariate analysis of risk factors for preeclampsia., JAMA; 266 : 237-241.

Fallis, N.E. and Langford, H.G. (1963) :

Relation of second trimester blood pressure to toxemia of pregnancy in the primigravida patient., Am. J. Obstet Gynecol; 87 : 123-125.

Fassbender, K.; Mielke, O.; Bersch, T.; Nafe, B.; Froschen, S. and Hennevi, M. (1999) :

Homocysteine in cerebral microangiopathy., Lancet; 353 : 1586-1587.

Finkelstein, J.D. (1990) :

Methionine metabolism in mammals., J. Nutr Biochem; 1 : 228-237.

Finley, P.R., Warnick, GP. And Albest N. (1983):

Selected Methods Clin. Chem. 10: 91-99.

Finnell, R.H.; Greer, K.A.; Barber, R.C.; Piedrahita, J.A.; Shaw, G.M. and Lammer, E.J. (1998) :

Neural tube and craniofacial defects with special emphasis on Folate pathway gene., Crit Rev Oral Biol. Med; 9 : 38-53.

Folsom, A.R.; Nieto, J.; Mc Govern P.G. et al., (1998) :

Prospective study of coronary heart disease incidence in relation to fasting total Homocysteine, related genetic polymorphism, and vitamins., Circulation; 98 : 204 – 210.

Francois Mouricean (1992) :

Hypertension In high-Risk pregnancy. Ed. Calded AA and Danlop W 98, Red Wood Press.

Frantzen, F.; Faaren, A.L.; Alfheim, I. and Nordhei, A.K. (1998) :

An enzyme conversion immunoassay determining total homocysteine in plasma or serum.
Clin Chem, 44 : 311.

Frausher, G.; Kamau Khova, E.; Muehl, Hoger, H. and Lubec, B. (1995) :

Oral administration of Homocysteine leads to increased plasma triglycerides and homocysteic acid: additional mechanisms in Homocysteine induced endothelial damage., Life Sci; 57 : 813 –817.

Friedman, E.A. and Neff, R.K. (1997) :

Pregnancy outcome as related to hypertension, edema and proteinuria in hypertension in pregnancy. Eds. Lindheim, M.D.K. at3, A.I. and Zuspan E.P., New York.

Garland, D. and Audrson, (1994) :

Pregnancy induced hypertension manual of clinical problems in Obstet and Gynecol Ed. Michel E.R., Rick W.M., Pub. Lihle Brown and company New York (4th). 33-42.

Garner, P.R.; D'Alton, M.E.; Dudley, D.K. et al. (1990) :

Preeclampsia in diabetic pregnancies., Am. J. Obstet Gynecol; 63 : 505-508.

Ghanem, A.; Nezar, M.; Abd El-Fatah O.; Ghanem, H. and Abd El-Hassieb, M.; (2003):

Second trimester plasma homocysteine assay in prediction of pre-eclampsia. Benha. M.J. Vol. 20. No. 1. 85-96.

Giles, W.H.; Kittner, S.J.; Croft, J.B. and Anda, R.F. (1998) :

Serum folate and risk of coronary heart disease: results from cohort of US adults.
Ann Epidemoil, 8 : 490.

Girling, J. and De-Swiet-M (1998) :

Inherited thrombophilia and pregnancy., Curr-opin-Obstet Gynecol; 10 (2) : 135-144.

Goddijn-Wessel, T.A.W.; Wouters, M.G.A.J. vdMolen, E.F.; Spuijbroek, M.D.E.H.; Stegers-Theunissen, R.P.M., Blom, H.J. et al. (1996) :

Hyperhomocysteinemia a risk factor for placental abruption or infarction., Eur J Obstet Gynecol Reprod Biol; 66 : 23-29.

Graham, I.M.; Daly, L.E.; Refsum, H.M.; Robinson, K.; Brattstrom, L.G.; Ueland, P.M.; Plama-Reis, R.J. et al., (1997) :

Plasma Homocysteine as a risk factor for vascular disease., The European concerted action project. JAMA; 277-281.

- Grandone, E.; Margalione, M.; Colaizzo, D.; Peters, J., Cameron, A. and Beltz, W. (1997) :**
Factor V Leiden C > MTHFR polymorphism and genetic susceptibility to preeclampsia., *Thromb Haemost*; 77 : 1052 – 1060.
- Green, R. and Miller, J.W. (1999) :**
Folate deficiency beyond megaloblastic anemia: Hyperhomocysteinemia and other manifestations of dysfunctional folate status., *Semin Hematol*; 36; 47-64.
- Guttormsen, A.B.; Mansoor, A.M.; Fiskerstrand, T. et al. (1993) :**
Kinetics of plasma Homocysteine in healthy subjects after pleural Homocysteine loading., *Clin Chem*; 39 : 1390 –1407.
- Guttormsen, A.B.; Svarstad, E.; Ueland, P.M. and Resfsum, H. (1995) :**
Elimination of homocysteine from plasma in subjects with end stage renal failure.
Irish J. Med. Sci, 164 (Suppl. 15) : 18.
- Guttormsen, A.B.; Ueland, P.M.; Svarstad, E. and Resfsum, H. (1997) :**
Kinetic basis of hyperhomocysteinemia in patients with chronic renal failure., *Kidney Int*; 52 : 495-502.
- Haagsma, C.J.; Blom, H.J.; Van Riel, Van't Hof, M.A.; Giesendorf, B.A. (1999):**
Influence of sulphasalazine, methotrexate, and the combination of both on plasma Homocysteine concentrations in patients with rheumatoid arthritis., *Ann Rheum Dis*.; 58 (2) : 79-84.
- Halligan, A.; Bonner, J.; Sheppard, B. et al. (1994) :**
Haemostatic, fibrinolytic and endothelial variables in normal pregnancy and preeclampsia., *Br. J. Obstet Gynecol*; 101 : (6) : 488 – 492.
- Hansen, J.P. (1986) :**
Older maternal age and pregnancy outcome : A Review of the literature., *obstet Gynecol Surv*; 41 : 726-728.
- Herrmann, W. (2001) :**
The importance of hyperhomocysteinemia as a risk factor for diseases. *Clin-Chem-Lab-Med. Aug*; 39)8) : 666-674.
-
-

- Hernanz, A.; Plaza, A.; Martin-Mola, E. et al., (1999) :**
Increased plasma levels of Homocysteine and other thiol compounds in rheumatoid arthritis women., Clin Biochem; 32 (1) : 65-70.
- Hietala, R.; Turpeinen U. and Laatikainen T. (2001):**
Serum Homocysteine at 16 weeks and subsequent preeclampsia: Obstet gynecol; 97 (4): 527-527.
- Hirano, K.; Ogiharat, Miki, M. et al., (1994) :**
Homocysteine induces iron catalyzed lipid peroxidation of low density lipoprotein that is prevented by alpha-Tocopherol., Free Radic Res; 21: 267-276.
- Hubel, G.A.; Mclaughlin, M.K.; Evans, R.W.; Hauth, B.A.; Sims, G.J.; Ronerts, J.M. et al. (1995) :**
Fasting serum Triglycerides, Free fatty acids, and Melanodialdehyde are increased in preeclampsia are positively correlated and decrease within 48 hr. post partum., Am J. Obstet Gynecol; 174 : 975-982.
- Ivan, J. Perry (1994):**
The definition of preeclampsia., Br. J. Obstet Gynecol, 101 : 587-591.
- Jancin, B. (1994) :**
Amino acid defect causes 20% of atherosclerosis in CHD., Family Pract News; 15 : 17.
- Kilby, M.D.; Pipkin, F.B. and Symonds, E.M. (1994) :**
Pharmacological prophylaxis in pregnancy induced hypertension. Progress in obstetrics and gynecology edited by John Studd : Vol. 11 : 53 – 74.
- Klonoff-Cohen, H.; Edelstein, S.; Savitz, D. (1993) :**
Cigarette smoking and preeclampsia., Obstet Gynecol; 81 : 541 – 544.

Klonoff-Cohen, H.S.; Savitz, D.A.; Cefalo, R.C. and McCann, M.R. (1989):

An epidemiology study of contraception of preeclampsia., JAMA; 262 : 3143 – 3147.

Lachmeijer, A.M.; Arngrimsson, R.; Bastiaans, E.J.; Pals, G.; TEN-Kate, L.P.; de-Vries, J.L.; Kostense, P.J.; Aarnoudse, J.G. and Dekker, G.A. (2001) :

Mutations in the gene for methylenetetrahydrofolate reductase, homocysteine levels, and vitamin status in women with a history of preeclampsia.

Am. J. Obstet. Gynecol. 2001 Feb. 184 – 202.

Laivuori, H.; Kaaja, R.; Turpeinen, U.; Viinikka, L. and Ylikorkala, O. (1999) :

Plasma Homocysteine levels elevated and inversely related to insulin sensitivity in preeclampsia., Obstet Gynecol; 93 (4) : 489 – 493.

Lewinsky, R.M. and Riskin, M.S. (1998) :

Autonomic imbalance in preeclampsia : evidence for increased sympathetic tone to the supine pressor test., Obstet Gynecol, 91 : 935 – 939.

Levine, Richard, J.; England, Lucinda, J. and Sibai, Baha, M. (2002) :

Elevated plasma homocysteine in early pregnancy : a risk factor for the development of severe preeclampsia.

Am-J. Obstet. Gynecol. May; 186 (5) : 1107; discussion 1108.

Lobo, A.; Naso, A.; Arheart, K. et al. (1999) :

Reduction of Homocysteine levels in coronary artery disease by low-dose Folic acid combined with vitamins B6 and B12. AM. J. Cardiol; 83 : 821-825.

Lopez-Quesada, E., Vilaseca M.A., Artuch, R., Gomez, E. Lailla J.M. (2003):

Homocysteine and other plasma amino acids in pre-eclampsia and in pregnancies without complications Clin Biochem. 36 (3): 185-192.

- Lorentzen, B.; Gndersen, M.J.; Clausen, T.; Henriksent, (1994) :**
Fasting serum fatty acids and triglycerides are increased before 20 wks of gestation in women who later develop pre-eclampsia., *Hypertens pregnancy*; 13 : 103-109.
- Loscalzo, J. (1996) :**
The oxidant stress of hyperhomocysteinemia., *J. Clin-Invest* : 98: 5-7.
- Mabie, W.C.; Pernoll, M.L. and Biswas, M.K. (1986) :**
Chronic hypertension in pregnacny., *Obstet Gynecol*; 67 : 197-205.
- Mac Gillivary (1961):**
Hypertension in pregnancy and its consequences., *J. Obstet Gynecol of the British common wealth*; 68 : 557 – 569.
- Mac Gillivary, J. (1983):**
Preeclampsia: The hypertensive disease of pregnancy. London : W.B. Saunders.
- Mahdy, E.; Abd, El-Kareem, A.; El-Hodaiby, K.H.; El-Maraghy, Khalifa, A. (1996) :**
Tumor necrosis alpha (TNF- α) in pregnancy induced hypertension., *The Egyptian Journal of Biochemistry*; 14 (1) : 109 – 114.
- Malinow, M.R.; Rajkovic, A.; Duell, P.B. Hess, D.L. and Upson, B.M. (1998) :**
The relationship between maternal and neonatal umbilical cord plasma homocysteine suggests a potential role for maternal homocysteine in fetal metabolism., *Am. J. Obstet Gynecol*; 178-180.
- Mandel, H.; Brenner, B.; Bernant, M.; Rosenberg, N.; Lanir, N.; Jacobs, C. (1999) :**
Coexistence of hereditary homocysteinuria and factor V leiden effect on thrombosis., *N. Engl. J. Med.*; 334 : 763-768.

- Marcoux, S.; Brisson, J. and Fabia, J. (1989) :**
The effect of cigarette smoking on the risk of preeclampsia and gestational hypertension., *Am J. Epidemiol*;130 : 950-957.
- Marnoni, E.; Menticoglou, S.; Harman, C.R. et al. (1994) :**
Cell mediated immunity imbalance in PIH., *Obstet Gynecol Invest.*; 83 : 236-240.
- Mayer, F.L.; Jacobson, D.W. and Robinson, K. (1996):**
Homocysteine and coronary atherosclerosis., *J. Am. Coll Cardiol*; 27 : 517-527.
- Mc-Cully, K.S. (1999):**
Vascular pathology of homocysteinemia : implication of pathogenesis of arteriosclerosis., *Am-J. Pathol*; 56 : 111-128.
- Mc Keever, M.P.; Weir, D.G.; Molloy, A. and Scott, J.M. (1991) :**
Betaine-homocysteine methyltransferase organ distribution in man, pig and rat, and subcellular distribution in the rat., *Clin. Sci*; 81 : 551-561.
- Medical Letter (1997) :**
Betaine for homocysteinuria., *The medical latter*; 39 (993) : 12.
- Meleady, R. and Graham, I. (1999):**
Plasma Homocysteine as a cardio-vascular risk factor : causal, consequential or of no consequence?., *Nutr Rev*; 57 (10): 299-305.
- Miller, J.W.; Nadeau, M.R.; Smith, D.; Selhub, J. (1994) :**
Vitamin B6 deficiency & Folate deficiency : Comparison of responses to methionine loading in rats., *Am, J. Clin. Nutr.*; 59 : 1033-1039.
- Miller, J.W.; Ribaya-Mercado, J.D.; Russell, R.M.; Shepard D.C.; Morrow, F.D. et al. (1992) :**
Total homocysteine in fasting plasma is not a good indicator of b6 deficiency., *Am J. Clin. Nut*; 55 : 1154-1160.
- Mills, J.L.; McParlin, J.M.; Kikke, P.N. et al. (1995) :**
Homocysteine metabolism in pregnancies complciated by neural-tube defects., *Lancet*; 345 : 149-151.

- Morgan, L.; Baker, P.; Broughton, F.; Pipkin, F.; Kalsheker, N. (1995):**
Preeclampsia and the angiotensinogen, Br. J. Obstet. Gynecol; 102 : 489-493.
- Morita, H.; Nappo F. and Olnegy (2000):**
Gentic poleomorphysim of 5.10- MTHFR as risk factor for coronary heart diseases circulated, 95: 2032-2036.
- Morris, N.H. Eaton, B.M. and Dekker, G. (1996) :**
Nitric oxide the endothelium, pregnancy and preeclampsia., Br. J. Obstet Gynecol; Jan. (103) : 4-15.
- Moutquin, J.; Rainvillec, Giroux, L. et al., (1985) :**
A prospective study of B1. Pr. In pregnancy : Prediction of preeclampsia., Am J. Obstet Gynecol; Jan. (103) : 4-15.
- Mudd, S.H.; Levy, H.L. and Skovby, F. (1989) :**
Disorders of transsulfuration. In : Scriver CR, Beaudet AL, Sly WS et al. editors. The metabolic basis of inherited disease, New York: McGraw-Hill, 693-700.
- Muller, T.; Werne, B.; Fowler, B. and Kuhnw, (1999) :**
Nigral endothelial dysfunction, homocysteine, and Parkinson's disease., Lancet; 354 (9173) : 126 –127.
- Nabhan, A.; Habib, S. and Tamara, T. (2000) :**
Plasma homocysteine concentration in preeclampsia., the new Egyptian journal of medicine; 22 (5) : 213-217.
- Nelen, W.L., (2002):**
Hyperhomocysteinemia and human reproduction., clin. Chem. Lab. Med. 39 (8): 758-763.
- Noto, R., Neri S., Noto, Z. Cilio, D. Abate, G. Noto P. Pepi F., Leanza A. Molino, G. (2003):**
Hyperhomocysteinemia in pre-eclampsia is associated to higher risk pressure profiles. Eur. Rev. Med. Pharmacol. 7 (3): 81-87.

- Nygard, O.; Vollset, S.E.; Refsum, H. and Ueland, P.M. (1998) :**
Major life style determinants of Plasma total Homocysteine distribution : The hordaland Homocysteine study., Am J. Clin Nutr; 67 : 263-270.
- Nygard, O.; Vollset, S.E.; Refsum, H.; Ueland, P.M. Nordrehang, J.E. and Farstad, M. (1997) :**
Plasma Homocysteine levels and mortality in patients with coronary artery disease., N Engl J. Med.; 337 : 230-236.
- Paul Zweifel (1916) :**
Quoted from William obstetrics 14th Ed. New York : Appleton-century-Crafts : 685 – 689.
- Parson, R.B.; Waring, R.H.; Ramsden, D.B.; Williams, A.C. (1998) :**
In vitro effect of the cysteine metabolites homocysteic acid, homocysteine and cysteic acid up on human neuronal cell lines., Neurotoxicology; 19 (4-5) : 599-603.
- Patrick, T.E. Powers R.W., Daftary, A.R., Ness, R.B. Roberts J.M. (2004):**
Homocysteine and folic acid are inversely related in black women with pre-eclampsia. Hypertension 43 (6): 1279-1282.
- Perales-Davila, J.; Martinez de-villarreal, L.E.; Triana-Saldana, H.; Saldivar-Rodrigues, D.; Valez-Leal, R.; zacarias-Villarreal-Perez, J. (2001):**
Folic acid levels, homocystein and polymorphism of methylenetetrahydrofolate reductase enzyme (MTHFR) in patients with pre-eclampsia and eclampsia. Ginecol. Obstet. Mex, 123: 696-711.
- Powers, R.W.; Evans, R.W.; Majors, A.K. et al. (1998) :**
Plasma Homocysteine concentration is increased in preeclampsia and is associated with evidence of endothelial activation, Am. J. Obstet Gynecol; 179 : 1605 – 1611.
- Powers, R.W.; Evans, R.W.; Ness, R.B.; Crombleholme, W.R. and Roberts, J.M. (2001) :**
Homocysteine and cellular fibronectin are increased in preeclampsia, not transient hypertension of pregnancy. Hypertens-Pregnancy. 20 (1) : 69-77.

- Prasmusinto, D.; Skrablin, S.; Hofstaetter, C.; Fimmers, R. and van-der-Ven, K. (2002) :**
The methylenetetrahydrofolate reductase 677 C--> T polymorphism and preeclampsia in two populations. Obstet-Gynecol. Jun; 99 (6): 1085-1092.
- Raijmakers, M.T.; Zusterzeel, P.L.; Steegers, E.A. and Peters, W.H. (2001) :**
Hyperhomocysteinaemia : a risk factor for preeclampsia? Eur. J. Obstet. Gynecol. Reprod. Biol. Apr; 95 (2): 226-228.
- Rajkovic, A.; Catalano, P.M. and Malinow, M.R. (1997) :**
Elevated Homocysteine levels with preeclampsia., Obstet Gynecol; 20 : 168-1671.
- Rajkovic, M.T.; Zusterzeel, P.L.; Roes, E.M.; Steegers, E.A.; Mulder, T.P. and Peters, W.H. (2001) :**
Oxidized and free whole blood thiols in preeclampsia. Obstet-Gynecol. Feb; 97 (2) : 272-276.
- Ray, J.G. and Laskin, C.A. (1999) :**
Folic acid and Homocysteine metabolic defects and the risk of placental abruption, preeclampsia and spontanous pregnancy loss : A systematic review., placenta; 20 (7) : 519-529.
- Redman, C.W. (2000) :**
Preeclampsia in : PM Shaighn O'brien (ed) The year book of obstetrics and gynecology 8th edition., London : RCOG Press, Henry Ling Limited, Dorchester Dorset, PP. 97.
- Refsum, H.; Ueland, P.M.; Nygard, O. and Vollset S.E. (1998):**
Homocysteine ans cardiovascular disease., Annu Rrv Med.; 49 : 31-62.
- Regland, B.; Andersson, M.; Abrahamsson, L.; Bagby, J.; Drehaag, L.G. and Gottfries, C.G. (1997):**
Increased concentrations of Homocysteine in the cerebrospinal fluid in patients with fibromyalgia and chronic fatigue syndrome., Scand, J. Rheumatol; 26 : 301-307.

Rey, E. and Couturier, A. (1994):

The prognosis of pregnancy in women with chronic hypertension., Am. J. Obstet Gynecol; 171 : 410 – 416.

Rippmaun, E.T. (1969) :

Preeclampsia., progress in Obstet Gynecol; 5 : 89-105.

Rizk, M.; Abdel-Hadi, M.; Al-Dessouky, M. and Abdel-Sattar, A. (2001):

Hyperhomocysteinemia at 16th week gestation and subsequent preeclampsia. Benha. M.J. Vol. 18. No. 2. 35-44.

Robert, W.; Power, Rhobert, W.; Evans, Alana, K.; Majors et al. (1998):

Plasma Homocysteine concentration is increased in preeclampsia and is associated with evidence of endothelial activation., Am. J. Obstet gynecol; 179 : 1605 – 1611.

Robert, J. (1998 a):

Possible ditary Measures in the prevention of preeclampsia and eclampsia. Clin. Obstet Gynacol. 15 :2-4.

Roberts, J.M. (1998 b) :

Endothelial dysfunction in preeclampsia., Semin Reprod Endocrinol; 16 : 5-15.

Roberts, J.M. and Redman, C.W.G. (1993) :

Preeclampsia : More than pregnancy induced hypertension., Lancet; 341 : 1451-1454.

Roberts, J.M. and Taylor, R.N. Goldfien, A. (1991) :

A clinical and biochemical evidence of endothelial cell dysfunction in the pregnancy syndrome pre-eclampsia., Am. J. Hyprtens; 4 : 700-708.

Robertson, E.G. (1971):

The natural history of edema during prengancy., J. Obstet Gynecol. Br. Commonw; 78 : 520-529.

Robillared, P.Y.; Dekker, G.A. and Hysley, T.C. (1998) :

Primipaternities in families : Is the incidence of pregnancy-induced hypertensive disorders in multigravidas an anthropological marker of reproduction? Aust NZJ., Obstet Gynecol, 38 (3) : 284-287.

Robillard, P.Y.; Hulse, T.C. Perionin, J. et al. (1994) :

Association of pregnancy induced hypertension with duration of sexual cohabitation before conception., Lancet; 344 : 973-975.

Robson, S.C. (1999) :

Hypertension in Pregnancy. In : Edmonds, DK (ed) Dewhurst's Text book of Obstetrics and Gynecology for postgraduates 6th edition., London : Blackwell Science, pp. 166 – 185.

Roeschlau P. et al., (1974):

Cholesterol importance assay; Klin Chem. Biochem. 12:226

Sargent, I.L. (1998) :

Maternal and fetal immune responses during pregnancy. Exp. Clin in immunoglobulins., 10 : 85-102.

Scoll, T.O. and Johnson, W.G. (2000) :

Folic acid influence on the outcome of pregnancy., Am. J. Clin. Nutr; 7 : 1295 -1303.

Scott, J.R. (1994):

Hypertensive disorders of pregnancy. In : Danforth's obstet & Gynecol; 7th ed., by Scott JR, Disaia PJ, Hammond CB and Spellacy WN, Pub. Lippincott Co., Philadelphia, P. 351 – 365.

Selhub, J. (1999) :

Homocysteine metabolism., Annu Rev Nutr; 19 : 217-246.

Selhub, J. and Miller, J.W. (1991) :

The pathogenesis of hyperhomocysteinemia : interruption of the coordinate regulation by S-adenosylmethionine of the remethylation and transulfuration of homocysteine., Am. J. Clin Nutr; 55 : 131-138.

Sheppard, B.L. and Bonnar, J. (1981) :

An ultrastructural study of utero-placental spiral arteries in hypertensive and normotensive pregnancy and fetal growth retardation., Br. J. Obstet Gynaecol; 88 : 695-705.

Shobel, H.P.; Fischer, T. Heuszerk, et al. (1996) :

Preeclampsia : a state of sympathetic over activity. N. Engl. J. Med., 335 : 1480 –1485.

Sibia, B.M. (1991) :

Immunologic aspect in preeclampsia., clin Obstet gynecol; 34 (1) : 27 –34.

Sibia, B.M.; Gordon, T.; Thom, E.; Caritis, S.N.; Klebanoff, M.; McNellis, D. and Paul, R.H. (1995) :

National institute of child-health and human development network of maternal-fetal medicine unites : Risk factors for preeclampsia in healthy nulliparous women : A prospective multicenter study., Am. J. Obstet Gynecol; 172: 53.

Sibia, B.M.; (1998) :

Prevention of pre-eclampsia: a big disappointment. Am. J. Obstet Gynecol; 179:1275-1278.

Sibia, B.M.; Lindheimer, M.; Hauth, J. et al. (1998) :

Risk factors for preeclampsia, abruptio placentae, and adverse neonatal outcomes among women with chronic hypertension., N. Engl. J. Med., 339 : 667-671.

Siddiqi, T.; Rosenn, B.; Mimouni, F. et al. (1991) :

Hypertension during pregnancy in insulin-dependant diabetic women., Obstet Gynecol; 77 : 514-519.

Silver, P.K.; Turbo, J.M.; Barid, J.A. et al. (1990) :

Soluble factors produced by isolated 1st trimester chorionic villi directly inhibit the immune system., Am. J. Obstet Gynecol; 163 : 84-88.

Smith, G.N.; Walker, M.; Tessier, J.L. et al. (1997) :

Increased incidence of preeclampsia in women conceiving by intrauterine insemination with donor versus partner sperm for treatment of primary infertility., Am J. Obstet Gynecol; 177 : 455 – 458.

- Snowdon, D.A. Tully, C.L.; Smith, C.D.; Perez Riley, K.; Markesbery, W.R. (2000) :**
Serum folate and the severity of atrophy of the neocortex in Alzheimer disease : findings from the Nun study., *Am Clin Nutr*; 71: 993.
- Sohda, S.; Arinami, T.; Hamada, H.; Yamada, N.; Hamaguchi, H. and Kubo, T. (1997) :**
Methylenetetrahydrofolate reductase polymorphism and preeclampsia., *J. Med. Genet*; 34 : 525-526.
- Stmaler, J.S.; Osborne, J.A.; Jaraki, O. et al. (1993) :**
Adverse vascular effects of Homocysteine are modulated by endothelium derived relaxing factor and related oxides of nitrogen., *J. Clin Invest*; 91 : 308 –318.
- Stampfer, M.J.; Malinow, R.; Willet, W.C. et al. (1992) :**
A prospective study of plasma Homocysteine and risk of myocardial infarction in US physicians., *JAMA*; 268 : 877-881.
- Steegers-Theunissen, R.P.; Boers G.H.; Blom, H.J.; Trijbelle, F.J.; Eskes, T.K. (1992) :**
Hyperhomocysteinemia and recurrent spontaneous abortion or abruptio placentage., *Lancet*; 339 : 1122-1123.
- Steegers-Theunissen, R.P.M.; Boers, G.H.J.; Trijbels, J.E.M. et al. (1994) :**
Maternal Hyperhomocysteinemia : a risk factor for neural tube defects., *Metabolism*; 43 : 1475-1480.
- Steegers-Theunissen, R.P.; Steegers, E.A. Thomas, C.M.; Hollanders, H.M. et al. (1993) :**
Study on the presence of Homocysteine in ovarian follicular fluid., *Fertil Steril*; 60 (6) : 1006-1010.
- Stein, J.H. and McBride, P.C. (1998) :**
Hyperhomocysteinemia and atherosclerotic vascular disease: Pathophysiology, screening, and treatment., *Arch intern Med*; 158 : 1301 –1306.

Stolzenberg-Solomon, R.Z.; Miller, E.R. 3rd, Maguire, M.G. et al. (1999) :

Association of dietary protein intake and coffee consumption with serum Homocysteine concentrations in an older population., *Am. J. Clin. Nutr*; 69 : 467-475.

Stone, J.L.; Lockwood, C.J.; Berkowitz, G.S. et al. (1994) :

Risk factors for sever preeclampsia., *Obstet Gynecol*; 83 : 357-361.

Symonds, E.M. (1995) :

Hypertension in pregnancy; *Arach. Did., Child-Fetal-Neonatal-Ed.*, 72 (2) : 139-144.

Tillman, A.J.B. (1955) :

The effect of normal and toxemic pregnancy on blood pressure., *Am. J. Obstet Gynecol*; 70 : 589-648.

Trudinger, B.J.; Cook, C.M. (1990) :

Doppler umbilical and uterine plow waveforms in severe pregnancy hypertension.
BJOG; 97 : 142-148.

Ueland, P.; Refsum, H.; Stabler, S.; Malinow, R.; Andersson, A. and Allen, R. (1993) :

Total Homocysteine in plasma or serum : Methods and clinical applications., *Clin Chem*; 39 : 1764-1779.

Upchurch, G.R.; Welch, G.N.; Fabian, A.J. et al. (1997) :

Homocysteine decreases bioavailable nitric oxide by a mechanism involving Glutathionine peroxidase., *J. Bio Chem*; 272 : 17012-17017.

Van de Berg, M.; Franken, D.G.; Boers, G.H.J. et al. (1994) :

Combined vitamin B6 plus folic acid therapy in young patients with arteriosclerosis and hyperhomocysteinemia., *J. Vasc Surg*; 20 : 933-940.

Vanderjagt, D.J., Patel, R.J., El-Nafaty A.U., Melah G.S., Crossey, M.J. Glew R.H. (2004):

High- density lipoprotein and homocysteine levels correlate inversely in pre-eclamptic women. *Acta Obste Gynecol. Scand* 83(6): 536-542.

Verhoef, P.; Kok, F.J.; and Krusysen DA. (1997):

Plasma total homocysteine, B vitamins, and risk of coronary atherosclerosis. *Arterioscler thromb. Vasc Biol.* 17:989.

Villar, M.A. and Sibai, B.M. (1989) :

Clinical significance of elevated mean arterial blood pressure in second trimester threshold increase in systolic or diastolic Bl. Pr. during third trimester., *Am. J. Obste. Gynecol*; 160 : 419-423.

Vollset, S.E.; Refsum, H.; Irgens, L.; Emblem, B.M.; Tverdal, A.; Giessing, H.; Monsen, A. and Ueland, M. (2000) :

Plasma total Homocysteine, pregnancy complications, and adverse pregnancy outcomes., *Am. J. Clin. Nutr*; 71 : 962-968.

Walker, J.J. (1998 a) :

Advances in the management of sever preeclampsia. In : John Bonnar (ed) *Recent advances in obstetrics and gynecology* 20th edition., Edinburgh : Churchill Livingstone, pp. 111.

Walker, J.J. (1998 b) :

Current thoughts on the pathophysiology of preeclampsia/eclampsia. In : *Progress in obstetrics and gynecology*, edited by John Studd, volume 13, Churchill Livingstone, Harcourt Brace and co., Egyptian Edition, P. 177-190.

Wang, J.; Mimuro, S.; Lahoud, R.; Trudinger, B. and Wang, X.L. (1998) :

Elevated levels of lipoprotein (a) in women with preeclampsia., *Am. J. Obstet Gynecol*; 178 : 146- 149.

Wang, J.; Trudinger, B.; Duarte, N.; Wilcken, D. and Wang, X.L. (2000) :

Elevated circulating Homocysteine levels in placental vascular disease and associated preeclampsia., *Br. J. Obstet & Gynecol.*, 107 : 935-938.

- Wang, Y.; Walsh, S.W.; Guo, J. and Zhang, J. (1991) :**
The imbalance between Thromboxane and Prostacyclin in preeclampsia is associated with imbalance between lipid peroxidases and vitamin E in maternal blood., *Am. J. Obstet Gynecol*; 165 : 1695-1700.
- Ward. Jata. A.; Jeunemaitre, X.; Helin, C.; Nilson, L.; Namikawa, C.; Farington, P.F. Ogasawara, M.; Suzumori, K.; Tomoda, S.; Berrebi, S.; Sasaki, M.; Corvol, P.; Lifton, R.P. and Laouel, J.M. (1993) :**
A molecular variant of angiotensinogen associated with preeclampsia (comment)., *Nat Genet*; 4 : 59 – 67.
- Warnick, GR, Benderson, And Alber S.N. (1983):**
Selected Methods., *Clin. Chem.* 10:91-99.
- Weiner, C.P. (1990):**
Evaluation of clotting disorder during pregnancy., *Obstet Gynecol*; 3 : 37-40.
- Welch, G.N. and Loscalzo, J. (1998) :**
Homocysteine and atherothrombosis., *N. Engl. J. Med.*; 338 : 1042-1050.
- Widschwendter, M.; Schrocksnadel, H. and Mortl, M.G. (1999) :**
Preeclampsia : a disorder of placental mitochondria? *Mol Med. Today*; 4 (7) : 286-291.
- William, J.W. (1912) :**
Obstet. Third New York, appleton Century crafts : 512.
- Wolfe, H.M.; Zador, I.; Gross, et al. (1991):**
The clinical utility of body mass index in pregnancy., *Am J. Obstet Gynecol*; 164 : 1306 – 1309.
- Woodworth, S.W.; Lix, Lei, Z.M.; Rao, G.V.; Yussman, M.A.; Spinnato, J.A.; Yokoyama, C.; Tanabe, T. and Ulrich, V. (1994) :**
Eicosanoid biosynthetic enzymes in placental and decidual tissue from preeclamptic pregnancies. Increased expression of thromboxane A2 synthesis gene., *J. Clin Endocrinal Metab*; 78 : 1225-1231.
- Worley, R.I. (1987):**
Pathogenesis of pregnancy induced hypertension., *Clin Obstet Gynecol*; 27 (4) : 821 – 835.
- Wouters, M.G.A.J.; Boers, G.H.J.; Blom, H.J. et al. (1993):**
Hyperhomocysteinemia: a risk factor in women with unexplained recurrent early pregnancy loss., *Fertil Steril*; 60: 820-850.

Zeeman, G.G. and Dekker, G.A. (1992) :

Pathogenesis of preeclampsia : A hypothesis., *Clinic Obstet Gynecol*; 34 (2) : 317-337.

Zeeman G.G., Alexander, J.M., Mcintire, D.D., Devaraj, S Leveno K.J. (2003):

Homocysteine plasma concentration levels for the prediction of preeclampsia in women with chronic hypertension.

Am. J. Obstet. Gynecol. 189 (2): 574-576.

Zhang, J.; Zeisler, J.; Hatch, M.C. and Berkowitz, G. (1997) :

Epidemiology of pregnancy induced hypertension, *Epidemiologic Rev*; 19 (2) : 218-232.