

REFERENCES

(In alphabetical order)

A

***Adair CD, Weeks JW, Barrilleaux S (1998):** Oral or vaginal misoprostol administration for induction of labor: a randomized, double-blind trial. *Obstet Gynecol*; 92:810–813.

***Allaire, A.D.; D'Andrea, N.; Truong, P.; McMahon, M.J. and Lessey, B.A (2001):** Cervical stroma apoptosis in pregnancy. *Obstet Gynecol*. 97: 399-115.

***Allot H.A. and Palmer, C.R (1993):** Sweeping of membranes. A valid procedure in stimulating the onset of labor. *Br. J. Obstet. Gynecol.*, 100: 898.

***American College of Obstetricians and Gynecologists (ACOG 1995a):** Induction of labor. ACOG technical bulletin no. 217. Washington DC.

***American College of Obstetricians and Gynecologists (ACOG 1995b):** Quality, evaluation and improvement in practice: post term pregnancy. Committee on quality assessment criteria set no. 10, August 1995.

***American College of Obstetricians and Gynecologists (ACOG 1997):** Induction and augmentation of labor. ACOG technical bulletin no. 216, Washington, DC, USA.

***American College of Obstetricians and Gynecologists (ACOG 1999):** Induction of labor with misoprostol. ACOG Committee Opinion 228 Washington DC.

***American College of Obstetricians and Gynecologists (ACOG 2000):** Response to Searle's drug warning on misoprostol. ACOG Committee Opinion 248 Washington DC.

***American College of Obstetricians and Gynecologists (ACOG 2003):** New U.S. Food and Drug Administration Labeling on Cytotec (Misoprostol) Use and Pregnancy. ACOG Committee Opinion 283 Obstet. Gynecol. 101: 1049 – 50.

***Arias, F. (1993):** Cervical ripening and induction of labor. In : Practical Guide to High Risk Pregnancy and Delivery, edited by Arias F, 2nd edition St. Louis, Baltimore, Ma, USA; 396 – 400.

***Arias, F. (2000):** Pharmacology of Oxytocin and prostaglandins. Clin Obstet Gynecol. 43: 455.

***Atad, J.; Hallak, M.; Auslender, R.; Zarfati, D. and Abramovici, H. (1996):** A randomized comparison of PGE₂, oxytocin and the double balloon device in inducing labor. Obstet. Gynecol., 87: 223.

B

***Barbosa RM, Arilha M. (1993):** The Brazilian experience with Cytotec. Stud Fam Plann; 24:236–240.

***Bennett BB. (1998):** Uterine rupture during induction of labor at term with intravaginal misoprostol. Obstet Gynecol; 89:832– 833.

***Bond GR, Van Zee A. (1994):** Overdosage of misoprostol in pregnancy. Am J Obstet Gynecol; 171:561–562.

***Bique C, Bugalho A, Bergstrom S. (1999):** Labor induction by vaginal misoprostol in grand multiparous women. Acta Obstet Gynecol Scand; 78:198–201.

***Bugalho A, Daniel A, Faundes A, (2001):** Misoprostol for prevention of postpartum hemorrhage. Int J Gynaecol Obstet; 73:1–6.

***Bergsjö, P.; Schmidt, E. and Pusch, D. (1983):** Differences in the reported frequencies of some obstetrical interventions in Europe. Br. J. Obstet. Gynecol., 90: 628.

***Bernstein, P.; Leyland, N.; Gurland, P. and Gare, D. (1991):** Cervical ripening and labor induction with prostaglandin E2 gel. A placebo controlled study. Am. J. Obstet. Gynecol. 156: 336 – 40.

***Bishop, E.H. (1964):** Pelvic scoring for elective induction of labor. Obstet. Gynecol. 24: 266 – 8.

***Bricker, L. and Luckas, M. (2001):** Amniotomy alone for induction of labor (Cochrane Review). In: Cochrane Library Issue 2. Oxford, Update Software Ltd., Summertown Pavilion, Middle way, Oxford OX2 7LG, UK.

***Bugalho, A.; Bique, C.; Machungo, F. and Faundes, A. (1994):** Induction of labor with intravaginal misoprostol in intrauterine fetal death. Am. J. Obstet. Gynecol. 171: 538 – 41.

***Bygdeman, M.; Gemzell Danielson, K.; Marions, L. and Swahn, M. (2000):** Pregnancy termination; Steroids 65: 801.

C

***Calder, A.A. and Embrey, M.P. (1973):** Prostaglandins and the unfavorable cervix. Lancet 2: 1322.

***Calder, A.A. (1986):** Cervical ripening. In Bygdeman M, Berger GS and Keith LG (eds.): Prostaglandins and their inhibitors in obstetric and gynecology. Lancaster MTP Press P. 145.

***Calder, A.A. and Greer, I.A. (1992):** Cervical physiology and induction of labor. In Bonner J. (ed.): Recent advances in obstetric and gynecology Edinburgh Churchill Livingstone P. 33.

***Casey, M.L. and MacDonald, P.C. (1988a):** Bimolecular process in the initiation of parturition: Decidual activation. Clin Obstet. Gynecol., 31: 533.

***Casey, M.L. and MacDonald, P.D. (1988b):** The role of a fetal maternal paracrine system in the maintenance of pregnancy and the initiation

of parturition. In: Jones (ed.): Fetal and neonatal development. Ithaca Perinatology, p. 521.

***Casey, M.L. and MacDonald, P.C. (1993a):** Human Parturition: Distinction between the initiation of parturition and the onset of labor. In: Ducusy CA (ed.): Seminar in Reproductive Endocrinology. New York Thieme p272.

***Casey, M.L. and MacDonald, P.C. (1993b):** Human Parturition: Prostaglandins and cytokines are sequel of labor. In: Marnex c, Forth, England, Parthenon p.638.

***Carlan SJ, Bouldin S, O'Brien WF. (1997):** Extemporaneous preparation of misoprostol gel for cervical ripening: a randomized trial. Obstet Gynecol; 90:911–915.

***Chong YS, Chua S, Shen L, et al. (2003):** Does the route of administration of misoprostol make a difference? The uterotonic effect and side effects of misoprostol given by different routes after vaginal delivery. Eur J Obstet Gynecol Reprod Biol;. In press.

***Chuck, F.J. and Huffaker, B.J. (1995):** Labor induction with intravaginal misoprostol versus intracervical PGE₂ gel (Prepidil gel): Randomized comparison. Am. J. Obstet. Gynecol., 173: 1137 – 42.

***Collins, O.W. (1990):** Misoprostol discovery, development and clinical application. Med. Res. Rev. 10: 149 – 72.

***Cunningham, P.G.; Leveno, K. and Gant, N.F. (2001a):** A normal labor: Dystocia due to abnormalities of the expulsive forces. In William's Obstetrics 21st edition, McGraw – Hill p. 763 – 817.

***Cunningham, F.G.; Leveno, K. and Gant, N.F. (2001b):** Abnormal labor: Hypertensive disorders in pregnancy. In William's Obstetrics 21st edition, McGraw – Hill, p. 763 – 817.

***Cunningham, F.G.; Leveno, K. and Gant, N.F. (2001c):** Parturition. In William's Obstetrics 21st edition, McGraw – Hill p. 262 – 87.

D

***Danforth, D.N. (1983):** The morphology of human cervix. Clin. Obstet. Gynecol., 26: 7.

***Danielson KG, Marions L, Rodriguez A (1999):** Comparison between oral and vaginal administration of misoprostol on uterine contractility. Obstet Gynecol; 93:275–280.

***Danielian, P.; Porter, b.; Feri, N.; summers, T. and Templeton, A. (1999):** Misoprostol for induction of labor at term: more effective than dinoprostone. Br. J. Obstet. Gynecol. Augu. 106 (8): 793 -7.

***Danforth, D.N.; Buckingham, J.C. and Monalo, P. (1974):** The effect of pregnancy and labor on the human cervix: Changes in collagen, glycoprotein and glycosaminoglycan. Am. J. Obstet. Gynecol. 120: 6f41 -9.

***Danforth, D.N. (1954):** The distribution and functional activity of cervical musculature. Am. J. Obstet. Gynecol., 68: 1261.

***Danforth, D.N.; Veis, A.; Breen, M.; Weinstein, H.G.; Buckingham, J.C. and Manale, O.P. (1947):** The effect of pregnancy and labor on the human cervix, Changes in collagen, glycoprotein and glycosaminoglycan. Am. J. Obstet. Gynecol., 120: 641 – 51.

***Den Hertog, C.E.; de Groot, A.N. and van Dongen, P.W. (2001):** History and use of oxytocics. Euro J. Obstet. Gynecol. Reprod. Biol. 94: 8.

***Don Shuwarger (1996):** Misoprostol for cervical ripening and induction of labor. Mater Fetal Med. Depart. Meeting in Sep. 1996, 1-3.

E

***Echeverria, E. and Rocha, M. (1995):** Randomized comparative study of induced labor with oxytocin and misoprostol in prolonged pregnancy. Rev. Clin. Obstet. Gynecol. 60 (2): 108 – 11.

***Edwards, R.K. and Richard, D.S. (2000):** Pre – induction cervical assessment. Clin. Obstet. Gynecol., 43: 440.

- *Egarter, C.H.; Hussein, P.W. and Rayburn, W.F. (1990):** Uterine stimulation after low dose PGE₂ therapy: Tocolytic treatment in 181 cases. Am. J. Obstet. Gynecol. 163: 794 – 6.
- *Ekman, F.; Forman, A.; Marshal, K. and Ulmsten, U. (1983):** Intravaginal versus intracervical application of PGE₂ in viscous gel for cervical priming and labor induction at term, in – patients with unfavorable cervical state. Am. J. Obstet. Gynecol., 147: 657 – 61.
- *Elliot, J.P.; Clewell, H.W. and Radin, T.G. (1992):** Intracervical PGE₂ gel: safety for outpatient cervical ripening before induction of labor. Reprod Med. 37: 713.
- *Elliot, J.P. and Flaherty, J.F. (1983):** The use of breast stimulation to ripen the cervix in term pregnancies. Am. J. Obstet. Gynecol., 145: 553.
- *El-Refaey, H.; Calder, L.; Wheatley, D.N. and Templeton, A. (1994):** Cervical ripening with prostaglandin E₁ analogue (misoprostol) and gemeprost. Lancet 343: 1207 – 9.
- *El-Refaey, H.; Rajasekar, D.; Abdullah, M.; Calder, L. and Templeton, A. (1995):** Induction of abortion with mifepristone (RU486) and oral or vaginal misoprostol. N. Engl. J. Med., 332: 983 – 7.
- *El-Refaey, H.; Pat O'Brien, Walemorafa and Charles Rodeck (1997):** Use of oral misoprostol in prevention of postpartum hemorrhage. Br. J. Obstet. Gynecol. 104: 336 – 39.
- *Embrey, M.P. and Calder (1978):** Induction of labor. In: Beard RW, Brundenell M, Dunn P. and Fairweather D (Eds.). The management of labor: London Collage of Obstetricians and Gynecologists p. 62.
- *EnKin, M.; Keirse, M.; Renfrew, M. and Neilson, J. (2000):** A guide to effective care in pregnancy and childbirth. 3rd ed. Oxford Univ. Press.

F

- *Farah, L.A.; Sanchez – Ramos, L.; Rosa, C.; Del Valle, g.; Gaudier, F.L. and Delke, I. (1997):** Randomized trial of two doses of misoprostol for labor induction. Am. J. Obstet. Gynecol., 177: 364.
- *Fletcher, H.M.; Mitchell, S.; Frederick, J.; Simon, D. and Brown, D. (1993):** Intravaginal misoprostol and cervical ripening agent: double blind clinical trial. Br. J. Obstet. Gynecol., 100: 641 – 4.
- *Fletcher, H.M.; Mitchell, S.; Frederick, J.; Simon, D. and Brown, D. (1994):** Intravaginal misoprostol versus dinoprostone as cervical ripening and labor inducing agents. Obstet. Gynecol. 83: 244 – 7.
- *Foong, L.G.; Vanaja, K.; Tan, G. and Chua, S. (2000):** Membrane sweeping in conjunction with labor induction. Obstet. Gynecol. 96: 539.
- *Friedman, E.A.; Niswander, K.R.; Bayonet, N.P. and Sachtleben, M.R. (1992):** Relation of pre – labor evaluation to inducibility and course of labor. Obstet. Gynecol. 28: 495.
- *Friedman, E.A. (1998):** Labor: clinical evaluation and management, 2nd Ed. New York: Appleton – Century – Crofts.
- *Fuchs, A.R.; Fuchs, S.; Hussein, P. and Soloff, M.S. (1984):** Oxytocin treatment in the human uterus during pregnancy and parturition. Am. J. Obstet. Gynecol. 150: 734.

G

- *Garfield, R.E. and Hayashi, R.H. (1981):** Appearance of gap junction in the myometrium of women during labor. Am. J. Obstet. Gynecol. 140: 254.
- *Garite, T.J.; Porto, M.; Carlson, N.J. and Romney, P.J. (1993):** The influence of elective amniotomy on fetal heart rate patterns and the course of labor in term patients: A randomized study. Am. J. Obstet. Gynecol. 168: 1827 – 31.

- *Gemzell Danielson, K.; Marions, L.; Rodriguez, B.W.; Wong, P.Y.K and Bygdeman, M. (1999):** Comparison between oral and vaginal administration of misoprostol on uterine contractility. J. Obstet. Gynecol. 93: 275 – 80.
- *Goldberg AB, Greenberg MB, Darney PD. (2001):** Misoprostol and pregnancy. N Engl J Med; 344:38–47.
- *Gulmezoglu AM, Forna F, Villar J (2001):** Prostaglandins for prevention of postpartum hemorrhage. Cochrane Database Syst Rev; 4:CD000494.
- *Gimpl, G. and Fahrwnholz, F. (2001):** The oxytocin receptor system: structure, function and regulation. Physiol Rev. 81: 629.
- *Gottschall, D.S.; Borgida, A.F. and Mihalek, J.J. (1977):** A randomized clinical trial comparing misoprostol with prostaglandin E2 gel for pre – induction cervical ripening. Am. J. Obstet. Gynecol. (177) 5: 1067 – 70.
- *Greer, I.A. and Calder, A.A. (1989):** Pre – induction cervical ripening with extra amniotic and vaginal PGE2. J. Obstet. Gynecol. 10: 18 – 22.

H

- *Halls, K.A.; Rayburn, W.F. and Turnbull, G.L. (1994):** Double blind comparison of intracervical and intravaginal PGE2 for cervical ripening and induction of labor. Am. J. Obstet. Gynecol., 171: 1087 – 91.
- *Hannah, M.E.; Ohlsson, A. and Farine, D. (1996):** Induction of labor compared with expectant management for pre – labor rupture of membranes at term. Term PROM study group. N. Engl. J. Med. 334: 1005.
- *Hofmeyr GJ, Gulmezoglu AM, Alfrevic Z. (1999):** Misoprostol for induction of labor: a systematic review. Br J Obstet Gynaecol; 106:798–803.
- *Hofmeyr, G.J. and Gulmezoglu, A.M. (2001):** Vaginal misoprostol for cervical ripening and induction of labor (Cochrane Review). In Cochrane Library, Issue 4. Update Software, Oxford.

***Hughesdon, P.E. (1952):** The fibromuscular structure of the cervix and its changes during pregnancy and labor. Quoted from: Stud J. (ed.): The cervix in pregnancy and labor. Progress in Obstet & Gynecol., Vol.12 Churchill Livingstone P. 108.

***Hunt JN, Smith JL, Jiang CL. (1983):** Effect of synthetic prostaglandin E1 analog on aspirin-induced gastric bleeding and secretion. Dig Dis Sci; 28:897–902.

J

***Johnson, M. and Bennett, P. (2001):** Endocrine control of labor. In Turnbull's Obstetrics, 3rd edition, 25: 417 – 25.

***James, Grassland (1980):** Prostaglandins, hormones and nucleotides, Lewis pharmacology, p. 387.

***Jarvelin, M.R.; Hartikaiaen – Sorri, A.L. and Rantakallio, P. (1993):** Labor induction policy in hospitals of different levels of specialization. Br. J. Obstet. Gynecol. 100: 310.

K

***Karim, S.M.M. and Sharma, S.D. (1971):** Oral administration of prostaglandin for induction of labor. Br. Med. J. 1: 260.

***Karim, A. (1987):** Anti – ulcer prostaglandin, misoprostol: Single and multiple doses Pharmacokinetic profile. Prostaglandins, 3 (suppl.): 40 – 50.

***Karim, A. (1988):** Pharmacokinetic profile of anti – ulcer prostaglandin, Misoprostol, in human (abstract). Postgraduate Med. J. 64 (suppl. 1): 80.

***Kavanagh, J.; Kelly, A.J. and Thomas, J. (2001):** Corticosteroids for induction of labor (Cochrane Review). In: Cochrane Library Issue 2, Oxford: Update Software.

***Keirse, M.J.N.C.; Mitchell, M.D. and Turnbull, A.C. (1977):** Changes in prostaglandin F and 13, 14-Dihydro–15–Keto–prostaglandin F

concentrations in amniotic fluid at the onset of and during labor. Br. J. Obstet. Gynecol. 84: 87 – 102, 743 – 6.

***Keirse, M.J.N.C. (1979):** Prostaglandins in parturition. In: Keirse M, Anderson A, Gravenhorst J. (eds.): Human Parturition; The Hague, Martinus Nijhof, P. 101.

***Keirse M.J.N.C. (1990):** Eicosanoids in human pregnancy and parturition. In: Mitchell (ed.): Eicosanoids in reproduction. Boca Raton FL: CRC Press, p. 199.

***Keirse, M.J.N.C. (1995):** Stripping / sweeping membrane sat term for induction of labor. [Revised 03 April 1992] In: EnKin MW, Keirse, MJNC, Renfrew MJ and Neilson JP (Ed.). Pregnancy and Childbirth Module In: The Cochrane Pregnancy and Childbirth Database. The Cochrane Collaboration; Issue 2, Oxford: Update Software.

***Kelly, A.J.; Kavanagh, J. and Thomas, J. (2001):** Castor oil, path and / or enema for cervical priming and induction of labor (Cochrane Review). In: Cochrane Library, Issue 2. Oxford: Update Software.

***Kemp, B.; Winkler, M.; Mitchell, M.D. and Rath, W. (2000):** Induction of labor by PGE 2 in relation to the Bishop score. Int. J. Obstet. Gynecol., 71: 13.

***Kramer, Ralph, L. and Coerge, J. (1997):** A randomized trial of misoprostol and oxytocin for induction of labor: Safety and Efficacy. Obstet. Gynecol., 89: 387- 91.

L

***Lange, A.P.; Secher, F.S. and Westergaard, J.D. (1982):** Pre-labor evaluation of inducibility. Obstet. Gynecol., 60: 137.

***Lawrie, A.; Penney, G. and Templeton, A. (1996):** A randomized comparison of oral and vaginal misoprostol for cervical priming before suction termination of pregnancy. Br. J. Obstet. Gynecol., 103: 1117 – 9.

***Lelaidier, C.; Baton, C.; Benifla, J.K. and Fernandez, H. (1994):** Mifeprestone for labor induction after previous cesarean section. Br. J. Obstet. Gynecol. 101: 501 – 3.

***Leppert, P.C. (1992):** Cervical softening, effacement and dilatation: A complex biochemical cascade Journal of Maternal and Fetal Medicine 1: 213 – 223.

***Leppert, P.C. and Yu, S.Y. (1991):** Three – dimensional structure of uterine elastic fibers, scanning electron microscopic studies. Connect. Tissue. Res., 26: 15.

***Leppert, P.C. (1986):** Uterine cervical elastin in normal and abnormal mammalian gestation and parturition. Graduate Schools of arts and science. Columbia University Ann Arbor M1: University Microfilms.

***Liu HS, Chu TY, Chang YK, et al. (1999):** Intracervical misoprostol as an effective method of labor induction at term. Int J Gynaecol Obstet; 64:49–53.

***Liggins, G.C. (1978):** Ripening of the cervix. In: Oliver TK and Kirschbaum TK (Eds.). New York, Sem Perinatol 2: 261 - 71.

***Lindmark, G.; Zador, G. and Nilsson, B.A. (1975):** The induction of labor with prostaglandin F₂ by intravenous infusion. Acta Obstet. Gynecol. Scand. 37: 17.

***Ludmir, J. and Sehdev, H.M. (2000):** Anatomy and physiology of human cervix. Clin. Obstet. Gynecol. 43: 433.

M

***MacDonald, D. (1970):** Surgical induction of labor. Am. J. Obstet. Gynecol., 107: 908 – 11.

***MacDonald, P.C. and Casey, M.L. (1996):** Preterm birth. Sci. Am. 3: 42.

***Mackenzie, Z. and Burns, E. (1997):** Randomized trial of one versus two doses of PGE₂ for induction of labor. Br. J. Obstet. Gynecol. 104: 1062.

- *Mainbrize, T.; Nimond, Dodd, G. and Persand, D. (1987):** Clinical utility of multiple dose administration of PGE₂ gel. *Am. J. Obstet. Gynecol.* 156: 341.
- *Manabe, Y.; Manabe, A. and Sagawa, N. (1982):** Stretch induced cervical softening and initiation of labor at term: a possible correction with prostaglandins. *Acta Obstet. Gynecol.* 61: 279.
- *Margulies, M.; Perez, G.C. and Voto, L.S. (1992):** Misoprostol to induce labor. *Lancet* 339: 64.
- *Majoko F, Nystrom L, Lindmark G. (2002):** No benefit, but increased harm from high dose (100 ug) misoprostol for induction of labor: a randomized trial of high vs. low (50 ug) dose misoprostol. *J Obstet Gynaecol*; 22:61614–61617.
- *McInnes, D.R.; Naftolin, F. and VanDer Rest, M. (1980):** Cervical changes in pregnant women. In: Naftolin F and Stubble Field PGE₂ (Eds.). *Dilatation of the uterine cervix* New York Raven. Press, P. 181.
- *McPherson, M. (1984):** Comparison of lamicel with PGE₂ gel as a cervical ripening agent before the induction of labor. *J. Obstet. Gynecol.* 4: 205.
- *Meydanli MM, Caliskan E, Burak F (2003):** Labor induction post-term with 25 micrograms vs. 50 micrograms of intravaginal misoprostol. *Int J Gynaecol Obstet*; 81:249– 255.
- *Michael, S. Rogers (2001):** Induction and augmentation of labor. In: *Turnbull's Obstetrics* edited by Chamberlain G and Steer P, 3rd edition, Harcourt Publishers limited 35: 563 – 80.
- *Miramoto, T.; Arai, K.; Hirakawa, S. and Ngai, Y. (1982):** Immunohistochemical studies on collagen types in the uterine cervix in pregnant and non – pregnant state. *Am. J. Obstet. Gynecol.* 156: 138 – 44.

N

***Nero, C.M.; Leao, C.J.; Baretto, E.; Kenj, G. and De Aquino, M.M. (1987):** Use of misoprostol for labor induction in intrauterine fetal death. Rev Paul Med. 105: 325 – 8.

***Ngai, S.W.; Tang, O.S.; Leo, T. and Ma, H.K. (1995):** Oral misoprostol versus vaginal gemeprost for cervical dilatation prior to vacuum aspiration in women in sixth to twelfth weeks of gestation. Contraception: 51: 347 – 50.

***Ngai, S.W.; To, K.O. and Lao, T. (1996):** Cervical priming with oral misoprostol in pre – labor rupture of membranes at term. Obstet. Gynecol. 87: 923 – 6.

***Ngai, S.W.; Chan, Y.M. and Liu, K.L. (1997):** Oral misoprostol for cervical priming in non – pregnant women. Human Reprod (12) ii: 2373 – 5.

***Ngai, S.W.; Chan, Y.M.; Tang, O.S. and Ho. P.C. (1999):** The use of misoprostol for preoperative cervical ripening before vacuum aspiration: A randomized trial. Hum Reprod. 14: 2139 – 42.

***Nopdonrattakoon L. (2003):** A comparison between intravaginal and oral misoprostol for labor induction: a randomized controlled trial. J Obstet Gynaecol Res; 29:87–91.

***Norman, J.E.; Thong, K.J. and Baird, D.T. (1991):** Uterine contractility and induction of abortion in early pregnancy by misoprostol and mifeprestone. Lancet. 338: 1233 – 6.

***Novy, M.J. and Liggins, G.C. (1980):** Role of prostaglandins, Prostacyclin and Thromboxanes in the physiologic control of the uterus and in parturition. Semin Perinatal 4: 45.

***Nuutila, M.; Hiilesmaa, V.; Karkkainen, T. and Ylikorkala, O. (1990):** Phosphorelated isoform of insulin like growth factor binding protein - 1 in the process of cervical ripening. Am. J. Obstet. Gynecol., 163: 157.

O

***O'Brien, W.F. (1995):** The role of prostaglandins in labor and delivery. In Knuppel RA (ed.): Clinics in Perinatology. Complicated labor and delivery, Philadelphia WB Saunders.

***O'Brien, P.; El-Refaey, H.; Gordon, A.; Geary, M. and Rodeck, C.H. (1998):** Rectally administered misoprostol for the treatment of postpartum hemorrhage unresponsive to oxytocin and ergometrine. Engl. Obstet. Gynecol., 1998 Aug. 92 (2): 212 – 4.

***O'Driscoll, K.; Foley, M. and MacDonald, D. (1984):** Active management of labor as an alternative to cesarean section for dystocia. Obstet. Gynecol., 63: 485.

***Olah, K.S. (1996):** The cervix in pregnancy and labor. In stud. J. (Eds): Progress in Obstet and Gynecol. Vol. (12).

***Orga, S.S.; Kirton, K.I.; Tomasi, T.B. and Lippes, G. (1974):** Prostaglandins in human fallopian tubes. Fertility and Sterility, 25: 250 – 5.

***Orioli IM, Castillo EE. (2000):** Epidemiological assessment of misoprostol teratogenicity. BJOG; 107:519–523.

P

***Paterson- Brawn, S.; Fisk, N.M.; Rodeck, C.H. and Rodeck, E. (1991):** Pre-induction cervical assessment by Bishop Score and transvaginal ultrasound. Euro. J. Obstet. Gynecol. Reprod. Biol. 40: 17.

***Pena, J.C.G. (1995):** Misoprostol for induction of labor (letter). Am. J. Obstet. Gynecol., 174: 797.

***Pepe, G.J. and Albrecht, E.D. (1995):** Actions of placental and fetal steroid hormones in primate pregnancy. Endocrine Rev. 16: 608.

***Phyllis, C. and Leppert, M.D. (1995):** Anatomy and physiology of cervical ripening. Clinical Obstet. Gynecol. 38: 267 – 79.

***Plaut MM, Schwartz ML, Lubarsky SL. (1999):** Uterine rupture associated with the use of misoprostol in the gravid patient with a previous cesarean section. Am J Obstet Gynecol; 180:1535–1542.

R, S

***Roberts, G. and Turnbull, A.C. (1971):** Uterine hypertonus during labor induced by prostaglandins. Br. Med. J. 1: 702 – 5.

***Sanchez – Ramos, L.; Kaunitz, A.M.; Delvalle, G.O.; Delke, I.; Schroeder, P.A. and Briones, D.K. (1993):** Labor induction with PGE1 methyl analogue (misoprostol) versus oxytocin. Obstet. Gynecol. 172: 687 – 90.

***Sanchez – Ramos, L.; Farah, L.A.; Kaunitz, A.M.; David, D.; Delvalle, G.O. and Fuqua, P. (1995):** Pre – induction cervical ripening with commercially available PGE2 gel: A randomized double – blind comparison with a hospital compounded preparation. Am. J. Obstet. Gynecol. 173: 1079 – 84.

***Sanchez – Ramos, L.; Kaunitz, A.M. and Wears, R.W. (1997):** Misoprostol for cervical ripening and labor induction. Obstet. Gynecol. 89: 633 – 42.

***Sanchez – Ramos, L.; Chen, A.H.; Kaunitz, A.M.; Gaudier, F.L. and Delke, I. (1997):** Labor induction with intravaginal misoprostol in term premature rupture of membranes. Obstet. Gynecol. Jun. 89 (6): 909 – 12.

***Sanchez – Ramos, L.; Peterson, D.E. and Delke, I. (1998):** Labor induction with PGE1 misoprostol compared with dinoprostone vaginal insert. Obstet. Gynecol., 91: 401 – 5.

***Schuler L, Pastuszak A, Sanseverino TV (1999):** Pregnancy outcome after exposure to misoprostol in Brazil: a prospective, controlled study. Reprod Toxicol; 13:147–151.

- *Schoenhard, G.; Fitzpatrick, R.J. and Dixon, W.C.W. (1985):** Metabolism and pharmacokinetic studies of misoprostol. Drug. Dis. Sci. 30 (suppl): 126 – 8.
- *Schonhofer, P.S. (1991):** Brazil: Misuse of misoprostol as an abortifacient may induce malformation. Lancet, 337: 1534 – 5.
- *Secher, N.J.; Lange, A.P. and Nielsen, F. (1981):** Induction of labor with or without primary amniotomy. A randomized study of PGE₂ tablet and intravenous oxytocin, Acta Obstet. Gynecol. Scand. 60: 237.
- *Seitchik, J. (1987):** The management of functional dystocia in the first stage of labor. Clinical Obstet. Gynecol. 30: 42.
- *Sennstorm, M.; Granstrom, L. and Ekman, G. (1998):** Increased cervical fetal fibronectin levels after PGE₂ – induced cervical ripening. Am. J. Obstet. Gynecol., 178: 540 – 5.
- *Sennstorm, M.; Ekman, G.; Westergren – thorsson, G.; Malmstrom, A. and Brauner, A. (2000):** Human cervical ripening, as an inflammatory process mediated by cytokines. Molecular Human Reproduction 6: 375.
- *Shetty A, Stewart K, Stewart G (2001):** Active management of term prelabor rupture of membranes with oral misoprostol. BJOG; 109:1354–1358.
- *Shetty A, Danielian P, Templeton A. (2002a):** Sublingual misoprostol for the induction of labor at term. Am J Obstet Gynecol; 186:72–76.
- *Shetty A, Mackie L, Danielian P (2002b):** Sublingual compared with oral misoprostol in term labor induction: a randomized controlled trial. BJOG; 109:645–650.
- *Shojo, H. and Kaneko, Y. (2000):** Characterization and expression of oxytocin and the oxytocin receptor. Mol. Genet. Metab. 71: 552.
- *Srisomboon J, Tongsong T, Tosiri V. (1996):** Pre-induction cervical ripening with intravaginal prostaglandin E₁ methyl analogue misoprostol: a randomized controlled trial. J Obstet Gynaecol Res; 22:119–124.

***Stefos, T.; Mouzakioti, E.; Sotiriadis, A.; Andronikou S. and Lolis, D. (2000):** Use of PGE2 for cervical ripening in pregnancies with oligohydramnios. *Obstet Gynecol. Invest.* 50: 58.

***Stewart, P.; Kennedy, J.H. and Calder, A.A. (1983):** A comparison between estradiol and PGE2 for ripening of the cervix. *Br. J. Obstet. Gynecol.*, 88: 236 – 9.

***Stubbs, T.M. (2000):** Oxytocin for labor induction. *Clin. Obstet. Gynecol.* Sep. 43 (3): 489 – 94.

***Stys, S.J. Dreser, B.L.; Otte, T.E. and Clark, K.E. (1981):** Effect of PGE2 on cervical compliance in pregnant Ewes. *Am. J. Obstet. Gynecol.*, 140: 415 -9.

***Summers, L. (1997):** Method of cervical ripening and labor induction. *J. Nurse Midwifery*, 42: 71 – 85.

T

***Tang, O.S.; Schweer, H.; Seyberth, H.W.; Lee, S.W. and Ho, P.C. (2002):** Pharmacokinetics of different routes of administration of misoprostol. *Hum. Reprod.* 2002 Feb 17 (2): 332 – 6.

***Thomas, F.V. and Dorflinger, G.T. (1978):** Mechanism of rapid antigonadotrophine action of prostaglandins on culture luteal cell proceeding of the national academy of science 75: 1344 – 8.

***Toppozada, M.K.; Anwar, M.Y.; Hassan, H.A. and El – Gzaerly, W.S. (1997):** Oral or vaginal misoprostol for induction of labor. *Int. J. Obstet. Gynecol.* 56 (2): 135 – 9.

***Toth, P.P. and Jayarani (2001):** J. University of Iowa, Family Practice Handbook, 3rd edition, Chapter 8, the Mosby Co.

***Trofatter, K.F. (1993):** Endo-cervical PGE1 for pre – induction cervical ripening: clinical trial results. *J. Reprod. Med.* 38: 78.

U

***Uldbjerg, N.; Ulmsten, U. and Ekman, G. (1983):** The ripening of the human uterine cervix in terms of connective tissue biochemistry. Clin. Obstet. Gynecol. 26: 14 – 26.

***Ulmstien, U.; Wingerup, L. and Ekman, G. (1983):** Local application of PGE2 for cervical ripening or induction of term labor. Clin. Obstet. Gynecol., 26: 95.

***Ulmstien, U. and Wingerup, L. (1988):** Clinical experience with new gel for intracervical application of PGE2 for therapeutic abortion. In: Prostaglandins 20: 533.

***Ulmstien, M.; Sagen, N. and Ekholm, S.N. (1979):** A comparative study of labor induced by PGE 2 and buccal tablet of demoxytocin. Int. J. Obstet. Gynecol., 17: 243.

***Urban R, Lemancewicz A, Urban J (2003):** Misoprostol and dinoprostone therapy for labor induction: a Doppler comparison of uterine and fetal hemodynamic effects. Eur J Obstet Gynecol Reprod Biol 2003; 106:20–24.

V

***Varaklis, K.; Mina, R. and Stubblefeild, P.G. (1995):** Randomized controlled trial of vaginal misoprostol and intracervical PGE 2 gel for induction of labor at term. Obstet. Gynecol., 86: 541 – 4.

***Ventura, S.J.; Martin, J.A.; Curtin, S.C. and Matthew, T.J. (1999):** Births: final data for 1997. National Center for Health Statistics, National Vital Statistics Reports 47: 1.

***Vincent du Vigneaud, V.; Ressler, C. and Swan, J.M. (1954):** The Synthesis of oxytocin. J. Am. Chem. Soc. 75: 4879.

***Vogel, I.; Salvig, J.D.; Secher, N.J. and Uldbjerg, N. (2001):** Association between raised serum relaxin levels during the eighteenth gestational week and very preterm delivery. Am. J. Obstet. Gynecol. 184: 390.

***Von Meir, C.A.; Ramirez, M.M.; Matthew, S.G.; Calder, A.A.; Keirse, M.J.N.C and Challis, J.R.C. (1997):** Chorionic prostaglandin catabolism is decreased in lower uterine segment with term labor. Placenta: 18: 109 – 14.

W

***Wilcott, H.D. and Conry (2000):** Normal labor. In Evans AT and Niswander KR (Eds.) Manual of Obstetrics: Sixth edition Lippincott Williams and Wilkins, p. 392 – 424.

***Windrim, R.; Bennett, R. and Mundle, W. (1997):** Oral misoprostol for labor induction: A randomized controlled trial. Obstet. Gynecol. 89: 721 – 8.

***Wing, D.A.; Jones, M.M.; Rahall, A.; Goodwin, T.M. and Paul, R.H. (1995):** Misoprostol: an effective agent for cervical ripening and labor induction Am. J. Obstet. Gynecol., 172: 1811 – 6.

***Wing, D.A. and Paul, R.H. (1997):** A comparison of differing dosing regimens of vaginally administered misoprostol for pre-induction cervical ripening and labor induction. Am. J. Obstet. Gynecol., 175: 158.

***Wing, D.A. and Paul, R.H. (1999):** Misoprostol for cervical ripening and labor induction: the clinician's perspective and guide to success. Contemp. Ob/Gyn. 44: 46.

***Wing, D.A.; Fassett, M.J. and Mishell, D.R. (2000):** Mifeprestone for pre – induction cervical ripening beyond 41 weeks' gestation: a randomized controlled trial. Obstet. Gynecol., 96: 543.

***Word, R.A.; Kamm, K.E.; Stull, J.T. and Casey, M.L. (1990):** Endothelin increases cytoplasmic calcium and myosin phosphorylation in human myometrium. Am. J. Obstet. Gynecol., 162: 1103.

***Word, R.A.; Stull, J.T.; Casey, M.L. and Kamm, K.E. (1993):** Contractile elements and myosin light chain phosphorylation in myometrial tissue from non – pregnant and pregnant women. J. Clin. Invest. 92: 29.

Y, Z

***Yawn, B.P.; Wollan, P.; McKeon, K. and Field, C.S. (1998):**

Temporal changes in rates and reasons for medical induction of term labor, 1980 – 1996. Am. J. Obstet. Gynecol., 184: 611.

***Zieman, M.; Fong, S.K. and Mitchell, S. (1997):** Absorption

kinetics of misoprostol with oral or vaginal administration. Obstet. Gynecol., 90: 88 – 92.

Thanks to God