

References

- Abbassi H., Aboulfalah A., Morsad F., et al. (2000):*** Maternal complications of cesarean section: retrospective analysis of 3,231 interventions at the Casablanca University Hospital. Morocco. Sante; 10 (6): 419.
- Aboulfalah A., Abbassi H., El Karroumi M., et al. (2000):*** Delivery of large baby after cesarean section: role of trial of labor. A propose of 355 cases. J Gynecol Obstet Biol Reprod (Paris); 29(4):409.
- ACOG practice bulletin (1999):*** Vaginal birth after previous cesarean delivery. Number 2, October 1998. Clinical management guidelines for obstetrician-gynecologists. American College of Obstetricians and Gynecologists. Int J Gynaecol Obstet; 64 (2): 201.
- Adair DC., Sanchez-Ramos L., Whitaker D., et al. (1996):*** Trial of labor in patients with a previous lower uterine vertical cesarean section. Am. J. Obst Gynecol; 174: 966.
- American College of Obstetrics and Gynecologists, Washington, DC 20090-6920, U.S.A. (2002):*** Induction of labor for vaginal birth after cesarean delivery. Committee on Obstetric Practice. Int J. Gynaecol Obstet; 77(3): 303.
- Amin AF., Misselwitz B. and Runzel W. (2001):*** Cesarean section rate as a measure of quality assurance in obstetric practice. The Journal of Egyptian Society of Obstetric and Gynecology; 27: 157.

- Araki T. and Inooka H. (1982):** The diagnostic value of ultrasonotomography with reference to previous cesarean section scars during full term pregnancy. *Acta Obstet Gynecol.*; 34: 738-44.
- Armstrong V., Hansen WF., Van Voorhis BJ., et al. (2003):** Detection of cesarean scars by transvaginal ultrasound. *Obstet Gynecol*; 101 (1): 61.
- Asakura H., Nakai A., Ishikawa G., et al. (2000):** Prediction of uterine dehiscence by measuring lower uterine segment thickness prior to the onset of labor: evaluation by transvaginal ultrasonography. *J Nippon Med Sch.*; 67 (5): 352. (pubmed English abstract).
- Ashoush S. (2004):** Trial of Scar. *ASJOG*; 1:115.
- Aslan H. Unlu E., Agar M., et al. (2004):** Uterine rupture associated with misoprostol labor induction in women with previous cesarean delivery. *Eur J Obstet Gynecol Reprod Biol.*; 113 (1): 45.
- Bedoya C., Bartha JL., Rodriguez I., et al. (1992):** A trial of labor after cesarean section in patients with or without a prior vaginal delivery. *Int J Gynaecol Obstet*; 39 (4): 285.
- Behrens O., Goeschen K., Jakob H., et al. (1994):** Induced labor with prostaglandin E2 gel after previous cesarean section. *Geburtshilfe Frauenheilkd*; 54 (3): 144. (English abstract)
- Berkowitz GS., Fiarman GS., Mojica MA., et al. (1990):** Effect of physician characteristics on the cesarean birth rate. *Am J Obstet Gynecol*; 162 (6): 1641.

- Biswas A. (2003):** Management of previous cesarean section. *Curr Opin Obstet Gynecol.*; 15 (2): 123.
- Brill Y. and Windrim R. (2003):** Vaginal birth after caesarean section: review of antenatal predictors of success. *J Obstet Gynaecol Can.*; 25 (4): 275.
- Brost BC., Goldenberg RL., Mercer BM., et al. (1997):** Association of cesarean delivery with increases in maternal weight and body mass index. *Am J Obstet Gynecol.*; 177 (2): 333.
- Bujold E., Hammoud AO., Hendler I., et al. (2004):** Trial of labor in patients with a previous cesarean section: does maternal age influence the outcome? *Am J Obstet. Gynecol.*; 190 (4): 1113.
- Bujold E., Mehta SH., Bujold C. et al. (2002):** Interdelivery interval and uterine rupture. *Am J Obstet Gynecol.*; 187 (5): 1199.
- Carroll CS., Magann EF., Cauhan SP., et al. (2003):** Vaginal birth after cesarean section versus elective repeat cesarean delivery: weight based outcome. *Am J Obstet. Gynecol.*; 188: 1516.
- Caughey AB., Shipp TD., Repke JT., et al. (1998):** Trial of labor after cesarean delivery: the effect of previous vaginal delivery. *Am J Obstet Gynecol.*; 179 (4): 938.
- Caughey AB., Shipp TD., Repke JT., et al. (1999):** Rate of uterine rupture during a trial of labor in women with one or two prior cesarean deliveries. *Am J Obstet Gynecol.*; 181 (4): 872.
- Chamberlain G. (2001):** Cesarean Section: In *Turnbull's Obstetrics*, Third edition, Chamberlain G. and Steer PJ Harcourt Publishers limited, p: 601.

- Chapman SJ., Owen J. and Hauth JC. (1997):** One versus two layer closure of a low transverse cesarean: The next pregnancy. *Obst Gynecol*; 89:16.
- Chauhan SP., Magann EF., Carroll CS., et al. (2001):** Mode of delivery for the morbidly obese with prior cesarean delivery: vaginal versus repeat cesarean section. *Am J Obstet Gynecol*; 185: 349.
- Chauhan SP., Martin JN., Henrichs CE., et al. (2003):** Maternal and perinatal complications with uterine rupture in 142,075 patients who attempted vaginal birth after cesarean delivery: review of the literature. *Am J Obstet Gynecol*; 189: 409.
- Chazotte C. and Cohen WR. (1990):** Catastrophic complications of previous cesarean section. *Am J Obstet Gynecol*; 163: 738.
- Chilaka VN., Cole MY., Habayeb OM., et al. (2004):** Risk of uterine rupture following induction of labor in women with a previous caesarean section in a large UK teaching hospital. *J Obstet Gynaecol.*; 24 (3): 264.
- Choy-Hee L. and Raynor BD. (2001):** Misoprostol induction of labor among women with a history of cesarean delivery. *Am J Obstet Gynecol.*; 184 (6): 1115.
- Clark SL., Scott JR., Porter TF., et al. (2000):** Is vaginal birth after cesarean less expensive than repeat cesarean delivery? *Am J Obstet Gynecol*; 182: 599
- Coleman TL., Randall H., Graves W., et al. (2001):** Vaginal birth after cesarean among women with gestational diabetes. *Am J Obstet Gynecol.*; 184 (6): 1104.

- Coltart TM., Davies JA. and Katesmark M. (1990):** Outcome of a second pregnancy after a previous elective caesarean section. Br J Obstet Gynaecol.; 97(12): 1140.
- Cowan RK., Kinch RA., Ellis B., et al. (1994):** Trial of labor following cesarean delivery. Obstet Gynecol.; 83 (6): 933.
- Cunningham F., Gant F., Leveno J., et al. (2001):** Abnormal labor and fetopelvic disproportion. In: Williams Obstetrics, twenty first edition, McGRAW-HILL; p:425.
- Cunningham F., Gant F., Leveno J., et al. (2001):** Cesarean delivery and postpartum hysterectomy. In: Williams Obstetrics, twenty first edition, McGRAW-HILL; p:54l.
- De Meeus JB., Ellia F. and Magnin G. (1998):** External cephalic version after previous cesarean section: a series of 38 cases. Eur J Obstet Gynecol Reprod Biol.; 81 (1): 65.
- Delaney T. and Young DC. (2003):** Trial of labor compared to elective Caesarean in twin gestations with a previous caesarean delivery. J Obstet Gynaecol Can.; 25 (4): 289.
- Depp (2002):** Cesarean section In: Gabbe Obstetrics-Normal and Problem Pregnancies, 4th edition, Gabbe SG, Niebyl JR and Simbson JL (ed.) Churchill Livingstone. P:573
- Drjfe J. (2002):** The start of life: a history of obstetrics. Postgrad Med J.; 78 (919): 311.
- Durnwald C. and Mercer B. (2003):** Uterine rupture, perioperative and perinatal morbidity after single-layer and double-layer closure at cesarean delivery. Am J Obstet Gynecol; 189: 925.

- E1-Kousy AM., Sammel M., Stevens E., et al. (2003):** The effect of birth weight on vaginal birth after cesarean delivery success rates. *Am J Obstet Gynecol*; 188: 824.
- Ecker JL., Chen KT., Cohen AP., et al. (2001):** Increased risk of cesarean delivery with advancing maternal age: indications and associated factors in nulliparous women. *Am J Obstet Gynecol.*; 185 (4): 883.
- Eltabbakh GH. and Watson JD. (1995):** Postpartum hysterectomy. *Int J Gynaecol Obstet.*; 50 (3): 257.
- Esposito MA., Menihan CA. and Malee MP. (2000):** Association of inter-pregnancy interval with uterine scar failure in labor: a case-control study. *Am J Obstet Gynecol.*; 183 (5): 1180.
- Falzone S., Chauhan SP., Mobley JA., et al. (1998):** Unengaged vertex in nulliparous women in active labor. A risk factor for cesarean delivery. *J Reprod Med.*; 43 (8): 676.
- Farmer-Richard M., Kirschbaum T., Daniel P., et al. (1991):** Uterine rupture during trial of labor after previous cesarean section. *Am J Obstet Gynecol*; 165: 996.
- Flamm BL. (1997):** Once a cesarean, always a controversy. *Obstet Gynecol*; 90: 312.
- Flamm BL. and Geiger AM. (1997):** Vaginal birth after cesarean delivery: an admission scoring system. *Obstet Gynecol*; 90: 907.
- Flamm BL., Goings JR., Lin Y., et al. (1994):** Elective repeat cesarean delivery versus trial of labor: A prospective multicenter study. *Obstet Gynecol*; 83: 927.

- Flamm BL., Newman LA., Thomas SJ., et al. (1990):** Vaginal birth after cesarean delivery: results of a 5-year multicenter collaborative study. *Obstet Gynecol.*; 76 (5): 750.
- Fraser W., Maunsell E., Hodnett E., et al. (1997):** Randomized controlled trial of a prenatal vaginal birth after cesarean section education and support program. *Am J Obstet Gynecol*; 176: 419.
- Fukuda M., Fukuda K. and Mochizuki M. (1988):** Examination of previous cesarean section scars by ultrasound *Arch Gynecol Obstet.*; 243: 221.
- Gotoh H., Masuzaki H., Yoshida A., et al. (2000):** Predicting incomplete uterine rupture with vaginal sonography during the late second trimester in women with prior cesarean. *Obstet Gynecol.*; 95 (4): 596.
- Gregory KD., Korst LM., Cane P., et al. (1999):** Vaginal birth after cesarean and uterine rupture rates in California. *Obstet Gynecol.*; 94 (6): 985.
- Haas DM. and Ayres AW. (2002):** Laceration injury at cesarean section. *J Matern Fetal Neonatal Med.*; 11(3): 196.
- Hamd ME. (2002):** Study of cesarean section delivery in Al-Mansoura University Hospital Msc, Thesis P:32.
- Heffner U., Elkin E. and Fretts RC. (2003):** Impact of labor induction, gestational age, and maternal age on cesarean delivery rates. *Obstet Gynecol.*; 102 (2): 287.
- Hem E. and Bordahi PE. (1998):** The first cesarean section in Norway. *Tidsskr Nor Laegeforen.* 10; 118 (30): 4648 (English abstract).

- Hem E. and Bordahi PE. (2003):** Max Sanger-father of the modern caesarean section. *Gynecol Obstet Invest.*; 55 (3): 127. (English abstract).
- Hendler I. and Bujold E. (2004):** Effect of prior vaginal delivery or prior vaginal birth after cesarean delivery on obstetric outcomes in women undergoing trial of labor. *Obstet Gynecol.*; 104 (2): 273.
- Hendricks MS., Chow YH., Bhagavath B., et al. (1999):** Previous cesarean section and abortion as risk factors for developing placenta previa. *J. Obstet Gynaecol Res.*; 25 (2): 137.
- Hibbard JU., Ismail MA., Wang Y., et al. (2001):** Failed vaginal birth after a cesarean section: how risky is it? *Am J Obstet Gynecol.*; 184 (7): 1365.
- Hoskins IA. and Gomez JL. (1997):** Correlation between maximum cervical dilatation at cesarean delivery and subsequent vaginal birth after cesarean delivery. *Am J Obstet Gynecol*; 89: 591.
- Huang WH., Nakashima DK., Rumney PJ., et al. (2002):** Interdelivery interval and the success of vaginal birth after cesarean delivery. *Obstet Gynecol*; 99 (1): 41.
- Impey L. and O'herlihy C. (1998):** First delivery after cesarean delivery for strictly defined cephalopelvic disproportion. *Am J Obstet Gynecol*; 92: 799.
- Ismail MA., Nagib N., Ismail T., et al. (1999):** Comparison of vaginal and cesarean section delivery for fetuses in breech presentation. *J Perinat Med.*; 27(5): 339.

- Johnson DP., Davis NR. and Brown AJ. (2003):** Risk of cesarean delivery after induction at term in nulliparous women with an unfavorable cervix. *Am J Obstet Gynecol.*; 188 (6): 1565.
- Jongen VH., Halfwerk MG. and Brouwer WK. (1998):** Vaginal delivery after previous caesarean section for failure of second stage of labor. *Br J Obstet Gynaecol.*; 105 (10): 1079.
- Khan KS. and Rizvi A. (1995):** The partograph in the management of labor following cesarean section. *Int J Gynecol Obstet.*; 50 (2): 151.
- Khan KS. and Rizvi A. (1995):** The partograph in the management of labor following cesarean section. *Int J Gynecol Obstet.*; 50 (2): 151.
- Khawaja M., Jurdi R. and Kabakian-Khasholian T. (2004):** Rising trends in cesarean section rates in Egypt. *Birth.*; 31 (1): 12.
- Kolas T., Hofoss D., Daitveit A.K., et al. (2003):** Indications for cesarean deliveries in Norway. *Am J Obstet Gynecol.*; (188): 864.
- Lao TT. and Leung BF. (1987):** Labor induction for planned vaginal delivery in patients with previous cesarean section. *Acta Obstet Gynecol Scand.*; 66 (5): 413. (English abstract)
- Lazarov L. (1993):** A breech presentation after a cesarean section--is cesarean section always needed? *Akush Ginekol (Sofia)*; 32 (1): 10. (English abstract)
- Lehmann M., Hedelin G., Sorgue C., et al. (1999):** Predictive factors of the delivery method in women with cesarean section scars. *J Gynecol Obstet Bioi Reprod (Paris)*; 28(4):358.

- Leung AS., Farmer RM., Leung EK., et al. (1993):** Risk factors associated with uterine rupture during trial of labor after cesarean delivery; a case-control study. *Am J Obstet Gynecol*; 168: 1358.
- Leung GM., Lam TH., Thach TQ., et al. (2001):** Rates of cesarean births in Hong Kong: 1987-1999. *Birth.*; 28 (3): 166.
- Levy AL., Gonzalez JL., Rappaport VJ., et al. (2002):** Effect of labor induction on cesarean section rates in diabetic pregnancies. *J Reprod Med.*; 47 (11): 931.
- Levy AL., Gonzalez JL., Rappaport VJ., et al. (2002):** Effect of labor induction on cesarean section rates in diabetic pregnancies. *J Reprod Med.*; 47(11): 931.
- Lieberman E., Lang JM., Heffner U., et al. (1998):** Assessing the role of case mix in cesarean delivery rates. *Obstet Gynecol.*; 92 (1): 1.
- Lin C. and Raynor BD. (2004):** Risk of uterine rupture in labor induction of patients with prior cesarean section: an inner city hospital experience. *Am J Obstet Gynecol.*; 190 (5): 1476.
- Ling C. and Xu L. (1995):** The prospective study of trial labor following previous cesarean section. *Zhonghua Fu Chan Ke Za Zhi.*; 30 (12): 721. (English abstract)
- Liu S., Rusen ID., Joseph KS., et al. (2004):** Recent trends in caesarean delivery rates and indications for caesarean delivery in Canada. *J Obstet Gynaecol Can.*; 26 (8): 735.
- Lieberman E. (2001):** Risk factors for uterine rupture during a trial of labor after cesarean. *Clin Obstet Gynecol*; 44 (3): 609.

- Lydon-Rochelle M., Holt VL., Easterling TR., et al. (2001):** First-birth cesarean and placental abruption or previa at second birth. *Obstet Gynecol.*; 97 (5): 765.
- Macones GA., Hausman N., Edeistein R., et al. (2001):** Predicting outcomes of trials of labor in women attempting vaginal birth after cesarean delivery: a comparison of multivariate methods with neural networks. *Am J Obstet Gynecol*; 184: 409.
- Maher HA. (1999):** Cesarean section rate in Kasr Elaini university Hospital during the past two years. Ms.Sc; Obsteric and gynecology Thesis; Cairo University Egypt.
- Makura H. and Myers SA. (1995):** More than one previous cesarean delivery: a 5-year experience with 435 patients. *Am J Obstet Gynecol*; 85: 924.
- Mankuta DD., Leshno MM., Menasche MM., et al. (2003):** Vaginal birth after cesarean section: trial of labor or repeat cesarean section? A decision analysis. *Am J Obstet Gynecol.*; 189 (3): 714.
- Marchiano D., Elkousy M., Stevens E., et al. (2004):** Diet-controlled gestational diabetes mellitus does not influence the success rates for vaginal birth after cesarean delivery. *Am J Obstet Gynecol.*;190 (3): 790.
- Martel MJ. and MacKinnon CJ. (2004):** Clinical Practice Obstetrics Committee of the Society of Obstetricians and Gynaecologists of Canada. Guidelines for vaginal birth after previous Caesarean birth. *J Obstet Gynaecol Can.*; 26 (7): 660.

- Martin J., Perry KG., Roberts WE., et al. (1997):** The case for trial of labor in the patient with a prior low-segment vertical cesarean incision. *Am J Obstet Gynecol.*; 177 (1): 144.
- McMahon MJ. (1998):** Vaginal birth after cesarean. *Clin Obstet Gynecol.*; 41(2): 369.
- McNally OM. and Turner MJ. (1999):** Induction of labour after 1 previous cesarean section. *J Obstet Gynaecol.*; 39 (4): 425.
- Meier PR. and Porreco R.P. (1982):** Trial of labor following cesarean section A two-year experience. *Am J Obstet Gynecol.*; 144: 671.
- Menacker F. and Curtin SC. (2001):** Trends in cesarean birth and vaginal birth after previous cesarean, 1991-99. *Natl Vital Stat Rep.* 27; 49 (13): 1.
- Mesleh RA., Asiri F. and Al-Naim MF. (2000):** Cesarean section in the primigravid. *Saudi Med J.*; 21 (10): 957.
- Miller DA., Diaz FG. and Paul RH (1994):** Vaginal birth after cesarean: 10-year experience. *Am J Obstet Gynecol.*; 84: 255.
- Miller DA., Mullin P., Hon D. and Paul RH. (1996):** Vaginal birth after cesarean section in twin gestation. *Am J Obstet Gynecol.*; 175(1):194.
- Mjcbals WH., Thompson HO., Boutt A., et al. (1998):** Ultrasound diagnosis of defects in the scarred lower uterine segment during pregnancy. *Obstet Gynecol.*; 71 (1): 112.
- Montanarj L., Alfei A., Drovanti A., et al. (1999):** Transvaginal ultrasonic evaluation of the thickness of the section of the uterine

wall in previous cesarean sections. *Minerva Ginecol.*; 51 (4): 107-12. (English abstract)

Mozurkewich EL. and Hutton EK. (2000): Elective repeat cesarean delivery versus trial of labor: a meta-analysis of the literature from 1989 to 1999. *Am J Obstet Gynecol.*; 183 (5): 187.

Myles T. (2001): Vaginal birth of twins after a previous Cesarean section. *J Matern Fetal Med.*; 10 (3): 171.

Naiden J. and Deshpande P. (2001): Using active management of labor and vaginal birth after previous cesarean delivery to lower cesarean delivery rates: A 10-year experience. *Am J Obstet Gynecol.*; 184 (7): 1535.

Nguyen TV., Dinh IV., Suresh MS., et al. (1992): Vaginal birth after cesarean section at the University of Texas. *J Reprod Med.*; 37 (10): 880.

Notzon FC. (1990): International differences in the use of obstetric interventions. *JAMA.*; 263 (24): 3286.

Ofir K., Sheiner E., Levvy A., et al. (2003): Uterine rupture: risk factors and pregnancy outcome. *Am J Obstet Gynecol.*; 189: 1042.

Ophir E., Oettinger M., Yagoda A., et al. (1989): Breech presentation after cesarean section: Always a section? *Am J Obstet Gynecol.*; 161: 25.

O'Sullivan MJ., Fumia F., Holsinger K., et al. (1981): Vaginal delivery after cesarean section. *Clin Perinatol.* Feb; 8 (1): 131.

- Parrish KM., Holt VL., Easterling TR., et al. (1994):** Effect of changes in maternal age, parity, and birth weight distribution on primary cesarean delivery rates. JAMA.; 271 (6): 443.
- Phelan JP. (1990):** Uterine rupture. Clin Obstet Gynecol; 33 (3): 432.
- Phelan JP., Ahn MO., Brar HS., et al. (1989):** Twice a cesarean: always a cesarean? Obstet Gynecol, 73: 161.
- Phelan JP., Clark SL., Diaz F., et al. (1987):** Vaginal birth after cesarean. Am J Obstet Gynecol.; 157 (6): 1510.
- Pickhardt MG., Martin JN., Meydrech EF., et al. (1992):** Vaginal birth after cesarean delivery: Are there useful and valid predictors of success or failure? Am J Obstet Gynecol; 166: 1811.
- Poma PA. (1999):** Effects of obstetrician characteristics on cesarean delivery rates. A community hospital experience. Am J Obstet Gynecol.; 180 (6): 1364.
- Poma PA. (2000):** Rupture of a cesarean-scarred uterus: a community hospital experience. J Nail Med Assoc.; 92 (6): 295.
- Raheem and Shalloum (2003):** Clinical audit: obstetric performance after cesarean section. J Obstet Gynecol.; 23 (5): 503.
- Raimer KA. and O'Sullivan MJ. (1992):** Cesarean section: History, incidence, and indications. In: surgical Obstetrics, ninth edition, Plauche WC., Morrison JC. And O'sullivan MJ., W.B. Saunders Company Harcourt Brace Jovanovich, Inc. Philadelphia, Pennsylvania. p: 405.

- Rasmussen OB., Pedersen BL., Wilken-Jensen C., et al. (2000):** Stratified rates of cesarean sections and spontaneous vaginal deliveries. Data from five labor wards in Denmark 1996. Acta Obstet Gynecol Scand.; 79 (3): 227.
- Ridgeway JJ., Weyrich DL. and Benedetti TJ. (2004):** Fetal heart rate changes associated with uterine rupture. Obstet Gyneeol.; 103 (3): 506.
- Robson MS., Scudamore IW. and Walsh SM (1996):** Using the medical audit cycle to reduce cesarean section rates. Am J Obstet Gynecol.; 174 (1): 199.
- Rooney BL., Thompson JE., Schauburger CW., et al. (1996):** Is a twelve-percent cesarean section rate at a perinatal center safe? J Perinatol.; 16 (3): 215.
- Rosen MG. and Dickinson J.C. (1990):** Vaginal birth after cesarean section a meta analysis of indicators of success. Obstet Gynecol; 76 (1): 865.
- Rosen MG., Dickinson JC. and Westhoff CL. (1991):** Vaginal birth after cesarean: A Meta-analysis of morbidity and mortality. Obstet Gynecol; 77: 465.
- Rozenberg P., Goffinet F., Phillippe HJ., et al. (1996):** Ultrasonographic measurement of lower uterine segment to assess risk of defects of scarred uterus. Lancet.; 347 (3): 281.
- Rozenberg P., Goffinet F., Phillippe HJ., et al. (1999):** Thickness of lower uterine segment :its influence in the management of

patients with previous cesarean sections. Eur J Obstet Gynecol Reprod Biol; 87 (1): 39.

Sachs BP., Kobelin C., Castro MA., et al. (1999): The risks of lowering the cesarean delivery rate. N Engl J Med; 340: 54.

Sansregret A., Bujold E. and Gauthier RJ. (2003): Twin delivery after a previous caesarean: a twelve-year experience. J Obstet Gynaecol Can.; 25 (4): 294.

Schachter M., Kogan S. and Blickstein I. (1994): External cephalic version after previous cesarean section--a clinical dilemma. Int J Gynaecol Obstet.; 45 (1): 17.

Scheller JM. and Nelson KB. (1994): Does cesarean delivery prevent cerebral palsy or other neurological problems of childhood? Obstet Gynecol; 83 (4), 624.

Scott JR. (1991): Mandatory trial of labor after cesarean delivery: An alternative viewpoint: Obstet Gynecol; 77: 811.

Sebhatu B. (1994): An experience with trial of scar in 66 Ethiopian women. East Afr Med J.; 71 (10): 676.

Sen S., Malik S. and Salhan S. (2004): Ultrasonographic evaluation of lower uterine segment thickness in patients of previous cesarean section. Int. J. Gynaecol Obstet.; 87 (3): 215-9.

Shipp TD., Zelop C., Cohen A., et al. (2003): Post-cesarean delivery fever and uterine rupture in a subsequent trial of labor. Obstet Gynecol.; 101 (1): 136.

- Shipp TD., Zelop CM. Repke JT., et al. (1999):** Intrapartum uterine rupture and dehiscence in patients with prior lower uterine segment vertical and transverse incisions. *Obstet Gynecol.*; 94 (5): 735.
- Shipp TD., Zelop CM., Repke JT., et al. (2002):** The association of maternal age and symptomatic uterine rupture during a trial of labor after prior cesarean delivery. *Am J Obst Gynecol*; 99: 585.
- Shipp TD., Zelop CM., Repke JT., et al. (2001):** Interdelivery interval and risk of symptomatic uterine rupture. *Obstet Gynecol.*; 97 (2): 175.
- Silberstein T., Wiznitzer A., Katz M., et al. (1998):** Routine revision of uterine scar after cesarean section: has it ever been necessary? *Eur J Obstet Gynecol Reprod Biol.*; 78 (1): 29.
- Silver RK. and Gibbs RS. (1987):** Predictors of vaginal delivery in patients with a previous cesarean section, who require oxytocin. *Am J Obst Gynecol*; 156 (1): 57.
- Smith GC., Pelt JP., Cameron AD., et al. (2002):** Risk of perinatal death associated with labor after previous cesarean delivery in uncomplicated term pregnancies. *JAMA.*; 287 (20): 2684.
- Smith JF., Hernandez C. and Wax JR. (1997):** Fetal laceration injury at cesarean delivery. *Obstet Gynecol.*; 90 (3): 344.
- Spaans WA., Sluijs MB., van Roosmalen J., et al. (2002):** Risk factors at caesarean section, and failure of subsequent trial of labor. *Eur J Obstet Gynecol Reprod Biol.*; 100 (2): 163.

- Suzuki S., Sawa R., Yoneyama Y., et al. (2000):** Preoperative diagnosis of dehiscence of the lower uterine segment in patients with a single previous caesarean section. *Aust N Z J Obstet Gynaecol.*; 40 (4): 402.
- Swaim LS., Hoiste CS. and Wailer DK. (1998):** Umbilical cord blood pH after prior cesarean delivery. *Obstet Gynecol.*; 92 (3): 390.
- Swaim LS., Hoiste CS. and Wailer DK. (1998):** Umbilical cord blood pH after prior cesarean delivery. *Obstet Gynecol.*; 92 (3): 390.
- Tanik A., Ustun C., Cil E. and Arslan A. (1996):** Sonographic evaluation of the wall thickness of the lower uterine segment in patients with previous cesarean section. *J. Clin Ultrasound.*; 24: 355-357.
- To WW. and Leung WC. (1995):** Placenta previa and previous cesarean section. *Int J Gynaecol Obstet.*; 51 (1): 25.
- Tucker JM., Hauth J., Hodgkins P., et al. (1993):** Trial of labor after a one- or two-layer closure of a low transverse uterine. *Am J Obstet Gynecol*; 168: 545.
- Turrentine MA. and Ramirez MM. (1999):** Adverse perinatal events and subsequent cesarean rate. *Obstet Gynecol.*; 94 (2): 185.
- Ventura SJ., Martin JA., Curtin SC., et al. (2000):** Final data for 1998 national vital statistical report. *Birth*; 48 (3): 1.
- Vinueza CA., Chauhan SP., Barker L., et al. (2000):** Predicting the success of a trial of labor with a simple scoring system. *J Reprod Med*; 45: 332.

- Wasef WR. (2000):** An audit of trial of labor after previous cesarean section. J Obstet Gynecol.; 20 (4): 380.
- Weinstein D., Benshushan A., Tanos V., et al. (1996):** Predictive score for vaginal birth after cesarean section. Am J Obstet Gynecol.: 174 (1): 192.
- Whiteside DC., Mahan CS. and Cook JC. (1983):** Factors associated with successful vaginal delivery after cesarean section. J Reprod Med.; 28 (11): 785.
- Wilkinson C., McIlwaine G., Boulton-Jones C., et al. (1998):** Is a rising caesarean section rate inevitable? Br J Obstet Gynecology.; 105 (1): 45.
- Yamani Zamzami TY. (2004):** Vaginal birth after cesarean section in grand multiparous women. Arch Gynecol Obstet.; 270 (1): 21.
- Zaideh SM., Abu-Heija AT. and El-Jallad MF. (1998):** Placenta praevia and accreta: analysis of a two-year experience. Gynecol Obstet Invest.; 46 (2): 96.
- Zelop CM., Shipp TD. Repke JT., et al. (1999):** Uterine rupture during induced or augmented labor in gravid women with one prior cesarean delivery. Am J Obstet Gynecol; 181 (4): 872.
- Zelop CM., Shipp TD., Repke JT., et al. (2000):** Effect of previous vaginal delivery on the risk of uterine rupture during a subsequent trial of labor. Am J Obstet Gynecol; 183: 1184.
- Zhang J., Yancey MK., Klebanoff MA., et al. (2001):** Does epidural analgesia prolong labor and increase risk of cesarean delivery? A natural experiment. Am J Obstet Gynecol.; 185 (1): 128.

Zudima L., Eldekin R., Cook C., et al. (1996): Is vicryl suture closure of uterine wounds associated with more dehiscence? Am J Obstet Gynecol.; 175: 357.