
REFERENCES

- Allolio, H.; Ziegler, G. and Klotz, U. (1997):** Midazolam kinetics. Clinical Pharmacology and Therapeutics. 30: 653 – 661.
- American Psychiatric Association (2000):** Diagnostic and Statistical Manual of Mental Disorders Washington, DC: American Psychiatric Association. 465 - 8.
- Amrein, R. and Hetzel, W. (1990):** Pharmacology of Dormicum (midazolam) and Anexate (flumazenil). Acta Anaesthesiol Scand, 92: 6.
- Andrews, P.R. and Mark, L.C. (1982):** Structural specificity of barbiturates and related drugs. Anesthesiology, 57:314.
- Armstrong, P.J. and Bersten, A. (1986):** Normeperidine toxicity. Anesth. Analg., 65: 536.
- AstraZeneca, Neuroscience (2004):** Target Controlled Infusion in anesthetic practice.
- Bacon, R.; Chandrasekan, V. Haigh, A. et al (1994):** Early extubation after open-heart surgery with total Intravenous anaesthetic technique. Lancet 345: 133 - 4.
- Bail, J.M. (2000):** A technique for approximately maintaining constant plasma levels of intravenous drugs. Anesthesiology; 78: 116 - 23.
- Baker, J.; Baker, A. and Mazer, C. (1995):** Determinants of postoperative hypothermia after normothermic cardiopulmonary bypass. Journal of Cardiothoracic and Vascular Anesthesia 9: 154 - 157.

-
- Bravo, M.T.; Chadam, M.V. and Ronnenberg, W.C. (2003):** Inhibitory effects of propofol on cytochrome P450 activities in rat hepatic microsomes. *Anesthesia and Analgesia*, 76: 817-21.
- Brown, G.W.; Patel, N. and Ellen, K. (1991):** Comparison of propofol and thiopental for laryngeal mask insertion. *Anaesthesia*, 46: 771.
- Calhoun, C.D. and Oparlis (2004):** Hypertension crisis. *N.E. Jour. Med.*, 33: 1038.
- Carlson, B.X.; Hales, T.O. and Olsen, R.V.V. (1998):** GABA_A receptors and anesthesia In Yaksh TL, Lynch C III. Zapol WM et al (eds): *Anesthesia: Biologic Foundations*. New York. Lippincott-Raven, p 250.
- Chaudhri, S.; White, M. and Kenny, G.N. (1992):** Induction of anaesthesia with propofol using a target-controlled infusion system. *Anaesthesia*. 47: 551 - 3.
- Chaudhri, S.; White, M. and Kenny, G.N. (1992):** Induction of anesthesia with propofol using a target controlled infusion system. *Anesthesia* 47: 551-3.
- Checketts, M.R.; Gilhooly, C.J. and Kenny, G.N.C. (1998):** Patient-maintained analgesia with target controlled alfentanil infusion after cardiac surgery: a comparison with morphine PC A. *Br J Anesthesia*, 80: 748-751.
-

-
- Choi, S.D.; Spaulding, B.C. Gross, J.B. et al (1985):** Comparison of the ventilatory effects of etomidate and methohexital. *Anesthesiology*, 62: 442.
- Clements, J.A. and Nimmo, W.S. (1981):** The pharmacokinetics and analgesic effect of ketamine in man. *Br. J. Anaesth.*, 53: 27.
- Cockshott, I.D.; Douglas, E.J.; Prys-Roberts, C.; Turtle, M. and Coates, D.P. (1990):** The pharmacokinetics of propofol during and after intravenous infusion in man. *European Journal of Anesthesiology*, 7: 265-75.
- Cockshott, L.D. (1985):** Propofol pharmacokinetics and metabolism - an overview. *Postgraduate Medical Journal*, 61 (Suppl. 3): 45-50.
- Coetzee, J.F.; Glen, J.B.; Wium, C.A. and Boshoff, L. (1995):** Pharmacokinetic model selection for target controlled infusions of propofol, Assessment of three parameter sets. *Anesthesiology*. 82: 1328 - 45.
- Collier, C. and Kelly, K. (1991):** Propofol and convulsions: the evidence mounts. *Anaesth. Intensive. Care* 19: 573 - 575.
- Collins, R. and Peto, R. (1990):** Blood pressure, stroke and coronary heart disease, part 2, short term reductions in blood pressure: overview, randomized drug trials in their epidemiologic context. *Lancet*, 335: 8693.
-

-
- Cook, D.R. and Wingard, L.B. (2004):** Pharmacokinetics of succinylcholine in infants, children and adults. Clin. Pharmacol Ther., 20: 493 - 498.
- Croughwell, N.; ReVes, J.G. and Hawkins, E. (1988):** Cardiovascular changes after midazolam in patients with aortic stenosis: Effects of nitrous oxide. Anesth. Analg., 67: S1.
- Dean andropoulos, K.B. (2005):** Closed malpractice claims for awareness during anesthesia. ASA Newsl 60:14-7.
- Domino, K.B.; Posner, K.L.; Caplan, R.A. and Cheney, F.W. (1999):** Awareness during anesthesia: a closed analysis. Anesthesiology 9-0:1053-61.
- Doyle, P.W. and Matta, B.F. (1999):** Burst suppression or isoelectric encephalogram for cerebral protection: Evidence from metabolic suppression studies. Br. J. Anaesth., 83: 580-4.
- Dundee, J.W.; Fee, J.P.H.; Moore, J.; McIlroy, P.D.A. and Wilson, D.B. (1980):** Changes in serum enzyme levels following ketamine infusions. Anesthesia 35: 12-18.
- Edwards Moss, (2003):** T.I.V.A. and sedation for neurosurgery. In: B. Kay (ed.), T.I.V.A., Elsevier Science Publishers, B.V., P. 247-84.
- Egan T.D. and Lemmens, H.J.M.; Fiset P. et al (1993):** The pharmacokinetics of the new short-acting opioid remifentanyl (G187084B) in healthy adult male volunteers. Anesthesiology 79:881-92.
-

-
-
- Egan, T.D. (1995):** Remifentanil pharmacokinetics and pharmacodynamics. A preliminary appraisal. Clin. Pharmacol., 29: 80.
- Egan, T.D. (2003):** Target Controlled Drug Delivery: progress toward an intravenous vaporizer and Automated Anesthetic Administration. Anesthesiology, 99: 1214 - 9.
- Egan, T.D.; Minto, C.F.; Hermann, D.J. et al (1996):** Remifentanil versus alfentanil: Comparative pharmacokinetics and pharmacodynamics in healthy adult male volunteers. Anesthesiology. 85:695.
- Egan, T.D.; Minto, C.F.; Hermann, D.J.; Barr, J.; Muir, K.T. and Shafer, S.L. (1996):** Remifentanil Versus Alfentanil: Comparative Pharmacokinetics and Pharmacodynamics in Healthy Adult Male Volunteers. Anesthesiology; 84 (4): 821-833.
- Engbers F. (1998):** Target controlled infusion in practice. European J. of Anaesthesia, 12, suppl (10): 88 - 90.
- Engbers, F. (1998):** Practical use of Diprifusor systems. Anesthesia, 53 (Supp. 1): 28 - 34.
- Epstein, R.H.; Dushane, T.E. and Okum, G. (2002):** Should monitoring the amnestic state during general anesthesia is standard of care. Am. J. Anesth., 27: 506 – 177.
-
-

-
-
- Forster, A.; Gardaz, J.P.; Suler, P.M. et al (2000):** I.V. midazolam as an induction agent for anaesthesia: A study in volunteers. Br. J. Anaesth., 52: 907.
- Fragen, R.J. and Avram, M.J. (1998):** comparative pharmacology of drugs used for the induction of anesthesia. In Stoelting RR, Barash PG, Gallagher TJ (eds): Advances in anesthesia. Chicago Yearbook Medical Publishers, p. 303.
- Fragen, R.J.; Diprivan (propofol) (1988):** A historical perspective. Semin Anesth., 7:1.
- Frances, B.; Gout, R.; Monsarrat, B. et al (1992):** Further evidence that morphine-6-glucuronide is a more potent opioid agonist than morphine. J. Pharmacol. Exp. Ther., 262:25.
- Frederick, E.; Sieber, (2005):** Evaluation of the patient with endocrine disease or diabetes mellitus. In: 1 Principles and practice of Anaesthesiology, 2nd edition, David, E.; John, H.; G. Edward (eds.), Mosby, 303-322.
- Freye, E.; Latasch, L.; and Schmidhammer, H. (1992):** Pharmacodynamic effects of S-(+)-ketamine on EEG, evoked potentials and respiration: A study the awake dog. Anaesthesist 41: 527.
- Gepts, E. (1998):** Pharmacokinetic concepts for TCI anaesthesia. Anaesthesia 53 (Suppl 1): 4 - 12.
- Gepts, E. (2005):** Pharmacokinetic concepts for "TCI" anaesthesia. Anaesthesia, 1: 4 - 12.
-
-

-
- Gepts, E.; Camu, F.; Cockshott, I.D. and Douglas, E.J. (1987):** Disposition of propofol administered as constant rate intravenous infusions in humans. *Anesthesia and Analgesia*, 66: 1256 - 63.
- Gepts, E.; Camu, F.; Cockshott, I.D. and Douglas, E.J. (1987):** Disposition of propofol administered as constant rate intravenous infusions in humans. *Anesth. Analg.*, 66 Suppl (12):1256 - 63.
- Gepts, E.; Jonckheer, K.; Maes, V.; Sonck, W. and Camu, F. (1988):** Disposition kinetics of propofol during alfentanil anaesthesia. *Anaesthesia*, 43: suppl, 8-13.
- Gepts, E.; Shafer, S.L.; Tamil, F. et al (1995):** Linearity of Lees, N.W. pharmacokinetics and model estimation of sufentanil. *Anesthesiology*, 83: 1194.
- Ghoneim,, P.S.; Glen, J.B.; Kenny, G.N.; Schuttler, J. and Shafer, S. (1997):** Nomenclature for computer assisted infusion devices. *Anaesthesiology*, 86:1430 - 1.
- Gilman, A.G. et al (1990):** Goodman and Gilman's the pharmacological basis of therapeutics, 8th ed. Pergamon.
- Glen, J.B. (1991):** Quality of anesthesia during spontaneous respiration: a proposed scoring system. *Anesthesia*; 46 (12): 1081-2, 1991.
- Goldman, L. (1995):** Cardiac risk in non cardiac surgery: an update. *Anesth. Analg.* 80: 810.
-

-
- Goodman C.; Mc Ewan, A.I.; Jhaveri, R.; Wilkinson, M.; D.; Smith, L.R.; Canada, A.T. and Glass, P.S. (2002):** The interaction of fentanyl on the Cp50 of propofol for loss of consciousness and skin incision. *Anesthesiology*, 8 1suppl (4): 820 - 8.
- Gray, J.M. and Kenny, G.N.C. (1998):** Development of the technology for diprifusor "TCI" system. *Anaesthesia*, 53 Suppl (1): 22 - 25.
- Gray, J.M. and Kenny, G.N.C.:** Development of the technology for Diprifusor TCI systems. *Anesthesia*, 53 (Supp. 1): 22 - 27.
- Guedes, P.A.; Park, G.R.; Cockshott, I.D.; Douglas, E.J.; Shuker, B. and Simons, P.J. (2003):** Propofol metabolism in man during the anhepatic and reperfusion phases of liver transplantation. *Xenobiotica*; 22 (1): 105 -114.
- Hensley, R.J.; Thomson, I.R.; Burgess, P.M. and Rosenbloom, M. (2007):** Alfentanil pharmacokinetics in patients.
- Hentgen, T.K.; Avram, M.J and Krejci, T.C. (2002):** The relationship of age to the pharmacokinetic of early drug distribution: the concurrent disposition of thiopental and indocyanine green. *Anesthesiology*. 72: 403 - 411
- Ian, D.V. (2006):** [TIVA in pediatric neurosurgery] - *Minerva Anesthesiol* – 59 (11): 631 - 4.
- Jacobs, J.R. and Hughes, M.A.; Glass, P.S. (2005):** Context-sensitive half-time in multicompartment pharmacokinetic
-

models for intravenous anesthetic drugs. *Anesthesiology*; 76: 334-341.

Keimy, G.N.C.; McFadzean, W. Mantzaridis, H. and Fisher, A.C. (1993): Propofol requirements during closed-loop anesthesia, *Anesthesiology*, 79: A329.

Kenny, G.N. and White, M. (1992): A portable target controlled propofol infusion system. *International J. of Clinical Monitoring & Computing*. 9: 179-82

Kenny, G.N. and White, M.A. (1990): portable computerized infusion system for propofol. *Anaesthesia*, 8: 692 - 3.

Kenny, G.N.C. (2002): "Intravenous versus Inhalational Anesthesia". Published in Italy by Italian SIVA foundation congress. Roma.

Kenny, G.N.C. (2002): "Target Controlled versus Manual Infusion of IV Anesthesia". Published in Italy by Italian SIVA foundation congress, Roma.

Kissin, I. and Wright, A.G. (2000): The introduction of halothane: a Russian contribution to intravenous anaesthesia. *Anesthesiology*, 69: 242 - 5.

Klees, T.M.; Sheffels, P.; Thummel, K.E. and Kharasch, E.D. (2005): Pharmacogenetic Determinants of Human Liver Microsomal Alfentanil Metabolism and the Role of Cytochrome P450 3A5. *Anesthesiology*; 102: 550 - 6.

-
- Kruger-Thiemer, E. (1968):** Continuous intravenous infusion and multicompartiment accumulation. *Eur. J. Pharmacol.*, 4: 317-324.
- Larson, M.D.; Kurz, A.; Sessler, D.I. et al (1997):** Alfentanil blocks reflex pupillary dilation in response to noxious stimulation but does not diminish the light reflex. *Anesthesiology*, 87: 849.
- Larsson, J.E. Wahlstrom (1994):** Optimum rate of administration of propofol for induction of anaesthesia in rats. *Br. J. Anaesth.*, 73: 692 - 4.
- Lee, G. and Antognini, J.F. (1994):** Complete recovery after prolonged bypass for hyperkalemia, cardiac arrest. *Anaesth. Analg.*, 79: 172.
- Lees, N.W. and Gillies, G.W. (1996):** The effects of speed of injection on induction with propofol, a comparison with etomidate. *Anaesthesia*, 44 Suppl (5): 386 - 8.
- Lepage J-YM.; Pinaud, M.L.; Heltas, J.H. et al (2006):** Left ventricular function during propofol and fentanyl anesthesia in patients with coronary artery disease: assessment with a radionuclide [approach]. *Anesthesia and Analgesia* 67: 949-55.
- Lepage J-YM.; Pinaud, M.L.; Heltas, J.H.; Cozian, A.Y.; Le Nonnanc, T.Y. and Souron, R.J. (1991):** Left ventricular performance during propofol or methohexital anesthesia:
-

isotopic and invasive cardiac monitoring. *Anesthesia and Analgesia*. 73: 3-9.

Lerot, J.J.; Boulieu, R.; Foussadier, A.; George, M.; Clerc, J.; Villard, J.; Chassignolle, J.; Ferry, S. and Estanove, S. (1993): Comparison of the pharmacokinetics of methohexital during cardiac surgery with cardiopulmonary bypass and vascular surgery. *Journal of Cardiothoracic and Vascular Anesthesia*, 7: 30 - 34.

Levin, G.M and Brown, L.A. . (2004): Role of propofol in refractory status epilepticus. *Ann. Pharmacother*, 32: 1053-1059.

Levy, J.K. and Rockoff, M.A. (1982): Anaphylaxis to meperidine. *Anesth. Analg.* 61:301.

Lin, C.C.; Chen, M.F.; Chen, H.M. et al (1996): Propofol anesthesia in a patient with myasthenia gravis - a case report. *Acta Anaesthesiol. Sin.*, 34: 89 - 92.

Lodes, U. (1999): Total intravenous anesthesia (TIVA) and balanced anesthesia with short-acting anesthetics for ENT surgery in children. *Anaesthesiol. Reanim.*, 24 (1): 13 – 8.

Lubin, M.F. (2005): Is age a risk factor for surgery? *Med. Clin. North, amer.* 77: 327.

MacClaurin, S.C. (2002): Awareness under TIVA. *Anaesth. Intensive Care*, 30 (6): 816.

Mallon, J.S. and Edelist, G. (2002): Total intravenous anaesthesia (editorial). *Can. J. Anaesth.* 37: 279 - 81.

-
-
- Manberg, P.J.; Zraket, D.; Kovitch, L. and Christman, L. (2000):** Awareness during anesthesia with BIS monitoring. *Anesthesiology*, 93: A1371.
- Mangano, D.T. (1990):** Preoperative cardiac morbidity. *Anesthesiology*, 72: 153.
- Mann, R.; Blobner, M. and Jelen-Bsselborn, S. (2000):** Preanesthetic train-of-four fade predicts the atracurium requirement of myasthenia gravis patients. *Anesthesiology* 95: 346 - 350.
- Mark, C. Roger; John, H. Tinker; Benjamin, G. Covino; David, E. Longnecker:** Principles and practice of anesthesiology; volume 1: 1155-1175, 1993.
- McKenzie, A.J.; Cousin, K.G. and Pollock, N. (2004):** Propofol is a safe anaesthetic agent in malignant hyperthermia susceptible patients. *Anaesth. Intern. Care*, 20:165 - 8.
- Metz, S.; Slojgoff, S. and Keats, A.S. (1990):** Single dose thiopental compared to infusion for central nervous system protection during cardiopulmonary bypass. *Anesthesia and Analgesia*, 70: S266.
- Millar, J.M. and Jewkes, C.F. (1988):** Recovery and morbidity after day case anaesthesia, a comparison of propofol with thiopentone with and without alfentanil. *Anaesthesia*. 43 Suppl (9): 738 - 43.
-
-

-
-
- Miller, D.E.; Cucchiara, F.; Gerald, R.; Savarese, J.J. (2006):** Anaesthesia. volume 1,463.
- Miller, D.R. (1994):** Intravenous infusion anaesthesia and delivery devices. *Can. Anaesth.*, 41: 639 - 52.
- Miller, E. and Atul, R. (1998):** Evaluation of the hypertensive patients In: *Principles and practice anaesthesiology*, 2nd edition. David John. H.; G. Edwards (eds.), Mosby, PP. 157 - 67.
- Miller, L.G. (1991):** Chronic benaodiazepine administration: From the patient to the gene. *J. Clin. Pharmacol.*, 31: 492.
- Miller, M.D. (1994):** Muscle relaxants and clinical monitoring. In: *Anaesthesia*, 4th edition, R. Millef, M.D. (Ed.), Churchill Livingstone.
- Miller, R.D. and Upton, R.A. (1982):** Renal and biliary elimination vecuronium and pancuronium. *Anaesth. Analg.*, 61: 313.
- Murray, Feley, R.N.; Parley, P.S.; Hartnett, J.D.; Kent, G.M.; D.C. and Barre, P.E. (2004):** The impact of anemia on cardiomyopathy, morbidity and mortality in end-stage renal disease. *Am. J. Kidney Dis.*, 28: 53 - 61.
- Mychaskiw, J.K. and Bishop, V.A. (1987):** Sinus arrest after alfentanil (letter). *Nr. J. Anesth.*, 59: 390.
- Myron, C.F.; Schnider, T.W.; Egan, T.D.; Youngs, E.; Lemmens, H.J.M.; Gambus, P.L.; Billard, V.; Hoke, J.F.; Moore, K.H.P.; Hermann, D.J.; Muir, K.T.; Mandema, J.W. and**
-
-

-
- Shafer, S.L. (2000):** Influence of age and gender on the pharmacokinetics and pharmacodynamics of remifentanyl. I. Model development. *Anesthesiology*; 86 (1): 10-23.
- Nicolas, F.; Bruhn, J. et al (2006):** Conventional cardiopulmonary bypass, A physiological approach 10 years of experience at Marie-Lamelongue Hospital. *Perfusion* 9: 41 - 8.
- Nilsson, A.; Lee, P.F.S. and Revenas, B. (1993):** Midazolam as induction agent prior to inhalation anesthesia: a comparison of propofol, etomidate and midazolam. *Anesthesist* 42: 435 - 40.
- Nilsson, E.; Paloheimo, M.; Muller, K. et al (1989):** Halothane-induced variability in the neuromuscular transmission of patients with myasthenia gravis. *Acta. Anesth. Scand* 33: 395-401.
- Nilsson, L.B. (2002):** Remifentanyl vs. alfentanil for direct laryngoscopy: a randomized study comparing two total intravenous anaesthesia techniques. TIVA for direct laryngoscopy. *Acta Anaesthesiol. Belg.*, 53 (3): 213 -9.
- Ogawa, A.; Uemura, M.; Kataoka, Y. et al., (1992):** Effects of ketamine on cardiovascular responses mediated by N-methyl-D-aspartate receptor in the rat nucleus tractus solitarius. *Anesthesiology*. 78: 163.
- Osterman, J.E.; Hopper, J.; Heran, W.J.; Keane, T.M. and Van der Kolk B.A. (2001):** Awareness under anesthesia and the
-

-
- development of posttraumatic stress disorder. *Gen Hosp Psychmtry*. 23: 198 – 204.
- Pack, A.L. and Millman, R.P. (1993):** Change in control of ventilation awake and a sleep in the elderly. *J. Am.* 34: 533.
- Passot, S.; Servin, F.; Allary, R.; Pascal, J.; Prades, J.; Auboyer, C. and Molliex, S. (2002):** Target-Controlled Versus Manually-Controlled Infusion of Propofol for Direct Laryngoscopy and Bronchoscopy. *Anesth Analg*, 94: 1212 - 6.
- Peduto, V.A.; Mezzeth, D. et al. (2000):** Sevoflurane provided a better recovery than propofol-fentanyl anaesthesia for day case surgery. *Eur. J. Anaesthesol.*, 17: 138.
- Philip, B.K.; Scuderi, P.E.; Chung, F. et al., (1997):** Remifentanil compared with alfentanil for ambulatory surgery using total intravenous anesthesia. *Anesth. Analg.*, 84: 515 – 21.
- Philip, G. and Boysen (1998):** Evaluation of the patient with pulmonary disease. In "Principle and Practice of Anaesthesiology, 2nd Edition, David, John, H.; G. Edwards (Eds.) Mosby.
- Philips, A.S. and Gibson, P.M. (1993):** Propofol-fentanyl anaesthesia in cardiac patients. *Anaesthesia*, 48: 661 - 3.
- Pilotti, L. (1999):** Total intravenous anesthesia in thoracotomy with one-lung ventilation. *Minerva Anesthesiol.*, 65 (7-8): 483 – 9.
-

-
- Pitsiu, M.; Wilmer, A.; Bodenham, A.; Breen, D.; Bash, V.; Bonde, J., Kessler, P.; Albrecht, S.; Fisher, G. and Kirkham, A. (2004):** Pharmacokinetics of remifentanyl and its major metabolite, remifentanyl acid, in ICU patients with renal impairment. *British Journal of Anesthesia*, 92 (4): 493 - 503.
- Price, H. (1966):** Effects of carbon dioxide on cardiovascular system. *Anesthesiology*, 21: 652.
- Randall, C.; Wetzel, Lyime, G. Maxwell, (2007):** Anaesthesia for children. In: *Principles and practice of anaesthesiology*, 2nd (Ed.), David, E.; John, H.; G. Edwards (eds.), Mosby, 2079-2112.
- Ranta, S.; Jussila, J. and Hynynen, M. (1999):** Recall of awareness during cardiac anaesthesia: influence of feedback information to the anaesthesiologists. *Acta Anaesthesiol. Scand.* 40: 554 - 60.
- Ranta, S.; Laurila, R.; Saario, J.; Ali-Melkkila, T. and Kynynen, M. (1999):** Awareness with recall during general anesthesia: incidence and risk factors. *Anesth. Analg.*, 86: 1084 - 9.
- Ravussin, P. and Wilder-Smith, O.H.G. (2003):** Intravenous anaesthetics in neurosurgery. *Curr. Opin. Anaesthesiol.*, 9: 381 - 4.
- Razvi, H.A.; Fields, D.; Vargas, J.C. et al. (1996):** Oliguria during laparoscopic surgery: evidence for direct renal parenchymal compression as an etiological factor. *J. Endourol*, 10: 1.
-

- Reddy, R.V.; Moorthy, S.S.; Dierdorf, S.F. et al (1993):** Excitatory effects and electroencephalographic correlation of etomidate, thiopental, methohexital, and propofol. *Anesth. Analg.*, 77: 1008.
- Restall, J.; Tully, A.N.; Ward, P.J. and Kidd, A.G. (1988):** Total intravenous anaesthesia for military surgery: a technique using ketamine, midazolam and vecronium. *Anaesthesia*, 43: 46.
- Reves, J.G. and Green, M.N. (2002):** Anaesthesiaology and the academic medical centre: place and promise at the start of the new millennium. *Int. Anaesthesiol Clin.* 38.
- Reves, J.G. and Hawkins, E. (1990):** Esmolol for treatment of intraoperative tachycardia and/or hypertension in patients having cardiac operation: bolus loading technique. *J. Thorac. Cardiovasc. Surg.*, 100: 211.
- Reves, J.G.: Benzodiazepines. In Prys-Roberts C. and Hugg, C.C. (1984):** Pharmacokinetics of Anesthesia Boston. Blackwell Scientific Publications, p. 157.
- Reves, J.G.; Glass, P.S.A. and Lubarsky, D.A. (2000):** "Nonbarbiturate Intravenous Anesthesia" edited by Ronald D. Miller, *Anesthesia*, Fifth edition. Published in USA by Churchill Livingstone.

-
-
- Roizen, M.F.; Hiromichi, B.; Morihiro, O. et al. (2002):** The effects of prolonged low-flow sevoflurane anaesthesia on renal and hepatic functions. *Anesth. Analg.*, 91: 1262.
- Russell, D. (1998):** Intravenous anaesthesia, manual infusion schemes versus "TCI" system. *Anaesthesia*, 53, suppl (1): 42 - 45.
- Russell, D. (1998):** Practical aspects of Target-Controlled Infusion. *Anaesthesia Rounds 1998* edited by Zeneca Pharmaceuticals.
- Russell, D.; Wilkes, M.P.; Hunter, S.C; Glen, J.B.; Hutton, P. and Kenny, G.N.C. (1995):** Manual compared with target-controlled infusion of propofol. *British Journal of Anaesthesia*; 75 (5): 562 - 6.
- Sand, J.B.; Bailey, P.L.; Eat, K.A. et al (1990):** Differences in magnitude and duration of opioid induced respiratory-depression and analgesia with fentanyl and sufentanil. *Anesth. Analg.*, 70: 8.
- Schuttler, J.; Kloos, S.; Schwilden, H. and Stoeckel, H. (2000):** Total intravenous anaesthesia with propofol and alfentanil by computer-assisted infusion. *Anaesthesia*, 43 Supp (1): 2 - 7.
- Schuttler, J.; Schuttler, M.; Kloos, S.; Nadstawck, J. and Schwilden, H. (1991):** Total intravenous anesthesia with ketamine and propofol with optimized dosing strategies. *Anesthesist.*, 40: 199 - 204.
-
-

-
-
- Schuttler, J.; Schwilden, H. and Stockel, H. (1983):**
Pharmacokinetics as applied to total intravenous anesthesia:
practical implications. *Anesthesia*, 38: 53 - 56.
- Schuttler, J.; Schwilden, H. and Stoeckel, H. (1983):**
Pharmacokinetics as applied to total intravenous anaesthesia.
Anaesthesia, 38: 53 - 56.
- Schuttler, J.; Schwilden, H. and Stoeckel, H. (1983):**
Pharmacokinetics as applied to intravenous anaesthesia.
Anaesthesia. 38 (suppl): 53.
- Schuttler, J.; Schwilden, H. and Stoeckel, H. (1985):** Infusion
strategies to investigate the pharmacokinetics and
pharmacodynamics of hypnotic drugs: Etomidate as an
example. *Eur. J. Anesthesiol.*, 2: 133.
- Schuttler, J.; Stoeckel, H. and Schwilden, H. (1985):**
Pharmacokinetic and pharmacodynamic modeling of propofol
in volunteers and surgical patients. *Postgraduate Medical
Journal*, 61 (Suppl3): 53 - 5.
- Schuttler, J.; Wilms, M.; Stoeckel, H.; Schwilden H. and Lauven,
P.M. (1983):** Pharmacokinetic interaction of etomidate and
fentanyl. *Anesthesiology*. 59: A247.
- Seitz, W.; Lubbe, N.; Schaps, D.; Haverich, A. and Kirchner, E.
(2001):** Propofol for induction and maintenance of anesthesia
in cardiac surgery. Results of pharmacological studies in
humans. *Anaesthesist*. 40: 145 - 52.
-
-

-
- Servin, E. (1998):** TCI compared with a manually controlled infusion of propofol: a multi-centre study *Anaesthesia*. 53 (suppl 1): 82-6.
- Servin, F.; Marchand, M.F. and Desmonts, J.M. (1998):** Influence of analgesic supplementation on the target propofol concentrations for anaesthesia with diprifusor "TCP". *Anaesthesia*, 53 Suppl (1): 72.
- Shacelford, D.P. and Hoollman, M.K. (1995):** Evaluation of preoperative cardiac risk index values. *Am. J. Obstet. Gynaeco.*, 173 (1): 80 – 84.
- Shafer, A.; Doze, V.A.; Shafer, S.L. et al (1988):** Pharmacokinetics and pharmacodynamics of propofol infusions during general anesthesia. *Anesthesiology*. 69: 348.
- Shafer, A.; Sun, M.L.; White, P.F. (1986):** Pharmacokinetics and pharmacodynamics of alfentanil infusions during general anesthesia, *Anesth. Analg.*, 65: 1021.
- Shafer, S.L. and Varvel, J.R. (1991):** Pharmacokinetics, pharmacodynamics, and rational opioid selection. *Anesthesiology*. 74:53-63.
- Short, T.G.; Lim, T.A. and Tarn, Y.H. (1996):** Prospective evaluation of pharmacokinetic model-controlled infusion of propofol in adult patients. *British Journal of Anesthesia*, 76: 313 - 315.
-

-
- Smith, L. and Thwaites, A.J. (2003):** Target-controlled propofol vs. sevoflurane: a double-blind, randomized comparison in day-case anesthesia. *Anesthesia*, 54 (8): 745 - 752.
- Steven, T.M.; Ramsay, M.A.E.; Curran, J.P.J. et al. (2000):** Intravenous anaesthesia by infusion technique using alfaxalone/alfadone (Althesin). *Anaesthesia*, 30: 757.
- Swinhoe, C.F.; Peacock, J.E.; Glen, J.B. and Reilly, C.S. (1998):** Evaluation of the predictive performance of a diprifusor "TCI" system. *Anaesthesia*, 53 Suppl (1): 61 - 7.
- Swinhoe, C.F.; Peacock, J.E.; Glen, J.B. and Reilly, C. S. (1998):** Evaluation of the predictive performance of a Diprifusor TCI system. *Anesthesia*, 53 (Suppl. 1): 61 - 67.
- Tackley, R.M.; Lewis, G.T.; Prys Roberts, C.; Boaden, R.W.; Dixon, J. and Harvey, J.T. (1989):** Computer controlled infusion of propofol. *Br. J. of Anaesth.*, 62 Suppl (1): 46 - 53.
- Taylor, I.; White, M. and Kenny, G.N. (1993):** Assessment of the value and pattern of use of a target controlled propofol infusion system. *Int. J. Clinical Monitoring and Computing*. 10 Suppl (3): 175 - 80.
- Taylor, I.; White, M. and Kenny, G.N.C. (1993):** Assessment of the value and pattern of use of a target controlled propofol infusion system. *International Journal of Clinical Monitoring & Computing*, 10 (3): 175 - 80.
-

-
- Taylor, I.N. and Kenny, G.N.C. (1998):** Requirements for target controlled infusion of propofol to insert the laryngeal mask airway. *Anesthesia*, 53 (3): 222 - 226.
- Todd, M.M.; Drummond, J.C. U HS (1984):** The hemodynamic consequences of high-dose methohexital anesthesia in humans. *Anesthesiology*. 61: 495 - 501.
- Tramer, M.; Moore, A. and McQuay, H. (1997):** Meta-analytic comparison of prophylactic antiemetic efficacy for postoperative nausea and vomiting: propofol anaesthesia vs omitting nitrous oxide vs total i.v. anaesthesia with propofol. *Br. J. Anaesth.*, 79: 256 - 9.
- Trinder, T.J. (1998):** Propofol and alfentanil total intravenous anaesthesia: a comparison of techniques for major thoracic surgery. *Acta Anaesthesiol. Scand.*, 42 (4): 452 – 9.
- Van den Nieuwenhuyzen, M.C.; Engbers, F.H.; Burm, A.G. et al (1993):** Computer-controlled infusion of alfentanil for postoperative analgesia. A pharmacokinetic and pharmacodynamic evaluation. *Anesthesiology* 79:481-92.
- Van den Nieuwenhuyzen, M.C.O.; Engbers, F.H.M.; Burm, A.G.L.; Vletter, A.A.; Van Kleef, J.W. and Bovill, J.G. (1999):** Target controlled infusion of alfentanil for postoperative analgesia: contribution of plasma protein binding to intra-patient and inter-patient variability. *British Journal of Anesthesia*, 82 (4): 580 - 5.
-

-
-
- Van den Nieuwenhuyzen, M.C.O.; Engbers, F.H.M.; Vuyk, J. and Burm, A.G.L. (2000):** Target-Controlled Infusion Systems: Role in Anesthesia and Analgesia. Clin Pharmacokinet, 38 (2): 181 - 190.
- Van Drmeersch, Maddineni and Ciarke, R.J. (1993):** Neuromuscular effects and intubation conditions following mivacurium: a comparison with suxamethonium. Anaesthesia. 48: 940 - 45.
- Van keer Wiele, B.; Rubinstein, E.; Peacock, W. et al (1999):** Propylene glycol toxicity caused by prolonged infusion of etomidate. J. Neurosurg. Anesthesiol., 7: 259.
- Vernon, Aj.; Major, E.; Vernon, Aj.; Yate, P.M. and Waddell, T.K. (1982):** Disoprofol and fentanyl for total intravenous anaesthesia. Anaesthesia, 37: 541 - 7.
- Vincent, G.E.; Brandon Bravo, L.J. and Shafer, S.L. (2004):** Target Controlled Infusions: Targeting the Effect Site While Limiting Peak Plasma Concentration. IEEE Transactions on Biochemical Engineering. 51 (11): 1869 – 1875.
- Vinik, J.; Engbaek, J.; Eriksson, L.I.; et al (2006):** Good clinical research practice (GCRP) in pharmacodynamic studies of neuromuscular blocking agents. Acta. Anaesthesiol. Scand., 40: 59- 74.

-
- Viuent, H.R. and Bradley, E.L. (1998):** Triple anaesthetic combination: Propofol, midazolam-alfentanil. *Anaesth. Analg.*, 78:354
- Vuyk, J. (2002):** Pharmacokinetic and pharmacodynamic interactions between opioids and propofol. *J. Clin. Anesth.*, 9: 23 – 6.
- Wagner, J.G. (1974):** A safe method for rapidly achieving plasma concentration plateaus. *Clin. Pharmacol.*, 16: 691 - 700.
- Wang, M. (2000):** The psychological consequences of explicit and implicit memories of events during surgery. In: Ghoneim, MM, editors. *Awareness during anesthesia* Woburn, MA: Butterworth Heinemann; p. 146-9.
- Ward, D.S. (2002):** Pharmacodynamics and pharmacokinetics of propofol in a medium-chain triglyceride emulsion. *Anesthesiology*. 97 (6): 1401 – 8.
- White, M. and Kenny, G.N. (2005):** Intravenous propofol anaesthesia using a computerised infusion system. *Anaesthesia*. 45 Suppl (3): 204 - 9.
- White, M.; Kenny, G.N.C. (1990):** Intravenous propofol anesthesia using a computerized infusion system. *Anesthesia*; 45: 204 - 9.
- White, P.F. (1989):** Clinical uses of intravenous anesthetic and analgesic infusions. *Anesth. Analg.*, 68: 161 - 71.
- Whitford, P.F. (1997):** Intravenous anaesthesia. Baltimore: Williams and Wilkins.
-

- Wood, P.F.; Marietta, M.P.; Pudwil, C.R. et al (2002):** Effects of halothane anesthesia on the biodisposition of ketamine in rats. J. Pharmacol. Exp. Ther., 196: 545.
- Yu-hong, L.; Jian-hong, X.; Jian-jun, Y.; Jie, T. and Jian-guo, X. (2005):** Predictive performance of Diprifusor TCI system in patients during upper abdominal surgery under propofol/fentanyl anesthesia. Journal of Zhejiang University SCIENCE, 6B (1): 43 - 48.
- Zamacona, M.K.; Suarez, E.; Gracia, E.; Aguirre, C. and Calvo, R. (1998):** The significance of lipoproteins in serum binding variations of propofol. Anesth. Analg., 87: 1147 - 51.