

REFERENCES

- **Abraham, E.P., Chain, E. (1940):** An enzyme from bacteria able to destroy penicillin. *Nature* 146, 837.
- **Ahmed A.S, Chowdhury M.A and Hoque M. (2002):** Clinical and Bacteriological Profile of Neonatal Septicemia in a Tertiary level Pediatric Hospital in Bangladesh *Epidemiol.* 113(2):297-306.
- **Akbar D.H., Mushtaq M.A., El-Tahawi A.T. and Bahnasy A.A. (2000):** Staphylococcus aureus bacteremia. *Saudi Med J.*, 21(2): 171-174.
- **Al-Asmary S.M., Al-Helali N.S., Abdel-Fattah M.M., Al-Jabban T.M., and Al-Bamri A.M. (2004):** Nosocomial urinary tract infection. Risk factors, rates and trends. *Saudi Med. J.*, 25(7):895-900
- **Anderson RLT Biand LA, Favero MS, (1985):** Factors associated with Pseudomonas intrinsic contamination of commercial respiratory therapy solutions marketed as sterile. *Appl Environ Microbiol*:50:1343-1348.
- **Anwer S.K., Mustafa S., Pariyani S., Ashraf S. and Taufiq K.M. (2000):** Neonatal sepsis; an etiological study; *JPM* Mar; 50 (3): 91-114.
- **Arisoy E.S., Corrcra A., Scilheimer O.K. and Kaplan S.L. (1993):** Candida rugosa central venous catheter infection in a child. *Ped. Infect. Dis. J.*, 12 (II): 961-983.

REFERENCES

- **Arnow PM, Flaherty JP. (1999):** Nonfermentative gram-negative bacilli. In: Mayliall CG, Hospital epidemiology and infection control. Philadelphia: *Lippincott Williams & Wilkins*; 431-451.
- **Avery G.B. (2000):** Sepsis Neonatorum. In Avery, G.B., Fletcher M.A. I and Mhairi, G.M. (eds.): Neonatology. Pathophysiology and Management of the Newborn, 5 ed., Lippincott Company Philadelphia. 1088-1096.
- **Ayliffe G. (2000):** Minimal Access Therapy Decontamination Workin Group. Decontamination of minimally invasive surgical endoscopes and Hosp. Infect.,45(4):263-277.
- **Bauchner H, Brown E and Peskin J. (2002):** Premature graduates on the newborn intensive care unit: A guide to follow up. Pediatric clinics of North America, 35:1207-1221.
- **Bearman G.M., Munro C., Sessler C.N., and Wenzel R.P. (2006):** infection control and the prevention of nosocomial infections in the; intensive care unit. Semin Respir Crit. Care. Med.,27(3):310-324 .
- **Bert F., Clarisson J., Durand F., Delefosse D., Chauvet C, Lefebvre P., Lambert N., and Branger C. (2003):** Prevalence, molecular epidemiology and clinical significance of heterogenous glycopeptide-intermediate *staphylococcus aureus* in liver transplant recipients. J. Clin. Microb-ol. 41:5147-5152.
- **Bhultu Z.A. and Yusuf K. (1997):** Early onset neonatal sepsis in 'Pakistan: a case control study of risk factors in birth cohort. American J. Perinatol, 14(9):577-581.

REFERENCES

- **Black SR and Wcinstein RA. (2004):** The pathogens of Hospital infections. In gorbaeh SI.. Barllctt JG and Blacklow RN.: Infections diseases. 3rd ed. Lippincolt Williams, and Wilkins, Philadelphia., P. 90-107.
- **Bochicchio G.V., Joshi M., Shih D., Bochicchio K., Tracy K., and Scalea T.M. (2003):** Reclassification of urinary tract infections in critically ill trauma patients: a time-dependent analysis. Surg. Infect. (Larchmt), 4(4):379-385..
- **Bolding B. (2003):** Flash sterilization (steam). Nurs J.,21(1):31-33.
- **Bolyard E.A., Tablan O.C., Williams W.W., Pearson M.L., Shapiro C.N., and Deitchmanii S.D. (1998):** Guideline for infection control in healthcare personnel, Hospital Infection Control Practices Advisory Committee. Infect. Control Hosp. Epidemiol., 19(6):407-463..
- **Bondi, A., Dietz, C.C. (1945):** Penicillin resistant staphylococci. Proc. Soc. Exp. Biol. Med. 60 (1), 55-58.
- **Briedigkeit H and Gbel U. (1997):** Anaerobic bacteria as the cause of endogenous infections. Z Arztl Fortbild Qualitatssich. Mar; 91 (2) : 165 - 70.
- **Bonten M.J., Gaillard, C.A. and Vantiel F.H. (1994):** The stomach is not a source for colonization of the upper respiratory and pneumonia in ICU patients. Chest, 105(3):878-884.
- **Bonten MJM. Austin DJ. Lipstich M. (2001):** Understanding the spread of antibiotic resistant pathogens in hospitals: mathematical models as tools for control. Clin Infect Dis ; 33: 1739-1746.

REFERENCES

- **Boubaker K., Diebold P., Blanc D.S., Vandenesch F., Praz G., Dupuis G., and Triollet N. (2004):** Panton-Valentine Leucocidin and staphylococcal skin infections in school children. *EID*. 10:21-124.
- **Boussaud V., Parroiti A., Mayaud C. Wislez M., Antoine M., Picard C, Delisle F., Etienne J. and Cadranel J. (2003):** Life-threatening hemoptysis in adults with community-acquired pneumonia due to Panton-Valentine Leucocidin-secreting Intensive Care Med. 29:1840-1843.
- **Boyle-Vavra S. and Daum R.S. (2007):** Community-acquired methicillin- resistant *Staphylococcus aureus*. The role of Panton-Valentine Leucocidin. *Lab. Investig.*; 87:3-9.
- **Boyle-Vavra S., Berke S.K., Lee J.C., and Daum R.S. (2000):** Reversion of the glycopeptide resistance phenotype in *Staphylococcus aureus* clinical isolates. *Antimicrob. Agents Chemother.* 44:272-277.
- **Bressler AM., Williams T., Culler E.E., Zhu W., Lonsway D., Patel J.B., and Nolte F.S. (2005):** Correlation of penicillin binding protein a detection with oxacillin resistance in *Staphylococcus aureus* and discovery of a novel penicillin binding protein a mutation. *J. Clin. Microbiol*, 43:4541-4544.
- **Briedigkeit H. and Gobel (1997):** Anaerobic bacteria as the cause of endogenous infections *Z. Arztl fortbild Qualitatssich*; 91: 165-170.

REFERENCES

- **Brook J. (1995):** Baeteroides infections in children. 43: 92-98.
- **Brown PD and Ngeno C. (2007):** Antimicrobial resistance in clinical isolates of Staphylococcus aureus from hospital and community sources in southern Jamaica. Int J Infect Dis; 11(3):220-5.
- **Brumfitt W and Miller H. (1989):** Methicillin – resistant : Staphylococcus aureus. N Engl J Med; 320:1188-96.
- **Bucknall T.E. (1995):** Factors affecting the development of surgical wound infection: a surgeon view. J. Hosp. Inf.,(6):1-8.
- **Burke J.P. (2003):** Infection control . A problem for patient safety. N. gl Med., 348(7):651-666.
- **Buxton AE, Anderson RL, Werdegar D, et at. (1978):** Nosocomial respiratory tract infection and colonization with Acinetobacter calcoaceticus. Am J Med; 65:507-513.
- **Byers J.F., and Sole M.L. (2000):** Analysis of factors related to the development of ventilator-associated pneumonia: use of existing databases. Am. J. Crit. Care, 9(5):344-349.
- **Calvacante SS, Mota E, Silva L R, et al. (2006):** Risk Factors for Developing Nosocomial Infections Among Prdiatric Patients. Pediatr Infect Dis 1; 25: 438-445.
- **Campana S., Taccetti G., Ravenni N., Masi I., Audino S., Sisi B., Repetto T., Doring G., and de Martino M. (2004):** Molecular epidemiology of Pseudomonas aeruginosa, Burkholderia cepacia complex and methicillin-resistant Staphylococcus aureus in a cystic fibrosis center, J. Cyst. Fibros., 3(3):159-163.

REFERENCES

- **Carapeti E.A., Andrews S.M. and Bentley B.G. (1996):** Randomized study of sterile versus non sterile urethral catheterization. *Annals of the Royal College of Surgeons of England*, (78):59-60.
- **Cardo D.M., and Simone P.M. (2005):** Monitoring and preventing healthcare associated infections. The CDC approach. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz*, 48(9):1043-1047.
- **Carvalho AMC, Giovanetti MCY, Rabelo LFD, Contijo Fitho PP. (2001):** Hospital-acquired infections in pediatric patients: a prevalence in three Brazilian government hospitals. *J. Hosp. Infect*; 47: 71-72.
- **Carven D.E. (2006): Preventing ventilator associated pneumonia in adults: sowing seeds of change.** *Chest*, 130(1):251-260.
- **Center KJ, Reboli AC, Hubler R, Rodgers GL, Long SS. (2003):** Decreased vancomycin susceptibility of coagulase-negative staphylococci in neonatal intensive care unit: evidence of spread of staphylococci warmer. *J Clin Microbiol.* Oct; 41(10):4660-4665.
- **Centers for Disease Control (CDC) (2002):** Sterilization or Disinfection of Medical Devices General Principle <http://www.cdc.gov/ncidod/hip/sterile/sterilgp.htm>.

REFERENCES

- **Chain, E., Florey, H.W., Adelaide, M.B., Gardner, A.D., Heatley, N.G., Jennings, M.A., et al. (1993):** Penicillin as a chemotherapeutic agent. 1940. Clin. Orthop. Relat. Res. 295, 3-7.
- **Chambers H.F. (1997):** Methicillin resistance in staphylococci: Molecular and biochemical basis and clinical implications. Clinical Microbiology Reviews 10:781-791.
- **Chastred and Fagon J. Y. (2002):** Ventilator-associated pneumonia.) 'AM. J. Respir. Crit. Care Med, (165):867-903.
- **Choban P.S., Heckler R., Burge J.C., Flancbaum L., and Gross P.A. (1991):** Striving for benchmark infection rates: progress in control for patient mix. Am. J. Med., 91(3B)165-205.
- **Chongtrakool P., Boyle-Vavra S., Draum R.S., and Hiramatsu K. (2002):** Novel type of Staphylococcal cassette chromosome mec identified in community-acquired methicillin-resistant *Staphylococcus aureus* strains Antimicrobiol Agents chemother. 46:1147-1152.
- **Control (CDC) (1997,2000, 2003, 2004):**
- **Coredero L., Ayers LAV. and Dams K. (1997):** Neonatal airway colonization with Gram negative bacilli: association with severity of bronchopulmonary dysplasia. 16: 18-23.
- **Couch R.B., Englund J.A. and Whimbey E. (1997):** Respiratory vira infections in immunocompetent and immunocompromised persons. An . Med., 102(3A):2-9.
- **Cox R.A. and Conquest C. (1995):** A major outbreak of methicillin-resistant staphylococcus aureus caused by a new phage

REFERENCES

- type (EMRSA 16). J. Hosp. Infect., (29):87-106.
- **Craven D.E. (2006):** What is healthcare-associated pneumonia, and how should it be treated?. Curr. Opin. Infect.Dis.,19(2):153-160.
 - **Cribier B., Prevost G., Couppie P., Finck-Barbancon V., Crosshans E., and Piemont Y., (1992).** *Staphylococcus aureus* leucocidin : a new virulence factor in cutaneous infections .An epidemiological and experimental study . Dermatology 185:175-180.
 - **Cui L., MaX., Sato K., Okuma K., Tenover F.C., Mamizuka E.M. Gemmell C.G., Kim M.N., Ploy M.C., El Solh N., Ferraz V., and Hiramatsu K. (2003):** Cell wall thickening is a common feature of vancomycin resistance in *Staphylococcus aureus*. J. Clin. Microbiol. 41:5-14.
 - **Cui L., Tominaga E., Neoh H.IVL, and Hiramatsu K. (2006):** Correlation between reduced daptomycin susceptibility and vancomycin resistance in vancomycin-intermediate *Staphylococcus aureus*. Antimicrob. Agents Chemother. 50:1079-1082.
 - **Cui., Murakami H., Kuwahara-Arai K., Hanaki H., and Hiramatsu K. (2000):** Contribution of a thickened cell wall and its glutamine nonamidated component to the vancomycin resistance expressed to *Staphylococcus aureus*. Antimicrob. Agents Chemother. 44:2276-2285.

REFERENCES

- **DeKema D.J., Beekmann S.E. and Chapin K.C. (2003):** Epidemiology outcome of nosocomial and community-onset blood stream infection. *J. Clin. Microbial*, 3655-3660.
- **Deveikis A., Schauf V. and Mizen M. (1977):** Antimicrobial therapy of experimental group B-streptococcal infection in mice antimicrob. *Agents Chemother. Infect Med* 15(8): 811-817.
- **Dinnen M.U., Wetter L.A. and May A.R. (1990):** Urethral strictures and aortic surgery. Suprapubic rather than urethral catheters. *Eur. J. Vase. Surg*, 4: 535-538.
- **Doebblebling B.N., Richard P. and Wenzel R.P. (1995):** Nosocomial viral hepatitis and infections transmitted by blood and blood products. In "Principle and practice of infectious diseases". 4th ed. Belgium. Chapter 284, 2616-2632.
- **Doebbling B.N. and Weizel R.P. (1990):** The direct costs of universal precautions in teaching hospitals. *JAMA* 264: 2083-2087.
- **Dohowitz LG. (2003):** Pediatric infection control. In long SS, Pickering LK and Prober CO: *Principles and Practice of Pediatric Infectious Diseases*, 2nd ed. Churchill Livingstone, Philadelphia, p17-27.
- **Donnio P.Y., Oliviera D.C., Faria N.A., Wilhelm N., Le Coustumier A., and de Lencastre H. (2005):** partial excision of the chromosomal cassette containing the methicillin-resistance determinant results in methicillin-susceptible *Staphylococcus aureus* *J, Clin Microbiol.* 43:4191-4193.

REFERENCES

- **Dreyfuss D., and Mier L. (2001):** Aspiration pneumonia. N. Engl. J. Med., 344(24): 1868-1869.
- **Dufour P., Gillet Y., Bes M., Lina G., Vandenesch F., Floret D., Etienne J. and Ricket. (2002):** Community-acquired methicillin-resistant *Staphylococcus aureus* infections in France: Emergence of a single clone that produces Panton-Valentine Leucocidin. Clin. Infect. Dis. 35 :819-824.
- **Dumont and Ralston (1993):** Persistence of multiresistant clone of staphylococcus epidermidis in neonatal intensive care unit. Clin. Infect. Dis., 20 (1): 294-299.
- **Dumpis U, Kovalova Z., Jansons J., Cupane L., Sominskaya I., Michailova M., Karayiannis P., Gardovska D., Viazov S., Ross S., Roggendorf M. and Pumpens P. (2003):** An outbreak of HBV and HCV infection in a paediatric oncology ward: epidemiological investigation and prevention of further spread. J. Med. Virol., 69(3):331-338.
- **Eastick, Lceming J.P, Bennett D, Nillar M.K. (1996):** Reservoirs of coagulase negative staphylococci in preterm infants. J. Hosp. Infect 74: 99-104.
- **Edmond R., Richard P. and Wenzel P.R. (1995):** "Isolation" principles and practice of infectious diseases. 4th ed. Chapt, 279 2575-2579.
- **Eggimann P and Pittet D. (2001):** Infection control in the ICU. Chest; 120:2059-2093.

REFERENCES

- **Emori T.G. and Gaynes R.P. (1993):** An overview of nosocomial infections, including the role of the microbiology laboratory. Clin. Microbial. Rev., (6):428-442.
- **Eriksen, K.R., (1961):** “Celbenin”-resistant Staphylococci. Ugeskr. Laeger 123, 384-386.
- **Esteban J.I., Gomez J., Martell M., Cabot B., Quer J., Camps J., Gonzalez A., Otero T., Moya A. and Esteban R. (1996):** Transmission of hepatitis C virus by a cardiac surgeon. N. Engl. J. Med., 334(9):555-560
- **Evert R.J., Murdoch D.R. and Chambers S.T. (2000):** Nosocomial pneumonia in adults general medicine and surgical patients at Christ Church Hospital. NY. Med. J., (113):221-224.
- **Eyer S., Brummitt C, Crossley K., Siegel R. and Cerra F. (1990):** Catheter-related sepsis: prospective, randomized study of three methods of long-term catheter maintenance. Crit Care Med;18:1073-1079.
- **Fagon J.Y. and Chastre J. (2005):** Antimicrobial treatment of hospital-acquired pneumonia. Clin. Chest Med., 26(1):97-104.
- **Fanaroff A.A, Korones S.B and Wright L.L. (1998):** Incidence, presenting features, risk factors, and significance of late onset septicemia in very low birth weight infants. Pediatr infect Dis J.;17:593-598.
- **Fanaroff A.A, Korones S.B and Wright L.L. (1998):** Incidence, presenting features, risk factors, and significance Of late onset septicemia in very low birth weight infants. Pediatr Infect

REFERENCES

Dic J.; 17 : 593 - 598.

- **Fayon M.J, Tucci M. and Lacroix J. (1997):** Nosocomial pneumonia and tracheitis in a pediatric intensive care unit: a prospective study. *Am J Respir Crit Care Med.*; 155:162-169.
- **Fernanda C.Lessa . MD MPH.Jonathan R. Edward Ms scottk. Fridkin MD (2009)** trend in incidence of late onset MRSA infection in NICU pediatric infection disease journal volume 58 number 7
- **Fierobe L, Lucet JC, Deere D. (2001):** An outbreak of imipenem-resistant *Acinetobacter baumannii* in critically ill patients. *Infect Control Hosp Epidemiol*; 22:35-40.
- **File T.M. (2000):** The epidemiology of respiratory tract infections, in. *Respir. Infect.*, 15(3):184-194.
- **Finan J., Rosato A., Dickinson T., Ko D., and Archer G. (2002):** Conversion of oxacillin-resistant staphylococci from heterotypic to homotypic resistance expression. *Antimicrob. Agents Chemother.* 46:24-30.
- **Fisher, J.F., Meroueh, S.O., Mobashery, S., (2005):** Bacterial resistance to beta-lactam antibiotics: compelling opportunism, compelling opportunity. *Chem. Rev.* 105 (2), 395-424.
- **Ford-Jones E T, Mindorff C M, Tangly J M. (1989):** Epidemiology study of 4684 hospital-acquired infections in pediatric patients. *Pediatr Infect Dis J.* 1989; 8:668-675.
- **Foster G., Ross H.M., Huston R.A., and Collins M.U. (1997):** Catalase-positive cocci. In: *Manual of Clinical Microbiology.* 9th ed.

REFERENCES

- Murray P.R., Baron E.J., Jorgensen J.H., Pfaller M.A., Tenover F.C., Tenover F.C. ASM Press American Society for Microbiology 1752 NST. N.W. Washington, DC 20036-2904.
- **Fowler V G., Jr, Miro J.M., A , Cobell C.H., Abretyn E., Rubinstein E., Corey G.R., Spleman D., Bradely S.F., (2003):** Catalase-positive cocci. In: Manual of Clinical Microbiology. 9th ed. Murray P.R., Baron E.J., Jorgensen J.H., Pfaller M.A., Tenover F.C., Tenover F.C. ASM Press American Society for Microbiology 1752 NST. N.W. Washington, DC 20036-2904.
 - **Fridkin S.K. (2001):** Vancomycin-intermediate and -resistant *Staphylococcus aureus*: what the infectious disease specialist needs to know. Clin. Infect. Dis. 32:108-115.
 - **Fryklund B, Tullus K and Burman L.G. (1995):** Epidemiology and attack index of gram-negative bacteria causing invasive infection in three special-care neonatal units and risk factors for infection. Infection. Mar-Apr;23(2):76-80.
 - **Fuda, C.C., Fisher, J.F., Mobashery, S., (2005):** Beta-lactam resistance in *Staphylococcus aureus*: the adaptive resistance of a plastic genome. Cell. Mol. Life Sci. 62 (22), 2617-2633.
 - **Garcia- Garmandia JL, Ortiz-Leyba C, Garnacho-Montero J. (2001):** Risk factors for *Acinetobacter baumannii* nosocomial bacteremia in critically ill patients: a cohort study. Clin Infect Dis 2001; 33:939-946.
 - **Gardner P. and Goldman D. (1992):** Nosocomial infection in: Feigen R.D. & Cherry J.D. (eds): Textbook of ped. Infectious

REFERENCES

- diseases, 3rd W.B. Saunders Co.. 2145-2156.
- **Garibaldi R A., (1999):** Residential care in the elderly: the burden of infection. J Hosp Infect; 43(Suppl) :9-18.
 - **Garner J.S. (1993):** The CDC Hospital Infection Control Practices Advisory Committee. Am. J. Infect. Control, 21(3):160-162.
 - **Giraham OR, Hand JD. (1981):** Citrobacter diversus brain abscess and meningitis in neonates. JAMA;245:1923-1925.
 - **Goldman RC and Capobianco JO. (1990):** Role of an energy-dependent efflux pump in plasmid PNE24-mediated resistance to 14 and 15 membered macrolides in staphylococcus epidermidis. Antimicrob agents chemother; 34:1973-1980.
 - **Gonzalez B.E., Martinez-Aguillar G., Hulten RG., Hammerman W.A., Cross-Bu J., Avalos-Mishaan A., Mason E.O., Jr, and Kaplan S.L.(2005):** Severe Staphylococcal sepsis in adolescents in the era of community-acquired methicillin-resistant *Staphylococcus aureus*. Pediatrics. 115:642-648.
 - **Graves N. (2004):** Economics and preventing hospital-acquired infection, g. Infect. Dis, 10(4):561-566.
 - **Griffith SJ, Nathan C, Sdander RK, (1989):** The epidemiology pseudomonas aeruginosa in oncology patients in a general hospital. J Infect Dis; 160:1030 – 1036.
 - **Guerina, (1998):** Bacterial and fungal infection. In Cloherty. J.P. and stark, A.R.(eds): Mamual of Neonatal Care, 4th ed, Lipincolt Raven Publisher, Philadelphia-New-York..271-300.

REFERENCES

- **Haley R, Culver, While J, (1985):** The nationwide nosocomial infection rate; A new need for vital statistics. *Am. J. Epidemiol*; 121:159-167.
- **Haley R.W., Culver D.H. and White J.W. (1995):** The rational wide nosocomial infection rate: a new need for vital statistics. *Am. J. Epidemiol.* 121: 159-173.
- **Hamaki H and Hiramatsu K. (2001):** Deletion methods of glycopeptide-resistant staphylococcus aureus: susceptibility testing. In: Gillespie SH, ed. *Antibiotic resistance methods*. New Jersey, Humana Press; 85-92.
- **Hansen U., Hussain ML, Villone D., Herrmann M., Robeneck H., Peters G., Sinha B. and Bruckner P. (2006):** The anchorless adhesin Eap (Extracellular adherence protein) from *Staphylococcus aureus* selectively recognizes extracellular matrix aggregates but binds promiscuously to monomeric matrix macromolecules. *Matrix Biol*; 25:252-260.
- **Hantedz H. and Arnaouth. (2000):** Anaerobic infection in neonatal sepsis. *J. Hosp. Infect.*, 43(5):689-698.
- **Harding G.K., Nicolle L.E. and Ronbald A.R. (1991):** How long should catheter acquired urinary tract infection? A randomized controlled study. *Ann. Intern. Med.*, 114: 713-719.
- **Harris J.A. (1997):** Pediatric nosocomial infections, children are not 4jttle adults. *Infect. Control Hosp. Epidemiol.*, 18 (11); 739-742.

REFERENCES

- **Hemming MD, James E and Michael RB. (1976):** Nosocomial infection in a newborn intensive care unit. N. Eng. J. Med; 294:263-271.
- **Henderson D., Faward J. and Montgomerie J. (1995):** Hematogenous Candida endophthalmitis in patients receiving TPN. Infect, Dis., 143: 655-661.
- **Hiramastuk. (1998):** Thermer gene of staphylococcus aureus with reduced susceptibility to vancomycin. Am J Med; 104: 75-105.
- **Hoffman D.J. and Harris M.C. (1996):** Diagnosis of neonatal sepsis. In Spitzer, A.R. (editor). Intensive Care of Fetus and Neonates. Mex., (60):201-204.
- **Holden M.T.G., Feil E.J., Lindsay J.A., Peacock S.J., Day N.P.J., (2004):** Complete genomes of two clinical *Staphylococcus aureus* strains: Evidence for the rapid evolution of virulence and drug resistance JPNAS. 101:9786-9791.
- **Horbar J.D, Rogowski J. and Plsek P.E. (2001):** Collaborative quality improvement for neonatal intensive care. Pediatrics.; 07: 14-22.
- **Hosseini Masoumi ASI Ali reza – Nateghian (2009):** Iranian journal of clinical infections diseases : 4 (2) : 83-86.
- **Hugonnet S, Harbarth S, Sax H, Duncan RA and Pittet D. (2004):** Nursing resources: a major determinant of nosocomial infection? Curr Opin Infect Dis. Aug; 17(4):329-33.
- **Hushins W.C and Goldmann D.A. (1998):** Nosocomial infections. In: Feigen RD, Cherry JD, eds. Textbook of Pediatric

REFERENCES

- Infectious Disease. Philadelphia, PA: WB Saunders Co; 1998:2545-2585
- **Hussain M., Von Eiff C, Sinha B., Joost I., Hermann M., Peters G., and V Becker K. (2008):** Eap gene as novel target for specific identification of *Staphylococcus aureus*. J. Clin. Microbiol. 46: 470-476
 - **Inweregbu K. (2005):** Nosocomial infections. Continuing Education in Anaesthesia, Critical Care & Pain 2005 5(1):14-17.
 - **Issacs , Barfield C.P, Jrimwood K., Mcphee A.J, Minntillo C. and Tudehope (1995):** Systemic bacterial and fungal infections in infants in Australian Neonatal Units.Australian study group for neonatal infections. I J.Hosp. Infect, 42(2): 135-141 .
 - **Issartel B., Tristan,. A., Lechevallier S., Bruijere F., Lina G., Garin B., Lacossin F., Bes M., Vandenesch F., and Etienne J. (2005):** Frequent carriage of Panton-Valentine Leucocidin genes by *Staphylococcus aureus* isolates form surgically drained abscesses. J. Clin. Microbiol. 43:3203-3207.
 - **Ito T., Katayama Y., Asada K., Mori N., Tsutsumimoto K., Tiensasiton C. and Hiramatsu K. (2001):** Structural comparison of three types of Staphylococcal cassette chromosome mec integrated in the chromosome in methicillin-resistant *Staphylococcus aureus*. 45:1323-1336.
 - **Jacob A., Lohr M.D. and Stephen M.D. (1994):** "Hospital acquired urinary tract infection". The pediatric patient; a prospective study. Infect. Dis. J., 13 (I); 8-12.

REFERENCES

- **Jan mordie (2008):** Lab tests for staph, staph culture, staph infection virulence factors sec17 /ch190/ch191.
- **Jarvis WR, Munn VP, Highsmith AK, (1985):** The epidemiology of nosocomial infections caused by klebsiella pneumoniae. Infect Control; 6:68-74.
- **Jin T., Bokarewa M., Foster T., Mitchell J., Higgins J., and Turkowski A. (2004):** *Staphylococcus aureus* resists human defensins by production of Staphylokinase, a novel bacterial evasion mechanism. J. Immunol 172:1169-1176.
- **Jonn C.C. and Schreiber J.R. (2006):** Therapies and vaccines for emergiti bacterial infections: learning from methicillin-resistant *Staphylococcus aureus*. Pediatr. Clin. N. Am. 53:699-713.
- **J. Vandepetitte, K. Engback P. Rohner, P. Piot. C.C Heuck. (2003):** Basic laboratory procedures in clinical bacteriology 2nd edition; p. 20:40.
- **Kaneko J., Kimura T., Kawakami Y., Tomita T., and Kamio Y. (1997):** Panton-Valentine Leucocidin genes in a, phage-like particle isolated from mitomycin C-treated *Staphylococcus aureus* V8 (ATCC49775). Biosci. Biotechnol. Biochem. 61:1960-1962
- **Kaneko J., Kimura T., Narita S., Tomita T., and Kamio Y. (1998):** Complete nucleotide sequence and molecular characterization of the temperate staphylococcal bacteriophage Phi PVL carrying Panton-Valentine Leucocidin genes. Gene. 215: 57-67.

REFERENCES

- **Kaszanyitzky E.J., Jaosi S., Somogyi P., Dan A., Van der Graaf-Van Bloois L., Van Duijkeren E., and Wagenaar J.A. (2007):** MRSA transmission between cows and humans.EID.13 : 1-4.
- **Katayama Y., Ito T., and Hiramatsu K. (2001):** Genetic organization of the chromosome region surrounding *mecA* in clinical staphylococcal strains: Role of IS431-mediated *mecI* deletion in expression of resistance in *mecA*-can-ying, low-level methicillin-resistant *Staphylococcus haemolyticus*. Antimicrob. Agents Chemother. 45:1955-1963.
- **Kaufman. D and Fairchild K.D. (2004):** Clinical Microbiology of Bacterial and Fungal Sepsis in Very-Low-Birth-Weight Infants. Clin Microbiol Rev.; 17(3): 638-680.
- **Kawagoe JY, Conceicao AM, Segre, Cresio RP, Maria FS, Cardoso, Claudia VS and Julia TF. (2001):** Risk factors for nosocomial infection in critically ill newborns: 5-year prospective cohort study. Am J Infect. Control; 29: 109-114.
- **Kenneth todar, Ph. D. (2008):** Text book of bacteriology microbes, university of Wisconsin microbiology, bacteria vol 304 : p. 1421.
- **Keyserling H. L., Sinkowitz R. L., Cochran J. M., Harris G. L., Levine J. D., Siegel B. H., Stover S. A., Lau and Jarvis W. R. (2003):** Vancomycin use in Hospitalized Pediatric Patients Pediatrics, August 1, 112(2): 104 – 111.

REFERENCES

- **Kim H.B., Park W.B., Lee R.D., Choi YJ, Park S.W., Oh M.D., Kim EC? and Chue K.W. (2003):** Nationwide surveillance for *Staphylococcus aureus* with reduced susceptibility to vancomycin in Korea. J .Clin. Microbiol. 41:2279-2281.
- **King M.D., Humphrey B.J., Wang Y.F., Kourbatova E.V., Ray S.M., and Blumberg H.M. (2006):** Emergence of community-acquired methicillin-resistant *Staphylococcus aureus* USA 300 clone as the predominant cause of skin and soft-tissue infections. Am. Inter. Med. 144:309-317.
- **Kipp F., Becker K., Peters G., and Von Eiff C. (2004):** Evaluation of different methods to detect methicillin-resistance in small colony variants of *Staphylococcus aureus*. J. Clin. Microbiol.42 :1277-1279.
- **Kipp F., Kahl B.C., Becker K., Baron E.J., Proctor R.A., Peter G., and Von Eiff C. (2005):** Evaluation of two chromogenic agar media for recovery and identification of *Staphylococcus aureus* small-colony variants. J. Clin. Microbial 43:1956-1959.
- **Kirby, W.M.M., (1944):** Extraction of a highly potent penicillin inactivator from penicillin resistant *Staphylococci*. Science 99 (2579), 452-453.
- **Kline MW, Kaplan SL, Hawkins EP. (1988):** Pathogenesis of brain abscess formation in an infant rat model of *Citrobacter diversus* meningitis. J. Infect. Dis; 157: 106-112.
- **Koontz C.S., Chang M.C., and Meredith J.W. (2000):** Effects of empiric antibiotic administration for suspected pneumonia on

REFERENCES

- subsequent opportunistic pulmonary infections. *Am. Surg.*, 66(121:1110-1114.
- **Krcmery V, Sykora P, Trupl., (2001):** Antibiotic use and development of resistance in blood culture isolates: 8 years of experience from a cancer referral center. *.1 Chemother*; 13: 133 -142.
 - **Kropec A, Huebner J, Wursthorn M and Daschner FD. (1993):** In vitro activity of vancomycin and teicoplanin against *Staphylococcus aureus* and *Staphylococcus epidermidis* colonizing catheters. *Eur J Clin Microbiol Infect Dis.* Jul;12(7):545-8.
 - **Kumarasinghe G, Goh H and Tan KN. Hospital acquired infections in a Singapore Hospital: (1985-1992):** *Malays J Pathol.* Jun;17(1):17-21.
 - **Ladhani S, Joannou C.L, Locherie D.P., Evans R.W., and Poston S.M. (1999):** Catalase-positive cocci. In: *Manual of Clinical Microbiology*. 9th ed. Murray P.R., Baron E.J., Jorgensen J.H., Tenover F.C., Tenover P.C. ASM Press American Society for Microbiology 1752 NST. N.W. Washington DC 20036-2904.
 - **Langley J.M. (2005):** Defining urinary tract infection in the critically ill child *Pediatr. Crit. Care Med.*, 6(3 Suppl.):525-529.
 - **Langley J.M., hanakowski M. and Bjrtiussi K. (1994):** Demand for isolation beds in a pediatric hospital. *Am. J. Infect. Control.*, 22 (4); 207-211.
 - **Leaper D. (2004):** Nosocomial infection. *Br J Surg.* May; 91(5): 526-7.

REFERENCES

- **Lessa FC, Edwards JR, Fridkin SK, Tenover FC, Horan TC and Gorwitz RJ. (2009):** Trends in Incidence of Late-Onset Methicillin-Resistant *Staphylococcus aureus* Infection in Neonatal Intensive Care Units. Data from the National Nosocomial Infections Surveillance System, 1995-2004. *Pediatr Infect Dis J*; 28: 577-581.
- **Levy, S.B., Marshall, B., (2004):** Antibacterial resistance worldwide: causes, challenges and responses. *Nat. Med.* 10 (12 Suppl.), S122-S129.
- **Liedberg H., Lundeberg T. and Jane, M. (1990):** Silver alloy coated catheter reduce catheter associated bacteriuria. *Br. J. Urol.*, 65:379-381.
- **Lim W.S., and Macfarlane J.T. (2001):** Hospital acquired pneumonia. *Zlin.Med.*, 1(3):180-184.
- **Lina G., Piemond Y., Godail-Gamot F., Bes M., Peter M.O., Gauduchon V., Vandenesch F., Etienne J. (1999):** Involvement of Panton-Valentine Leucocidin producing *Staphylococcus aureus* in primary skin infections and pneumonia. *Clin. Infect. Dis.*29: 1128-1132.
- **Lipp A., and Edwards P. (2005):** Disposable surgical face masks. *Nurs.J.*, 23(3):20-1, 24-5, 33-38.
- **Lowy F.D. (1998):** The Staphylococci. In: *Medical Microbiology*. 23 Brooks G.F., Butel J.S., Morse S.A. McGraw-Hill Companies.
- **Lu PL., Chin L.C., Peng C.F., Chiang Y.H., Chen T.P., Ma L., and Siu L.K. (2005):** Risk factors and molecular analysis of

REFERENCES

- community methicillin-resistant *Staphylococcus aureus* carriage. J. Clin. Microbiol. 43:132-139.
- **Ma X., Ito T., Tiensasitorn E., Jamklang M., ChongtVakool P., Boyle-Vavra S., Draum R.S., and Hiramatsu K (2002):** Novel type of Staphylococcal cassette chromosome mec identified in community-acquired methicillin-resistant *Staphylococcus aureus* strains. 46: 1147-1152.
 - **Madeo, M. and Barlow, G. (2008):** Reducing blood-culture contamination rates by the use of a 2% chlorhexadine solution applicator in acute admission units Journal of Hospital Infection 69, 207-309.
 - **Maki D.G. (1992):** Infections due to infusion therapy In Bennet J.V. and Brachman, P.S. (eds). Hospital infection, edition Boston: Little brown 849-898.
 - **Maple PAC, Hamilton GMT, Brumfitt W. (1989):** World-wide antibiotic resistance in methicillin-resistant, *Staphylococcus aureus*. Lancet; 1:537-40.
 - **Markowitz N, Quinn EL, Saravolatz LD. (1992):** Trimethoprim-sulfamethoxazole compared with vancomycin for the treatment of *Staphylococcus aureus* infection. Ann Intern Med; 117: 390-8.
 - **Martin M, Dinges, Paul M, Orwin and Patrik M. (2009):** Exotoxins of *staphylococcus aureus* clinical microbiology review P. 16-34. 0893-8512.
 - **Massanari RM, Pfaller MA, Wakefield DS. (1988):** Implication of acquired oxacillin resistance in the management and control of

REFERENCES

Staphylococcus aureus infections. J Infect Dis; 158:702-9.

- **Massey R.C., Scriba T.J., Brown E.L., Phillips R.E., and Sewell A.I. (2007):** Use of peptide-major histocompatibility complex tetramer technology to study interactions between *Staphylococcus aureus* proteins and human cells. *Infect. Immun.* 75:5711-5715.
- **McClure J.A., Conly J.M., Lau V., El-Sayed S., Lowie T., Hutchins W., and Zhang K. (2006):** Novel multiplex PCR assay for detection of the staphylococcal virulence marker Panton-Valentine Leucocidin genes and simultaneous discrimination of methicillin-susceptible from resistant staphylococci. *J. Clin. Microbiol.* 44:1141-1144.
- **McDonald LC, Walker M, Carson L, (1998):** *Outbreak of Acinetobacter* spp. bloodstream infections in a nursery associated with contaminated aerosols and air conditioners. *Pediatr Infect Dis J*; 17:716-722.
- **Mcdougal L.K., Steward CD., Killgore G.E., Ckairam J.M., McAllister S.K., and Tenover FC. (2003):** Pulsed-field gel electrophoresis typing of oxacillin-resistant *Staphylococcus aureus* isolates from the United States: establishing a national database. *J. Clin. Microbiol* 41:5113-5120.
- **Mehrotra M.; Wang G. and Johnson W.M. (2000):** Catalase-positive cocci. In: *Manual of Clinical Microbiology*. 9th ed. Murray P.R., Baron E.J., Tenover J.C., Tenover F.C. ASM Press American Society for Microbiology 1752 N.W. Washington, DC 20036-2904.

REFERENCES

- **Mempel M., Lina G., Hojka M., Schopp C, Seidl P., Schafer T., Ring J., Vandenesch F., and Abeck D. (2003):** High prevalence of superantigens associated with the *egc* locus in *Staphylococcus aureus* isolates from patients with atopic eczema. Eur. J. Clin. Microbiol. Infect. Dis. 22:306-309.
- **Merighi A, Contato E, Scaglianrini R. (1996):** Quality improvement in gastrointestinal endoscopy : microbiologic surveillance of disinfection Gastrointest Endosc; 43:457-462.
- **Mermel L.A. (2000):** Prevention of intravascular catheter-related infections Ann Intern Med 2000;132:391-402.
- **Mermel L.A., Farr B.M and Sherertz R.J. (2001):** Guidelines for the management of urinary catheter-related infections. Infect Control Hosp Epidemiol . 22:222-242.
- **Merrerr J, Santoli F, Appcre-De Vecchi C. (2000):** Colonization pressure and risk of acqusition of methicillin-resistant *Staphylococcus ourcus* ht a medical intensive care unit. Infect Control I losp Kpidemiol ; 2 1:718-723.
- **Metchock J.E. (1995):** Changing pharyngeal flora of hospitalized patient. J. Hosp. Infect. 30 (Supp.): 272-282.
- **Miller L.G., Perdreau-Remington F., Rieg G., Mehdi S., Periroth J., Bayer A.S., Tang A.W., Phung T.O., and Spellberg B. (2005):** Necrotizing fasciitis caused by community-associated. methicillin-resistant *staphylococcus aurous* in Los Angeles. NEJM. 352: 1445-1453.

REFERENCES

- **Mimoz, O., Karim, A., Mercat, A. Cosseron, M., Falissard, B., Parker, F., Richard, C., Samii, K. and Nordmann, P. (1999):** Chlorhexidine compared with providone-iodine a skin preparation before blood culture *Annals of Internal Medicine* 131(11), 834-837.
- **Mitchell D.H., Swift G., and Gilbert G.L. (1999):** Surgical wound infection surveillance: the importance of infections that develop after hospital discharge. *Aust. J. Surg.*, 69(2): 117-120.
- **Moar Y., Rahav G., Belausov N., Ben-David D., Smollan G., and Keller N. (2007):** Prevalence and characteristics of heteroresistant vancomycin-intermediate *Staphylococcus aureus* bacteraemia in a tertiary care center. *J. Clin. Microbiol.* 45:1511-1514.
- **Mobley HLT, Chippendale GW, Tenney JH, (1988):** Hemagglutination of *Providencia stuartii* correlates with adherence to catheters and with persistence in catheter-associated bacteriuria. *J Infect Dis* 1988;157:264-271.
- **Moellering. QC. JR., (2009):** Recommended treatments for community acquired MRSA infections *Journal of American Academy of Family Physicians* 64 (7) : 1032 – 1037.
- **Mongko Trattanotliai KTT Boyle S., Kahana M.D., and Daum R.S. (2003):** Severe *Staphylococcus aureus* infections caused by clonally related community-acquired methicillin-susceptible and methicillin-resistant isolates. *Clin. Infect. Dis.* 37:1050-1058.

REFERENCES

- **Myianthefs P.M., Kalafati M., Samara I. and Baltopoulos G.J. (2004):** Nosocomial pneumonia. Crit. Care Nurse, 27(3):241-257.
- **Mylotte J.M and Tayara A. (2000):** Blood cultures: clinical aspects and controversies. Eur J Clin Microbiol Infect Dis;19: 157-163.
- **Narita S., Kaneko J., Chiba J., Piemont Y., Jarraud S., Etienne J., and Kamio Y. (2001):** Phage conversion of Panton-Valentine leucocidin in *Staphylococcus aureus* molecular analysis of a PVL converting phage, phiSLT. Gene. 268 : 195-206.
- **Neoh H.M., Cui L., Yuzawa H., Takenchi F., Matsuo M., and Hiramatsu K. (2007):** Mutated response-regulator graR is responsible for pherotypic conversion of *Staphylococcus aureus* from heterogenous vancomycin - intermediate resistance tovancomycin-intermediate resistance. Antimicrob. Agents Chemother.52:45-53.
- **Neu, H.C., (1992):** The crisis in antibiotic resistance. Science 257 (5073), 1064-1073.
- **Niven DJ, Laupland KB, Gregson DB and Church DL. (2009):** (The *S aureus* screening initiative group). Epidemiology of *Staphylococcus aureus* nasal colonization and influence on outcome in the critically ill.
- **Novick R.P., Schilevert P.,and Ruzin A. (2001):** The Staphlococci. In: Medical Microbiology. 23rd ed. Brook G.F., Butel J.s., Morse S.A. McGraw-Hill Companies.

REFERENCES

- **Nrtrita S., Kaneko J., Chiba J., Piemont Y., Jarraud S., Etienne J., and Kamio Y. (2001):** Phage conversion of Panton-Valentine leucocidin in *Staphylococcus aureus* molecular analysis of a PVL converting phage, phiSLT. *Gene*. 268 : 195-206.
- **O'Brien F.G., Price C, Grubb W.B., and Gustafson J.E. (2002):** *Antimicrob. Chemother.* 50: 313-321.
- **Ogrady N.P., Alexander M., Dellinger E.P. (2002):** Guidelines for the Prevention of Intravascular Catheter-Related; 51-5 Infections. *Pediatrics*;110:51-55.
- **Okuma K,Iwakawa K., Turnidge J.D., Grubb , Tenover F.C., Kapi M.. Ticensatoru C, Ito T., and Hiramatsu K. (2002):** Dissemination of new methicillin-resistant *Staphylococcus aureus* clones in the community. I. *Clin. Microbiol.* 40:4289-4294.
- **Oliveira G.A., Aquila A.M., Masiero R.A., Levy C, Gomes S.G., Cui L., Hiramatsu K., and Mamizuka E.M. (2001):** Isolation in Brazil of nosocomial *Staphylococcus aureus* with reduced susceptibility to vancomycin. *Infect. Control Hosp. Epidemiol* 22:443-448.
- **Orlan D., Arne O. and Cynthia K. (1995):** Accuracy of leukocyte indices and c-reactive protein for diagnosis of neonatal sepsis: a critical review. *Ped. Infect, dis.*, 14: 362-366.
- **Orrett FA and Land M. (2006):** Methicillin-resistant *Staphylococcus aureus* prevalence: Current susceptibility patterns in Trinidad. *BMC Infectious Diseases*, 6:83-88.

REFERENCES

- **Pai M., Kalantri S., Aggarwal A.N., Menzies D., and Blumberg H.M. (2006):** Nosocomial tuberculosis in India. *Emerg. Infect. Dis.*, 511-1318.
- **Palazzo I.C.V., Araujo M.L.C., and Darini A-L.C. (2005):** First report of vancomycin-resistant staphylococci isolated from healthy carriers in Brazil. *J Clin Microbiol.* 43:179-185.
- **Paradisi F., Corti G. and Mangani V. (1998):** Urosepsis in the critical "care unit. *Crit Care Clin* 14J65-180.
- **Parienti J.J., Thibon P., Heller R., Le Roux Y., von Theobald P., Bensadoun H., Bouvet A., Lemarchand F., and Le Coutour X. (2002):** Antisepsie Chirurgicale des mains Study Group. Hand-rubbing with an aqueous alcoholic solution traditional surgical hand-scrubbing and 30-day surgical site infection rates: a randomized equivalence study. *JAMA*, 288(6):722-727.
- **ParkerL. J. (1999):** Infection control 1: A practical guide to glove usage." *Br.J.NUTS.*, 8(7): 420-424.
- **Parry MF, Hutchinson JH, Brown NA, (1980):** Cram-negative sepsis in neonates: A nursery outbreak due lo hand carriage of *Citrobacter diversus*. *Pediatrics*; 65: 1105-1109.
- **Peacock S.J., Moore C.E., Justice A., Kantzanou M., Story L., Mackie K., O'Neill G., and Day N.P.J. (2002):** Virulent combinations of adhesion and toxin genes in natural populations of *Staphylococcus aureus* *Infect. Immun.* 70:4987-4996.
- **Peleg A.Y. and Munckhof W.J. (2004):** Fatal necrotizing pneumonia due to community-acquired methiillin-resistant

REFERENCES

- isolates. Clin. J. Aust. 181:228-229.
- **Pennington J.R. (1995):** Nosocomial respiratory infections. Principle; and practice of infectious diseases 4th ed. N. Engl. J. Med.,Chapter 282 2599-2607.
 - **Peter G. and Cashore W.J. (1995):** Infections acquired in the nursery : Epidemiology and control. Infectious diseases of the fetus and newborn infant Fourth edition, UK practices. J. Hosp. Infect., 48(3):180-185.
 - **Pierce G. and Joseph J.K. (1991):** Hospital acquired infections. In Harrison's principles of internal medicine. Vol. I (12th ed). LCDC. Health Canada. ,22:1188-4169.
 - **Pittet D., Harbarth S. and Ruef C. (1999):** Prevalence and risk factors for nosocomial infections in four university hospitals in Switzerland. Infect Control Hosp Epidemiol.; 20: 37-42.
 - **Pittet D., Hugonnet S. and Harbarth S. (2000):** Effectiveness of a hospital-wide programmed to improve compliance with hand hygiene. Infection Control Programmed. Lancet.;3 56:1307-1312.
 - **Plsek P.E. (1999):** Quality improvement methods in clinical medicine. Pediatrics. 103 (suppl):203-214.
 - **Powell K.R., Marcy S.M. and Jones K.T. (1994):** Laboratory aids for diagnosis of neonatal sepsis. In Remington J.S., Klein (ed). Infectious diseases of fetus and newborn infants. Philadelphia: W.B. Saunders. 4th ed.1223-1240.
 - **Pratt, (2007):** National Evidence Guidelines for preventing healthcare associated infections in NHS hospitals in England

REFERENCES

Journal of Hospital Infection 65 (1) S14

- **Qi W., Ender M., O'Brien F., Imhof A., Ruef C., McCallum N., and Berger-Bachi B. (2005):** Molecular epidemiology of methicillin-resistant *Staphylococcus aureus* in Zurich, Switzerland (2003): Prevalence epidemiology of methicillin-resistant *Staphylococcus aureus* in Zurich, Switzerland (2003): Prevalence of type IV SCC mec and a New SCC mec element associated with isolates from intravenous drug users. J. Clin. Microbiol 43:5164-5170.
- **Raad H., Hohn D.C. and Gilbreath B.J. (2000):** Prevention of central venous catheter-related infections by using maximal sterile barrier precautions during insertion. Infect Control Hosp Epidemiol.; 15:231-23.
- **Ramsy CM., Rcidy J. and Richard H. (1990):** Selective decontamination in intensive care practice. Intern. Care. Med,, 16:S217- 23.
- **Raygada JL and Levine DP. (2009):** Managing CA-MRSA Infections: Current and Emerging Options. Infect Med; 26:49-58.
- **Raymond J. and Aujard Y. (2002):** Nosocomial infections in pediatric patient. A European multicenter study group. Infect Control Hosp Epidemiol; 21(4): 260-263.
- **Rebollo MH, Bernal JM, Llorca J, Rabasa JM and Revuelta JM. (1996):** Nosocomial infections in patients having cardiovascular operations: a multivariate analysis of risk factors. J Thorac Cardiovasc Surg. Oct;112(4):908-13.

REFERENCES

- **Renzoni A., Francois P., Li D., Kelly W.L., Lew D.P., Vaudaux P., and Schrenzel J. (2004):** Modulation of fibronectin adhesins and other virulence factors in a teicoplanin-resistant derivative of methicillin-resistant *Staphylococcus aureus*. Antimicro. Agents chemother. 48 : 2958-2965.
- **Richard E.D. and George P.M. (1992):** Nosocomial infections In maxcy-Roscnan Public health and preventive medicine, 12ed by Appleton century-Crofls, Norwalk, Connecticut. Med. Bull., 73-74:123 -137.
- **Richards M.J., Edwards J.R., Culver D.H., and Gaynes R.P. (2000):** Nosocomial infections in combined medical-surgical intensive care units in the United States.infect. Control Hosp. Epidemiol, 21(8):510-515.
- **Robinson DA., and Enright M.C. (2004):** Multilocus sequence typing and the evolution of methicillin-resistant *Staphylococcus aureus*. Clin. Microbiol. Infect. 10:92-97.
- **Rojo D, Pinedo A., Clavijo E., Rodriguez G. A., and Garcia V. (1999):** Analysis of risk factors associated with nosocomial bacteraemias " Lancet, 1(8123):951-953.
- **Rossney A.S., Shore A.C., Morgan P.M., Fitzgibbon M.M., O'Connell B.,and Coleman D.C. (2007):** The emergence and importation of diverse genotypes of methicillin-resistant *Staphylococcus aureus* (MRSA) harboring the Panton-Valentine Leucocidin gene (PVL) reveal that PVL is a poor marker for community-acquired MRSA strains in Ireland. J. Clin. Microbiol.

REFERENCES

45:2554-2563.

- **Rubin, P. J. Sanchez, J. Siegel, G. Levine, L. Saiman, and W. R. Jarvis. (2002):** Evaluation and Treatment of Neonates With Suspected Late-Onset Sepsis: A Survey of Neonatologists' Practices Pediatrics; 110(4): 42-45.
- **Ruiz M.E., Guerrero I.C., and Tuazon C.U. (2002):** Endocarditis caused by methicillin-resistant *Staphylococcus aureus*: Treatment failure with linezolid. Catalase-positive cocci. In: Manual of Clinical Microbiology. 9th ed. Murray P.R., Baron E.J., Jorgensen J.H., Tenover F.C., Tenover P.C. ASM Press American Society for Microbiology 1752 N.W. Washington, DC 20036-2904.
- **Rutala W.A., and Weber D.J. (2004):** Disinfection and sterilization in health care facilities: what clinicians need to know. Clin. Infect. Dis., 39(5): 702-709.
- **Sadoyama G and Gontijo Filho PP.(2003):** Comparison between the jugular and subclavian vein as insertion site for central venous catheters: microbiological aspects and risk factors for colonization and infection. Braz J Infect Dis.;7(2):142-148.
- **Said-Salim B., Mathema B., Braughton K., Davis S., Sinsimer D., Eisner W., Likhoshvay Y., Deteo and Krciswirth B.N. (2005):** Differential distribution and expression of Panton - Valentine Crobiol Leucocidin among community - acquired methicillin - resistant *Staphylococcus aureus* strains. J. Clin. Microbiol. 43:3373-3379.

REFERENCES

- **Saiman L, Luirigton E. and Pfaller M. (2000):** Risk factors for bacteraemia in neonatal intensive care unit patients. The National Epidemiology of Bacterial Survey study group. *Pediatr Infect DisJ.*;19:319-324.
- **Savolainen K., Paulin L., Westerlund-Wikstrom B., Foster T.J., Korhonen T.K., and Kuusela P. (2001):** Expression of *pls*, a gene closely associated with the *mecA* gene of methicillin-resistant *Staphylococcus aureus*, prevents bacterial adhesion in vitro. *Infect. Immun.* 69:3013-3020.
- **Seifert H., Wisplinghoff H., Schnabel P., and von Eiff C. (2003):** Small colony variants (SCVs) of *Staphylococcus aureus* as a cause of acemaker-related infection. *EID.* 9:1316-1318.
- **Severin A, Wu S.W., Tabei K., and Tomasz A. (2004):** Penicillin-binding protein is essential for expression of high-level vancomycin resistance and cell wall synthesis in vancomycin-resistant *Staphylococcus aureus* carrying the enterococcal *vanA* gene complex. *Antimicrob. Agent Chemother.* 48:4566-4573.
- **Sherertz RJ, Sullivan ML. (1985):** An outbreak of infections with *Acinetobacter calcoaceticus* in burn patients: contamination of patients' mattresses. *Infect Dis*, 151-.252-258.
- **Shore A., Rossney A.S., Keane C, Enright M C, and Coleman D.C. P. (2005):** Seven novel variants of the staphylococcal chromosomal cassette *mec* in methicillin-resistant *Staphylococcal aureus* isolates from Ireland. *Anitimicrob. Agents Chemother.* 49:2070-2083.

REFERENCES

- **Showsh S.A., De Boever E.H., and Clewell D.B. (2001):** S vancomycin- resistant plasmid in *Enterococcus faecalis* that encodes sensitivity to a sex pheromone also produced by *Staphylococcus aureus*. *Antimicrob Agents Chemother.* 45 : 2177-2178.
- **Siegel J.,Rhinehart E., Jackson M., Chiarello L. the healthcare Infection Control Practices Advisory Committee. (2006):** Management of multidrug-resistant organisms in healthcare setting. DC.<http://www.cdc.gov/ncidod/dhqp/pdf/ar/mdroGuideline2006.pdf>
- **Sieradzki K., LeskiTVDick J., Borio LaBd-TomaSz A. (2003):** Evolution of a vancomycin-intermediate *Staphylococcus aureus* strain in vivo: multiple changes in the antibiotic resistance phenotypes of a single lineage of methicillin-rjsistant 5". *aureus* under the impact of antibiotics administered for chemotherapy. *J. Clin. Microbil.* 41: 1687-1693.
- **Silva HA, Pereira EM, Schuenck RB, Pinto RCM, Abdallah VOS, Santos KRN and Gontijo-Filho PP. (2009):** Molecular surveillance of methicillin susceptible *Staphylococcus aureus* at a neonatal intensive care unit in Brazil. *Am J Infect Control*; 36; article in press.
- **Sinclair R.C., Maxfield A., Marks E.L., Thompson D.R., and Gershon R.R. (2002):** Prevalence of safer needle devices and factors associated with their adoption: results of a national hospital survey. *Public Health Rep.*, 117(4)340-349.

REFERENCES

- **Skinner, D., Keefer, C.S., (1941):** Significance of bacteremia caused by *Staphylococcus aureus*. Arch. Intern. Med. 68, 851-875.
- **Skow A., Mangold A., Tajuddin M., Huntington A., Fritz B., Thomson RB.. Jr, and Kaul K.L. (2005):** Species level identification of *Staphylococcus* isolates by real-time PCR and melt curve analysis. J. Clin. Microbiol. 43:2876-2880.
- **Sobke A.C., Selimovic D., Orlova V., Hassan M., Chavakis T., Athanasopoulos A.N., Schubert V., Hussain M., (2006):** The extracellular adherence protein from *Staphylococcus aureus* abrogates angiogenic responses of endothelial cells by blocking Ras activation. FASEBJ. 20:2621-2623.
- **Solnim A.D, Kurtines H.C, Sprague B.M. and Singh N. (2001):** The associated with nosocomial bloodstream infections in the pediatric intensive care unit. Pediatr Crit Care Med; 2(2): 170-174.
- **Speller D. and Humphreys H. (1998):** Hospital-acquired infections In and Wilson's Microbiology and Microbial Infections, Edited by L Collier, A Balowsy and Y Mussman, 9th edition. Great Britain, P. 188-229, P. 188-229.
- **Spelman D.W. (2002):** Hospital acquired infections. Med. J. Aust, 176(6): 286-291.
- **Stamm, W.E. (1997):** Catheter associated urinary tract infection epidemiology, pathogenesis and prevention. Am. J. Med. 91 .(Suppl. 3D): 655-715.
- **Stephenson J. (1998):** Emerging infections on center stage at first major international meeting. JAMA, 279(14):1055-1056.

REFERENCES

- **Stoll, Hansen N, Fanaroff A. A, Wright L. L, Carlo W. A, Ehrenkranz R. A, Lemons J. A, Donovan E. F, Laptook A. R, Stevenson D. K , Papile L. A. and Poole W. K. (2002):** Late-Onset Sepsis in Very Low Birth Weight Neonates: The Experience of the NICHD Neonatal Research Network Pediatrics; 110(2): 285 – 291.
- **Susanj. Rehm. M.D. (2008):** Staphylococcus aureus : the new adventures of legendary pathogen Cleveland clinical journal of medicine volume 75 number 3.
- **Takano T., Higuchi W., Otsuka T., Baranovich T., Enany S., Saito K., Isobe H., Dohmae S., Ozaki K., Takano M., Iwao Y., Shibuya M., Okubo T., Yabe S., Shi D., Reva I., Teng L.J., and Yamamoto T. (2008):** Novel characteristics of community-acquired methicillin-resistant *Staphylococcus aureus* strains belonging to multilocus sequence type 59 in Taiwan. Antimicrob. Agents Chemother.52: 837-845.
- **Tambyah P.A, Halvorson K.T and Maki D.G. (1999):** A prospective study of pathogenesis of catheter-associated urinary tract infections. Mayo ClinProc 74,131-136.
- **Tokars J.I, Cookson S.T, McArthur M.A, Boyer C.L, McGeer A.J. and Jarvis W.R. (1999):** Prospective evaluation of risk factors for bloodstream infection in patients receiving home infusion therapy; 131 :340 -347.
- **Trindale P.A., Pacheco R.L., Costa S.F., Rossi F., Barone A.A., Mamizuka E.M., and Levin A.S. (2005):** Prevalence of SCC mec

REFERENCES

- type IV nosocomial blood stream isolates of methicillin-resistant *staphylococcus aureus*. J. Clin. Microbiol. 43:3435-3437.
- **Ultman RF, Gurevich I, Schock PE and Cumnha BA. (2002):** The role of understaffing and over crowding in recurrent out breaks of staph. Infection in pediatric special care area. J. infect dis; 145: 845-880.
 - **Umesh S Kamat, Agnelo, Ferrira Dilip Amonkar, Dilip D Motghare Manoj, Kulkarni. (2009):** Epidemiology of hospital acquired infection in a medical college hospital in goa issue 1 volume 28 page 76-80.
 - **Urrea M., Pons M., Serra ML, Latorre C. and Palomeque A. (2003):** Prospective incidence study of nosocomial infections in a pediatric intensive care unit. Pediatr Infect Dis J.;22(6):490-494.
 - **Vandenesch F., Naim T., Enright M.C., Lina G., Nimmo G.R., Henernan H., Liassine N., BesM., Green land T., Reverdy M.E., and Etienne J. (2003):** Community – acquired methicilli- resistant *Staphylococcus aureus* carrying panton – Valentine Leucocidin genes:Word Wide emergence EID. 9: 978-984.
 - **Vaudaux P., Francois P., Lew D.P., and Waldvogel F.A. (2000):** Host factors predisposing to and influencing therapy of foreign body infections, p. 1-26. In: Waldvogel F.A. and Bisno A.L. (ed.), Infections associated with indwelling medical devices. American Society for Microbiology, Washington. D.C.

REFERENCES

- **Viglione A, Nottebart VF, Bodman HA and Platt R. (1991):** Recurrent group A streptococcal carriage in a health care worker associated with widely separated nosocomial outbreaks. *Am J Med.* Sep 16;91(3B):329S-333S.
- **Vincent JL, Chierego M, Struelens M and Byl B. (2004):** Infection control in the intensive care unit. *Expert Rev Anti Infect Ther.* Oct; 2(5):795-805.
- **Waldvogel F.A. (2000):** The Staphylococci. In: *Medical Microbiology*. 23rd ed. Brooks G.F., Butel J.S., Morse S.A. McGraw-Hill Companies.
- **Weber DJ, Raasch R and Rutala WA. (1999):** Nosocomial infections in the ICU: the growing importance of antibiotic-resistant pathogens. *Chest.* Mar;115(3 Suppl):34S-41S.
- **Webster, (2009):** Webster's new world 2009 nosocomial infection definition in medical dictionary third edition 480 pages.
- **Weernink A, Severin WPJ, Tjernberg I, (1995):** Pillows, an unexpected source of *Acinetobacter*. *J Hosp Infect*; 29:189-199.
- **Wendt C, Dietze B, Dietz E, (1997):** Survival of *Acinetobacter baumannii* on dry surfaces. *J Clin Microbiol* ; 35:1394-1397.
- **Wenzel RP and Edmond MB. (2006):** Team-based prevention of catheter-related infections. *N Engl J Med.* Dec 28;355(26):2781-3.
- **Wilbrandt B, Wischniewski N, Geffers C, Daschner F, Rüdén H AND Gastmeier P. (2001):** Reduction of nosocomial infections by public health quality management in surgical patients. 2: No reduction in low initial infection rate. *Zentralbl Chir.*

REFERENCES

- Sep;126(9):691-5.
- **Wilcox MH, Fawley WN. (2000):** Hospital disinfectants and spore formation by *Clostridium difficile*. Lancet: 356: 2098-2099.
 - **Wooton M., MacGowan A.P., Walsh T.R., and Robin R.A. (2007):** A multicenter study evaluating the current strategies for isolating *Staphylococcus aureus* with reduced susceptibility to glycopeptides. J. Clin. Microbiol. 45: 329-332.
 - **Yamasaki O., Kaneko J., Morizane S., Akiyama H., Aratta J., Narita S., Chiba J., Kamio Y., and Iwatsuki K. (2005):** The association - between *Staphylococcus aureus* strains carrying Panton-Valentine Leucocidin genes and the development of deep-seated follicular infection. Clin. Infect. Dis. 40:381-385.
 - **Yologlu S, Durmaz B and Bayindir Y. (2003):** Nosocomial infections and risk factors in intensive care units. New Microbiol. Jul; 26(3):299-303.
 - **Zhang K., McClure J., Elsayed S., Louie T. and Conly J. (2005):** Novel multiplex PCR assay for characterization and subtyping of staphylococcal cassette chromosome mec types I to V in methicillin-resistant *Staphylococcus aureus*. J. Clin. Microbiol. 43:5026-5033 .
 - **Zimakoff J, Hoiby N, Rosendal K, (1983):** Epidemiology of *Pseudomonas aeruginosa* infection and the role of contamination in a cystic fibrosis clinic. . J Hosp Infect 1983; 4:31-40.

This document was created with Win2PDF available at <http://www.win2pdf.com>.
The unregistered version of Win2PDF is for evaluation or non-commercial use only.
This page will not be added after purchasing Win2PDF.