

References

AAP Subcommittee on Hyperbilirubinemia (2004): Clinical practice guideline. Management of hyperbilirubinemia in the newborn infant 35 or more weeks of gestation. Pediatrics, 114: 297-316.

Adélia Jehan Nasser Bernaldo, and Conceicao a parecida de Matoos Segre (2004): Bilirubin dosage in cord blood: could it predict neonatal hyperbilirubinemia. Saopaulo Med. J Vol, 122, No. 3 Sao Paulo.

Agarwal R, Kaushal M, Aggarwal R, Paul VK and Deorari AK (2002): Early neonatal hyperbilirubinemia using first day serum bilirubin level. Indian Ped: 39: 724-30.

Akaba K, Kimura T, Sasaki A, Tanabe S, Ikegami T, Hashimoto M, Umeda H, Yoshida H, Umetsu K, Chiba H, Yuasa I and Hayasaka K (1998): Neonatal hyperbilirubinemia and mutation of the bilirubin uridine diphosphate glucuronosyltransferase gene: a common missense mutation among Japanese, Koreans and Chinese. Biochem Mol Biol Int; 46 (1): 21-6.

Alcock GS and Liley H (2003): Immunoglobulin infusion for isoimmune haemolytic jaundice in neonates. (Cochrane Review). In: The Cochrane Library, Issue 2. Oxford: Update Software.

Alcock GS and Liley H (2002): Immunoglobulin infusion for isoimmune hemolytic jaundice in neonates. Cochrane-Database – Syst – Rev; (3): 1469-1493.

Alpay FA, Sarici SU, Tosuncuk HD, Serdar MA, Inanc N and Gokcay E (2000): The value of the first day bilirubin measurement in predicting the development of significant hyperbilirubinaemia in healthy term newborns. *Peadiatrics*: 106: E16.

American Academy of Pediatrics (1994): Committee on Quality Improvement and Subcommittee on Hyperbilirubinemia. Practice parameter: management of hyperbilirubinemia in the healthy term newborn. *Pediatrics*; 94: 558-65.

Aono S, Yamada Y and Keino H (1994): A new type of defect in the gene for bilirubin uridine 5-diphosphate glucuronosyl transferase syndrome type 1. *Pediatr. Res*; 35: 629-39.

Bhutani VK, Johnson L and Sivieri EM (1999): Predictive ability of a predischarge hour-specific serum bilirubin for subsequent significant hyperbilirubinemia in healthy term and near-term newborns. *Pediatrics*, 103: 6-14.

Bhutani VK, Johnson L and Sivieri EM (2000): Predictive ability of a predischarge hour specific bilirubin for subsequent

significant hyperbilirubinemia in healthy term and near term newborns. *Pediatrics*; 103: 6-14.

Bhutani VK, and Johnson LH (2000): Managing the assessment of neonatal jaundice: importance of timing. *Indian J Pediatr*; 67: 733-7.

Bhutani VK, Johnson LH, and Stevenson DK (on behalf of the Jaundice Multi National Study Group) (2001): prospective pre-discharge (30 ± 6 hours) diagnostic evaluation strategies for severe neonatal hyperbilirubinemia. *J Perinatol*; 21: 488-9.

Bhutani VK, and Johnson LH (2003): Newborn jaundice and kernicterus health and social perspectives. *Indian J Pediatrics*; 70: 407-16.

Blackmon LR et al. (2004): Research on prevention of bilirubin induced brain injury and kernicterus: national institute of child health and human development conference summary. *Pediatrics*, volume 114, number 1, pages 229-233.

Boo NY and Lee HT (2002): Randomized controlled trial of oral versus intravenous fluid supplementation on serum bilirubin level during phototherapy of term infants with

severe hyperbilirubinemia. J Paediatr Child Health; 38 (2): 151-5.

Braveman P, Egerter S, Pearl M, Marchi K, and Miller C (1995): Problems associated with early discharge of newborn infants: early discharge of newborns and mothers: a critical review of the literature. Pediatrics; 96: 716-726.

Britton JR, Britton HL and Beebe SA (1994): Early discharge of the term newborn: a continued dilemma. Pediatrics; 94: 291-295.

Burtis CA, Ashwood ER, Joan EA and Ldrich MS (1996): Tietz fundamentals of clinical chemistry; 4th edition. W.B. Saunders Company Biochemical assessment of liver function; 543-48.

Brown AK, and Johnson L (1996): Loss of concern about jaundice and the reemergence of kernicterus in full term infants in the era of managed care. In: Fanaroff AA, Klaus MH, editors. Yearbook of neonatal and perinatal medicine. St. Louis: Mosby Yearbook; P. 17-28.

Buchan PC (1997): pathogenesis of neonatal hyperbilirubinemia after induction of labour with oxytocin. Br Med J, Nov. 17; 2(6200): 1255-7.

Campbell AGM and McIntosh N (1998): Forfar and Arneil' textbook of pediatrics; fifth edition. Churchill Livingstone. Neonatal Jaundice; 214-226.

Campbell O, et al. (2004): Perinatal / Neonatal Mortality Study. Journal of Perinatology, 24(5): 284-9.

Cash and Sheryl (2004): Guideline Offers Direction for Prompt Diagnosis, treatment of hyperbilirubinemia. AAP New, American Academy of Pediatrics.

Children's National Medical Center (1998): NICU Protocol. Washington D.C.

Chou SC and Palmer RH (2003): Management of hyperbilirubinemia in newborns: measuring performance by using a benchmarking model. Pediatrics, 112: 1264-1273.

Cloherty and AR Stark, eds. (1998): Manual of Neonatal Care, 4th ed. Philadelphia: Lippencott-Raven.

Danic, Mortelli E, Bertini G, Pezzati M, Filippil, Rossetti , Rizzuti G and Rubaltelli FF (2003): Plasma bilirubin level and oxidative stress in preterm infants. Arch Dis Child Fetal Neonatal; 88(2): F119.

DeJonge MH, Khuntia A, Maisels MJ and Bandagi A (1998):

Bilirubin (BR) and severe retinopathy of prematurity (ROP) in 23-26 week estimated gestational age (EGA) infants. *Pediatr Res*; 43: 171A.

Dennery PA, Seidman DS, and Stevenson DK (2001): Neonatal

hyperbilirubinemia. *New Eng J Med*, 344: 581-590.

Driscoll JM, KF Schuetz and JH Wung (1999): Neonatal care a

Practical Approach, 10th Annual Course. New York: Columbia Presbyterian Medical Center and Columbia University College of Physicians and Surgeons.

ECRI (1995): Fiberoptic phototherapy systems. *Health Devices*; 24:

134.

Fanaroff AA and RJ Martin, eds (1997): Neonatal Perinatal

Medicine: Diseases of the fetus and infant, 6th ed. ST. Louis: Mosby-Year Book.

Fanaroff AA and RJ Martin, eds (2002): Neonatal Perinatal

Medicine: Diseases of the fetus and infant, 7th ed. ST. Louis: Mosby-Year Book.

Fenton TR (1992): Neonatal Jaundice in: Campbell AGM and MC

Intosh NM (eds.): *L For Far and Arneil's textbook of pediatrics*; 14th ed., Churchill Livingstone; 240-5.

Fenton (2001): Jaundice in the healthy newborn infant: a new approach to an old problem. *Pediatrics*, 81: 505-511.

Gartner LM (1994): Neonatal Jaundice. *Pediatrics in Review*; 15: 422-432.

Gathwala G and Sharma S (2002): Phototherapy induces oxidative stress in premature neonates. *Indian J Gastroenterol*; 21 (4): 153-4.

Gopinathan V, Miller NJ, Milner AD and Rice Evans CA (1994): Bilirubin and ascorbate antioxidant activity in neonatal plasma. *FEBS Lett*; 349: 197-200.

Halamek LP and Stevenson DK (1997): Neonatal jaundice and liver disease. In: *Neonatal and perinatal medicine disease of the fetus and infant*. 6th ed. Edited by Fanaroff AA and Martin RJ, Mosby-Boston, New York; 1345-1389.

Halamek LP, and Stevenson DK (2002): Neonatal jaundice and liver disease. Chapter in: Fanaroff AA, Marin RJ, eds. *Neonatal Perinatal Medicine: Diseases of the fetus and infant*. 7th ed. St. Louis, MO: Mosby, Inc 1309-1350.

Han JL and Yan JM (1993): [Treatment of ABO blood type incompatibility caused early abortion with traditional

Chinese medicine]. [Article in Chinese]. Zhongguo Zhong Xi Yi Jie He Za Zhi, 13(1): 13-5,4.

Hansen TW (2002): Kernicterus: An international perspective. Semin neonatol. 7(2): 103-9.

Hansen TWR (2002): Mechanisms of bilirubin toxicity: clinical implications. Clin Perinat, 29: 765-778.

Hinkes M and Cloherty JP (1998): Manual of Neonatal Care Textbook; 4th Edition. Lipincott Raven. Philadelphia. New York. John P. Cloherty and Ann R Stark (eds.); Chapter 18: 175-210.

Ho NK (2002): Using a spectral reflectance technique to measure transcutaneous bilirubin in neonates: a new device. Issues Emerge Health Technol; 33: 1-4.

Holt LE (1897): The disease of infancy and childhood: for the use of students and practitioners of medicine. New York: D. Appleton (quoted from Dennery et al., 2001).

Holtzman NA (2004): Management of hyperbilirubinemia: Quality of evidence and cost. Pediatrics 114: 1086-1088.

Hosono S, Ohno T, and Kimoto H (2002): Follow up study of auditory brainstem responses in infants with high

unbound bilirubin levels treated with albumin infusion therapy, *Pediatr Int*; 44(5): 488-92.

Ip S, et al (2004): Hyperbilirubinemia and kernicterus: 50 years later. *Pediatrics*, 114: 263-264.

Ives NK (1999): Neonatal Jaundice. In: *Textbook of Neonatology*; Edited by Janet M. Rennie and NRC Robertson. Third Edition. Churchill Livingstone; 715-730.

Jackson CL, Tudehope D, Willis L, Law T, and Venz J (2000): Home phototherapy for neonatal jaundice technology and teamwork. *Aust Health Rev*; 23(2): 162-8.

Jendrassik L and Grof P (1938): *Biochem. Z.*, 297: 81.

Johnson LH, and Bhutani VK (2002): System based approach to management of neonatal jaundice and prevention of kernicterus. *Jour Peds*, 140: 396-403.

Joint Commission on Accreditation of Healthcare organizations (2004): Revised guidance to help prevent kernicterus. Sentinel Event Alert. August, Issue 31.

Joint Commission on Accreditation of Healthcare Organizations (2001): Kernicterus threatens healthy newborns. Sentinel Event Alert. April, Issue 18.

Knudsen A (1996): Prediction and non-invasive assessment of neonatal jaundice in the term healthy new born infant. Acta Paediatr; 85: 393-7.

Knudsen A (1998): Predicting the need of phototherapy in healthy mature neonates using transcutaneous bilirubinometry on the first postnatal day. Boil Neonate; 68: 398-403.

Knudsen A (1999): Prediction of the development of neonatal jaundice by increased umbilical cord blood bilirubin. Acta Paediatr Scand; 78: 217-21.

Knupfer M, Pulzer F, GeBauer C and Vogtman C (2005): Predictive value of umbilical cord blood bilirubin for post natal hyperbilirubinaemia. Acta Paediatrica; 94: 581-587.

Liu YJ, Chen W, Wu KW, Broadberry RE and Lin M (1996): The development of ABO isohemagglutinins in Taiwanese. Hum Hered; 46(4): 181-4.

Locham KK, Kaur K, Tandon R, Kaur M and Garg R (2002): Exchange blood transfusion in neonatal hyperbilirubinemia role of calcium. Indian Pediatr; 39(7): 657-9.

Maisels MJ and Newman TB (1995): Kernicterus in otherwise healthy, breast fed term newborn. Peadiatircs, 96: 730-3.

Maisels MJ (1996): Why use homeopathic doses of phototherapy?
Pediatrics; 98: 283.

Maisels MJ and Kring E (1998): Length of stay, jaundice, and
hospital readmission. Pediatrics; 101: 995-998.

Maisels MJ (1999): Neonatology: pathophysiology and
management of the Newborn, fifth edition, edited by
Aveny GB, Fletcher MA and MacDonald MG, fifth
edition. Lippincott Company, Philadelphia; 38: 765-919.

Maisels MJ, Newman TB (1999): Predicting hyperbilirubinemia in
newborns: the importance of timing pediatrics; 103: 493-
494.

Maisels, MJ (2001): Phototherapy traditional and nontraditional. J
Prenatol; 1: S93-7; discussion S 104-7.

**Matthias Knupfer, Ferdinand Pulzer, Corinna Gebauer, Ova
Robel Tilig and Christoph Vogtmann (2005):**
Predictive value of umbilical cord blood bilirubin for
postnatal hyperbilirubinemia. Acta Paediatrica; 94: 581-
587.

McDonagh AF (1998): Is bilirubin good for you? Clinics in
Perinatology; 17: 259-269.

Mireles LC, Lum MA, and Dennery PA (1999): Antioxidant and Cytotoxic effects of bilirubin on neonatal erythrocytes. *Pediatric Res*; 45: 355-62.

Monaghan G, McLellan A, McGeehan A, Li Volti S, Mollica F, Salemi I, Din Z, Cassidy A, Hume R, and Burchell B (1999): Gilbert's syndrome is a contributory factor in prolonged unconjugated hyperbilirubinemia of the newborn. *Pediatr*; 134(4): 441-6.

Moyer VA, Ahn C and Sneed B (2000): Accuracy of clinical judgment in neonatal jaundice. *Arch Pediatr Adolesc Med*; 154: 391-4.

Neonatal Care, Protocols for physicians. Cairo-Egypt (2004): Neonatal Hyperbilirubinemia; chapter 14: 223-234.

Newman TB, and Maisels MJ (1990): Does hyperbilirubinemia damage the brain of healthy full term infants? *Clin Perinatol*; 17: 331-358.

Newman TB, and Maisels MJ (1992): Evaluation and treatment of jaundice in the term newborn: A kinder, gentler approach. *Pediatrics*, 89: 809-818.

Newman TB, and Klebanoff MA (1999): Neonatal hyperbilirubinemia and long-term outcome: Another

look at the collaborative perinatal project. *Pediatrics*, 92: 651-657.

Newman TB, Liljestrnd P and Escobar GJ (2002): Jaundice noted in the first 24 hours after birth in a managed care organization. *Arch Pediatr Adolesc Med*; 2002, 156: 1266-50.

North Shore University Medical System (1998): NICU policy and Procedures. Bayshore, NY: Southside Hospital.

Olcay Oran MP, Simit Sarici, MD, Muhittin A, Serdar MD, Aysekorkmaz MD, Gulsen Erden MD, Gulsevin Tekinalp MD, Murat Yurdakok MD and Suleyigit, MD (2004): Incidence, course, and prediction of hyperbilirubienmia in near-term and term newborns. *Pediatrics*, Vol. 113, No. 4, April 2004, pp. 775-780.

Peter F, Whittington, Estello M and Alonso (1998): Disorders of bilirubin metabolism in: Nathan and Oski's Hematology of infancy and childhood; David G, Nathan, Stuart H. Orkin and David Ginsburg; (eds.) 5th edition, W.B. Saunders Company, 79-113.

Poland RL (2002): Preventing kernicterus: Almost there. *Jour Pediatr*, 140: 385-386.

Risemberg HM, Mazzi E, MacDonald MG, Perlta M, and Heldrich F (1997): Correlation of cord bilirubin level with hyperbilirubinemia in ABO in Compatibility. Arch Dis Child; 52(3): 219-22.

Riskin A, David M, Peskin B, Tamir A, Vafsi O, Leivovitz Z, Riskin Mashiah S, Israel N, Merksamer R and Bader D (2004): The role of umbilical cord alpha fetoprotein as a screening tool for neonatal hyperbilirubinemia. Am J perinatol 21(2): 93-8.

Rosenfeld J (1996): Umbilical cord bilirubin levels as a predictor of subsequent hyperbilirubinemia. J Fam Pract; 23: 556-8.

Rubaltelli FF (1998): Current drug treatment options in neonatal. Drugs; Jul; 56(1): 23-30.

Sarici SU, Yurdakok M, and Serdar MA (2002): An early (sixth-hour) serum bilirubin measurement is useful in predicting the development of significant hyperbilirubinemia and severe ABO hemolytic disease in a selective high risk population of newborns with ABO incompatibility. Pediatrics; 109: e53.

Schaffer and Abvery's (1991): Diseases of the newborn, 6th ed. Philadelphia, W.B. Saunders.

Schmorl G (1904): Zur kenntniss des Ikterus neonatorum, insbesondere der dabei auftretenden Gehirnveränderungen. Verh Dtsch Pathol Ges; 6: 109-15 (quoted from Dennery et al., 2001).

Segal M, Adachi Y and Koiwai O (1996): The genetic basis of Gilbert's Syndrome. Lancet; 374: 557-8.

Sheridan SE (2002): Kernicterus parents' group. Jour Peds, 141: 597-598.

Shibuya A, Hirono A, Ishii S, Fujii H, and Miwa S (1999): Hemolytic crisis after excessive ingestion of fava beans in a male infant with G6PD canton. Int J Hematol; 70(4): 233-5.

Smith JF and Baker JM (1994): Crigler Najjar disease in pregnancy. Obstet Gynecol; 84: 670.

Stephenson T, Zuccolo J and Mohajen M (1994): Diagnosis and management of non immune hydrops in the newborn. Arch Dis Child; 70: F151.

Stevenson DK and Vreman HJ (1998): Bilirubin production in healthy term infants as measured by carbon monoxide in breath. Clinical Chemistry; 40: 1934-1939.

Stoll BJ and Kliegman RM (2000): Nelson text book of pediatrics: 16th edition, edited by Behrman RE, Kliegman RM and Jenson HB, W.B Saunders company volume 2; 513-519.

Sung RY, Pang CP, Lyon A, Loong PL and Davies DP (1996): Urinary D-glucaric acid excretion in idiopathic neonatal jaundice. Chin Med J; 109(3): 201-5.

Suresh GK, and Clark RE (2004): Cost-effectiveness of strategies that are intended to prevent kernicterus in newborn infants. Pediatrics.

Suresh GK, Marin CL and Soll RF (2003): Metalloporphyrins for treatment of unconjugated hyperbilirubinemia in neonates (Cochrane Review). In: The Cochrane Library, Issue 2. Oxford: Update Software.

Taeusch W and RA Ballard eds (1998): Schaffer and Avery's Diseases of the Newborn, 7th ed. Philadelphia: W.B. Saunders Co.

Tan KL, Chia HP and Koh BC (1996): Transcutaneous bilirubinometry in Chinese, Malay and Indian infants. Acta Paediatr; 85(8): 986-90.

Thibeault DW (2000): The precarious ant: Oxidant defuses of the pretorm infant. Amgperinatal; 17: 167-81.

Tsai FJ, Tsai CH, Peng CT, and Wang TR (1998): Sister chromatid exchange in Chinese newborn infants treated with phototherapy for more than five days. *Zhonghua Min Guo Xiao Er Ke Yi Xue Hui Za Zhi*; 39(5): 327-9.

Tsang RC et al eds (1997): Nutrition during Infancy: Principles and Practice, 2nd ed. Cincinnati: Digital Educational Publishing.

Van Bel F, Dorrepaal CA, Benders MJNL, Zeeuwe PEM, Van de Bor M and Berger HM (1994): Changes in cerebral hemodynamics and oxygenation in the first 24 hours after birth asphyxia. *Pediatrics*; 92: 365-372.

Vinod K, Bhutani, Lois Johnson and Emidio M and Sivieri (1999): predictive ability of a predischage hour specific serum bilirubin for subsequent significant hyperbilirubinemia. In healthy term and near-term newborns. *Pediatrics*; Vol 103, No. 1 January, pp. 6-14.

Volpe JJ (2001): Bilirubin and brain injury. Chapter in: Volpe JJ. *Neurology of the Newborn*. 4th ed. Philadelphia, PA: WB Saunders Co, 521-546.

Watchko J and Classen D (1994): Kernicterus in premature infants: current prevalence and relationship to NICHD phototherapy study exchange criteria. *Pediatrics*; 93: 996.

Whittington PF, and Alonso EM (2003): Disorders of bilirubin metabolism. Chapter in: Nathan DG, Ginsburg D, Orkin SH, Look AT, eds. Nathan and Oski's Hematology of Infancy and Childhood. 6th ed. Philadelphia, PA: WB Saunders Co, 86-120.

Wong KF, Tse KT, Lee AW, Mak CS and So CC (1997): Is antenatal antibody screening worthwhile in Chinese? Br J Haematol; 97(4): 917-9.

Yeo KL, Perlman M and Hao YMP (1998): Outcomes of extremely premature infants related to their peak serum bilirubin concentrations and exposure to phototherapy. Pediatrics; 102-1426.

Yesilkaya A, Altinayak R and Kipmen D (2000): The antioxidant effect of free bilirubin on cumene-hydroperoxide treated human leukocytes. General Pharmacology: the vascular system; 35: 17-20.

Yigi TS, Yurdakok M, and Kilinc K (1999): Serum malondialdehyde concentrations in babies with hyperbilirubinemia. Arch Des Child Fetal Neonatal; 80: P235-237.

Zweig MH, Campbell G (1993): Receiver-operating characteristic (ROC) plots: a fundamental evaluation tool in clinical medicine. Clin Chem 1993; 39(4); 561-77.