

References

Abaci A, Kiranati B and Ozkutlu S (1999):

Are the left ventricular diastolic tissue Doppler diastolic filling velocities independent from preload? Arch Turk Soc Cardiol 1999; 27:578-582

Agmon Y, Oh JK and Mc Carthy JT., (2000):

Effect of volume reduction on mitral annular diastolic velocities in hemodialysis patients Am J Cardiol 2000, 85:665-668.

Amman K, Ritz E and Wiest G. (2001):

A role of parathyroid hormones for the activation of fibroblasts in uremia J Am Soc Nephrol 1994; 4:1814-1819

Appleton C.P., (1993):

Doppler assessment of left ventricular diastolic function: the refinements continue. (Editorial)

J. Am. Coll. Cardiol., 21(7):1697-700.

Appleton CP and Redfield MM. (2000):

Clinical utility of Doppler echocardiography and tissue Doppler echocardiography in the estimation of left ventricular filling pressure: A comparative simultaneous Doppler catheterization study. Circulation 2000; 102:1788-1794

American Echocardiography Society. (2003) :

Heart Diseases and stroke statistics.

Dallas Tex.: American heart association, 2003.

Appleton C.P, Tugcine. And Hatle L.K. (1992):

The natural history of left ventricular filling abnormalities: Assessment by two-dimensional and Doppler echocardiography.

Echocardiography; 9(4):548-568

Ayabakan C and Ozkutlu S. (2004):

Left ventricular diastolic velocities by tissue Doppler and relation to the characteristics of filling ventricles. Young. Apr:14(2):156-63.

Baker D.W. (1980).

Application of pulsed Doppler.

Rad. Clin. North. Am., 18(1, 4):79-103.

Bauer F, Jamal J and Douillet R, (2007):

Acute change in load: Effects of myocardial velocities measured by TDI

.Arc Mal Coeur Vaiss 2001:94:1155-1160

Bergehin H,Daugirdas JT and Ing TS (Eds.). (2000):

Complications during hemodialysis handbook of dialysis .3rd edition.

lippincot Williams and wilkins, 2000; p.148-68

Bergstein, (2000):

Renal failure, In:Berhrman et al., Nelson textbook of pediatrics, 16th (ed.).Saunders company.pp 1605-1612.

Bess RL, Khan S and Rosman HS,(2007):

Technical aspects of diastology: why mitral inflow and tissue Doppler imaging are the preferred parameters? Echocardiography 2006:23:332-339.

Bora Polat, Tugcine and Uraginci T, (2006):

Journal of pediatric Gastroenterology and nutrition.43 (2):222-227, August 2006

Choong CY and Thomas JD, (1998):

Combined influence of ventricular loading and relaxation on the transe mitral flow velocity profile measured by Doppler echocardiography. Circulation 1998; 78:672-683.

Daugirdas JT, (2001):

Pathophysiology of dialysis hypotension: An update: Am J kidney disease 2001; 38(4):1-65.

David, Alpert MA and Ravens craft Md (, 2003):

Pericardial involvement in end stage renal disease .In: Department of medicine, st John Mercy Medical Center's Louis ,Missouri. Am J med Sci: 325(4):228.

Derumeaux G, Douillet R, Troniou A, Jarnal F, Litzler PY and Pontier G (1999):

Distinguishing between physiologic hypertrophy in athletes and primary hypertrophic cardiomyopathies. Importance of tissue color Doppler. Arch Mal Coeur Vaiss 1999; 92:197-199

Derumeaux G., Michel O. and Loufoua J., (2000):

Assessment of non uniformity of transmural myocardial velocities by color-coded tissue Doppler imaging. Characterization of normal, ischemic and stunned myocardium. Circulation, 101:1390-95.

Derumeaux G, Mulder P, Richard V and Bauer F. (2002):

Tissue Doppler imaging differentiates physiological from pathological pressure overload left ventricular hypertrophy in rats. Circulation 2002; 105:1602-1608

Desai A, Kern MJ., and J Fuller CJ, (2006):

Patent foramen ovale and cerebrovascular diseases. Nat Clin Pract Cardiovascular Med. 2006; 3:446-455.

Doherty AM, Patt WC and Berryman KA. (1997):

Gastrointestinal complication in chronic renal failure. Nephrol 1997; 26:358-61.

Dowborn JK, (1996):

CRF IN: Textbook of renal disease 1st Ed (Judith) Witworth JR, Lawrence AO, ed. Churchill Livingstone-Edinberg. p.343.

Dursunoglu D., Polat B. and Evrengul Hl. 2004 :

Assessment of systolic function by atrioventricular plane displacement in patients with diastolic dysfunction. Acta. Cardiol. Aug;59(4):409-15.

Edvardsen T, Urehim S, Ihlen H and Smiseth OA (2002):

Quantification of left ventricular systolic function by tissue Doppler echocardiography: added value of measuring pre and post ejection velocities in ischemic myocardium. Circulation 2002; 105:2071-2077

Eidem BW, Mc Mahon CJ, Ayres NA and Kovachin JP, (2005):
Impact of chronic left ventricular preload and after load on Doppler tissue imaging .J Am Soc Echocardiography .Aug:18(8):830-8.

El-Nahas AM, Richard V and Winerals CG, (1996):
Chronic renal failure in: David AW, Edward J (Eds).oxford textbook of medicine 1996.3rd edition .oxford university press, London; chapter 20:3294-306.

Erbel R., Nesser H and Drozd J., (1995):
Atlas of tissue Doppler echocardiography.
Darmstadt: Steinkopf 1995.

Eric HY and Robert Z (2006):
Evaluation of cardiac function in the dialysis patients –a primer for the non –expert. Nephrology Dialysis Transplantation 21(6):1474-1481.

European Study Group on Diastolic Heart Failure, (1998):
Working Group Report: How to diagnose diastolic heart failure.
Eur. Heart J. 19:990-1003.

Farid F, Russhan L, Badawy A and Yossef D., (2005):
Effect of hemodialysis on carnitine levels in children with chronic renal failure. Geget 5:11-21.

Farid FA, (1992):
Eighteen months experience in the pediatric dialysis unit (PDU). Ain Shams University children Hospital .Proceedings of the annual congress of the Egyptian Society of nephrology 1992 ;(1):p.121

Fauchier L and Mario O, (2003):
Reliability of QRS duration and morphology on surface electrocardiogram to identify ventricular desynchronize in patients with idiopathic dilated cardiomyopathy .Am J Cardiol.2003; 92:341-344.

Fine RN, Davidson DM. and Gentile DE. (1990):
Growth in children with chronic renal insufficiency .Clinical dialysis 2nd edition 1990; 9:667.

Fishbane S and Paganini EP. (2000):

Hematologic abnormalities .In Daugirdas JT, Blake PG, (eds.) Handbook of dialysis, 3rd edition, Philadelphia 2000;p.477-94.

Francisco L, (2005):

Renal Failure: Diagnosis and management, In: Wilcox CS, Tisher CC (eds.).Handbook of nephrology and hypertension, 5th edition, Philadelphia 2005; 267-293.

Firstenberg MS and Greenberg NL (2001):

Determinants of diastolic myocardial tissue Doppler velocities: influence of relaxation and preload Appl Physiol 2001; 90:299-307.

Foley RN, Parfrey PS, Harnett JD, Kent GM, Murray DC and Barre PE. (1995):

The prognostic importance of left ventricular geometry in uremia cardiomyopathy.J Am Soc Nephrol 1995; 5:2024-2031

Galiuto L., Ignone G. and De Maria A.N., (1998):

Contraction and relaxation velocities of the normal left ventricle using pulsed wave tissue Doppler echocardiography.
Am. J. Cardiol., 67:222-224.

Garcia MJ and Thomas JD, (1998):

New Doppler echocardiography applications for the study of diastolic function. J Am Coll Cardiol 1998; 32:865-875.

Gardin JM, Rohan MK., and Davidson DM., (1987) :

Doppler transmitral flow velocities parameters: Relationship between age, body surface area, blood pressure and gender in normal subjects. Am. J. Non-invasive Cardiol 1:3

Gillber JC. And Glantz SA., (1989):

Determinants of left ventricular filling and of the diastolic pressure volume relation. Circ. Res. 64:827-832

Golderisi M., Benjamin EJ. Evans JC. and Fuller DL., (1997):

Intra and inter observer reproducibility of Doppler assessed indexes of left ventricular diastolic function in a population based study (the framingham heart study). Am. J. Cardiol. 70:1341.

Gorcsan J., Gulati V.K. and Mandarino W.A., (1996):

Color-coded measures of myocardial velocities throughout the cardiac cycle by tissue Doppler imaging to quantify regional left ventricular function.

Am. Heart J., 131:1203-1213.

Hamed NA. (1999):

Serum carnitine levels in hemodialysis patient's .Master thesis, Pediatrics .Supervised by, Farid FA, EL-Hakim IZ. Faculty of medicine Ain shams University, Cairo.

Harizi RC. Bianco JA. and Alpert JS.,(1988):

Diastolic function of the heart in clinical cardiology. Arch. Intern. Med. 148:99-109

Hatle and Angeleson, (1985):

Doppler Ultrasound in Cardiology, 2nd Ed.

Philadelphia: Lea and Febiger.

Hayashi SY, Brodin LA, Alvestrand A and Mazza Do. (2004) :

Improvement of cardiac function after haemodialysis .Quantitative evaluation by color tissue velocity imaging. Nephrol Dial Transplant 2004; 19:1497-1506

Hiraishi S and Di Sesha TG (1983);

Two dimensional echocardiography assessment of left atrial size in children. J Am Cardiol 1983; 52:1249.

Huysman and Kramer W, (1998):

Cardiac structure and function in continuous ambulatory peritoneal Dialysis: influence of blood purification and hyper circulation Am Heart J.119:pp.52-344

Keren G, Nielson JC. and Sherez R., (1985):

Pulmonary venous flow pattern.Its relationship to cardiac dynamics. A pulsed Doppler echocardiography study.

Circulation, 71:1105.

Kowari V and Roguljic M. (2003):

Metabolic acidosis of chronically hemodialyzed Patient's .Am J Nephrol 2003; 23(3):158-64

Lawrence MT and Stephen JM, (1999):

Chronic renal failure. Current medial diagnosis and treatment 38th edition printed in the united state of America 1999; p.873-80

Levin a, Singer J, Thompson CR, Ross H and Lewis M, (1996):

Prevalent left ventricular hypertrophy in the predialysis population: identifying opportunities for intervention. Am J Kidney Disease 1996; 27:347-354?

Locatelli F and Marcelli D. (2000):

Cardiovascular disease in chronic renal Failure: the challenge continues .Nephrol Dial Transplant 2000; 15(5):569-80

London GM, (2003):

Cardiovascular disease in chronic renal Failure: path physiologic aspects .Semin Dial 2003; 16:85-94

London GM, (1997):

Cardiac disease in chronic uremia: pathogenesis.Adv Ren Replace Ther 1997; 4:194-211

Lytrivi ID, Lai WW, Ko HH, Nielson JC and Strivasyava S., (2005):

Assessment of ventricular function by tissue Doppler in children with end stag renal disease. Oct: 18(10):1099-104.

Mackenzie HS, (2001):

Prevention of progressive renal failure. Brenner and rector therapy in nephrology and hypertension .5th edition .philadilphia 2001; 463-73.

Maron MS and Olivotto I, (2003):

Effect of left ventricular outflow tract obstruction on clinical outcome in hypertrophy cardiomyopathy.N Engl J Med.2003; 348:295-303.

Matsumura Y and Elliott, (2002):

Left ventricular diastolic function assessed using Doppler tissue imaging in patients with hypertrophy cardiomyopathy: relation symptoms and exercise capacity. *Heart*. 2002;87:347-251

McDicken W.N., Sutherland G.R. and Moran C.M. (1992):

Color Doppler velocity imaging of the myocardium. *Ultrasound Med.BIOL.*, 18(6\7):651-654.

Moe SM and Sprango SM. (1994): Uremic

Encephalopathy.*Clin Nephrol* 1994; 41(4):521-6.

Morris ST, Edwards W.D. and Jardine AG. (2000):

The vascular endothelium in chronic renal failure. *J Nephrol* 2000; 13:96-105.

Moudgil A and Bagg A. (1999):

Evaluation and treatment of chronic renal failure .*Indian J Pediatric* 1999; 66(2):241-53.

Movillie E, Zani R and Sangalli L. (1998):

Correction of metabolic acidosis increases serum albumin concentration .Kinetically evaluated protein intake in hemodialysis patien: a prospective study .*Nephrol Dial Tranplant* 1998; 73:1719-22.

Mueller X., Stauffer J.C. and Jaussi A., (1991):

Subjective visual echocardiographic estimate of left ventricular ejection fraction as an alternative to conventional echocardiography methods; Comparison with contrast angiography.
Clin. Cardiol. 14:898.

Nagueh SF, and Kopelen HA, (1997):

Doppler tissue imaging: A non invasive technique for evaluation of left ventricular relaxation and estimation of filling pressure.
J. Am. Coll. Cardiol., 30:1527-1533.

Nandia N.C., Pinheiro L. and Schwartz S.I., (1992):

Multiplane transoesophageal imaging and three-dimensional reconstruction.
Echocardiography, 9:677.

Naqvi TZ, Neyman G and Broyde A (2001):

Comparison of myocardial tissue Doppler with transmitral flow Doppler in left ventricular hypertrophy. *J Am Soc Echocardiography* 14:1153-1160

Nishimura RA, (1984):

Non-invasive measurement of cardiac output by continuous wave Doppler echocardiography: initial experience and review of the literature. *Mayo.Clin.Proc.*, 59(7):484-489.

Nishimura R.A., (1985):

Doppler echocardiography: Theory, instrumentation, technique, and application *Mayo. Clinic. Proc.*, 60(5):321-343.

Nishimura R.A.and Schwartz C., (1993):

Assessment of diastolic function of the heart: Background and current applications of Doppler echocardiography. Part II. Clinical study .*Mayo Clin Proc* 1989; 64:181-204.

Nishimura R.A. and Tajik J., (1997):

Evaluation of diastolic filling of left ventricle in health and disease: Doppler echocardiography is the clinicians Rosetta stone. *J. Am. Coll. Cardiol.*, 30:8-18.

Nishimura R.A., Housmans P.R. and Hatle L.K. (1989):

Assessment of diastolic function of the heart: Background and current applications of Doppler echocardiography (part 1). *Mayo. Clinic. Proc.*, 64:71-81.

Nishimura R.A., Edwards W.D. and Warnes C.A., 1990):

Intravascular ultrasound imaging: In-vitro validation and pathologic correlation. *J. Am. Coll. Cardiol.*, 16(1)145-154.

Nissen S.E., Gurley J.C and Grines C.L, (1991):

Intravascular ultrasound assessment of lumen size and wall morphology in normal subjects and patients with coronary heart disease. *Circulation*, 84:1087.

Nissenson AR and Steinwald B. (1996):

The effects of recombinant human erythropoietins on functional health and well-being in chronic dialysis patients Am Soc Nephrol 1996; 7(5):763-73.

Obrador GT, Appleton C and Kasusz AT, (2002) :

Chronic kidney diseases in united states: an under recognized problem.Semin Nephrol.2002; 22:441-8.

Oguzhan A.,Arinic H.,Abaci A and Topsakal R., (2005):

Preload dependence of Doppler tissue imaging derived indexes of left ventricular diastolic function.

Echocardiography: A journal of cardiovascular ultrasound and altered technique, volume 22.No 4:320-324

Oki T,Tabata T, Yamada H andNishikado A (1998):

Clinical application of pulsed tissue Doppler imaging for assessing abnormal left ventricular relaxation .Am J Cardiol 1998,79:921-928

Okubo M and Takahashi K, (2001) :

Age dependency of stiffness of the abdominal aorta and the mechanical properties of the aorta in Kawasaki disease in children.Pediatr Cardiol 22:198-203

Ommen SR, Nishimura RA, Appleton C and Redfield MM (2000):

Clinical utility of Doppler echocardiography and tissue Doppler echocardiography in the estimation of left ventricular filling pressure: A comparative simultaneous Doppler catheterization study. Circulation 2000; 102:1788-1794

Olivotto I and Betocchi S (2003):

Effect of left ventricular outflow obstruction on clinical outcome in hypertrophic cardiomyopathy.N .Engl J Med.2003:348:295-303.

Parfery PS,Harnett JD and Barre PE. (1996)A:

The natural history of myocardial disease in dialysis patients, J Am Soc Nephrology 1996; 2:2-12.

Parfery PS, Foley RN and Harnet JD. (1996)B. :

Impact of hypertension on cardiomyopathy. Morbidity and mortality in end-stage renal disease .Kidney Int 1996; 49:1379-1385

Plotnick GD., (1989):

Changes in diastolic function difficult to measure, harder to interpret. Am. Heart J. 118:637

Ramalanjaona G (editor) (2005). :

Diagnosis and management. In: chronic renal failure in children, 1st edition, ever new york 2005; p.2.

Ritz E, Wiest G and Mall G (1994):

A role of parathyroid hormone for the activation of cardiac fibroblasts in uremia .J Am Soc Nephrol 4:1814-1819

Rokey R., Kuo LC. and Zoghbi WA., (1985):

Determination of parameters of left ventricular diastolic filling with pulsed Doppler echocardiography in comparison with cineradiography. Circulation 71:543-50

Ross JR and Tokars JE.(2005).:

Successful treatment of a hemodialysis access system pocket infection with large volume kanamycin solution irrigation .Adv Ren Replace Ther 2005;10:248-53.

Sabbah HN. and Stein PD., (1981):

Negative diastolic pressure in the intact canine ventricle: Evidence of diastolic suction. Circ. Res. 49:108.

Salski IB and Goodman W. (1994).:

Disorders of bone and mineral metabolism chronic renal failure .In: Trachtman H (editor). Barrats textbook of pediatrics nephrology, 3rd edition, 1994:p.390-483.

Sagie A. and Benjamin E.J., (1993):

Reference values for Doppler indexes of left ventricular diastolic filling in the elderly.

J.Am.Soc.Echocardiogr. 6(6):570-576.

Salaymeh K, Kern MJ. and Banerjee A,(2007).:

Evaluation of atrial stiffness in children with Williams syndrome : does it play a role in evolving hypertension?. Am Heart J 142:549-555

Senior R., SomanP., and KhattabRS., (1999):

Improved endocardial visualization with second harmonic imaging comparing with fundamental two dimensionalechocardiographic imaging. Am Heart Journal 138:163-168?
Philadelphia: Lea and Febiger.

Shapiro SM., Laks MM. and Bersohn MM. (1991):

In search of the Holy Grail: The studies of diastolic ventricular function by the use of Doppler echocardiography. J. Am.Coll. Cardiol. 17:1517.

Shiller N.B. and Shah P.M., (1989):

Recommendations for quantification of the left ventricle by two dimensional echocardiography.
J.Am. Soc.Echocardiogr., 2(5):358-367.

Smith MD. , (1997):

Left ventricular diastolic function:
Clinical utility of Doppler Echocardiography. Quoted from: The practice of clinical echocardiography: Otto, C.M. WB Saunders company.

Smith K.S., Stein PD. And Traber MG.(1997).:

Vitamin E supplementation increase circulating vitamin E metabolites ten fold in end state renal disease patients. Lipids1997; 38(8):813-9.

Sohn D.W., Chai I.H. and Lee D.J., (1997):

Assessment of mitral annulus velocity by tissue Doppler imaging in the evaluation of left ventricular diastolic function.
J.Am. coll.Cardiol. 30:474-480.

Srivastava R and Abuelo JG (Eds.). (2006).:

Diagnosis and treatment .In: Chronic renal failure in children, 2nd edition, Harvard, USA 2005; PP1-284.

Stoddard MF. Pearson AC. And Kern MJ. (1989):

Left ventricular diastolic function: comparison of pulsed Doppler echocardiography and hemodynamic indexes in subjects with and without coronary heart disease. J Am.Coll. Cardiol.13-327.

Takahashi K, (2001).:

Age dependency of stiffness of the abdominal aorta and the mechanical properties of the aorta in Kawasaki disease in children.Pediatr Cardiol 22:198-203

The Task Force on Heart Failure of the European Society of cardiology (1995):

Guideline for the diagnosis of the heart failure Eur Heart J 1995; 16:741-751

Vasan RS and Larson MG, 1999):

Congestive heart failure in subjects with normal versus reduced left ventricular ejection fraction: prevalence and mortality in a population – based cohort. J Am Coll Cardiol 1999; 33:1948-1955.

Warady BA and Harmon WE. (1999):.

Optimal care of the pediatric end stage renal disease patient on dialysis. Is J Kidney disease 1999; 33:567-583?

Wierzbowska-Drabik. K, Drozd J and Plewka M., (2006):

Pulsed tissue Doppler echocardiography parameters for the diagnosis of advanced left ventricular hypertrophy .Echocardiography Mar, 23(3):189-96

Yamada H., Oki T. and Tabata T., (1999) :

Effect of aging on diastolic left ventricular myocardial velocities measured by pulsed Doppler tissue imaging in healthy subjects.

J.Am. Soc.Echocardiogr., 12:574-581.

Yamamoto K. and Redfield MM. and Nishimura RA., (1996):

Analysis of left ventricular diastolic function. British Heart Jornal 72(supll II):27-35

Zile MR., (1991):

Homodynamic determinants of echocardiographically-derived indices of left ventricular filling echocardiography9:289.

Zile M, Larson MG, and Gaasch WH, (2001):

Heart failure with a normal ejection fraction, is measurement of diastolic function necessary to make the diagnosis of diastolic heart failure? Circulation 2001:104:779-782.

Ziolkowska H, Tabata T.and Debinki A.,(2000):.

Serum bone turnover parameters in uremia children. 89(6):666-71

Zoghbi W.A. and Bolli R., (1991):

The increasing complexity of assessing diastolic function from ventricular filling dynamic.

J.Am. coll.Cardiol., 17:1237.