

REFERENCES

Agha HM (1993):

Cardiopulmonary assessment of the newborn with perinatal hypoxia
M.D. Thesis. Faculty of Medicine Cairo University.

Allen MC, Cauter AJ. (1986):

Assessment of early auditory and visual abilities of extremely premature infants. Dev. Med. Child. Neurol. 28: 458-466.

American academy of pediatrics and American college of Obstetricians (2002):

Guidelines for Perinatal Care.

Amiel-Tison C, Grenier A. (1986):

Neurological Assessment During the First Year of Life. 1st ed. New York: Oxford University Press.

Anand NK, Gupta AK, and Lampa IM (1994):

Neurosonographic abnormalities in neonates with hypoxic ischemic encephalopathy. Indian Pediatrics; Jul;31(7):767-74.

Anderson RB, Rosenblith JF. (1968):

Photic sneeze reflex in the human newborn. Dev Psychobiol. 1: 65.

Apgar and James, LS (1962):

Further observations on the newborn scoring system Am. J. Dis. Child., 104-419.

Azzopardi D, Wyatt JS, Cady EB et al (1989):

Prognosis of newborn infants with hypoxic-ischemic brain injury assessed by phosphorus magnetic resonance spectroscopy Pediatr. Res. 25:445.

Babcock, D.S. and Ball, W. (1983):

Postasphyxial Encephalopathy in full-term infants: Ultrasound Diagnosis. Radiology 148: 417-423.

Babcock D.S and Han.B.K. (1981):

The accuracy of high resolution real – time ultrasonography of the head in infancy. Radiology: 135: 665 – 676.

Bagenholm R., Andine P and Hagberg H (1991);

Effects of 21 aminosteroid U 7400F on brain damage edema following perinatal hypoxia-ischemia in the rat J. Cereb. Blood Flow Metab., 11:134.

Bozynski M, Dipietro M and Claflin C (1993):

Cranial sonography and neurological examination at term and motor performance through three months of age. J. Dev. Behav. Pediatr. Apr; 14 (2):112-6.

Brann AW (1985):

Factors during neonatal life that influence brain disorders. In: Feeman J. (ed.) Prenatal and perinatal factors associated with brain disorder. NIH Pub. No. 85-1149, p. 263, Bethesda. Maryland. National Institutes of Health.

Brazelton TB. (1973):

Neonatal Behavioral Assessment Scale. 1st ed. Philadelphia: Spastics International Medical Publications with JB Lippincott; (Clin. Dev. Med., Vol. 50).

Buonocore G, Perrone S and Bracci R (2001):

Free radicals and brain damage in the newborn. Biol. Neonate, 79(3-4); 180-6

Bussion A, Plotkine M .and Boulu RG (1992):

The neuroprotective effect of nitric oxide inhibitor in arat model of focal cerebral ischemia. Br. J. Pharmacol; 106-766.

Buttke TM, Sandstorm PA (1994):

Oxidative stress as a mediator of apoptosis. Immunology Today 15:7-10.

Carlo and Chatburn (1988):

Oxygen therapy. In Neonatal Respiratory Care P-197:204.

Capute AJ, Shapiro BK, Accardo PJ, Wachtel RC, Ross A, Palmer FB/ (1982):

Motor functions: associated primitive reflex profiles. Dev. Med. Child. Neurol. 24: 662-669.

Chrion C, Raynaud and Maziere (1992):

Changes in regional cerebral blood flow during maturation in children and adolescents. Journal of Nuclear Medicine 33:696-703.

Chao, C. P, Christopher G. and Alice C. (2006):

Neonatal Hypoxic-Ischemic Encephalopathy: Multimodality Imaging Findings RadioGraphics 2006;26:S159-S172

Chugani H.T. (1993):

Positron emission tomography scanning:

Applications in newborn. In: Shankaran S (guest ed.) Perinatal asphyxia., Clin. Perinatol; 20(2) 395, W.B. Saunders Co., Philadelphia.

Cruits PD, Mathews TG, Clarke TA et al., (1988):

Neonatal seizures; the Dublin collaborative study. Arch. Dis. Child., 63:1065-1068.

Dammann O and Leviton A, (2000):

Brain damage in preterm newborns: biological response modification as a strategy to reduce disabilities, J. Pediatr; 136: 433-8.

Denkhaus H, and winsberg F. (1979):

Ultrasonic measurements of fetal Ventricular system;Radiology 11:781-786

Dewbury K.C, and Alu wihare A.P.R.(1980)

The anterior as an ultrasound window for the study of the brain;

British journal of radiology; 53:81-84

Dubowitz LMD, Dubowitz V. (1981):

The Neurologic Assessment of the Preterm and Full-Term Newborn Infant. London: Spastics International Publications; Clin. Dev. Med. Vol. 79.

De Vries LS (1993):

Somatosensory-evoked potentials in term neonates with post-asphyxial encephalopathy. Classification of cranial ultrasound findings. Clin. Perinatol. 20:472-475.

Edwards AD, Mehmet H. (1996):

Apoptosis in perinatal hypoxic – ischemic cerebral damage. Neuropathol Appl Neurobiol. 22: 494-498.

Eicher DJ, Wagner CL, Katikaneni LP, Hulsey TC, Bass WT, Kaufman DA, et al. (2005):

Moderate hypothermia in neonatal encephalopathy: safety outcomes. Pediatr Neurol: 32: 18-24.

Ekert P, Perlman M and Hao Y (1997):

Predicting the outcome of postasphyxial hypoxic ischemic encephalopathy within 4 hours of birth. J. Pediatr Oct., 131(4): 613-7.

Ellison P. (1990):

The infant neurological examination. Adv. Dev. Behav Pediatr. 9:75138.

Evans JD. and Levene IM. (2004):

Hypoxic-ischemic injury. In: Rennie MJ. and Robertson CN., 3rd (ed). Textbook of Neonatology. Churchill. Livingstone. Harcourt Brace and Company, Edinburgh, p. 1231-1251.

Fenichel G. (1993):

Hypoxic ischemic encephalopathy in the newborn. Arch. Neurol.,

40-261.

Fischer A., Anderson J, Shuman K., and Stinson W. (1985):

Pediatric neurosonography clinical, tomography and neuropathologic correlates; Chapter 5, p. 33.

Fiske C.E Filly R.A., Callen P.W. (1981):

Sonographic measurement of lateral ventricular width in early ventricular dilatation; journal of clinical ultrasound; 9: 303 -307.

Fletcher Mary Ann (1998):

physical diagnosis in neonatology. 441-501.

Forfar and Arniel (1998):

Textbook of Pediatrics, Campbell A.G.M., McIntosh N., New York.

Gluckman PD, Wyatt JS, Azzopardi D, Ballard R, Edwards AD, Ferriero DM, et al. (2005):

Selective head cooling with mild systemic hypothermia after neonatal encephalopathy: multicenter randomized trial. Lancer: 365: 663-70.

Gonzalez D, Dios J and Moya M (1996):

Perinatal asphyxia, hypoxia ischemic encephalopathy and neurological sequelae in full term newborns. Rev. Neurol. Agu., 24(132):969-76.

Gonzalez D, Dios J and Moya M (1997):

Perinatal difference in asphyxic full term newborns in relation to the presence of hypoxic ischemic encephalopathy. Rev. Neurol. Agu., 25(144), 1189-94.

Gordon B, Avery and Mary Ann Fletcher (2004):

Neonatology: pathophysiology and management of the newborn, 4th edition, 248-268.

Govaert P, Matthys E, Zecic A et al. (2000):

Perinatal cortical infarction within middle artery trunks, Arch. Dis. Child. Fetal. Neonatal. ed. 82: F59-F63.

Gunn AJ, Parer JT, Mallard EC, et al. (1992):

Cerebral histological and electrophysiological changes after asphyxia in fetal sheep, Pediatr. Res., 31:486.

Gunn AJ, Gluckman PD, Wyatt JS, Thoresen M, Edwards AD, (2005):

On behalf of the CoolCap study Group. Selective head cooling after neonatal encephalopathy. Lancet: 365: 1619-20.

Hadlock F.P., Deter R.Lpark S.K (1981):

Real time sonography. Ventricular and vascular anatomy of the fetal brain in utero; American journal of radiol; 136: 133 -137.

Hard AL, Niklasson A, Svensson E, Hellstrom A (2000):

Visual function in school-aged children born before 29 weeks of gestation: a population-based study, Dev. Med Child Neurol; 40: 100-105.

Hegyi T, Carbone T, Anwar M, Ostfeld B, Hiatt M and Paneth N. (1998):

The Apgar score and its components in the preterm infant. Pediatr, (101).77-81.

Hill A and Volpe JJ (1992):

Hypoxic-ischemic cerebra! injury. In: Robertson NCR. (ed) Textbook of Neonatology. P. 1061. Churchill Livingstone, London.

Hoekelman RA. (1991):

The physical examination in infants and children, In: Bates B; ed. A Guide to physical Examination and History Taking. 5th ed. Philadelphia: JB Lippincott; 561-635.

Huang C. C.(1994):

Recent advances in perinatal asphyxia: Pathogenesis, clinical

assessment and potential treatment. *Acta Pediatr. Sin.*; 35 (5): 356.

Illingworth RS. (1974):

The Development of the infant and Young Child, 5th ed. Edinburg, Scotland: Churchill Livingstone.

Imaizumi S, Wollworth V, and Lishman RA (1989):

Superoxide dismutase activities and their role in focal cerebral ischemia. *J. Cereb. Blood Flow Metab*; 9:5217.

Jacobs SE, Stewart M, Inder TE, Doyle L, Morley C. (2002):

Feasibility of a pragmatic randomized controlled trial of whole body cooling for term newborns with hypoxic-ischaemic encephalopathy. December 9-10, Hot Topics in Neonatology Meeting Proceedings, Washington, D. C.

Jeffery M. Perlman, MB Rollins, MD (2000)

Surveillance protocol for the detection of intracranial abnormalities in premature neonates.

Arch Ped. Adolesc. Med. vol. 154 (522-826), Aug 2000.

Jepson HA, Tala Shek ML and Tichy AM (1991):

The Apgar score: evaluation, Limitations and scoring guidelines. *Birth* (18): 83-92.

Johnson M.L., Dunne M.G., Mack L.A, Kashbaum C.M. (1980):

Evaluation of fetal intracranial anatomy by static and real time ultrasound *journal of clinical ultrasound*, 8: 311 -318.

Kalifa G. (1986):

Pediatric ultrasonography springer-verlag, Berlin H., Delburg, New York, Tokyo.

Kirimi, E.; Tunecur, O.; Atasa, B. et al. (2002):

Clinical value of Doppler ultrasonography measurements in full term neonates with perinatal asphyxia, and hypoxic ischemic encephalopathy in the first 12 hours of life and long term prognosis *Tohoku. J.Exp. Med.* 197, 27-33.

Kossoff G., Garrett W.J., Kadovanovich G., (1974):

Ultrasonic atlas of normal brain of infant, ultrasound in Medicine and biology; 1: 259 – 266.

Lawrence AJ and Jarrott B (1993):

Nitric oxide increases interstitial excitatory amino acid release in the rat dorsomedial medulla oblongata. Neuroscience Letters 151:126-129.

Leksell L., (1956):

Echencephalography: Detection of intracranial complications following head injury. Acta chin.soud 110:301-315.

Levene MI, Gibson NA, Fenton AC, Papathoma E, Barneti D. (1990):

The use of a calcium-channel blocker, nicardipine, for severely asphyxiated newborn infants, Dev Med Child Neurol. 32: 567-574.

Levene MI (1993):

Management of the asphyxiated full term infant. Arch. Dis. Child; 68:612.

Levene MI (1995):

The asphyxiated newborn infant: The clinical diagnosis of birth asphyxia. In: Levene. MI, Lifford RJ, Bennett MJ, and Punt J (Eds). Fetal and Neonatal Neurology and Neurosurgery. Churchill Livingstone. Edinburg. (ed.).427-442.

Malcolm I. Levene, Jonattan L.W., Claire and Lise fawer, (1985):

Ultrasound of the infant brain: Scanning techniques and normal anatomy. Clinics in developmental medicine no. 92, Philadelphia J.B Lippincott co.

Mallard EC, Gunn AJ, Williams CE et al. (1992):

Umbilical cord occlusion causes cerebral damage in the fetal sheep. Am. J. Obstet. Gynecol; 167: 1423-30.

Marilyn J. S, Gary D. S, Jeffrey M, and Keith H. F (1984):

Hypoxic-ischemic Encephalopathy in Term Infants: Diagnosis and Prognosis Evaluated by Ultrasound' Radiology; 152: 395-399

Marinelli PV. (1983):

The asymmetric tonic neck reflex. Clin. Pediatr. 22:544-546.

Marret S, Gressens P, Gadisseux JF, Evrard P, 1995:

Prevention by magnesium of excitotoxic neuronal in the developing brain: an animal model of clinical intervention studies Dev Med Child Neurol. 37:473-484.

Martian-Ancel A., et al. (1995):

Multiple organ involvement in perinatal asphyxia. J Pediatr. 127:786.

Medoff-Cooper B, Verklan T. Carlson S. (1993):

The decelopment of sucking patterns and physiologic correlates in very-low-birth-weight infants. Nurs. Res. 42: 100105.

Nagdyman N., et al (2001):

Early biochemical indicators of hypoxic-ischemic encephalopathy after birth asphyxia. Pediatr REs. 49: 502.

Ny Boo, V Chandran, Ma Zulfiqar, Sm Zamratol, Mk Nyein, Ms Haliza, Ms Lye (2000)

Early cranial ultrasound changes as predictors of outcome during first year of life in term infants with perinatal asphyxia Journal of Paediatrics and Child Health 36 (4), 363–369.

Palmer C and Vannucci RC (1993):

Potential new therapies for perinatal cerebral hypoxia-ischemia. In: Shankaran S. (ed.) Perinatal asphyxia. Clin. Perinatol.; 20(2): 411, W.B. Saunders Co., Philadelphia

Patt A, Horesh IR and Berger EM (1990):

Iron depletion or chelation reduces ischemia reperfusion-induced edema in gebril brains. J. Pediatr. Surg; 25:224.

Perlman JM, Tack ED, Martin T et al, (1989):

Acute systemic organ injury in term infants after asphyxia. Am. J. Dis. Child., 143:617.

Peroutka SJ, Peroutka LA. (1984):

Autosomal sominant transmission of the “photic sneeze reflex,” N Engl. J. Med. 599-600.

Prechtl H, Beintema D, (1964):

The Neurologic Examination of the Full-Newborn Infant. London: SIMP Heinemann; Cli. Dev. Med. Vol. 12.1.

Prechtl H, (1977):

The neurological Examination of the Full-Term Newborn Infant. 2nd ed. Philadelphia: JB Lippincott. Clin. Dev. Med. Vol. 63.1.

Rennie JM and Robertson NRC (2002):

Resuscitation of the newborn. In: Rennie JM and Robertson NRC (eds.) Manual of Neonatal Intensive Care, 4th ed. p68.

Rhine W. D., and Blankenberg F. G, (2001):

Cranial Ultrasonography
NeoReviews Vol.2 No.1 3-11

Rivikin MJ and Volpe JJ (1996):

Asphyxia and brain injury. In: Spitzer AR. (ed.) Intensive Care of the Fetus and Neonate; 59:685. Mosby-Year Book, Inc; USA.

Robertson C and Finer N (1993):

Long term follow up of term neonates with perinatal asphyxia. Clin. Perinatol Jun., 20(2):483-500.

Roberts DS, Haycock CB, Dalton RN et al (1990):

Prediction of acute renal failure after birth asphyxia. Arch. Dis. Child. 65:1021-8.

Rosenkrantz T, Zalneraitis L and Napolitano E (1991):

Prediction of survival in severely asphyxiated infants. Pediatr

Neurol Nov. 7(6): 446-51.

Sarnat H and Sarnat M (1976):

Neonatal encephalopathy following fetal distress. Arch. Neurol.,
33:696.

Scalais E, Francois-Adant A, Muttin C, Bachy A and Cuesrit JM (1998):

Multimodality evoked potentials as a prognostic tool in term asphyxiated newborns. *Electroencephalography & Clinical Neurophysiology*, 108(2): 199-207.

Schenk-Rootlieb AJ, van Nieuwenhuizen O, van der Graaf Y, et al, (1992):

The prevalence of cerebral visual disturbance in children with cerebral palsy *Dev. Med Child Neurol* 34: 473-480

Schneider H and Herrmann U (1990):

Prolonged pregnancy Pathophysiology and clinical aspects. *Gebunshilfe Frauenheilkd*; 50: 3.

Shankaran S, Laptook A, Wright LL, Ehrenkranz RA, Donovan EF, Fanaroff AA, et al. (2002):

Whole-body hypothermia for neonatal encephalopathy: animal observations as a basis for a randomized, controlled pilot study in term infants. *Pediatrics*: 110: 377-85.

Shankaran S, Laptook AR, Ehrenkranz RA, Tyson JE, McDonald SA, Donovan EF, et al. (2005):

Whole-body hypothermia for neonates with hypoxicischemic encephalopathy. *N Engl J Med*: 353: 1574-1584.

Siesjo BK (1992):

Pathophysiology and treatment of focal cerebral ischemia. Part 2: Mechanisms of damage and treatment. *Journal of Neurosurgery* 77:337-354.

Singh M (1994):

Diagnosis and Management of perinatal asphyxia: Current concepts. *Indian Pediatr*: 31(10): 1169.

Snyder EY and Cloherty JP (2005):

Neurology: Perinatal asphyxia In: Cloherty JP and Stark AR (eds.) Manual of Neonatal Care. 4th ed., 27: 515. Little. Brown and Co, Boston.

Stanton WR, McGee R and Silva PA (1991):

Indices of perinatal complications, family background, child rearing and health as predictors of early cognitive and motor development. *Pediatr.*; 88: 954.

Stark JE, Seibert JJ. (1994):

Cerebral artery Doppler ultrasonography for prediction of outcome after perinatal asphyxia. *J.*

Steel, R.G.D. & Torrie, J.H. (1960):

Principles and procedures of statistics, with special reference to the biological science McGraw Hill Book company Inc., New York, Toronto, London.

TOBY Trial, <http://www.npeu.ox.ac.uk/toby/>**Van Bel F, Dorrepaal CA, Benders MJ, Zeeuwe PE, Van Debor M, Berger HM (1993):**

Changes in cerebral hemodynamics and oxygenation in the first 24 hours after birth asphyxia. *Pediatrics* 92:365-372.

Van Bel F and Walther FJ (1990):

Myocardial dysfunction and cerebral blood flow velocity following birth asphyxia. *Acta Paediatrica Scandinavia.* 79:756-762.

Vannucci RC (1990):

Experimental biology of cerebral hypoxia-ischemia: relation to perinatal birth damage. *Pediatr. Res.* 27:317.

Volpe JJ (1992):

Perinatal hypoxic-ischemic brain injury. In: Fukuyama Y, Sazuki V, and Komoshita S et al. (eds.) *Fetal and Perinatal Neurology*: p. 232, Basel, Karger

Volpe J.J. (1995):

Hypoxic-ischemic encephalopathy. In: Neurology of the newborn, 3rd ed.; p. 250. W.B. Saunders Co., Philadelphia.

Volpe J.J. (1995):

Neurology of the Newborn. 3rd ed. Philadelphia: WB Saunders.

Volpe JJ (2001):

Hypoxic ischemic encephalopathy. In: Neurology of the newborn, 4th ed. p.217- 394, W.B. Saunders Co; Philadelphia.

Williams CE, Mallaard C, Tan W et al., (1993):

Pathophysiology of perinatal asphyxia. In: Shankaran S. (ed.) perinatal asphyxia. Clin. Perinatol.; 20(2): 305-25.

Wyatt JS, Edwards AD and Azzopardi D (1989):

Cerebral hemodynamic during failure of oxidative phosphorylation following birth asphyxia. Pediatr. Res., 26:511.

Wyatt JS, Cope M and Delpy DT (1990):

Measurement of optical length for cerebral near-infrared spectroscopy in newborn infants. Developmental Neuroscience, 12: 140-144.