

REFERENCES

- ❖ *Adams J, Bodor GS and Davla R(1993)* : Cardiac troponin-I circ., 88:101-106 .
- ❖ *Adams J, Schechtmann KB and Landty S (1994)* : Compatible detection of acute myocardial infarction. Clin chem , 40 : 1291 – 95 .
- ❖ *Agarwal R, Klair-el Din, Graybusn PA and Victor RG (1993)* : Effects of cooler temperature on heamodynamic ability in “problem” dialysis patients. kidney Int. ; 44 : 606-612 .
- ❖ *Agnes Fogo and Valentina Kon (1999)* : Pathophysiology of progresssive renal disease, from pediatric nephrology, 4th Ed: T. martin, Ellis D., Harmon P : 1183 –85 .
- ❖ *Alfery AC and Chan L (1992):* Pathogenesis of CRF, In renal and electrolyte disorders, 4th ed., Schrier RW, Brown and Company, Boston, P:568.
- ❖ *Anand IS, Ferrarir A and Kalra GS (1989):* Edema of cardiac origin, studies of body water, sodium, renal function, heamodynamics, and plasma hormones in untreated congestive heart failure. Cric .;30:299.
- ❖ *Antemann EM, Gradizenc and Sacks D(1995):* Evaluation of rapid assay or detection of serum CtnI . J-Am-Chem, 273-1279.
- ❖ *Antila K, Forsstornn J and Valimaki L.(1986):* Effects of heamodialys is an heart rate variability in chronic renal failure . scand J clin Lab Invest ; 46: 546 – 550 .
- ❖ *Apajasale M, Rautomes J and Sintones H (1997):*Health related quality of life after organ transplantation in children. Ped. Transplant, P:B 0-7 .
- ❖ *Apple FS(1999):* The specificity of biochemical markers of cardiac damage; A problem solved. Clin. Chem. Lab. Med. 37(11-12): 1085-9 .
- ❖ *Balland N, Saint-Remy M and Collignon P.(1982):* Heamo dynamic study of hypotension during heamodialysis, Kidney Int.; 21:868-76 .
- ❖ *Bauer JD (1982):* Enzymology in clinical laboratory medicine. Mosby company st. louis, Toronto, loudon, 9th ed., P: 442-504.

- ❖ **Bennett WM, Muther RS, Parker PA and Fegi P (1980):** Drug therapy in renal failure: Dosing guidelines. *Ann Intern Med*; 93: 286-325.

- ❖ **Bodor GS, Porter S and Landty D (1992):** Development of monoclonal antibodies for and assay of cardiac troponin I *clin chem.*, 38:2203-2205

- ❖ **Brenner BM(1998):** Neurological complications of chronic renal Failure. In: *kid. Int.* p: 1520-36, W.S. Saunders, Philadelphia.

- ❖ **Brenner BM. and Meyer TW;(1986):** The distribution of glomerular hemodynamic alterations to progressive renal disease, New York: churchill living store,: 1-16 .

- ❖ **Carl Leier and Harisios B(1997):** Renal disorders and heart disease. From Braunwald's text book of cardiovascular disease, 5th ed., Eds: Eugene Braunwald, Saunders Company, Philadelphia.

- ❖ **Chanard J, Haguet M, Basile C and Asson R (1980):** Carnitine improves lipid anomalies in hemodialysis patients. *Lancet*; 2:763-64 .

- ❖ **Chenseny RW and Avioli LV(1992):** Childhood renal osteodystrophy. In: *Ped. Kid. Dis.*, 2nd ed, Eds, Edelman CM, Brenstien J and Meadow, Brown and Company, Bosron; P:647-74 .

- ❖ **Cohn IN and Burke LP(1980):** Nitroprusside. *Ann intern Med*; 91:752-757 .

- ❖ **Creasman MR, Burrel DE and Tolins PS (1992):** Hypertension associated with CRF (pathology and management), *Am J cardiol*; 9:215-17 .

- ❖ **Cummins B (1987):** Cardiac specific troponin I release in canine experimental myocardial infarction. *J Moll cell cardiol*; 19:994-1010 .

- ❖ **Cupta BX, Spinowitz BS, Charytan C and Wahl SJ(1993):** Cholesterol crystal embolization – associated renal failure after therapy with recombinant tissue-type plasminogen activator. *Am J kidney Dis.*, 21:659 .

- ❖ **Daugiradas JT, Leehy DJ and Mechary GM (1966):** subxiphoid pericardiostomy for hemodialysis associated pericardial effusion, *Arch intern Med*; 146: 1113-1115.

- ❖ **David G.W.(1996):** Diagnosis and management of chronic renal Failure. In cecil text book of medicine, 20th ed. Eds, claude Bennet and Fred plum.
- ❖ **Davis I., Bundsman TE and Grionon PC(1998):** Pediatric rend Trans: indications and special consedirations Ped. Transplant; 2: 117-129 .
- ❖ **De winter RJ and Sanders GT(2001):** Determination of cardiac troponins for diagnosis of Acute myocardial infarction Ned Tijdschr Geneeskd 10; 145 (10): 461-6 .
- ❖ **Eidelson B., Kim KE, Onseti G, Greco J and Fernandes M(1991):** Sequential heamodynamic changes in end-stage renal disease during volume expansion and hypertension; 2:102-110 .
- ❖ **Eknoyan G and Lister BJ(1985):** Renal complications of bacterial endocarditis. Am J Nephroll; 5:477 .
- ❖ **El-Nahas AM.(1996):** Chronic renal failure and its treatment, in oxford text book of medicine, 3rd ed. Eds: Dj weatherall, G.ledingham, D.A.warrell 20:3294-96 .
- ❖ **Ersleu AJ and Besarab A(1997):** Erythropoiten in the pathogenesis and treatment of anemia of chronic renal failure. Kid. Int; 51:622 (production, action) and therapentic use of erythro poitien in anemia of CRF.
- ❖ **Feinstien EI, et al.(1985):** Renal complications of heart disease, Am J Nephrol.
- ❖ **Feld LG,Stablein D and Fivash B. (1997):**Annual report of the north America pediatric renal and transplantation cooperative study 1:146-162.
- ❖ **Fennel EB, Fennel RS and Mings E(1986):** The effects of various modes of therapy for ESRD on cognitine performance in ped.population. Int J Ped Nephrol; (2): 107-12 .
- ❖ **Ferrori R, Kalra GS and Colly K(1989):** Edema of cardiac origin cirDculation; 30:299 .

- ❖ **Foreman JW and Chan JCM (1988):** Chronic renal failure in infants and children. J Pediatrics: 113:793-800 .
- ❖ **Franz Schafeer and Otto Mehls (1998):** Endocrine and growth disturbance in children with CRF, from Ped. Neph. Text book, 4th ed; 79:1197-1000 .
- ❖ **Friedmann HS ,Shah BS ,Kin HG ,Govel A and Smith JA(1981):**Clinical study of cardiac findings in CRF(abstract).East Africa Med J,70(2);107-111.
- ❖ **Fromme RE, Meyer D, and Zimmorman J(1998):** comparison of CK-MB subforms cardiac troponin T and I as alternative diagnostic standards to CK-MB mass in a randomized prospective blinded trial, circ. 98:1-423 .
- ❖ **Fujimoto S,Kogoshima T,Hashimoto T and Dahn K (1994):**Left ventricular diastolic function in patients on regular HD.Clin Nephrol ;42(2):109-116.
- ❖ **George B, and Dalton R(1998):** Laboratory investigations of renal disease, from text book of Ped.Neph., 4th de, Eds; T.martin, Ellis, William Harman, 20: P : 343-365 .
- ❖ **Gerhard T, Katus H and Ravkilde J (1991):** serum troponin T in suspected ischemic myocardial damage compared with mass and catalytic concentration of serum CK-MB. Clin Chem, 37:1405-11 .
- ❖ **Gilli G, Mehls O and Scharner R (1984):** Final hight of children with CRF, proc. Eur. Dial Transp. Association, 21: 830-36 .
- ❖ **Gold Blum and Read WP (1980):** Host defense and immunologic alternations associated with chronic heamodialysis. Am Intern Med ; 93:547-613 .
- ❖ **Grayburn PA, Victor RG and Anrich WL (1993):** Reduction of the incidence of hypotension during heamodialysis, Am J Kidney Dis; 5: 712-18 .
- ❖ **Green EL, Raji L.and Makw(1992):** Cardiovascular risk factors in chronic renal failure and heamodialysis patients-Am J Kidney Dis, 19: 505-513 .

- ❖ **Gretz N (1983):** Predictability of the progression of CRF. *Kid. Int.*; 24(suppl 15): S2- S5.
- ❖ **Grooms P, Dusso A, Windus DW and Tindira C (1991):** Parathyroid hormone suppression by intravenous 1.25 -dihydroxy vit. D. *J clin Invest*; 83:1349-55 .
- ❖ **Guerin A(2000):** Principal cardiovascular complications in end-stage renal disease. *Prees-Med.*; 29(5): 274-80 .
- ❖ **Guler N, Bilge M, Eryonucu B, uzum K, Avci ME and Dulger H(2001):** Cardiac troponin I levels in patients with left heart failure and cor pulmonale, Departement of cardiology, school of medicine, Turkey, *Angiology*; 52(5): 317-22 .
- ❖ **Harisios Boudolas and Lier CV(1997):** Cardio-renal disorders and diseases, from Brounwalds text book of cardiac medicine, 4th ed.
- ❖ **Harman WE and Jabs KL (1999):** Heamodialysis: In. pediatric nephrology Eds: Martin T, Ellis D, william E. 4th ed. Lippincott, Williams and Wilkins. Baltimore, Maryland, P:1267-87 .
- ❖ **Harmiller JB (1984):** Association between central heamodynamics and blood flow in patients with congestive heart failwa.
- ❖ **Herrmann J, Eggebrecht H and Malyar N.(2001):** "Biochemical markers of ischenic and non ischemic myocardial damoge", *Med klin* 15 ; 96 (3): 144-156 .
- ❖ **Hokken AC, Van Zeal and Van Bergen (1999):** Final hight and its prediction factor after renal transplantation in childhood, *pediatrics* 36:323-28 .
- ❖ **Houston MC ,Mcchesney JA and Chaterejje K(1981):** Pericordial effusion associated with minoxidil therapy .*Arch Inter Med*;141:69-71.
- ❖ **Huting J(1993):** Course of left ventricular hypertrophy and function in end stage renal disease after renal transplantation. *Am.J. Cardiol.* 70:1481 .
- ❖ **James L and William E (2000):** Patho physiology of anemia: in : The kidney diseases, 8th ed., Eds, Bany M, Sammul A, WB Sanclers Company, Philadelphia, London, Tranto, Tokyo, P:2062 .

- ❖ *Jsky Gordon and Rose de burgn (1998)*: Diagnostic imaging of renal disease; from text book of Ped. Meph., 4th edy 22:P:377-391 .
- ❖ *Katus HA, Remppis S and Neumann FJ (1991)*: Diagnostic defeciency of troponin T measurements in acute myocardial damage. Circulation, 83:902-912 .
- ❖ *Katus H, Scheffold T and Remppis A (1992)*: Proteins of troponin complex. Lab Med, 23:311 .
- ❖ *Kelly RA, O'hara DS, Mitch WE and Solmon HS (1986)*: Endogenous digitalis-like factors in hypertension and elmonic renal insufficiency. Kidney Int.; 30:723-729 .
- ❖ *Klahr S., Levy A. and Beck WE(1999)*:The effects of diatary protein restriction and blood pressure control on the progression of CRF, N.Engl J Med, 330:877.
- ❖ *Koo W and Kaplan LA (1988)* : Aluminum and bone disorders with specific referance on aluminum contamination of infants nurtients, J.AM. Nut; 7(3): 199-214
- ❖ *Kuch W and George ER (1982)*: Clinical pharmakokinetics of Atenolol. Eur J Drug Metab pharmacokinet: 7:81
- ❖ *Lai KN ,Whitford J ,Fassett RG and Mathew (1993)*:Left ventricular function in uremia;Echocardiographic assesment in patients on regular HD.Clin Nephrol .23(3);125-133.
- ❖ *Lazarus JM (1990)*: Diagnostic imaging of renal disease; from text book of Ped. Meph., 4th ed, 22:P:377-391 .
- ❖ *Leir C.V. (1992)*: Cardio-renal disorders; from Brawnwald's text book of Medecine, 3rd ed., Eds:Eugene Brawnwald, Saundus Company,Philadelphia.
- ❖ *Leir MF and Magorien RD (1984)*: Relationship between central heamodynamics and regional blood blos in normal subjects and in patients with congestive heart failure, circulation 69:57 .

- ❖ **Levinsky NG (1993):** The organisation of medical case, lessons from the medicase End-stage renal disease programm, New England Journal of Medecine 329:1395-90
- ❖ **Levy AS (1990):** Measurements of renal function in chronic renal disease, Kid. Int; 38:167-184 .
- ❖ **Levy A, Beck GJ and weerts E (1994):** Factors effects the progression of chronic renal failure. N Engl J Med., 329:150-160 .
- ❖ **Llach F and Bover J (2000):** Renal osteodystrophy. In: The kidney diseases, Eds; Brenner BM, 6th ed., Philadelphia. P:2103-66 .
- ❖ **London GM, Marchais SJ, Gurein AP and Pannier B(1993):** Cardiac hypertrophy and arterial alternations in ESRD. Kid Inter ,43:supple41;S42-S49.
- ❖ **Logan MG (1990):** Renal effects of atrial natriuretric factor, Annual report; 52:699 .
- ❖ **Luscher TF, wenzel RR and Moreu P. (1995):** Vascular protection effects of ACE inhibitors and Ca antagonists therapeutic basis for a combination therapy in hypertention and other CVD; 9:509-523 .
- ❖ **Macdouglass IC (1998):** Treatement of renal anemia with rh EPO. Curr opin Nephrol; 1:210-219 .
- ❖ **Maher RE, Robertson LM and Curtis JR (1985):** Calcific Aortic stenosis in chronic renal failure . Lancet; 2:1007 .
- ❖ **Maher JF ,Bryan CW and Ahearn PS (1975):** Prognosis of chronic renal failure and factors affecting survival. Arch Inter Med, 135:273-8.
- ❖ **Mair J, Dient LF and puschendorf B. (1992):** Cardiac specific troponin T in the diagnosis of myocarial injary. Clin lab Med, 29:37-35
- ❖ **Mann JFE (1995):** Hypertension and cardiovascular effects-long term safety and potential long term benefits or r-HuEpo. Nephrol Dial Toansplant; 10(suppl 2): 80-84 .

- ❖ *Martin M, Ralf S and Ludwig K (1999):*Prognostic values of cardiac troponin T and I elevations in renal disease.Nephrol Dial Transp ;14:1489-95.
- ❖ *Marchais. S (2000):* Principal cardiovascular complications in end-stage renal disease. Press-Med; 29(5): 274-80 .
- ❖ *Mathew P ,Suarez W,Kip K and Bayar E (2001):*Is there a potential role for serum CTnI as a marker for myocardial dysfunction ?Cancer Invest ;19(4):352-9.
- ❖ *Meeus F, Kourilsky O, Gurein AP, Growdry C and Marchais SJ (2000):* Pathophysiology of cardiovascular disease in heamodialysis patients. Kidney Int.; 58 (76) : 5140-7 .
- ❖ *Meinerts T and Hamm CW (1998):* Rapid testing for cardiac troponin in patients with a cute chest pain in Emegency room. European heart J.:19;973-4 .
- ❖ *Methenge RN ,Mcligeyo AK, Mutia AK and Otimo LS (1993):*The specificity of ECHO findings in CRF.East Arican Med J,70;107-117.
- ❖ *Miache H,StephenG and Edwin AR (1981):*Cardiac diseases in dialysis patients .clinical dialysis (2nd ed.)Eds:Allen RN ,Richard N and Lona P.p:405.
- ❖ *Mitch WE (1991):* Dietary protein restriction in patients with CRF, Kid. Int.; 40:326-391 .
- ❖ *Morris KP ,Skinner JF, and Coulthand MG (1993):*Cardiac abnormalities in ESRD and anemia .Archeves of disease in childhood .66;637-642.
- ❖ *Neugarten and Baldwin DS (1994):*Glomerulonephritis in bacterial endocarditis ,Am J Med .77:297.
- ❖ *Norwood K, Slatapolsky E and Weerts(1998):*Long term effects of Ca carbonate on mineral metabolism.kid int ;36:897.
- ❖ *Osanloo E, parker RH and shalhoub RJ (1989):*Viral pericarditis in patients receiving heamodialysis. Arch Inter Med, 139:301-303 .

- ❖ **Packer M and Lee WH (1987):** Effects of diabets mellitus on changes in ventricular performance and renal function produced by ACE inhibitors in patients with sever chronic heart failure. Am.J. Med.82:1119.
- ❖ **Pacher M, Lee WH and Kessler PD (1988):** Preservation of glomerular filtration in human heart failure by activation of renin angiotensin system. Circ.
- ❖ **Pannier B and Guerin AP(1994):** Cardic hypertrophy, Aortic compliance, peripheral resistance and wave reflection in end stage renal disease; comparative effects of ACE inhibitors and calcium.channel blochage. Circ 90:278 .
- ❖ **Pannier B (2000):** Cardiac anomalies in chronic renal failure. Press – Med; 29(5): 274-80
- ❖ **Parfery PS and Harnett JD (1994):** Long term cardiac morbidity and mortality during dialysis therapy .Arch Nephrol Necker Hosp.;23:311-330.
- ❖ **Persand J, chan MK and Varghese Z (1991):** pathogenic roles of post-heparin lipases in lipid abnormalities in heamodialysis patients. Kidney Int. 25:812-818 .
- ❖ **Peterson JC(1995):** Chronic renal failure In:Pediatric Nephrology. 3rd edition , Ticher CC and Wilcox CS. Eds. Mass publishing Co., William and Wilkins, London. P:443 .
- ❖ **Pickering TG, Devereux RB and James GD (1988):** Recurrent pulmonary oedema in hypertension due to bilateral renal artery stenosis; Treatment by angioplasty or surgical revascularization. Lancet 2:551 .
- ❖ **Plebani M and Zaniomotto M(1999):** Cardiac markers ,present and feature. Int-J-Clin-lab, 20:56-63 .
- ❖ **Prasad AS (1963):** Zinc metabolism in patients with Fe defeciency anemia, HSM, dwarfism. Journal of clin. Med, 61:537 .
- ❖ **Purkerson MI and Cole BR (1990):** CRF in ped., text book of fluid and electrolyte; Eds; William and Wilkins P:242.

- ❖ **Reach J, Dennies C, Rouby JJ, Di Giulio and Aime F (1981):** Epidemiology of dialysis induced hypotension, Proc Eur Dial Transplant Assoc 18:133-138 .
- ❖ **Rector TS, Kubo SH, Bank AJ and S.M (1991):** Endothelium dependent vasodilation in attenuated in treatment of heart failure . circ; 88:55 .
- ❖ **Rees L., Ridgen SP and Word GM (1989):** CRF and Growth, Arch. Dis. Chil. 64; 573-77 .
- ❖ **Rigeal W. (1993):** Complications of systemic arterial hypertension specially on renal system., Am J cardiol, 5; 110-119 .
- ❖ **Robert WC, Perloff J and Baylar EN (1972):**Mitral valve disease ;surgery of the condition causing the mitral valve to function abnormally .Ann Int Med,77:939-975.
- ❖ **Robinson RM, Formann MB and Viramanir D (1984):** Mitral annular calcification in chronic renal failure. Chest; 85:367-371 .
- ❖ **Rodeheffer RJ, Naruse M and Atkinson JB (1993):** Molecular atrial natriuetric factor in normal and failing human myocardiam circ.; 88:364 .
- ❖ **Ross EA and Barri M (1995):** Heamodialysis In: Nephrology. Eds: Tisher CC, Wilcox CS. 3rd ed. Mass publishing co., Williams and Wilkins. London. Tokyo, P:1070-1120 .
- ❖ **Rostand SG, Brown G and Kirk K (1989):** Renal insuffeciency in treated essential hypertension. N Engl J Med 320:684 .
- ❖ **Rostand SG, Gretes JG and Kirk KA (1990):** Ischemic heart disease in patients with uremia undergoing maintanance heamodialysis. Kid-Int; 16:600-611 .
- ❖ **Rostand SG, Brunzell JD, Cannan KO, and victor RG (1991):** Cardiovascular complications in renal failure . J Am soc Nephrol ; 2:1053-62 .
- ❖ **Rudnick MR (1994):** Nephrotoxic risks of renal Angiography,contrast media-associated nephrotoxicity. Am J kid D is, 24:713-727 .

- ❖ **Rutsky EA (1990):** Pericarditis in end-stage renal disease, *cardiol clin*; 8:701-707 .
- ❖ **Sacks DB (1999):** Acute coronary ischemia. *Vasc Med*, 49:253-259 .
- ❖ **Schoefer F., Weicek A and Ritz E (1998):** Endocrine disorders, In: *Oxford text book of clin. Neph.*, 2nd ed., Eds, Alexander M, J.stewants, Jean pearre; Oxford-New York Tokyo, Oxford University, P:1854-65 .
- ❖ **Schrier RW (1988):** Pathogenesi of sodium and water retention in high low output cardiac failure, nephrotic syndrome, cirrhosis and pregnancy. *N Engl J Med*.
- ❖ **Shichiri M, Hirata Y, Ando K, Ohta K and Kimoro S (1990):** Plasma endothelin levels in patients with uremia, and hypertension; 15:493-496.
- ❖ **Sobki SH, saadeddin SM and Habbab MA (2000):** "Cardiac markers used in the detechion of myocardial injury"; *saudi Med J*; 21(9): 843-6
- ❖ **Sohn HJ, stokes Gs and Johnston H (1983):** A controlled study of the effect of indomethacin in uremic pericaritis. *Kidney Int*; 24:663-669 .
- ❖ **Stephen O. pastan (1998):** The heart and kidney diseases, from *Hursts text book of cardiac diseases* 94:2410-21.
- ❖ **Steven J., and Michelle Baum (1998):** Physiology and management of CRF in pediatric age group , from *pediatric nephrology* 72:1156-58 .
- ❖ **Tabak MA, Christenson PC and Fine RN (1986):** Prediction of progression of CRF in children are current models accurate ? *ped.*; 78:1007-1012 .
- ❖ **Teitlebaum SL, Auska KA and George R (1995):** Renal osteodystrophy. *N Engl J Med*; 33:166-74.
- ❖ **TermanJ ,Ruedin P,Monin C and Lerch R (1971):**Effects of altered loading conditions during HD on left vent.filling pattern. *Eur Heart J*,14;p:655.

- ❖ **Tonshoff and Fine (1995):** Growth and growth hormone treatment in children with chronic renal insufficiency . In: clinical dialysis, 3rd ed., Allen R, Nissenson, Richard, N. Fine. P 567-79.
- ❖ **Turner C, Copston J. and Mack RH (1998):** Bone turnover and 1.25 dihydroxy cholecalciferol with phosphate binders, Kid. Int.; 3:189-59 .
- ❖ **USRDS** (United states Renal Data System); Annual report, Am J kidney Dis, 1994; 24(suppl 2):48-56 .
- ❖ **Vatikus PT (2000):** Current status of prevention, diagnosis and management of coronary artery in patients with CRF Am. Heart J; 139 (6): 1000-8 .
- ❖ **Volbracht L, Haude M, Mann K and Erbel R (2001):** "Biochemical markers of ischemic and non-ischemic myocardial damage", Mcd Kli 51; 96(3):144-156 .
- ❖ **Walkers K (1997):** General management of end stage-renal disease (Description and management of non-renal complications of end-stage renal disease, this includes cardiovascular disease, anemia, diet, hypertension, hyper lipedemia and hyper parathyroidism) .
- ❖ **Warady BA, Herbet D and Sullivan EK. (1997):** Renal transplantation and chronic renal failure in children and adolescents, pediatric nephrology 11(1):49-64.
- ❖ **Warnock PG (1996):** Uremic acidosis . Kid. Int. 34:278, A review of the renal responses to chronic acidosis, the importance of chronic acidosis in the development of renal osteodystrophy .
- ❖ **Warnock PG (1996):** CRF in: Cecil textbook of medecine, 20th ed., Bennetle JC and Alen F, Eds, WB Sauders Co., Philadelphia, P:586 .
- ❖ **Wassner SJ (1982):** The role of nutrition in the case of children with CRF, ped. Clinics of North America; 9:973 .
- ❖ **Wassner SJ and Alexander S (1996):** Annual report, vol. 1: Dialysis and Transplantation . candidian organ replacement register, candidian institute for health information: 3-16 .

- ❖ **Wassner SJ and Baum M (1999):** Physiology and management of CRF, from ped.neph., Eds: Barrat TM, Avener ED, Harman WE, 4th ed., Lippincott William and Wilkins P: 1155-82 .
- ❖ **Weber J and Campese VM (1984):** Renal Hemodynamic effects of vasodilatation with nifedipine and hydralazine in patients with heart failure . J Am coll. Cardiol .
- ❖ **Weber H, Joskocs G and Schwarzer C (1984):** Chronic hemodialysis high risk patient for arrhythmias ? Nephrol ; 3:180-185 .
- ❖ **Whilehead VM (1979):** Renal dialysis treatment and hematological system In: Hematological problems in renal disease. Eds; Jepson M, Addison wesley publishing Co., California USA P:145.
- ❖ **William E Mitch (1998):** The heart and kidney disease. From Hursts text book of cardiology 94:2410-21
- ❖ **William E. Harmon, Ellis D. and T. Martin (1999):** Overview of chronic renal failure, from text book of pediatric nephrology, lippincott williams and wilkins, Baltimore, Maryland- P:1151-1154 .
- ❖ **Wilmore DW (1991):** Management of dialysis associated pericarditis in children with regular hemodialysis . J Am soc Nephrol ; 2:130-35 .
- ❖ **Windus DW, Tindira C and Slatopolsky K (1991):** Parathyroid hormone suppression by intravenous 1.25-dihydroxy vit. D. J clin Invest; 83:1349-55 .
- ❖ **Wing AJ, Brunners FP, Brynger H, Jacobs C and Karmer P (1984):** Cardiovascular related causes of death and the fate of patients with renovascular disease. Contrib Nephrol; 41:306-3110 .
- ❖ **Winters KJ and Eisberge PR (1998):** Ischemic heart diseases, The Washington manual of medical therapeutics carey H (ed)Lippincott-Raven-Philadelphia, New York, 29th edition, P 85-94 .
- ❖ **Zdon MJ and Crawford (1993):** Renal micro embolization syndrome. A cause for renal dysfunction after abdominal aortic .
- ❖ **Zeigler TR, Lazarus JM and young LS (1991):** Uremic pericarditis in acute and chronic renal failure 16: 156-79