

REFERENCES

- Adib M., Ostadi V., Navaei F., et al., (2007):** Evaluation of CD11b Expression on Peripheral Blood Neutrophils for Early Detection of Neonatal Sepsis. Iran J Allergy Asthma Immunol, June 2007; 6(2): 93-96
- Agerberth B., Charo J., Werr J., et al. (2000) :** The human antimicrobial and chemotactic peptides LL-37 And alpha defensins are expressed by specific lymphocyte and monocyte populations. Blood; 96:3086-93.
- AL-Mulla ZS. And Chrisensen RD. (2005):** Neutropenia in the neonates. Cli. Perinatal, 22:711-739
- Apgar B.S., Greenberg G., and Yen G. (2005):** Prevention of Group B Streptococcal Disease in the Newborn. Am fam Physician; 71 (5): 903-10. Arch Dis Child, in press.
- Artman M, Mahony L, Teitel D:** Approach to the infant with excessive pulmonary blood flow. Neonatal Cardiology 2002.
- Avroy A, Fanaroff RY. (2002):** Neonatal- Perinatal Medicine: Diseases of the Fetus and Infant. USA, Lowis: Mosby, 2002: 360-91.
- Baird T., Martin R. and Abu-Shaweesh J.:** Clinical associations, treatment, and outcome of apnea of prematurity. NeoReviews 2002; 3
- Balkundi DR, Nycyk JA and Cooke RW (1994):** Immunisation and C reactive protein in infants on neonatal intensive care units [letter]. Arch Dis Child. 1994; 71:F149
- Ballard JL, Khoury JC, Wedig K, Wang L, Eilers-Walsman BL, Lipp R. (1991):** New Ballard Score, expanded to include extremely premature infants. J Pediatr.; 119(3):417-23.
- Ballard JL, Novak KK, and Driver M. (1979):** A simplified score for assessment of fetal maturation of newly born infants. J Pediatr.; 95(5 Pt 1):769-74.
- Baltimore R.S. (2003):** Neonatal species: Epidemiology and management. Pediatr Drugs; 5(11): 723-40.

References

- Banks-Randall B, Ballard R (2005):** Bronchopulmonary dysplasia. Avery's Diseases of the Newborn. 8th ed.2005.
- Baltimore R.S. (2002):** perinatal bacterial and fungal infection In:Hal B.Jenson and Robert S.Baltimore ,editors.Pediatric infectious diseases principals and practice.Philadelphia 96:1119-33.
- Benitz WE, Gould JB, Druzin ML.1999.**Antimicrobial prevention of early-onset group B streptococcal sepsis Estimates of risk reduction based on r Pediatrics; 103:e78.a critical literature.
- Benitz WE, Han MY, Madan A, et al.,(1998):** Serial serum C-reactive proteinlevels in the diagnosis of neonatal infection. Pediatrics 102:E41
- Benuck L. and David RJ. (2003):**Sensitivity of published neutrophil indices in identifying newborn infants with sepsis J.Pediatr.,138-144.
- Blackmon L:** Biologic limits of viability: implications for clinical decision making. NeoReviews 2003
- Blaymore-Bier J, Pezzullo J, Kim E, et al.(1994):** Outcome of extremely low-birth-weight infants: 1980-1990. Acta Paediatr 1994 Dec
- Borgnolo G, Barbone F, Guidobaldi G,et al.,(1996):**C-reactive protein in viral and bacterial gastroenteritis in childhood. Acta Paediatr. 1996;85:670–674
- Brodie SB, Sands KE, Gray JE, et al.(2000)**Occurence of nosocomial blood-stream infection insix neonatal intensive care units.Pediatr Infect Dis J;19:56–65.
- Brooks S., Khan A., Stoica D., et al. (1998):**Reduction in vancomycin-resistant enteroccus and Clostridium difficile infections following change to tympanic thermometers. Infect Control Hosp Epidemiol; 19:333-
- Brown GE, Silver GM, Reiff J, et al., (1999):** Polymorphonuclearneutrophil chemiluminescence in whole blood from blunt trauma patients withmultiple injuries. J Trauma 46:297–305).
- Buras-Buisson C., Legrand P., Philippon A., Montravers F., et al. (1987):**Transferable enzymatic resistance to third-generation

References

cephalosporins during nosocomial outbreak of multiresistant klebsiella pneumoniae. Lancet; ii:302-6.

Butler J, Pathi VL, Paton RD, et al.(1996): Acute-phase responses to cardiopulmonary bypass in children weighing less than 10 kilograms. Ann Thorac Surg. 1996;62:538-542

Cairo M.S.(1989): Neonatal neutrophil host defense. Prospects for immunologic enhancement during neonatal sepsis .Am J Child;143(1):406.

Caplan M and Jilling The(2001): The pathophysiology of necrotizing enterocolitis. NeoReviews; 2001

Cloherly J.P. Stark .R.and Eichenwald.R. (2004): Bacterial and fungal infections. In :Manual of Neonatal Care :Lippincott-Raven.23:271-300.

Carr R.(2000): Neutrophil production and function in newborn infants.Br J Haematol ;110:18-28

Carrigan S.D., Scott G., and Tabrizian M.(2004): Toward Resolving the Challenges of Sepsis Diagnosis. Clinical Chemistry. 2004;50:1301-1314.

Ceccon ME, Vaz FA, Diniz EM, et al.,(2006): Interleukins 6 and C-reactive protein for the diagnosis of late onset sepsis in the newborn infant. Rev Assoc Med Bras. 2006 Mar-Apr;52(2):79-85. Epub 2006 Jun.

Chapman, R. L., and faix R.G. (2003): Candida infections in the neonate. Curr Opin pediatr ; 15:97-102. Chemistry; 50:1301- 14.

Chwals WJ, Letton RW, Jamie A, et al.,(1995): Stratification of injury severity using energy expenditure response in surgical infants. J PediatrSurg. 1995;30:1161-1164

Chiesa C., Panero A., Osborn J., et al. (2004): Diagnosis of neonatal sepsis : And clinical and laboratory challenge. Clinical Chemistry; 50: 279- 87. Child Fetal;87:F25-8

Collee F (1996): Acute phase reactant. Clinical Microbiology; 2:93.

Cui YB, Du LZ, Chen YZ, Yu YB, Wang FM, Mao QQ.(2003): Expression of neutrophil adhesion molecule CD11b as an early diagnostic marker for neonatal sepsis. Zhonghua Er Ke Za Zhi 2003; 41(5):348-51.

References

- Da Silva O., Ohlsson A., and Kenyon C. (1995):** Accuracy of leukocyte indices and Cell-reactive protein for diagnosis of neonatal sepsis: a critical review. *Pediatr Infect Dis J*; 14:362-6.
- Dollberg S, Hoath S: (2001)** Temperature regulation in preterm infants: role of the skin-environment interface. *NeoReviews* 2001
- Dubowitz LM, Dubowitz V, Goldberg C. (1970):** Clinical assessment of gestational age in the newborn infant. *J Pediatr.*; 77(1):1-10
- ✓ Dyck RF, Bingham W, Tan L and Rogers SL(1984):L.** Serum levels of C-reactiveprotein in neonatal respiratory distress syndrome. *Clin Pediatr (Phila).*1984;23:381-3-83
- ✓ Edwards M.S. and Baker C.J.(2004):**Sepsis in the newborn. In *Diseases of children*, 11th ed : 545-61. Edited by Gershon A.A., Hotez P.J., and Katz S.L. By Mosby.
- Eisenthal A.,Hassner A.,Shenav M ., et al.(2003):** Phenotype and function of lymphocytes from the neonatal umbilical cord compared paired maternal peripheral isolated during delivery .
- El Badawy A., El-sebaei D., Khairat S., and shible S.F.(2003):**A study of microbiological pattern of neonatal sepsis. M.Sc. Thesis.
- Engel W.D.,Rosenfeld C.R. and Mouzinho A.(2001):** Circulating neutrophils in septic preterm neonates :Comparison of two reference ranges ,*Pediatrics*,99:110.
- Erman, Wayan Retayasa, and Soetjiningsih (2006):**Clinical gestational age assessment in newborns using the new Ballard score. *Paediatrica Indonesiana*, 46, (5-6), 97-101.
- Eskander R.M., Badrawi N.H., El-Atter S. et al. (2003):**Epidemiological study of neonatal sepsis in a neonatal intensive care unit. M.D. Thesis.*Exp Mol Pathol*;75(1):45-52.
- Espinosa EL, Gonzalez LFP, Montes AT. (2002):** Expression of CD64 as a potential marker of neonatal sepsis. *Pediatr Allergy Immunol* 2002; 13:319-27.
- Falconer A.F.Carline R.A.,and Edward S.W. (1995):**Impaired neutrophil phagocytosis in preterm neonates :lack of correlation with expression of immunoglobulin or complement receptors .*Bio Neonate*;68(4):264-9.

References

- Feldman m. and Marini j.O.(2002):** Cell corporation in the antibody response .in immunology Roitt,Brostoff,Male,6th ed:131-47 .Edited by Roitt,Brostoff J ,Male D. By Mosby
- Fjaertoft G, Hakansson L, Ewald U, et al(1999).** Neutrophils from term and preterm newborn infants express the high affinity Fc[gamma]-receptor I (CD64) during bacterial infection *Pediatr Res* ;45:871–6.
- Freij B.J. and McCracken G.H.(1999):**Acute Infections. In the textbook of Neonatology: Pathophysiology and Management of the Newborn, 5th ed: 1189-230. Edited by Avery G.B., Fletcher M.A., and MacDonald M.G. By Lippincott Williams & Wilkins.
- Gabriel J. Escobar, Marie C. McCormick, John A. Zupancic, et al (2006)** Outcomes of moderately premature infants in the NICU," by Archives of Disease in Childhood: *Fetal and Neonatal Edition* 91(4), pp. F238-F244.
- Garcia-Prats JA, Cooper TR, Schneider VF,et al (2000):**Rapid detection of microorganisms in blood cultures of newborn infants utilizing an automated blood culture system.
- Georgeson G.D.,Szony B.J.,Streitman K ,et al.(2001):** Natural killer cell cytotoxicity is deficient in newborns with sepsis and recurrent infections.*Eur J Pediatr*;160(8):478.82.
- Gerdes G. (2004):**Diagnosis and management of bacterial infections in the neonates. *Ped.Clin.N.Am*; 51: 939-59.
- Goldenberg RL and Rouse DJ. (1998):** Prevention of premature birth. *N Engl J Med.*; 339(5):313-20.
- Gottoff S.P. (2000):**Infections of the newborn infant. In Nelson Textbook of Pediatrics, 16th edition : 538-52. Edited by Behrman R.E., Kliegman R.M., and Jenson H.B. By W.B. Saunders.
- Griffin, M.P., and Moorman J. R. (2001):**Toward the early diagnosis of neonatal sepsis and sepsis-like illness using novel heart rate analysis. *Pediatrics*; 107:97-104.
- Griffin, M.P., O'shea T. M., Bissonette E. A., et al. (2003):**Abnormal heart rate characteristics preceding neonatal sepsis and sepsis-like illness. *Pediatr. Res*; 53:920-26.
- Guida J.D., Kunig A.M., Leef K.H. et al.,(2003):**

References

Platelet Count and Sepsis in Very Low Birth Weight Neonates: Is There an Organism-Specific Response? *PEDIATRICS* Vol. 111 No. 6 June 2003, pp. 1411-1415

Hammerman C, Kaplan M : (2005) American Academy of Pediatrics guidelines for detecting neonatal hyperbilirubinaemia. *Arch. Dis. Child. Fetal Neonatal* Ed.2005; 90: F448-F449.

Harris J.S. and Goldmann D.A. (2001):Infections acquired in the nursery: Epidemiology and control. In *Infectious Diseases of the fetus and Newborn Infant* 5th ed: 1371-418. Edited by Remington J.S. and Klein J.O. By W.B. Saunders.

Hillier SL, Nugent RP, Eschenbach DA, et al. (1995): Association between bacterial vaginosis and preterm delivery of a low-birth-weight infant. The Vaginal Infections and Prematurity Study Group. *N Engl J Med.*; 333(26):1737-42.

Hoath S, Pickens W.(2003) : The biology of the vernix; in Hoath SB, Maibach HI (eds): *Neonatal Skin: Structure and Function*. New York, Marcel Dekker, 2003.

Horns KM.(2000): Neoteric physiologic and immunologic methods for assessing early-onset neonatal sepsis. *J Perinat Neonatal Nurs* 2000; 13(4):50-66.

Iams JD, Johnson FF, Sonek J, et al., (1995):Cervical competence as a continuum: a study of ultrasonographic cervical length and obstetric performance. *Am J Obstet Gynecol.*; 172(4 Pt 1):1097-103; discussion 1104-6.

Jiang J.H., Chiu N.C., Huang F.Y., et al. (2004):Neonatal sepsis in the neonatal intensive care unit: characteristics of early versus late onset. *J Microbiol Immunol Infect*; 37(5): 301-6.

Karen M.(2004): bacterial and fungal infections . In :*Manual of Neonatal Care* .(2004), Cloherty J.P. Stark .R.and Eichenwald.R. eds:Lippincott-Raven.23:287-313

Karlowicz MG, Buescher ES, Surka AE:(2000):Fulminant late-onset sepsis in a neonatal intensive care unit, 1988- 1997, and the impact of avoiding empiric vancomycin therapy. *Pediatrics* 2000 Dec

References

- Kaufman . And Fairchild K.D. (2004):** Clinical microbiology of bacterial and fungal sepsis in very-low-birth-weight infants .Clin microbiology Rev;17(3):638-80.
- Kennon C.,Overturf G., Bessman S., et al., (1996):**Granulocyte colony-stimulating factor as a marker for bacterial infection in neonate .J Pediatr;128:765-9.
- Khalil A, Kaufmann RC, Wortsman J, Winters SJ, and Huffman DG. (1995):** Inhibin in normal and abnormal pregnancy: maternal serum concentration and partial characterization. Am J Obstet Gynecol.; 172(3):1019-25.
- Kidokoro A., Iba T., Fukunaga M., and Yagi Y. (1996):**Alterations in coagulation and fibrinolysis during sepsis. Shok; 5:223-8.
- Klein J. O. and Remington J.S. (2001):**Current concepts of infections of the fetus and newborn infant. In Infectious Diseases of the fetus and Newborn Infant 5th ed: 1-24. Edited by Remington J.S. and Klein J.O. By W.B. Saunders.
- Klein J.O. (2001):**Bacterial sepsis and meningitis. In Infectious Diseases of the fetus and Newborn Infant 5th ed: 943-98. Edited by Remington J. S. and Klein J. O. By W. B. Saunders.
- Kuster, H., Weiss M., Willeitner A. E., Detlefsen S., et al. (2003):**Interleukin-1 receptor antagonist and interleukin-6 for early diagnosis of neonatal sepsis 2 days before clinical manifestation. Lancet; 352:1271-7.
- La Pine T.R., Joyner J.L.,Augustine N.H.,et al (2003):** Defective production of IL-18 and IL-12 by blood mononuclear cells influences the T helper-1 interferon gamma response to group B Streptococci.Pediatr Res;54(2)276-81.
- Landers C.D.,Chelvarajan R.L.,Bondada S.(2005):** The role of Bcells and accessory cells in the neonatal response to TI-2 antigens. Immunol Res;31(1):25-36.
- Lehr HA, Krombach F, Munzing S, et al(1995):** In vitro effect of oxidized low density lipoprotein on CD11b/CD18 and L-selectin presentation on neutrophils and

References

- Levy O.(2002):** Impaired innate immunity at birth: bactericidal/permeability –increasing protein in the neutrophils of newborns. *Pediatr Res*;51(6)667-9.
- Levy O.,Martin S., Eichenwald E.,et al.(1999):** Impaired innate immunity in newborns :Newborns neutrophils are deficient in bactericidal /permeability –increasing protein .*PEDIATRICS*;104(6):1327-33.
- Levy O.,Sisson R., Kenyon J.,et al (2000):** Enhancement of neonatal innate defenses: Effect of adding an N-terminal recombinant fragment of bactericidal/permeability - increasing protein (Rbpi21)on growth and TNF-inducing activity of gram negative bacteria tested neonatal cord blood ex vivo.*Infect Immun*;68:5120-5.
- Lewis D.B. and Wilson C.B.(2001):** Developmental immunology and role of host defense in fetal and neonatal susceptibility to infection. In *infectious diseases of the fetus and Newborn infant 5th ed*:25-138.Edited by Remington J.S. and Klein J.O. By Saunders.
- Macones G.A. (2005):** Prematurity: causes and prevention. In: Taeusch H. A., Ballard R. A., and Gleason C. A. (ed), *Avery's diseases of the newborn. 8th ed*, Elsevier Saunders, p139-45.
- Madan A, Hamrick SEG, Ferriero D(2005):** Central nervous system injury and neuroprotection. In: *Avery's Diseases of the Newborn. 8th ed. 2005*
- May J.T. (1994):** Antimicrobial factors and microbial contaminants in human milk:recent studies .J. *Pediatr Child Health* :30(6):470-1995.
- McKenney WM(2001):** Understanding the neonatal immune system: high risk for infection. *Crit Care Nurse* 2001; 21(6):35-47.
- Meis PJ, Michielutte R, Peters TJ, Wells HB, Sands RE, Coles EC, Johns KA. (1995):** Factors associated with preterm birth in Cardiff, Wales. II. Indicated and spontaneous preterm birth. *Am J Obstet Gynecol.*; 173(2):597-602.
- Mentzer W, Glader B: Erythrocyte disorders in infancy(2005):.** In: *Avery's Diseases of the Newborn. 8th ed. 2005:*

References

- Neubert R.T., Delgado I., Webb J.R., et al (2000):** Assessing lymphocyte functions in neonates for revealing abnormal paternal development of the immune system. *Teratog Carcinog Mutagen*; 20:171-93.
- Ng PC. (2006):** Diagnostic markers for neonatal sepsis. *Curr Opin. Pediatr* 2006; 18(2):125-31.
- Ng P.C. (2004):** Diagnostic Markers of infection in neonates. *Arch Dis Child Fetal Neonatal Ed*; 89:F229.
- Ng PC, Li K, Wong RPO, et al, (2002):** Neutrophil CD64 expression: a sensitive diagnostic marker for late-onset nosocomial infection in very low birthweight infants. *Pediatr Res* ;51:296-303.
- Nicaise C., Gire C., Fagianelli P., et al (2002):** Neonatal consequences of preterm premature rupture of membrane (PPROM) at 24-34 WG: 118 singleton pregnancies. *J Gynecol Obstet Biol Reprod (Paris)*. 2002 Dec;31(8):747-54.
- Nupponen I, Andersson S, Jarvenpaa AL et al, (2001):** Neutrophil CD11b expression and circulating interleukin-8 as diagnostic markers for early-onset neonatal sepsis. *Pediatrics*; 108:E12.
- Ohlsson A. and Lacy J.B., (2004):** Intravenous immunoglobulin for suspected or subsequently proven infection in neonates. *Cochrane Database Syst Rev*; (1)CD001239
- Pablo J. and Jane D., (1999):** Bacterial And viral infections. In *Oski's sepsis pediatrics' principles and practice 3rd ed*:404 By Lippincott Williams and Wilkins.
- Philip A.G. (2000):** Acute phase proteins in neonatal sepsis. *Pediatrics*, 65:1036-41.
- Polin R, Parravicini E, Regan J (2005):** Bacterial sepsis and meningitis. In: *Avery's Diseases of the Newborn*. 8th eds. 2005:
- Pourcyrous M, Bada HS, Korones SB, et al, (1991):** Acute phase reactants in neonatal bacterial infection. *J Perinatol*. 1991; 11:319-325
- Powell K.R. AND & Marcy S.M. (2000):** Laboratory aids for diagnosis of neonatal sepsis. In: Klein J.O. and Remington J.J.S. (eds): *Infectious diseases of fetus and newborn infants*, W.B. Saunders, Philadelphia, P. 1223-1240.

References

- Pruchniewski D, Pawlowski T, Morkowski J, et al., (1987):** C-reactive protein in management of children's burns. *Ann Clin Res.* 1987;19:334-338
- Pursely and Cloherhy J.P. (1998):** Identifying the high risk newborn and evaluating the gestational age, prematurity, postmaturity. *In: Cloherty J.P., and Stark A.R. (ed), Manual of Neonatal care 4th ed Lippin Cott. Rawan Company, Chapter III Section 2, 37-49.*
- Remington J.S. , & Klein J.O. (2001):** Infectious diseases of the fetus and newborn infant ,5th ed.WB. Saunders ,Philadelphia.
- Rennie J.M., and Robertson N.R. , (2002):** Infection .InA Manual of Neonatal Intensive Care 4th ed:223-67.By Arnold and member of the Holder Headline Group.
- Remington J.S. , & Klein J.O. (2001):** Infectious diseases of the fetus and newborn infant ,5th ed.WB. Saunders ,Philadelphia.
- Rennie J.M., and Robertson N.R. , (2002):** Infection .InA Manual of Neonatal Intensive Care 4th ed:223-67.By Arnold and member of the Holder Headline Group.
- Rodwell RL, Leslie AL, and Tudehope DI (1988):** Early diagnosis of neonatal sepsis using a hematologic scoring system. *J Pediatr;* 112(5):761-7.
- Roilides E., Holmes A., Blake C., et al.(1995):** Effect of granulocyte colony-stimulating factor and interferon gamma on antifungal activity of human polymorphnuclear neutrophils against pseudohyphae of different medically important candida species.*J Clin Microbiol;*57:651-6.
- Roitt I.M., AND Delves P.J.(2001):** Lymphocyte activation .in Roitte's Essential Immunology,10th ed:164-76.By Blackwell Science Ltd.
- Rønnestad A., Abrahamsen G., Medbø S. ,et al.,(2005):** Late-Onset Septicemia in a Norwegian National Cohort of Extremely Premature Infants Receiving Very Early Full Human Milk Feeding *PEDIATRICS* Vol. 115 No. 3 March 2005, pp. e269-e276
- Rowen J.L., Rench M.A., Kozinetz C.A., et al. (1999):** Endotracheal colonization with candida enhances risk of systemic candidiasis in very low birth weight neonates.*J Pediatr;*124:789-94.

References

Rudnick C.M. and Hoekzema G.S. (2002):

Neonatal Herpes Simplex Virus Infections, Am Fam Physician. 2002 Mar 15;65(6):1138-42.

Saez-Llorens X and McCracken G.H. (1998): Sepsis syndrome and septic shock in pediatrics:current terminology,pathophysiology,and management.J Pediatr123:497-508.

Schelonka R.L.and Infant A.J. (1998): Neonatal immunology.semin perinatol.;22(1):2-14.

Seri I, Ramanathan R, Evans J(2005):Acid-base, fluid and electrolyte management. In: Avery's Diseases of the Newborn. 8th ed. (2005)

Shang S.,Chen Z.,and Yu X. (2001): Detection of bacterial DNA by PCR and reverse hybridization in the 16s r RNA gene with particular reference to neonatal septicaemia .Acta Pediatr ;90:179-83

Schelonka R.L.and Infant A.J. (1998): Neonatal immunology.semin perinatol.;22(1):2-14.

Seri I, Ramanathan R, Evans J(2005):Acid-base, fluid and electrolyte management. In: Avery's Diseases of the Newborn. 8th ed. (2005)

Shang S.,Chen Z.,and Yu X. (2001): Detection of bacterial DNA by PCR and reverse hybridization in the 16s r RNA gene with particular reference to neonatal septicaemia .Acta Pediatr ;90:179-83

✓ **Schober PH and Nassiri J.(1994):** Risk factors and severity indices in necrotizingenterocolitis. Acta Paediatr Suppl. 1994;396:49-52

Short M.A.(2004):Linking the sepsis triad of inflammation ,coagulation,and suppressed fibrinolysis to infants ,Adv Neonatal care;;4(5)258-73.

Schouten-Van Meeteren NY, et al.,(1992):Influence of perinatal conditions on C-reactive protein production. J Pediatr.1992;120:621-624

✓ **Singh B, Merchant P, Walker CR, et al.,(1996):**Interleukin-6 expression in cord blood of patients with clinical chorioamnionitis.Pediatr Res. 1996;39:976-979

Simms HH and D'Amico R(1995): Lipopolysaccharide induces intracytoplasmic migration of the polymorphonuclear leukocyte CD11b/CD18 receptor Shock;3:196-203.

References

- Smyth SS, Joneckis CC and Parise LV(1993).** Regulation of vascular integrins. *Blood* ;81:2827-2843.
- Stagno, S. (2004):**Cytomegalovirus .in Nelson textbook of pediatrics ,17th edition:1066-9edited by Behrman R.E. Kliegman R.M.,And Jenson H.B. by W.B. Saunders.
- Steel, R.G.D. & Torrie, J.H. (1960):** Principles and procedures of statistics, with special reference to the biological sciences. McGraw-Hill Book Company, Inc., New York – Toronto – London.
- Stoll B.J.(2004):** Infectious of the neonatal infant .in Nelson Text Book of Pediatrics.17th ed:623-40.Edited by Behrman R.E.,Kliegman R.M.,andJenson H.B. By Saunders
- Suri M.,Van de Ven C ., and Cairo M.S.(2003):** Immunotherapy in the prophylaxis and treatment of neonatal sepsis .*Curr Opin Pediatr*;15:155-60.
- Takala A, Jousela I, Jansson SE, et al(1999):** Markers of systemic inflammation predicting organ failure in community- acquired septic shock. *Clin Sci* ;97:529-538.
- Thornton Catherine A.; Capristo et al., (2003):**The Effect of Labor on Neonatal T-Cell Phenotype and Function **Pediatric Research. 54(1):120-124, July 2003.**
- Tirodker U.H., Nataro J.P., Smith S., et al.(2003):** Detection of fungemia by polymerasechain reaction in critically ill neonates and children . *J Perinatol* ; 23(2):117-22.
- Thurlbeck,S.M. and McIntoch n.,(2002):** Preterm blood counts vary with sampling site.*Arch.Dis.Child*,62:74.
- Tollner,(1982):** Early diagnosis of septicaemia in the newborn,clinical studies and sepsis score.*European J.Pediatrics*.,183:331
- Trappe U.a Riess H.(2005):** Basics in the pathophysiology of sepsis.*Hamostaseologie*;25(2):175-82.
- Tsiotou A.G., Sakorafas G.H., Anagnostopoulos G.,and Bramis J. (2005):** Septic shock; current pathogenetic concepts from and clinical perspective . *Med Sci Monit* ;11(3):RA 76-85.

References

- Wintrobe N(1990):** Clinical hematology. In: Wintrobe N ,clinical hematology ,Lea &Febgin,pp:154
- WHO 2003:**Recommendations for first line empiric therapy of serious neonatal Infections
- Yaron Y, Bryant-Greenwood PK, Dave N.et al., (1999):** Multifetal pregnancy reductions of triplets to twins: comparison with nonreduced triplets and twins. Am J Obstet Gynecol.; 180(5):1268-71.
- Zaidi A.K.,Huskins W.C., Thaver D., Bhutta Z. et al.,(2005):** Hospital-aquired neonatal infections in developing countries.Lancet;365:1175-88.