

References

A

Abdel-Fatah Mohd Abu-Haweleh (1998) : Acute Renal Failure in Newborn: Etiology and Mortality Rate in Jordan Patients Year : 1998 , Volume : 9 , Issue : 1 , Page : 18-21

Abitbol CL, Buer CR, Montae .B (2003): long term follows up of low birth weight with acute renal failure. *Pediatr nephrol.* 18:887-893

Adams-Chapman I, Stoll BJ (2006): Neonatal infection and long-term neurodevelopmental outcome in the preterm infant. *Curr Opin Infect Dis.* Jun 2006; 19(3):290-7.

Aggarwal A, Kumar P, Chowdhary G, Majumdar S, Narang A. (2005): Evaluation of renal functions in asphyxiated newborns. *J .Trop. Pediatr.* Oct; 51(5):295-9. Epub 2005 Jul 6.

Agras PI, Tarcan A, Baskin E, Cengiz N, Gürakan B, Saatci U.(2004) :Acute renal failure in the neonatal period. *Ren Fail.* May; 26 (3):305-9.

Airede A, Bello M, Weerasinghe HD(1997) :Acute renal failure in the newborn: incidence and outcome. *J Paediatr Child Health* . Dec; 33(6):550.

Al-DahanJ, Haycock GB, Chantler C et al., (1983): sodium homeostasis in term and preterm neonates; *Arab dis child*; 1983; 58; 335-342.

Al-Idressy HM, Abdel Basit OB, Haque KN, Abdurrahman MB (1991): Acute renal failure in neonates: A prospective study *Ann Saudi Med.* May;11(3):297-301.

American Academy of Pediatrics. Red Book (2003):26th ed. 117-123, 237-43, 561-73,584-91.

Amoo-Lamptey A, Dickman P, Carcillo JA. (2001): Comparative pathology of children with sepsis and MOF, pneumonia withoutMOF, andMOFwithout infection [abstract]. *Pediatr Res.*49:46A.

Anand SK (1995): acute renal failure in neonates *Ped.Clin.North, Am.J*; 9; 361-368.

Anderson, M.R. & Blumer, J.L et al., (1997): Advances in the therapy of children with sepsis. *Ped.Clin.North, Am.J.Perinatol.* 44(1): 179-205.

Andreoli SP (2004): Acute renal failure in the newborn. *Semin Perinatol.* Apr; 28(2):112-23

Andreoli SP,Bergestin JM,Sherrad DC (1984) : Aluminum toxicity in acute renal failure : *N Eng J Med* 1984 ;310 :1079-1084.

Andreoli SP,Dunson JW (1987):calcium carbonate as effective phosphorus binder in chronic renal failure;*Amj Kidney Dis* .9206-210.

Angus DC, Wax RS (2001): Epidemiology of sepsis: an update. *Crit Care Med* .29(Suppl):S109-S116.

Ann L Anderson &Berry (2006): e-medicine, Neonatal sepsis, *Pediatrics: Cardiac Disease and Critical Care Medicine Neonatology.*

Annabelle N. Chua & Minne M, Sarwel (2005): acute renal failure management in neonates, *Neoreviews* .6; 369-e376

Anne Stormorken & Kleith R. Powell (2004): Sepsis & Shock: chpter 163, *nelson text book of pediatrics*, Saunders 17thed; 846.

Anonymous(1998) : Adoption of hospital policies for prevention of Perinatal Group B streptococcal disease. *MMWR Morb Mortal Wkly Rept*; 47(32):665– 70.

Apgar V (1953): A proposal for a new method of evaluation of the newborn infant. *Curr Res Anesth Analg* .32:260-267.

Arnold W, Holliday M. (1976): Renal failure in the newborn. Read before the American Society of Pediatric Nephrology, St. Louis, April

Arvind Shenoï and Kishore D. Phadke (2000): Uric Acid Nephropathy as an Unusual Cause of Acute Renal Failure in a Neonate. *Indian Pediatrics* : Mar;37:322 – 324

Assumma M, Signore F, Pacifico L, Rossi N, Osborn JF, Chiesa .C. (2000): Serum procalcitonin concentrations in term delivering mothers and their healthy offspring: a longitudinal study. *Clin Chem*; 46:1583-1587.

B

- Bailie, M.D. (1992):** Renal function and disease. Clin.Perinatol. 19(1).
- Baker CJ. (1997):** Group B streptococcal infections. Clin Perinatol; 24:59-70.
- Balasubramaniam J. (2000):** Nimesulide and neonatal renal failure (Comment). *Lancet* 355:575,
- Balasubramaniam J. (2000):** Selective cox -2 inhibitors and nephrotoxicity. *American Journal of Kidney Diseases*, Vol 36, No 3 (September), pp 675-676
- Balasubramaniam J. (2001):** COX 2 inhibitors and nephrotoxicity. 2nd International Congress of Nephrology in Internet
- Balasubramaniam J. (2003):** Neonatal Renal Failure, Kidney Care centre. Tirunelveli, Tamilnadu. India.3; 76-78.
- Balows, A., and Sonnenwirth, A.C. (1983):** Bacteriemia: laboratory and clinical aspects .Charles C Thomas, Springfield, III.
- Baltimore.R.S. (1998):** Neonatal nosocomial infections. Semin Perinatol; 22:25-32.
- Barr M Jr. (1994):** Teratogen update: angiotensin-converting enzyme inhibitors. Teratology Dec; 50(6):399-409
- Behrman RE, Shiono PH. (2002):** Neonatal risk factors. Fanaroff AA Martin RJ eds. Neonatal-perinatal medicine: 17-26 St. Mosby Louis, MO.
- Bell M, Jackson E, Mi Z, et al. (1998):** Low-dose theophylline increases urine output in diuretic dependent critically ill children. Intensive Care Med 1998; 24(10):1099–105.
- Benitz WE, Han MY, Madan A, Ramachandra P. (1998):** Serial serum C-reactive protein levels in the diagnosis of neonatal infection. Pediatrics; 102:e41.
- Bennet, C.O.; Leonard, A.A.; & Quse, P. (1996):** Laminar air flow protection in bone marrow transplantation.Clin, inf.Dis 22(6):900-74.

Berger C, Uehlinger J, Ghelfi D, Blau N, Fanconi S. (1995): Comparison of C-reactive protein and white blood cell count with differential in neonates at risk for septicemia. Eur J Pediatr;154:138-144.

Bernard GR, Vincent JL, Laterre PF, et al. Safety and efficacy trial of activated protein C in

Bernirschke K. (1960): Routes and types of infection in the fetus and newborn. Am J Dis Child; 99:714-721.

Beth A, Vogt & Eills D, Anver (2004): acute renal failure, chapter 527 nelson text book of pediatrics, saunders 17th ed ;1767-1768.

Bick RL. (1996): Disseminated intravascular coagulation: objective clinical and laboratory diagnosis, treatment, and assessment of therapeutic response. Semin Thromb Hemost; 22:69-88.

Bick RL. (1998): Disseminated intravascular coagulation: pathophysiologic mechanisms and manifestations. Semin Thromb Haemost; 24.

Blanc WA. (1961): Pathways of fetal and early **neonatal** infection: viral placentitis, bacterial and fungal chorioamnionitis. J Pediatr; 59:473-496.

Blesha CW, Kohaut EC, Warady BA (1995): Dialytic management of childhood renal failure Pediatr nephrol .9:361-368

Bone RC. (1992): Modulators of coagulation: a critical appraisal of their role in sepsis. Arch Intern Med; 152:1381-9.

Booy R, Habibi P, Nadel J, et al. (2001): Reduction in case fatality rate from meningococcal disease associated with health care delivery. Arch Dis Child 2001; 85(5):386 –90.

Bourquia A, Chakib F, Jennah A, Boughnama .A.(2002): Acute renal failure in moroccan children. Saudi .J. Kidney .Dis Transpl. Jan-Mar;13(1):66-70.

Brilli RJ, Jacobs BR, Lyons K, et al. (1999): Severe protein C deficiency in children who do not survive purpura fulminans. Crit Care Med; 27(12):A127.

Brion LP, Bernstein J, Spitzer A. (1997): Kidney and urinary tract. In Neonatal- perinatal medicine: Diseases of the Fetus and Infant, Fanaroff AA, Martin RJ eds. St. Louis, Mosby-Year Book Inc., 1997, pp 1586.

Buck C, Bundschu J, Gallati H, Bartmann P, Pohlandt F. (1994): Interleukin-6: a sensitive parameter for the early diagnosis of neonatal bacterial infection. *Pediatrics*; 93:54-58.

Byington CL, Enriquez FR, Hoff C. (2004): Serious bacterial infections in febrile infants 1 to 90 days old with & without viral infections. *Pediatrics*. Jun 2004; 113(6):1662-6.

C

Cairo, M.S.; & Worcester, C.C. (1992): Randomized trials of granulocyte transfusion versus immunoglobulin administration for neonatal neutrophilia & sepsis .J. *Pediatr*, 120: 281-285.

Carcillo J.A., Cunnion RE. (1997): Septic shock. *Crit Care Clin*; 13(3):553– 74.

Carcillo J.A., Davis A.L., Zaritsky. (1991): A. Role of early fluid resuscitation in pediatric septic shock. *JAMA*; 266(9):1242– 5.

Carcillo J.A., Fields A.I, and Task Force Members. (2000): Clinical practice parameters for hemodynamic support of pediatric and neonatal patients in septic shock. *Crit Care Med*; 30(6):1– 13.

Chantler C. (1979): Renal failure in childhood. In. *Renal disease*, Black DAK, Jones NF eds. ed4, Oxford, Blackwell, 1979; 825.

Chapman RL, Faix RG. (2003): Persistent bacteremia and outcome in late onset infection among infants in a neonatal intensive care unit. *Pediatr Infect Dis J*. Jan; 22(1):17-21.

Charles D, Edwards WR. (1981): Infectious complications of cervical cerclage. *Am J Obstet Gynecol* 1981; 141:1065-1071.

Chevalier RL, Campbell F, Brenbridge AG. (1984): Prognostic Factors in neonatal acute renal failure. *Pediatr*; 74: 265-272

Chiesa C, Pellegrini G, Panero. A, Osborn JF, Signore F, Assumma M, et al. (2003): C-Reactive protein, interleukin-6, and procalcitonin in the immediate postnatal period: influence of illness severity, risk status, antenatal and Perinatal complications, and infection. Clin .Chem. 49:60-68.

Chiesa C, Signore F, Assumma M, Buffone E, Tramontozzi P, Osborn JF, et al. (2001): Serial measurements of C-reactive protein and interleukin-6 in the immediate postnatal period: reference intervals and analysis of maternal and Perinatal confounders. Clin Chem; 47:1016-1022.

Christensen RD, Rothstein G. (1979): Pitfalls in the interpretation of leukocyte counts of newborn infants. Am. J. Clin .Pathol.72:608-611.

Churchwell KB, McManus ML, Kent P, et al. (1995): Intensive blood and plasma exchange for treatment of coagulopathy in meningococemia. J. Clin .Apheresis; 10(4):171– 7.

Clarkson AR, MacDonald MK, Fuster V et al. (1970): Glomerular coagulation in acute renal failure. Q J Med 1970; 39: 585.

Claudio Chiesa Alessandra Panero, John F. Osborn, Antonella F. Simonetti and Lucia Pacific (2004): Diagnosis of Neonatal Sepsis: A Clinical and Laboratory Challenge; Clinical Chemistry. 50:279-287.

Csaicsich D, Russo-Schlaff N, Messerschmidt A, Weninger M, Pollak A, Aufrecht C. (2008): Renal failure, co morbidity and mortality in preterm infants Wien Klin Wochenschr.; 120(5-6):153-7

D

D. S. Roberts, G. B. Haycock, R. N. Dalton, C. Turner, P. Tomlinson, L. Stimmler, and J. W. Scopes. (1990): Prediction of acute renal failure after birth asphyxia. Arch Dis .Child. 1990 October; 65(10 Spec No): 1021–1028.

David Moskowitz, (1997): Personal communication. Rajesh, V.T. Manual of neonatal care / Joint program in Neonatology, Harvard Medical School, – 4th editions.

Davis, D.J., Kennedy A. & Roberts C. (1970): Renal medullary necrosis in infancy. *J.Path* 99: 125.

Dawodu, A.S. Abue-Rteish, F. & Obeidate, A. et al., (1994): A case control study in neonatal sepsis, Saudi. Arabia. *J.Trop.Ped.* 43(2):84-8.

Dehne M.G., Sablotzki A., Mühling J., Papke G., Kuntzsch U., Hempelmann G.(2001): from energy failure in a cell culture model of sepsis. *Res Commun Mol Pathol Pharmacol*; 110(1&2):107– 16.

Delves P.J., Reitt.I.M. (2000): The immune system. *N .Engl. J Med* 2000; 343(1):37– 49.

Despond O, Proulx F, Carcillo J, et al. (2001) Pediatric sepsis and multiple organ dysfunction syndrome. *Curr Opin Crit Care* 2001; 13(3):247– 53.

Diamond JR, Yoburn DC. (1982): Nonoliguric acute renal failure. *Arch Intern Med.* Oct; 142(10):1882-4.

Donohoe, J.F.; Venkatachalam, A.M. et al., (1978): Tubular leakage & obstruction after renal ischaemia. *Kid. Int.*13:208.

Dorman HR, Sonhemier JH, Cadnapaphornchai P (1990): Mannitol induced acute renal failure; *Medicine* .1990; 60:153-159.

Doughty L, Carcillo JA, Kaplan S, et al (1998): Plasma nitrite and nitrate concentrations and multiple Organ failure in pediatric sepsis. *Crit Care Med* 1998; 26(1):157 –62.

Douglas A. Lind (1994): Basic Statistics, Two samples test about proportion McGraw-Hill, 326.

Drukker A,Guignard JP (2002): Renal aspects of term &preterm infants:*Curr Opin pediatr* ; 14:175-182.

DuPont HL, Spink WW. (1968): Infections due to gram negative organisms: an analysis of 860 patients with bacteremia at University of Minnesota Medical Center. 1958– 1966. *Medicine (Baltimore)*; 48(4):307– 11.

E

Edwards WR. (1981): Infectious complications of cervical cerclage. *Am J Obstet Gynecol*; 141:1065-1071.

Edwards, M.S. and Baker, C.J.I (1997): Bacterial infection in neonates in: *pediatric infectious diseases* chapter 98 page 606 – 614, Churchill Livingstone, New York.

El- Maraghi, S.: El-Safi, A. and Abou Alam, M. (1984): diagnosis of neonatal septicemia by gastric aspirate .*J. Egypt, Soc. Ob. Gyn.*, 10 (1):43.

Ellis EN, Arnold WC. (1982): Use of urinary indices in renal failure in the newborn. *Am. J. Dis. Child* 1982; 136: 615-617.

EL-Najjar, M. (2002): practical pediatric diagnosis, neonatal septicemia. University Book Center

Escobar GJ, Fischer A, Li DK, Kremers R, Armstrong MA. (1995): Score for neonatal acute physiology: validation in three Kaiser Permanente neonatal intensive care units. *Pediatrics*; 96:918-922.

Escobar GJ. (1999): The neonatal "sepsis work-up": personal reflections on the development of an evidence-based approach toward newborn infections in a managed care organization. *Pediatrics*; 103(Suppl 1):S360-S373.

F

Fanaroff, AA, Korons S.B., Wright, L.L., Veter .J. (1998): incidence, presenting features, risk factors and significance of late onset sepsis in very low birth weight. The national institute of Child Health & Human Development, Neonatal research network. *Pediatr. Infect.Dis. J.*, 17(7):593-8.

Faust SN, Levin M, et al. (2001): Dysfunction of endothelial protein C activation in severe meningococcal sepsis. *N .Engl. J .Med*; 345(6):408–16.

Fawcett, J. and Scott J. (1970): A rapid precise method for determination of BUN and creatinine.

Fejgin M, Amiel A, Kaneti H, Ben-Nun I, Beyth Y. (1993) :Fulminant sepsis due to group B β -hemolytic streptococci following transcervical chorionic villi sampling [Letter]. Clin Infect Dis 1993; 17:142-143.

Fetter, W.PF ; Bearts , W. & Vonlingen, R.A.(1995):Surfactant replacement therapy in neonates with respiratory distress due to bacterial sepsis. Acta. Ped. 84:14-16.

Forest JC, Larivière F, Dolcé P, Masson M, Nadeau L. (1986): C - reactive protein as biochemical indicator of bacterial infection in neonates. Clin. Biochem 1986; 19:192-194.

Fourrier F, Chopin C, Huart JJ, et al. (1993): Double blind placebo controlled trial of antithrombin III concentrate in septic shock with disseminated intravascular coagulation. Chest; 104(3): 882– 8.

Fowlie PW, Schmidt B. (1998): Diagnostic tests for bacterial infection from birth to 90 days—a systematic review. Arch Dis Child Fetal Neonatal Ed; 78:F92-F98.

Franz AR, Kron M, Pohlandt F, Steinbach G. (1999): Comparison of procalcitonin with interleukin 8, C-reactive protein and differential white blood cell count for the early diagnosis of bacterial infections in newborn infants. Pediatr. Infect. Dis. J; 18:666-671.

Franz AR, Steinbach G, Kron M, Pohlandt F. (1999): Reduction of unnecessary antibiotic therapy in newborn infants using interleukin-8 and C-reactive protein as markers of bacterial infections. Pediatrics; 104:447-453.

Freedman RM, Baltimore R. (1990): Fatal *Streptococcus viridans* septicemia and meningitis: a relationship to fetal scalp electrode monitoring. J. Perinatol 1990; 10:272-274.

Freedman RM, Baltimore R. Fatal (1990): *Streptococcus viridians* septicemia and meningitis: a relationship to fetal scalp electrode monitoring. J. Perinatol; 10:272-274.

Fridrich JO, Adhikari N, Herridge M, Beyene M (2005): low dose dopamine increase urine out put but not prevent renal dysfunction. Ann. intern .Med. 142:510-542.

G

Garges HP, Moody MA, Cotten CM. (2006): Neonatal meningitis: what is the correlation among cerebrospinal fluid cultures, blood cultures, and cerebrospinal fluid parameters, *Pediatrics*. Apr; 117(4):1094-100.

Gendrel D, Assicot M, Raymond J, Moulin F, Francoual C, Badoual J, et al. (1996): Procalcitonin as a marker for the early diagnosis of neonatal infection. *J Pediatr*; 128:570-573.

Gerdes, J.S. (1991): Clinico-pathological approach for diagnosis of neonatal sepsis. *Clin. Perinatology*, 18(2):361.

Gharehbaghi MM, Peirovifar A. (2007): evaluation of causes of ARF in newborn Pakistan *Journal of Medical Sciences*, 2007 Oct-Dec;23(6):877-880

Gibbs RS, Duff P. (1991): Progress in pathogenesis and management of clinical intraamniotic infection. *Am.J Obstet. Gynecol* 1991; 164:1317-1326.

Giroir BP, Scannon PJ, Levin M. (2001): Bactericidal permeability increasing protein; lessons learned from the phase III randomized clinical trial of rBPI21 for adjuvant treatment of children with severe meningococemia. *Crit Care Med* 29(7 Suppl):S130 –5.

Gladder BE. (1985): Bleeding disorders in the newborn infant. In Schaffer's Diseases of the newborn, Avery ME, Taeusch HW eds. 5th Edn, Philadelphia, WB Saunders Co, 1985, pp562.

Golej J, Kitzmueller E, et al. (2002): Low-volume peritoneal dialysis in 116 neonatal and paediatric critical care patients. *Eur. J. Pediatr*. Jul; 161(7):385-9. Epub 2002 May 09.

Gotoff (1992): sepsis in newborn, infectious diseases of children chapter, 24 pages 402 -5.

Gotoff SP. (2000): Infections of the **neonatal** infant. Behrman RE Kliegman RM Jenson HB eds. *Nelson textbook of pediatrics* 2000:538-552, WB Saunders Philadelphia.

Gotoff, S.P. (1996): infections of neonatal infants. In: Behram R. & Kliegman RM, Nelson Text Book of pediatrics (15th Ed). W.B. Saunders Co.

Gouyon JB, Guignard JB (2000): Management of acute renal failure in newborn. *Pediatr Nephrol.*; 14:1037-1044.

Gouyon JB, Guignard JB. (1986): drugs & acute renal insufficiency in neonates: *Biol neonate*; 50:177-181.

Graham PL, Begg MD, Larson E. (2006): Risk factors for late onset gram-negative sepsis in low birth weight infants hospitalized in the neonatal intensive care unit. *Pediatr Infect Dis J.* Feb 25(2):113-7.

Gray, D. (1992): Immunological memory. *Ann. Rev. Immunol.*, 11:49.

Green J, Dougherty L, Kaplan SS, et al. (2002): The tissue factor and plasminogen activator inhibitor type 1 response in pediatric sepsis induced multiple organ failure. *Thromb Haemost*; 87(2): 218– 23.

Grella MJ. (1997): C - reactive protein to determine the duration of antibiotic therapy in infants with suspected sepsis [Letter]. *Pediatrics*; 100:900.

Griffin NK, McElena J, Baratt JM. (1976): Acute renal failure in early life. *Arch. Dis. Child*, 1976; 51: 459-462.

Grylack L, Medani C, Hultzen C, Sivasubramanian K, Davitt MK, Jose P, Scanlon JW. (1982): Nonoliguric acute renal failure in the newborn. *Am. J. Dis. Child*, 1982; 136: 518-520.

Guerina , N.B. (1991) : Bacterial & Fungal infection of neonates . in : manual of neonatal care (4th ed) . (Eds) Cloherty, JP & Stark, AR., 271 Lippin Cott –Raven- Publ. Philadelphia, New York.

Gupta BD, Sharma P, Bagla J, Parakh M, Soni JP. (2005): Renal failure in asphyxiated neonates *Indian Pediatr. Sep*; 42(9):928-34.

Gutteberg TJ, Haneberg B, Jorgensen T. (1984): Lactoferrin in relation to acute phase proteins in sera from newborn infants with severe infections. *Eur.J. Pediatr* 1984; 142:37-39.

H

Hall MW, Carcillo JA. (2000): Immunoparalysis and the state of immunologic dissonance in pediatric multiple organ dysfunction syndrome. Crit Care Med 2000; 29(12):A20.

Hall RT, Wheeler RE, Rippetoe LE (1993) : Calcium & phosphorus supplementation after initial discharge in breast fed infants . J. perinatol. 13:272-278.

Han Y, Doughty L, Carcillo JA. (2003): Procalcitonin is increased bacterial sepsis-induced multiple organ failure. Pediatr Crit Care Med 2003; 41:21–5.

Han YY, Huang L, Jackson EK, et al. (2001): Liposomal ATP or NAD protects human endothelial cells from energy failure in a cell culture model of sepsis. Res. Commun Mol. Pathol. Pharmacol .2001; 110(1&2):107– 16.

Harris JAS, Goldmann DA. (2001): Infections acquired in the nursery: epidemiology and control. Remington JS Klein JO eds. Infectious diseases of the fetus and newborn infant: 1371-1418 WB .Saunders Philadelphia. .

Harris MC, Costantine A.T., & Sukivan A.S. (1994): Cytokines elevation in critically ill infants with sepsis, J.pediatr 124:105.

Hauth JC, Merenstein GB. Infection control. Hauth JC Merenstein GB (1997): eds. Guidelines for Perinatal care: 251-277 American Academy of Pediatrics and American College of Obstetricians and Gynecologists Elk Grove Village, IL. .

Hazeltet JA, de Kleijn ED, de Groot R. (2001): Endothelial protein C activity in meningococcal sepsis. N Engl J Med; 345(24):1776– 7.

Hentschel R, Loddgi B, Bulla M(1996) : Renal insufficiency in neonatal period. Clin Nephrol.1996; 49:54-58.

Hentschel R, Lodige B, Bulla M. (1996): Renal insufficiency in the neonatal period. Clin. Nephrol. 1996; 46: 54-58.

Hickman ME, Rench MA, Ferrieri P, Baker CJ. (1999): Changing epidemiology of group B streptococcal colonization. *Pediatrics*. Aug; 104(2 Pt 1):203-9.

Hillier SL, Krohn MA, Kiviat NB, Watts DH, Eschenback DA. (1991): Microbiologic causes and neonatal outcomes associated with chorioamnion infection. *Am .J. Obstet Gynecol*; 165:955-961.

I

Inbender E, Cabatu EE, Guzman DM, Sweet AY. (1982): Serum C-reactive protein and problems of newborn infants. *J Pediatr*; 101:438-440.

International Neonatal Network. The CRIB (1993): (clinical risk index for babies score)—a tool for assessing initial neonatal risk and comparing performance of neonatal intensive care units. *Lancet*; 342:193-198.

Ishibashi M, Takemura Y, Ishida H, Watanabe K, Kawai T. (2002): C-reactive protein kinetics in newborns: application of a high-sensitivity analytic method in its determination [Technical Brief]. *Clin. Chem*; 48:1103-1106.

J

Jacobs RF, Sowell MK, Moss MM, et al. Septic shock in children: bacterial etiologies and

Jayashree G, Dutta AK, Sarna MS, Saili A. (1991): Acute renal failure in asphyxiated newborns. *Ind J Pediatr*; 28: 19-23.

Jayashree G, Saili A, Sarna MS, Dutta AK. (1991): Renal dysfunction in septicemic newborns. *Indian Pediatr*. Jan; 28(1):25-9

Jepson HA, Talashek ML, Tichy AM. (1991): The Apgar score: evolution, limitations, and scoring guidelines. *Birth*; 18(Suppl 2):S83-S92.

John E. (1981): Renal physiology and acute renal failure. *Ind. J. Pediatr*, 48: 313-321.

John P. and Ann R.Starket, et al., (1997): Manual of neonatal care / Joint program in Neonatology, Harvard Medical school 4th edition,.

John V.Dacie (1991): bleeding tendency investigations, practical hematology Churchill Livingstone 293:295

Joseph A. Carcillo, AG. (1984): Prognostic Factors in neonatal acute renal failure. *Pediatr*, 1984; 74: 265-272. Arnold W, Holliday M. Renal failure in the newborn. Read before the American Society of Pediatric Nephrology, St. Louis, and April 1976.

Joseph A. Carcillo, (2003): Pediatric septic shock and multiple, *Crit Care Clin*. 19: 413– 440

K

Kellum JA. (1998): Use of diuretics in critical care setting. *Kid .int*. 49:4-14.

Kemp AS, Campbell DE. (1996): The neonatal immune system. *Seminar in Neonatology* ; 1:67-75.

Khashu M, Osiovich H, Henry D. (2006): Persistent bacteremia and severe thrombocytopenia caused by coagulase-negative *Staphylococcus* in a neonatal intensive care unit. *Pediatrics*. Feb ;117(2):340-8.

Klein JO, Marcy SM. (1995): Bacterial Sepsis and Meningitis. *Infectious Diseases of the Fetus & Newborn Infant. 4th ed.* ;835-878.

Klein JO. (2001): Bacterial sepsis and meningitis. Remington JS Klein JO eds. *Infectious diseases of the fetus and newborn infant* .943-998 WB Saunders Philadelphia. .

Klien J.O.& Marcy S.M (1990): Bacterial sepsis & meningitis in Remington J.S., Klien J.O , (eds), *infectious disease of newborn* 4th ed ,Philadelphia ,Suanders,835-90.

Knaus WA, Zimmerman JE, Wagner DP, Draper EA, Lawrence DE. (1981): Acute physiology and chronic health evaluation: a physiologically based classification system. *Crit Care Med*; 9:591-597.

Kocherlakota P, La Gamma EF. (1998): Preliminary report: rhG-CSF may reduce the incidence of neonatal sepsis in prolonged preeclampsia-associated neutropenia *Pediatrics*. Nov; 102(5):1107.

Kofos D, Frye R, Burckart G, et al. (1998): Diminished oxidative metabolizing capacity of specific cytochrome P450 isozymes in sepsis induced pediatric multiple organ failure. *Crit Care Med* 1998; 143:A79.

Kutko MC, Calarco MP, Ushay M, et al. (2000): Mortality of pediatric septic shock may be less than previously reported. *Crit Care Med*; 28(12):T212.

L

L. Catald, R Leone, U Morett , B De Mitri, V Fanos, L Ruggeri, G Sabatino, F Torcasio, V Zanardo.(2005): Potential risk factors for the development of acute renal failure in preterm newborn infants: a case-control study *Archives of Disease in Childhood - Fetal and Neonatal Edition*;90:F514-F519;

Lackland DT, Bendall HE, Osmond C, Egan BM, Barker DJ. (2000): Low birth weights contribute to high rates of early-onset chronic renal failure in the Southeastern United States. *Arch Intern Med*. May 22; 160(10):1472-6.

Lapillonne A, Basson E, Monneret G, Bienvenu J, Salle BL. (1998): Lack of specificity of procalcitonin for sepsis diagnosis in premature infants [Letter]. *Lancet* 351:1211-1212.

Lauterbach R, Pawlik D, Kowalczyk D, et al. (1999): The effect of the immunomodulatory agent, pentoxifylline in the treatment of sepsis in prematurely delivered infants; placebo controlled, double blinded trial. *Crit Care Med*;27(4):807– 14.

LeDoux D, Astiz ME, Carpati CM, et al. (2000): Effects of perfusion pressure on tissue perfusion in septic shock. *Crit Care Med*. 28(8):2729– 32.

Levi M, van der Poll T, ten Cate H, van Deventer SJH. (1997): The cytokine-mediated imbalance between coagulant and anticoagulant mechanisms in sepsis and endotoxemia. *Eur J Clin Invest.* 27:3-9.

Levy MM, Fink MP, Marshall JC, Abraham E, Angus D, and Cook D, et al. (2001): SCCM/ESICM/ACCP/ATS/SIS International Sepsis Definitions Conference. *Crit Care Med* 2003; 31:1250-1256.

Liaño F, Pascual J. (1996): Epidemiology of acute renal failure: a prospective, multicenter, community-based study. Madrid Acute Renal Failure Study Group. *Kidney Int.* 1996 Sep; 50(3):811-8

Liberman .E. (1937): Management of acute renal failure in infants & children. *Nephron* 11; 193.

M

Macintoch, K. (1984): Bacterial infection of the newborn, *Biol neonate*, 38:114.

Maire F, Héraud MC, Loriette Y, Normand B, Bègue RJ, Labbé A. (1999): Intérêt de la procalcitonine dans les infections néonatales. *Arch Pédiatr*; 6:503-509.

Malik A, Hui CPS, Pennie RA, Kirpalani H. (2003): Beyond the complete blood cell count and C-reactive protein. A systematic review of modern diagnostic tests for neonatal sepsis. *Arch. Pediatr Adolesc. Med*; 157:511-516.

Manroe BL, Weinberg AG, Rosenfeld CR, Browne R. (1979): The neonatal blood count in health and disease. I. Reference values for neutrophilic cells. *J Pediatr*; 95:89-98.

Marcel Levi M.D., & Hugo Ten Cate (1999): DISSEMINATED INTRAVASCULAR COAGULATION; *Current Concepts*; 19, 586-592

Mathers NJ, Pohlandt F. (1987): Diagnostic audit of C-reactive protein in neonatal infection. *Eur J Pediatr*; 146: 147-151.

Mathew O.P. et al. (1980): Neonatal renal failure: Usefulness of diagnostic indices. *Pediatrics* 65:57, (Modified)

Mathew OP, Jones AS, James E, et al., (1980): Neonatal acute renal failure usefulness of diagnostic indices. *Pediatr.* 65:57-60.

Mathur N.B., Himmanshu S, Agarwal and Arti Maria.(2006): Acute renal failure in neonatal sepsis. *Ind J, peditr*, 73, 199-500-501.

Matlin, R.A. & Gary, N.E. (1984): Acute cortical necrosis. *Am. J. Med.* 56: 110.

McKenna DS, Iams JD. (1998): Group B streptococcal infections. *Seminars in Perinatology.* ; 22:267-76.

Mehr S, Doyle LW. (2000): Cytokines as markers of bacterial sepsis in newborn infants: a review. *Pediatr Infect Dis J*; 19:879-887.

Members of the American College of Chest Physicians/Society of Critical Care Medicine Consensus Conference Committee. (1992): American College of Chest Physicians/Society of Critical Care Medicine Consensus Conference: definitions for sepsis and organ failure and guidelines for the use of innovative therapies in sepsis. *Crit Care Med*; 20:864-874.

Merril, J.P. (1971): Acute renal failure. *Int. Diseases of the kidney*, Little brown & Co. Boston.

Messer J, Eyer D, Donato L, Gallati H, Matis J, Simeoni U. (1996): Evaluation of interleukin-6 and soluble receptors of tumor necrosis factor for early diagnosis of neonatal infection. *J Pediatr*; 129:574-580.

Minelli L, Bernardi M, Locatelli G (1994): Acute tubular necrosis in a newborn suffering from neonatal necrotizing enterocolitis *Acta Paediatr Suppl.*; 396:84-5.

Mizuno S. (1997): Granulocyte colony-stimulating factor and neonatal infection [Letter]. *J Pediatr*; 130:844-845.

MMWR Morb Mortal Wkly. (2005): Early-onset and late-onset neonatal group B streptococcal disease--United States, 2005. *Rep.* Dec 2 2005; 54(47):1205-8.

MMWR. Morb Mortal Wkly (2002): CDC. Prevention of Perinatal group B streptococcal disease: revised guidelines from CDC. 2002; 51 :(No. RR-11).

Monneret G, Labaune JM, Isaac C, Bienvenu F, Putet G, Bienvenu J. (1997): Procalcitonin and C-reactive protein levels in neonatal infections. *Acta Paediatr*; 86:209-212.

Morales WJ, Dickey SS, Bornick P, Lim DV. (1999): Change in antibiotic resistance of group B streptococcus: impact on intrapartum management. *Am J Obstet Gynecol*. Aug ;181(2):310-4.

Mouhammad El- Najjar (2002): neonatal septicemia, pediatric clinical diagnosis 4th Ed, university book center.

Muller-Berghaus G, ten Cate H, Levi MM (1988): Disseminated intravascular coagulation. In: Verstraete M, Fuster V, Topol EJ, eds. *Cardiovascular*

N

Nguyen T, Hall M, Han Y, et al. (2001): Microvascular thrombosis in pediatric multiple organ failure: is it a therapeutic target? *Pediatr Crit Care Med* 2001; 2(3):187 – 96.

Norman Jones AS, James E, Bland H, Groshong T. (1979): Renal failure in the newborn. *Clint. Ped..* 1979; 18: 286-290.

Norman ME, Asadi FK. (1979): A prospective study of acute renal failure in the newborn infant. *Pediatrics*, 1979; 63: 475-479.

Nupponen I, Andersson S, Järvenpää A-L, Kautiainen H, Repo H. (2001): Neutrophil CD11b expression and circulating interleukin-8 as diagnostic markers for early-onset neonatal sepsis. *Pediatrics*;108:e12.org failure MDCrit Care Clin 19 (2003) 413– 440.

P

Panero A, Pacifico L, Rossi N, Mancuso G, Stegagno M, Chiesa C. (1997): Interleukin 6 in neonates with early and late onset infection. *Pediatr Infect Dis J*; 16:370-375.

Pejović B, Peco-Antić A, Dunjić R (2002) Acute oliguric renal failure in hypoxic neonates born at full term] p *Arh Celok Lek*. Nov-Dec;130(11-12):367-70.

Pepys, M.B... (1981): Lancet 1:653

Pereira S, Pereira BJG, Bhakoo ON, Narang A, Sakhuja VS, Chugh KS. (1988): Peritoneal dialysis in neonates with acute renal failure. *Ind. J. Pediatr*, 58: 973-978.

Petridou E, Richardson DK, Dessypris N, Malamitsi-Puchner A, Mantagos S, Nicolopoulos D, et al. (1998): Outcome prediction in Greek neonatal intensive care units using a score for neonatal acute physiology (SNAP). *Pediatrics*; **101:1037-1044**.

Petzold L, Guibourdenche J, Boissinot C, Benoist JF, Luton D, Demelier JF, et al, (1998): Apport de la procalcitonine dans le diagnostic des infections materno-foetales. *Ann Biol Clin (Paris)*;56:599-602.

Philip AGS, Hewitt JR. (1980): Early diagnosis of neonatal sepsis. *Pediatrics*; 65:1036-1041.

Philip AGS, Mills PC. (2000): Use of C-reactive protein in minimizing antibiotic exposure: experience with infants initially admitted to a well-baby nursery. *Pediatrics*; 106:e4.

Pierce, J.M; Merensten, G.B. & Stocker, J.T. (1984): Immediate post mortem culture in intensive care nursery. *Pediatric INF: Dis.*, 2:80-6.

Pietrement C, Malot L, et al. (2003): Neonatal Acute Renal Failure Secondary to Maternal Exposure to Telmisartan, Angiotensin II Receptor Antagonist. *J Perinatol*. May;23(3):254-5.

Pollack MM, Koch MA, Bartel DA, Rapoport I, Dhanireddy R, El-Mohandes AA, et al. (1985): A comparison of neonatal mortality risk prediction models in very low birth weight infants. *Pediatrics* 105:1051-1057.

Pollack MM, Ruttimann UE, (1988): Getson RR. Pediatric risk of mortality (PRISM) scores. *Crit Care Med*; 16:1110-1116.

Pollard AJ, Britto J, Nadel S, et al. (1999): Emergency management of meningococcal disease. *Arch Dis Child*; 80(3):290– 6.

Poole, J.C.F...(1952): Correction of ESR in anemia .*British medical journal*, i, 353.

Pourcyrous M, Bada HS, Korones SB, Barrett FF, Jennings W, Lockey T. (1991): Acute phase reactants in neonatal bacterial infection. *J Perinatol*; 11:319-325.

Pourcyrous M, Bada HS, Korones SB, Baselski V, Wong SP. (1993): Significance of serial C-reactive protein responses in neonatal infection and other disorders. *Pediatrics*; 92:431-435

Prober CG. (1997): Clinical approach to the infected neonate. Long SS Pickering LK Prober CG eds. *Principles and practice of pediatric infectious diseases*:603-606 Churchill Livingstone New York. .

Prober CG. Clinical approach to the infected neonate. Long SS Pickering LK Prober CG eds. (1997): Principles and practice of pediatric infectious diseases 1997:603-606 Churchill Livingstone New York.

R

Racine A , Abribat D, Ensergueix G, Lucas Y, Poux JB. (1989): Value of C-reactive protein assay for the early diagnosis of neonatal infection in a general hospital maternity ward and pediatric department. *Ann Pediatr (Paris)* 36:253-257.

Richardson D, Tarnow-Mordi WO, Lee SK. (1999): Risk adjustment for quality improvement. *Pediatrics*; 103(Suppl 1):S255-S265.

Richardson DK, Corcoran JD, Escobar GJ, Lee SK. (2001): SNAP-II and SNAPPE-II: simplified newborn illness severity and mortality risk scores. *J Pediatr*; 138:92-100.

Richardson DK, Gray JE, McCormick MC, Workman K, Goldmann DA. (1993): Score for neonatal acute physiology: a physiologic severity index for neonatal intensive care. *Pediatrics*; 91:617-623.

Rintala E, Kauppila M, Seppala O, et al. (2000): Protein C substitution in sepsis-associated Purpura fulminans. *Crit Care Med*; 28:2373–8.

Rivers E, Nguyen B, Havstad S, et al. (2001): Early goal directed therapy in the treatment of severe sepsis and septic Shock. *N. Engl .J. Med*; 346(19):1368–77.

Robert L. Chevalier MD, Fern Campbell RNFNP, and A. Norman A. G. Brenbridge MD (1984), Prognostic Factors in Neonatal Acute Renal Failure PEDIATRICS Vol. 74 No., pp. 265-272

Rodriguez- Cervilla, J.;Farag J.M.; Gracia-Restira, C.;Fernandez J.R. , Martinez-Soto, I.(1998):Neonatal sepsis epidemiologic indicators & relation to birth weight and length of hospitalization time. An. Esp.Pediatr., 48(4):401-8.

Roger FR, Li SC, (2001): Acute colonic necrosis associated with sodium polystyrene in critically ill patient. J. Truman. 51:395-397.

Rose BD (1987). Pathophysiology of renal diseases, 2d ed ,mac Graw Hill, 1987 ,pp.63-70.

Rudd, P.T, Hughes, E.A. et al. Reference ranges for plasma creatinine during the first month of life. Arch. Dis. Child. 58:212, 1983

S

Sabel K-G, Wadsworth C. (1979): C-reactive protein (CRP) in early diagnosis of neonatal septicemia. Acta .Paediatr. Scand; **68:825-831**.

Sachse C, Dressler F, Henkel E. (1998): Increased serum procalcitonin in newborn infants without infection [Technical Brief]. Clin Chem; 44:1343-1344.

Saez X, McCracken GH. (1995): Clinical Pharmacology of Antibacterial Agents. Infectious Diseases of the Fetus & Newborn Infant. 4th ed. ;1294-1327.

Sáez-Llorens X, Lagrutta F. (1993): The acute-phase host reaction during bacterial infections in children. Pediatr Infect Dis J 1993;12:83-87.

Sáez-Llorens X, McCracken GH, Jr. (1993): Sepsis syndrome and septic shock in pediatrics: current concepts of terminology, pathophysiology, and management. J Pediatr 1993; 123:497-508.

Sáez-Llorens X, McCracken GH, Jr. (1995): Perinatal bacterial diseases. Feigin RD Cherry JD eds. Textbook of pediatric infectious diseases 1998:892-926 WB Saunders Philadelphia.

Sagripanti A, Carpi A, Rosaia B, et al. Iloprost in the treatment of thrombotic microangiopathy: report of thirteen cases. *Biomed Pharmacother* 1996; 50(8):350– 6. 438 J.A. Carcillo / *Crit Care Clin* 19 (2003) 413–440.

Sanghvi, K.P. (1996): Neonatal bacterial infection in neonatal intensive care Unit.*J.Pediatr*, 32(4):333 -8.

Sarkar S, Bhagat I, DeCristofaro JD. (2006): A study of the role of multiple site blood cultures in the evaluation of neonatal sepsis. *J Perinatol*. 26(1):18-22.

Schaison G, Eden OB, Henze G, et al. (1998): Recommendations on the use of colony-stimulating factors in children: conclusions of a European panel. *Eur. J. Pediatr* .157:955– 66.

Schelonka RL, Yoder BA, desJardins SE, Hall RB, Butler TJ. (1994): Peripheral leukocyte count and leukocyte indexes in healthy newborn term infants. *J Pediatr*; 125:603-606.

Schelonka RL, Yoder BA, Hall RB. (1995): Band neutrophil counts in neonates [Letter: Reply]. *J Pediatr*; 126:505-506.

Schelonka RL, Yoder BA. The WBC count and differential: its uses and misuses. *Contemp. Pediatr*. **1996**; 13:124-141.

Schiffel H, Lang SM, Fischer R. (2002): Daily hemodialysis and the outcome of acute renal failure. *N .Engl. J .Med*; 346(5):305– 16.

Schmidt B, Kirpalani H, Rosenbaum P, Cadman D. (1988): Strengths and limitations of the Apgar score: a critical appraisal. *J Clin Epidemiol*;41:843-850.

Schmidt BK, Kirpalani HM, Corey M, Low DE, Philip AGS, Ford-Jones EL. (1987): Coagulase-negative staphylococci as true pathogens in newborn infants: a cohort study. *Pediatr Infect Dis J*; 6:1026-1031.

Schouten-Van Meeteren NYN, Rietveld A, Moolenaar AJ, Van Bel F. (1992): Influence of perinatal conditions on C-reactive protein production. *J Pediatr*;120:621-624

Schouten-Van Meeteren NYN, Rietveld a, Moolenaar AJ, Van Bel F. (1992): Influence of perinatal conditions on C-reactive protein production. *J Pediatr* 1992; 120:621-624.

Schrag SJ, Zell ER, Lynfield R, Roome A, Arnold KE, Craig AS, et al. (2002): A population-based comparison of strategies to prevent early-onset group B streptococcal disease in neonates. *N. Engl. J. Med.*; 347:233-239.

Schrag SJ, Zywicki S, Farley MM, Reingold AL, Harrison LH, Lefkowitz LB, et al. (2000): Group B streptococcal disease in the era of intrapartum antibiotic prophylaxis. *N Engl J Med*;342:15-20

Seaward PG, Hannah ME, Myhr TL, et al. (1998): International multicenter term PROM study: evaluation of predictors of neonatal infection in infants born to patients with premature rupture of membranes at term. *Premature Rupture of the Membranes. Am J Obstet Gynecol.* Sep; 179(3 Pt 1):635-9.

Seigle J.D. &MacCracken, C.H. (1981): single dose penicillin prophylaxis against group B-streptococcal infection. *New Eng., J. Med.*, 303: 769-776. severe sepsis. *N Engl J Med* 2001;344(10):699– 709.

Shine B, Gould J, Campbell C, Hindocha P, Wilmot RP, Wood CBS. (1985): Serum C-reactive protein in normal and infected neonates. *Clin Chim Acta*;148:97-103.

Speer CH, Bruns A, Gahr M. (1983): Sequential determination of CRP, α_1 -antitrypsin and haptoglobin in neonatal septicemia. *Acta Paediatr Scand*;72:679-683.

Spitzer A. (1978): Renal physiology and functional development. In *Pediatric Kidney Disease*, Edelmann CM (Ed): Boston, Little Brown, , pp25.

Squire E, Favara B, Todd J. (1979): Diagnosis of neonatal bacterial infection: hematologic and pathologic findings in fatal and nonfatal cases. *Pediatrics*;64:60-64.

St. Geme JW, Jr, Murray DL, Carter J, Hobel CJ, Leake RD, Anthon BF, et al. (1984): Perinatal bacterial infection after prolonged rupture of amniotic membranes: an analysis of risk and management. *J Pediatr*; 104:608-613.

Stapleton FB, Jones DP, Green RS. (1987): Acute renal failure in neonates: incidence, etiology and outcome. *Pediatr. Nephrol.* 1987 Jul; 1(3):314-20

Steele BT, Paes B, Towell ME, Hunter DJ. (1988): Fetal renal failure associated with intrauterine growth retardation Am J Obstet Gynecol. Nov; 159(5):1200-2.

Stoll BJ, Gordon T, Korones SB, Shankaran S, Tyson JE, Bauer CR, et al. (1996): Early-onset sepsis in very low birth weight neonates: a report from the National Institute of Child Health and Human Development Neonatal Research Network. J Pediatr;129:72-80.

Stoll BJ, Gordon T, Korones SB, Shankaran S, Tyson JE, Bauer CR, et al. (1996): Late-onset sepsis in very low birth weight neonates: a report from the National Institute of Child Health and Human Development Neonatal Research Network. J Pediatr;129:63-71.

Stoll BJ, Hansen N, Fanaroff AA, Wright LL, Carlo WA, Ehrenkranz RA, et al. (2002): Changes in pathogens causing early-onset sepsis in very-low-birth-weight infants. N Engl J Med; 347:240-247

Stoll BJ, Holman RC, Shuchat A. (1998): Decline in sepsis-associated neonatal and infant deaths 1974–1994. Pediatrics; 102:E18.

T

Taketomo CK, Hoddings JH, Kraus DM. (1996): Pediatric Dosage Handbook. 3rd ed. 1996-97.

Tausch WH, Ballard RA, Avery. Schaffer & Avery's (1998): Diseases of the Newborn. 7th ed. 1998; 435-452, 490-512.

Taylor FB, Chang A, Ruf W. Lethal E.(1991) coli septic shock is prevented by blocking tissue factor with a monoclonal antibody. Circ Shock 1991;33:127– 34.
temporal relationships. Pediatr Infect Dis J 1990;(3):196– 200.

Thomas NJ, Carcillo JA, Doughty LA, et al. (2002): Plasma concentrations of defensins and lactoferrin in children with severe sepsis. Pediatr Infect Dis J 2002; 21(1):34– 8.

Tollner U (1982):Early diagnosis of sepsis in newborn:clinical studies &sepsis score Eur. J.Pediatr.,138:331.

Toth-Heyn P ,Drukker A,Guignard JP. (2000): the stressed neonate kidney: *Pediatr. Nephrol.* 14:227-239.

Towers CV, Briggs GG. (2002): Antepartum use of antibiotics and early-onset neonatal sepsis: the next 4 years. *Am J Obstet Gynecol*; 187:495-500.

Trachtman H,:Sodium &water (2004) : pediatric Nephrology 5th ed Williams &Wilkins; 2004

Tulassy T,Vasarhely B .(2002): Birth weight &Renal functions. *Curr. Opin .Nephrol.* 14:347-352.

U

Unni JC, Date A, Raghupathy P, Shastry JCM. (1986): Medical renal disease in South Indian infants. *Acta. Pediatr. Scand* 1986; 75: 1030-1031.

V

Van Lente F, Pippenger CE. (1987): The pediatric acute care laboratory. *Pediatr Clin North Am*; 34:231-246.

Volk HD, Reinke P, Krausch D, et al. (1996): Monocyte deactivation – rationale for a new therapeutic strategy in sepsis. *Intensive Care Med*; 22:S4

W

Wasunna A, Whitelaw A, Gallimore R, Hawkins PN, Pepys MB. (1990): C-Reactive protein and bacterial infection in preterm infants. *Eur J Pediatr*;149:424-427.

Watson RS, Carcillo J, Linde-Zwirble WT, et al. (2003): The epidemiology of severe sepsis in children in the United States. *Am J Respir Crit Care Med*;167:695– 703.

Wedekin M, Ehrich JH, Offner G, Pape L. (2008): Aetiology and outcome of acute and chronic renal failure in infants. *Nephrol Dial Transplant*. 2008 May; 23(5):1575-80. Epub .

Weinberg GA, Powel KR. (2001): Laboratory aids for diagnosis of neonatal sepsis. Remington JS Klein JO eds. *Infectious diseases of the fetus and newborn infant* 2001:1327-1344 WB Saunders Philadelphia.

Wendel GD, Jr, Leveno KJ, Sánchez PJ, Jackson GL, McIntire DD, Siegel JD. (2000): Prevention of neonatal group B streptococcal disease: a combined intrapartum and neonatal protocol. *Am J Obstet Gynecol*; 186:618-626.

Whalen MJ, Doughty LA, Carlos TM, et al. (2000): Intercellular adhesion molecule– 1 and vascular Cell adhesion molecule– 1 are increased in the plasma of children with sepsis induced multiple Organ failure. *Crit Care Med* 2000; 28:2600–7.

Wilkins I, Mezrow G, Lynch L, Bottone EJ, Berkowitz RL. (1989): Amnionitis and life-threatening respiratory distress after percutaneous umbilical blood sampling. *Am J Obstet Gynecol* 1989; 160:427-428.

Y

Yoder PR, Gibbs RS, Blanco JD, Castaneda YS, St. Clair PJ. A (1983): prospective, controlled study of maternal and perinatal outcome after intra-amniotic infection at term. *Am J Obstet Gynecol*; 145:695-701.

Young, TE, Mangum, B. (2006): Neofax 2006. *Neofax 2006*. 2006;19th edition:2-75.

Z

Zager RA, (1992): renal ischemia .*Circ .Res*; 70:20

Zenz W, Bodo Z, Zobel G. (2000): Recombinant tissue plasminogen activator restores perfusion in meningococcal purpura fulminans. *Crit. Care Med*; 26(5):969– 71.