

# Reference

**Abdalla MGK, Beattie B and Alfirevic Z.** Intrauterine transfusion technique in fetuses with Rhesus incompatibility. In The Cochrane Library, John Wiley and Sons L, eds. Issue 2, 2004.

**Adin CA, Croker BP and Agarwal A.** Protective effects of exogenous bilirubin or ischemia-reperfusion injury in the isolated perfused rat kidney. *Am J Physiol Renal* 2005; 288(4): F778-F784.

**Agarwal RK and Deorari AK.** Neonatal Jaundice. *Indian Pediatr* 1999; 36: 110-112.

**Agarwal V, Singh V, Goel SP, et al.** Maternal and neonatal factors affecting physiological jaundice in western VP. *Indian J Physiol Pharmacol* 2007 Apr-Jun; 51(2): 203-6.

**Avent ND and Reid ME.** The Rh blood group system: In Neil D and Marion E, eds. A review, *Blood* 2000; 95(2):375-387.

**Bazlul Karim ASM1 and Kamal M2.** Cholestatic jaundice during infancy: experience at a tertiary-care center in Bangladesh. *Indian J Gastroenterol* 2005; 24: 52-4.

**Bertini G, Dani C, Tronchin M and Rubaltelli FF.** Is breastfeeding really favoring early neonatal jaundice? *Pediatrics* 2001; 107(3): E41.

**Bertini G and Rubaltelli FF.** Non-invasive bilirubinometry in neonatal jaundice, *Semin Neonatol* 2002; 7:129-33.

- Bertini G, Pratesi S, Cosenza E and Dani C.** Transcutaneous bilirubin measurement: evaluation of Bilitest. *Neonatology* 2008; 93(2):101-5.
- Bhutani VK, Johnson L and Sivieri EM.** Predictive ability of a predischage hour-specific serum bilirubin for subsequent significant hyperbilirubinemia in healthy term and near-term newborns, *Pediatrics*.1999; 103 :6 –14
- Bhutani V, Gourley GR, Adler S, Kreamer B, et al.** Non invasive measurement of total serum bilirubin in a multiracial predischage newborn population to asses the risk of sever hyperbilirubinemia, *Pediatrics* 2000; 106 (2).
- Bhutani VK, Johnson LH, Maisels MJ, et al.** Kernicterus: epidemiological strategies for its prevention through system-based approaches. *J Perinatol* 2004; 24:650-62.
- Bhutani VK and Johnson LH.** Urgent clinical need for accurate and precise bilirubin measurements in the United States to prevent kernicterus, *Clin Chem* 2004, 50: 477-80.
- Billing BM.** Intestinal and renal metabolism of bilirubin including enterohepatic circulation In: Ostrow JD, editor. *Bile Pigments and Jaundice; Molecular, Metabolic and Medical Aspects*. New York: Marcel Dekker, 1986: 255-269.
- Boo NY and Lee HT.** Randomized controlled trial of oral versus intravenous fluid supplementation on serum bilirubin level

during phototherapy of term infants with severe hyperbilirubinaemia, J Paediatr Child Health 2002; 38:151-5

**Bosch AM.** Classical galactosemia revisited. J Inherit Metab Dis, Jul(11) 2006.

**Bowman JM.** ABO hemolytic disease. In: Creasy RK, Resnik R, eds. Maternal and Fetal Medicine. 4<sup>th</sup> eds. Philadelphia, PA: WB Saunders; 1999:736-767.

**Briscoe L, Clark S and Yoxall CW.** Can transcutaneous bilirubinometry reduce the need for blood tests in jaundiced full term babies? Archives of Disease in Childhood Fetal & Neonatal Edition, 2002; 86(3), F190-192.

**Buchan PC.** The pathogenesis of neonatal hyperbilirubinemia after induction of labour with oxytocin. Br Med J 1979 Nov17: 1255-7.

**Buiter HD, Dijkstra SS, et al.** Neonatal jaundice and stool production in breast or formula-fed term infants. Eur J pediatr 2008 May; 167(5):501-7.

**Burgos AE, Schmitt SK, Stevenson DK, et al.** Readmission for neonatal jaundice in California 1991-2000: Trends and Implications. Pediatrics 2008 Apr; 121(4) e86-9.

**Carbonell X, Botet F, Figueras J, et al.** Prediction of hyperbilirubinaemia in the healthy term newborn, Acta Paediatr 2001; 90:166 –170.

**Crowther CA and Middleton P.** Anti D administration after childbirth for preventing Rhesus alloimmunization. In The Cochrane Library, John Wiley and Chichester UK, eds. Issue 2, 2004.

**Chou SC, Palmer RH, Ezhuthachan S, et al.** Management of hyperbilirubinemia in newborns: measuring performance by using a benchmarking model. Pediatrics 2003; 112:1264-73.

**Cui Y, Konig J, Leier I, Buchholz U, et al.** Hepatic uptake of bilirubin and its conjugates. J Biochem 2001.

**Dennery PA, Seidman DS and Stevenson DK.** Neonatal hyperbilirubinemia. In Johnson LH, Bhutani VK, Brown AK, et al, (eds). Case Based Pediatrics For Medical Students and Residents, N Engl J Med 2001; 344:581-90.

**Dennery PA, Stevenson DK and Hintz SR.** Understanding neonatal jaundice. J Perinatol 2001; 21: 521-4.

**Diane J and Madlon-Kay D.** Maternal assessment of neonatal jaundice after hospital discharge- Original Research-Statistical Data Included, Journal of Family Practice, May 2002.

**Dsouza SW, Black P, MacFarlane T and Richards B.** The effect of oxytocin in induced labour on neonatal jaundice. Br J Obstet Gynaecol 1979 Feb 86(2): 133-8.

**Ebbesen F, Rasmussen LM and Wimberley PD.** A new transcutaneous bilirubinometer, BiliCheck, used in the neonatal intensive care unit and the maternity ward, Acta Paediatr 2002; 91:203-11.

- Edwards MC and Fletcher MA.** Exchange transfusions. In: Fletcher MA, MacDonald MG, eds. Atlas of Procedures in Neonatology. 2<sup>nd</sup> ed. Philadelphia, PA: JB Lippincott Co; 1993: 363-372.
- Engmann C and Schumacher RE:** In phototherapy, Michigan Manual of Neonatal Intensive Care 3<sup>rd</sup> Edition Ed. Donn SM, Hanley and Belfus 2003 – Pg 351 – 354.
- Eyssette-Guerreau S, Bader-Meunier B, Garconl L, et al.** Infantile Pyknocytosis: a cause of hemolytic anemia of the newborn: Br J Haematol 2006 May; 133(4): 439-42.
- Farrant P, Meire HB and Mieli-Vergani G.** Ultrasound features of the gall bladder in infants presenting with conjugated hyperbilirubinemia. Br J Radiol 2000; 73: 1154.
- Fernandes A, Falcao AS, Silva RF, et al.** Inflammatory signalling involved in astroglial activation by unconjugated bilirubin. J Neurochem 2006; 96(6): 1667-1679.
- Foley TP.** Congenital hypothyroidism. In: Werner and Ingbars the thyroid, 7<sup>th</sup> ed. J Clin Endocrinal Metab 2000; 85: 3975-3987.
- Fox IJ, Roychowdhury J, Kaufman SS, et al.** Treatment of Crigler-Najjar Syndrome type 1 with hepatocyte transplantation. N Engl J Med 1998; 38: 1422-1428.
- Funato M, Teraoka S, Tamai H, et al.** Follow-up study of auditory brainstem responses in hyperbilirubinemic newborns treated with exchange transfusion, Acta Paediatr Jpn 1996; 38: 17-21.

- Gale R, Seidman DS, Dollberg S, et al.** Epidemiology of neonatal jaundice in the Jerusalem population, J Pediatr Gastroenterol Nutr 1990;10(1): 82-6.
- Gallagher pG and Lux SE.** Disorders of the Erythrocyte Membrane.In: Nathan DG and Orkin SH, eds. Hematology of infancy and childhood 2003; 617-632.
- Garcia FJ and Nager AL.** Jaundic as an early diagnostic sign of urinary tract infection in infancy. Pediatrics 2002; 109: 846-851.
- Garden AS and Davidson DC.** Recommendations for the management of galactosemia, Arch Dis Child, Mar 2000; 82(3): 266.
- Gartner LM and Hershel M.** Jaundice and breastfeeding. Pediatr Clin North Am. 2001; 48: 389-99.
- Gault DT.** Extravasation Injuries. Br J Plast Surg 1993; 46(2): 91-6.
- Geiger AM, Petitti DB and Yao JF.** Rehospitalization for neonatal jaundice: Risk factors and outcomes. Pediatr Perinat Epidemiol 2001 Oct; 15(4):353-8.
- Gennuso F, Ferentti C, Tirolò C, et al.** Bilirubin protects astrocytes from its own toxicity by inducing up-regulation and translation of multidrug resistance-associated protein 1 (Mrp1). Proc Natl Acad Sci USA 2004, 101(8): 2470-2475.
- Goessling W and Zucker SD.** The Role of Apolipoprotein D in the Transport of Bilirubin in Plasma. Am J Physiol Gastrointest, Liver Physiol 2000; 279(2) G356-365.

- Gonzalez-Quintero VH, Istwan NB, Rhea DJ, Rodriquez LI, et al.** The impact of glycemic control on neonatal outcome in singleton pregnancies complicated by gestational diabetes. *Diabetes Care*, 2007 Mar; 30(3):467-70.
- Gourley GR, Li Z, Kreamer BL, et al.** A controlled, Randomized, Double-Blind Trial of prophylaxis against jaundice among breastfed newborn. *Pediatrics* 2005; 116: 385-391.
- Grimmer I, Berger-Jones K, Buhrer C, et al.** Late neurological sequelae of non-hemolytic hyperbilirubinemia of healthy term neonates. *Acta Paediatr* 1999; 88: 661-3.
- Grompe M.** The pathophysiology and treatment of hereditary tyrosinemia type 1. *Semin Liver Dis*, 2001: Nov; 21(4): 563-71.
- Guglielmi FW, Boggio-Bertinet D, Federico, et al.** Total parenteral nutrition-related gastroenterological complication. In: Palmo A, Panella C, et al (eds). *Digestive and Liver Disease*, Vol 38, Issue9, Sep 2006, P 623-642.
- Gupta AK and Mann SB.** Is auditory brainstem response a bilirubin neurotoxicity marker? *Am J Otolaryngol* 1998; 19: 232-236.
- Gupta R and Parashar Y.** Crigler-Najjar syndrome type II. *Indian J Pediatr*, Nov 2004; 71(11): 1043.
- Halamek LP and Stevenson DK.** Neonatal Jaundice and Liver disease; In Fanaroff AA, Martin RJ (eds): *Hepatology* 2002; 35:905-911.

**Hansen TW.** Bilirubin Oxidation in Brain. *Mol Genet Metab* 2000; 71(1-2) 411-417.

**Herschel M and Beutler E.** Low glucose 6-phosphate dehydrogenase enzyme activity level at the time of hemolysis in a male neonate with the African type of deficiency. *Blood Cells, Molecules and Diseases* 2001; 24,918-923.

**Hirschfield GM and Alexander GJ.** Gilberts syndrome: an overview for clinical biochemists. *Ann Clin Biochem*, Sep 2006; 43: 340-3.

**Holton JB, Walter JH and Tyfield LD.** Galactosemia. In: Scriver CR, Kaufman S, Eisensmith E, (eds). *The Metabolic and Molecular Aspects of Inherited Disease*, Mc Mill 2001, New York Ch 72.

**Horvath A, Koletzko B, Kalisz M and Szajewska H.** The effect of supplemental fluids or feedings during the first days of life on the success and duration of breast-feeding: A systematic review of randomized controlled trials, *Arch Pediatr Adolesc Med* 2005; 159:597-8.

**Huang MJ, Kua KE, Teng HC, et al.** Risk factors for severe hyperbilirubinemia in neonates. *Pediatr Res* 2004 Nov; 56(5):682-9.

**Hwang S and Shulman R.** Approach to Neonatal Cholestasis, *Semin Liver Disease* 2002; 18: 281-285.

**Iinuma Y, Narisawa R, Iwafuchi M, et al.** The role of endoscopic retrograde cholangiopancreatography in cholestatic infants. *J Pediatr Surg* 2000; 35:545.

- Immenschuh S and Schroder H.** Heme oxygenase 1 and cardiovascular disease. *Histol Histopathol* 2006; 21(6): 679-685.
- Iolascan A, Miraglia Del Giudice E, Perrotta S, et al.** Hereditary Spherocytosis: *Haematologica* Vol 83, Issue 3, 1998: 240-257.
- Iranpour R, Akbar MR and Haghshenas I.** Glucose-6-phosphate dehydrogenase deficiency in neonate. *Indian J Pediatr* 2003 Nov; 70(11) 855-7.
- Jackson JC.** Adverse events associated with exchange transfusion in healthy and ill newborns, *Pediatrics* 1997; 99(5).
- Jansen PL.** Diagnosis and Management of Crigler-Najjar Syndrome. *Eur J Pediatr*, Dec 1999; 158 Suppl 2: S 89-94.
- Johnson JD, Aldrich M, Angelus P, et al.** Oxytocin and neonatal hyperbilirubinemia. Studies of bilirubin production. *Am J Dis Child* 1984 Nov; 138(11):1047-50.
- Johnson L and Bhutani VK.** Guidelines for management of the jaundiced term and near-term infant, *Clin Perinatol* 1998; 25(3): 555-74.
- Kadakol A, Ghosh SS and Sappal BS.** Genetic lesions of bilirubin uridine-diphosphoglucuronate glucuronosyl transferase (UGT1A1) causing Crigler-Najjar and Gilbert syndrome: Correlation of genotype to phenotype, *Pediatrics* 2000 oct; 106(4): 297-306.

**Kanno H.** Hexokinase: gene structure and mutations. Baillieres Best Pract Res Clin Haematol 2000; 13:83-86.

**Kaplan M, Algur N and Hammerman C.** Onset of jaundice in glucose 6-phosphate dehydrogenase-deficient neonate. Pediatrics 2001 Oct; 108(4): 956-9.

**Kaplan M, Herschel M, Hammerman C, et al.** Hyperbilirubinemia among African American, Glucose-6-phosphate dehydrogenase deficient neonate. Pediatrics 2004; 114. e213-e214.

**Kaplan M, Muraca M, Hammerman C, et al.** Imbalance between production and conjugation of bilirubin. A fundamental concept in the mechanism of neonatal jaundice. Pediatrics 2002;110(4).

**Keffler S, Kelly DA, Powell JE, et al.** Population screening for neonatal liver disease: a feasibility study. J Pediatr Gastroenterol Nutr 1998, Sep; 27(3): 306-11.

**Keren R, Bhutani VK, Luan X, Nihtianova S, et al.** Identifying newborns at risk of significant hyperbilirubinemia: a comparison of two recommended approaches. Arch Dis Child 2005; 90; 415-421.

**Kilic S, Texcan S, Tascilar E, et al.** Morbidity and Mortality characteristics of infants hospitalized in the pediatrics department of the largest Turkish military hospital. Mil Med 2005; 170 (1): 48-51.

**Keren R, Luan X, Friedman S, et al.** A comparison of alternative risk-assessment strategies for predicting significant neonatal

hyperbilirubinemia in term and near term infants.

Pediatrics, 2008 Jan; 121(1): e 170-9

**Khatun N, Latif SA and Vddin MM.** Infant outcomes of gestational diabetes mellitus. Mymensingh Med J 2005 Jan; 14(1):29-31.

**Kolman KB, Mathieson KM and Frias C.** A comparison of transcutaneous and total serum bilirubin in neonate. Hispanic infants at 35 or more weeks of gestation. J Am Board Fam Med 2007 May-Jun; 20(3):260-71.

**Kotal P, Van Der Veer CN, Sinaasappel M, et al.** Intestinal excretion of unconjugated bilirubin in man and rats with inherited unconjugated hyperbilirubinemia. Pediatr Res 1997; 42(2): 195-200.

**Kuroda M, Kobayashi Y, Tanaka Y, et al.** Increased hepatic and renal expressions of multidrug resistance-associated protein 3 in hyperbilirubinemia in rats. J Gastroenterol Hepatol 2004; 19(2):146-153.

**Lee CH, wang PW, Lee TT, et al.** The significance of functioning gall bladder visualization on hepatobiliary scintigraphy in infants with persistent jaundice, J Nucl Med 2000; 41:1209.

**Leslie ND.** Insights into the pathogenesis of galactosemia, Ann Rev Nutr 2003; 23: 59-80.

**Linder N, Yatsir T, Tsur M, et al.** Unexplained neonatal jaundice as an early diagnostic sign of septicemia in the newborn. *J Prenatal* 1988; (8): 325-7.

**Macasai MS, Schwartz TL, Hinkle D, et al.** Tyrosinemia type II: nine cases of ocular signs and symptoms, *Am J Ophthalmol* 2001; 132: 522-7.

**Maisels MJ: Jaundice In: Avery GB, Fletcher MA, Mac Donald MG,** eds: *Neonatology: Pathophysiology and Management of the newborn*, 4<sup>th</sup> edition, Philadelphia: JB Lippincott Company, 1994:630-725.

**Maisels MJ and Kring EA.** Length of stay, jaundice, and hospital readmission, *Pediatric* 1998; 101:995 –998.

**Maisels MJ and Newman TB.** Predicting hyperbilirubinemia in newborns:the importance of timing. *Pediatrics* 1999; 103:493-495.

**Maisels MJ, Baltz RD, Bhutani V, et al.** Management of hyperbilirubinemia in newborn infant 35 or more weeks of gestation, *Pediatrics* 2004 (A); 114: 297-316.

**Maisels MJ, Ostrea EM, Touch S, et al.** Evaluation of a new transcutaneous bilirubinometer. *Pediatrics*, 2004 (B); 114: (6), 1628-1635.

**Maisels MJ.** A primer on phototherapy for the jaundiced newborn, *Contemp Pediatr* 2005 (A): 768-846.

- Maisels MJ.** Jaundice In: Mac Donald MG, Seshia MMK, Mullett MD, eds. Neonatology: Pathophysiology and Management of the Newborn. Philadelphia, Pa: Lippincott Co; 2005 (B): 768-896.
- Marcuello E, Altes A, Menoyo A, Del Rio E, et al.** UGT1A1 gene mutations and irinotecan treatment in patients with metastatic cancer; Journal of Autoimmunity 2004.
- Martin TC, Shaa M, Alexander D, et al.** Did exclusive breast feeding and early discharge lead to excessive bilirubin levels in newborns in Antigua and Barbuda? Pediatrics 2002.
- Martinez JC, Maisels MJ, Otheguy L, et al.** Hyperbilirubinemia in the breast-fed newborn: A controlled trial of four interventions. Pediatrics 1993; 91:470-3.
- Martinez JC, Maisels MJ, Otheguy L, et al.** Hyperbilirubinemia in the breast-fed newborn: a controlled trial of four interventions. Pediatrics 1993; 91: 470-473.
- Maruo Y, Nishizawak K, Sato M, et al.** Prolonged unconjugated hyperbilirubinemia associated with breast milk and mutations of the bilirubin uridine diphosphate glucucosyl transferase gene. Pediatrics 2000; 106: 59-63.
- McKiernan PJ.** Neonatal Cholestasis, Semin Neonatol 2002; 7: 153-165.
- McLin VA and Balistereri WF.** Approach to neonatal cholestasis. In: Pediatric Gastrointestinal Disease: Pathophysiology, Diagnosis, Management, 4<sup>th</sup> ed, Walker WA, Goulet O, Kleinman RE, et al (eds), 2004.p1079.

- Mehta S, Kumar P and Narang A.** A randomized controlled trial of fluid supplementation in term neonates with severe hyperbilirubinemia, J Pediatr 2005; 147:781-5.
- Melton K and Akinbi HT.** Neonatal jaundice strategies to reduce bilirubin-induced complications. Postgrad Med 1999; 106: 167-8,171-4,177-8.
- Mercier CE, Barry SE and Paul K.** Improving Newborn Preventive Services at the Birth Hospitalization: A collaborative, Hospital-Based Quality-Improvement Project. Pediatrics: 2007 Sep; 120(3): 481-488.
- Mills JF and Tudehope D.** Fiberoptic therapy for neonatal jaundice. Cochrane Database Syst Rev 2001.
- Miqdad AM, Abdelbasit OB, Shaheed MM, et al.** Intravenous immunoglobulin G (IVIG) therapy for significant hyperbilirubinemia in ABO hemolytic disease of the newborn, J Matern Fetal Neonatal Med 2004; 16:163-6.
- Mishra S, Agarwal R, Deorari AK, et al.** Jaundice in the newborn, Indian J Pediatr 2008; 75:157-63.
- Moise KJ.** Red blood cell alloimmunization in pregnancy. Semin Hematol 2005 Jul; 42(3) 169-78.
- Morita T.** Heme Oxygenase and Atherosclerosis. Arterioscler Thromb Vasc Biol 2005; 25(9): 1686-1795.

- Mowat AP, Davidson LL and Dick MC.** Earlier identification of biliary atresia and hepatobiliary disease: Selective Screening in the third week of life, Arch Dis Child 1995; 72(1): 90-92.
- Moyer V, Freese DK, Whittington PF, et al:** Guideline for the evaluation of cholestatic jaundice in infants, J Pediatr Gastroenterol Nutr 2004, Aug; 39(2): 115-28.
- Moyer VA, Ahn C and Sneed S.** Accuracy of clinical judgment in neonatal jaundice. Arch Pediatr Adolesc Med 2000; Apr 154:391-4.
- Murray NA and Roberts IA.** Haemolytic Disease of the Newborn. Arch Dis Child Fetal Neonatal Ed.2007 Mar; 92(2): F83-89.
- Nanjundaswamy S, Petrova A, Mehta R, et al.** The accuracy of transcutaneous bilirubin measurements in neonates: a correlation study. Biol Neonate 2004; 85(1):21-5.
- Newman TB, Xiong B, Gonzales VM, et al.** Prediction and prevention of extreme neonatal hyperbilirubinemia in a mature health maintenance organization, Arch Pediatr Adolesc Med. 2000; 15 :1140 –1147.
- Newman TB, Escobar GJ, Liljestrand P, et al.** Combining laboratory and clinical data for prediction of hyperbilirubinemia in newborns. Pediatr Res 2003; 53:2259.
- Nicoll A, Ginsburg R and Tripp JH.** Supplementary feeding and jaundice in newborn. Acta Pediatr Scand 1982; 71: 759-761.

**Norton KI, Glass RB, Kogan D, et al.** MRCholangiography in the Evaluation of Neonatal Cholestasis: Initial Results, Radiology 2002; 222-687.

**Oral E, Gezer a, Cagdas a, et al.** Oxytocin infusion in labour: the different indications and the use of different diluents on neonatal bilirubin level. Arch Gynecol Obstet 2003 Jan; 267(3):117-20.

**Ostrow JD, Pascolo L, Brites D, Tiribelli C, et al.** Molecular basis of bilirubin induced toxicity. Trends Mol Med 2004; 10(2): 65-70.

**Ostrow JD, Pascolo L, Shapiro SM, Tiribelli C, et al.** New Concepts in Bilirubin Encephalopathy. Eur J Clin Invest, 2003; 33(11): 988-997.

**Perlstein MA.** The late clinical syndrome of posticteric encephalopathy; Pediatr Clin North Am 1960; 7:665-87.

**Phillips MJ, Azuma T, Meredith SL, et al.** Abnormalities in villin gene expression and canalicular microvillus structure in progressive cholestatic liver disease of childhood, Lancet 2003, 362: 1112-1119.

**Prchal JT and Gregg XT.** Red cell enzymopathies. In: Hoffman R, Benz E, eds. Hematology: Basic Principles and Practice, 4<sup>th</sup> ed 2005: 653-659.

**Prickles CJ, Symonds EM and Broughton Pipkin F.** The fetal outcome in randomized double-blind controlled trial of labetalol versus placebo in pregnancy-induced hypertension, Br J Obstet Gynaecol 1989 Jan; 96(1):38-43.

- Raijmakers MT, Jansen PL, Steegers EA, et al.** Association of human liver bilirubin UDP glucuronyl transferase activity with apolipomorphism in the promoter region of the UGT1A1 gene, *Journal of Hepatology* 2000; 33(3): 348-351.
- Riskin A, Abend-weinger M and Bader D.** How Accurate are Neonatologists in Identifying Clinical Jaundice in Newborn? *Clin Pediatr* 2003; 42: 153-158.
- Rodrigues CMP, Solo S and Brites D.** Bilirubin induce apoptosis via the mitochondrial pathway in developing rat brain neurons. *Hepatology* 2002; 35: 1185-1195.
- Rosenkrantz TS.** Polycythemia and hyperviscosity in the newborn. *Semin Thromb Hemost*, Oct 2003; 29(5): 515-27.
- Roth P and Polin RA.** Controversial Topics in Kernicterus. *Clin Perinatol* 1988; 15(4): 965-990.
- Rubaltelli FF, Gourley GR, Loskamp N, et al.** Transcutaneous bilirubin measurement: a multicenter evaluation of a new device, *Pediatrics* 2001; 107: 1264 –1271.
- Russo PA, Mitchell GA and Tanguay RM.** Tyrosinemia: a review, *Pediatr Dev Pathol* 2001, May-Jun; 4(3) 212-21.
- Samanta S, Tan M, Kissack C, et al.** The value of Bilicheck as a screening tool for neonatal jaundice in term and near-term babies. *Acta Paediatr* 2004; 93:1486-90.

- Sanpavat S& Nuchprayoon I.** Noninvasive transcutaneous bilirubin as a screening test to identify the need for serum bilirubin assessment. *Journal of the Medical Association of Thailand*, 2004; 87: (10), 1193-1198.
- Seidman DS, Ergaz Z, Paz I, et al.** Predicting the risk of jaundice in full term newborns: a prospective population based study, *J Perinatol* 1999 Dec; 19:564-7.
- Seidman DS, Moise J, Ergaz Z, et al.** A new blue light-emitting phototherapy device: A prospective randomized controlled study, *J Pediatr* 2000; 136: 771-4.
- Semmekrot BA, Devries MC, et al.** Optimal breast feeding to prevent hyperbilirubinemia in healthy term newborns. *Ned Tijdschr Geneesk* 2004 Oct9; 148(41):2016-9.
- Shapiro SM.** Definition of the clinical spectrum of kernicterus and Bilirubin-Induced Neurological Dysfunction (BIND). *J Perinatol* 2005; 25(1): 54-59.
- Snyder EL and Shoos Lipton K.** Prevention of hemolytic disease of the newborn due to anti-D. In: *Bullet in 98-2*, Bethesda (MD): American Association of Blood Banks; February16, 1998 p 1-6.
- Stevenson DK, Fanaroff AA, Maisels MJ, et al.** Prediction of hyperbilirubinemia in near-term and term infants, *Pediatrics* 2001; 108 :31 –39.

**Stoll BJ and Kliegman RM.** Digestive System Disorder. In: Behrman RE, et al (eds). Nelson Textbook of Pediatrics, 17<sup>th</sup> ed 2004; pp.588-599.

**Stoll BJ and Kliegman RM.** The Foetus and Neonatal Infant. In: Nelson Textbook of Pediatrics, 16<sup>th</sup> , Behrman RE, Kliegman RM, Jenson HB, eds. Philadelphia, WB, Saunders 2000; pp 451-551.

**Szabo P, Wolf M, Bucher HU, et al.** Detection of hyperbilirubinaemia in jaundiced full-term neonates by eye or by bilirubinometer? Eur J Pediatr 2004; 163:722-7.

**Tiberi E, Latella C, Parenti D and Romagnoli C.** Predictive ability of a pre discharge hour-specific serum bilirubin for hyperbilirubinemia in full term infants. Minerva Pediatr, 2007 Jun; 59(3): 183-9.

**Trevett TN, Dorman K, Lamvu G and Moise KJ.** Antenatal maternal administration of phenobarbital for the prevention of exchange transfusion in neonates with hemolytic disease of the fetus and newborn. Am J Obstet Gynecol. 2005; 192(2):478-82.

**Trioseco JA, Aly H, Milner J, Patel K, et al.** Dose gender affect the neonatal hyperbilirubinemia in low birth weight infants? Pediatr Crit Care Med 2005 Mar; 6(2):171-4.T

**Van Mil SW, Houwen RH and Klomp LW.** Genetics of Familial Intrahepatic cholestasis syndromes. J Med Genet, Jun 2005; 42(6): 449-63.

- Venigalla S and Gourley GR.** Neonatal Cholestasis, *Semin Perinatol* 2004, Oct; 28(5): 348-55.
- Verlinsky Y, Rechitsky S, Schoolcraft W, et al.** Preimplantation diagnosis for fanconi anemia combined with HLA matching. *J AMA* 2001; 285, 3130-3133.
- Vitek L, Majer F, Muchoval L, Zelenka J, et al.** Identification of bilirubin reduction products formed by clostridium perfringens isolated from human neonatal rectal flora. *Analyt Technol Biomed Life Sci* 2006; 833(2) 149-159.
- Wananukul S, Praisuwanna P and Kesorncam K.** Effects of clear topical ointment on transepidermal water loss in jaundiced preterm infants receiving phototherapy. *J Med Assoc Thai* 2001; 84(6):837-41.
- Watchko JF.** Exchange transfusion in the management of neonatal hyperbilirubinemia. In: Maisels MJ, Watchko JF, eds. *Neonatal Jaundice*. London, United Kingdom: Harwood Academic Publishers; 2000: 169-176.
- Wennberg RP, Ahlfors CE, Bhutain VK, et al.** Toward Understanding Kernicterus: A challenge to improve the management of jaundiced newborn. *Pediatrics* 2006; 117: 474-485.
- Willems WA, Van Den Berg LM, De Wit H, et al.** Transcutaneous bilirubinometry with the Bilicheck in very premature newborns, *J Matern Fetal Neonatal Med* 2004; 16: 209-14.
- Woyton J, Agarwal P and Zimmer M.** Evaluation of the effect of oxytocin use for labour induction or frequency of occurrence and

severity of neonatal jaundice. Ginekol Pol 1994 Dec: 65(12):682-5.

**Zanella A, Fermo E, Bianchi P, et al.** Red cell pyruvate kinase deficiency: molecular and clinical aspects. Br J Haematol. Jul 2005; 130(1): 11-25.