

REFERENCES

REFERENCES

- Adinolfi, M.; (1977)**
Human complement; Onset and site of synthesis during fetal life.
Am. J. Dis. Child.; 131:1015-23.
- Adler, S.M.; Denton, R.L.; (1975)**
The erythrocyte sedimentation rate in the newborn period.
J. Pediatr.; 86 (6): 942-8.
- Ainbender, E.; Cabatu, E. E.; Guzman, D.M.; Sweet, A.Y.; (1982)**
Clinical and laboratory observations: Serum C-reactive protein and problems of newborn infants.
J.Pediatr.; 101 (3): 438-40.
- Akenzua, G.I.; Hui, Y.T.; Milner, R; (1974)**
Neutrophil and band counts in the diagnosis of neonatal infections.
Pediatr.; 54:38-42.
- Avery, G.B.; (1987)**
Bacterial and viral infections of the newborn. In: Avery, G.B. (Ed.);
Neonatology, Pathophysiology and Management of the Newborn.
3 rd edition; 922-31.
JB Lippicott.
- Badr-El-Din, A.A.; Hammad, S.; Abdel-Latif, M.T.; Abdel-Aal, A.A.; Massoud, M.Z.; (1986)**
Prediction of sepsis among high-risk newborns.
The Gaz. Egypt. Paed. Ass.; 34 (1-4): 149-54.
- Bahaa El-Din, H.K.; Sound, L; Kamel, M.A.; (1991)**
Acute phase reactants in neonatal sepsis.
M.Sc. Thesis, Pediatr., Cairo University.
- Baker, C.J.; Webb, B.J.; Jackson, C.V.; (1980)**
Contercurrent immunoelectrophoresis in the evaluation of infants with group B streptococcal disease.
Pediatr.; 65:1110.

References

- Battisti, O.; Mitchison, R.; Davies, P.A.; (1981)**
Changing blood culture isolates in a referral neonatal intensive care unit.
Arch. Dis. Child.; 56:775-78.
- Becker, W.; Rapp, W.; Schwick, H.G.; Storiko, K.; (1968)**
Z. Klin. Chem.; 6:113.
- Belohradsky, B.H.; Roos, R.; Margret, W.; (1978)**
Exchange transfusion in neonatal septicemia.
Infection; 6:139-42.
- Bennet, R.; Bergdahl, S.; Eriksson, M.; Zetterstrom, R.; (1989)**
The outcome of neonatal septicemia during fifteen years.
Acta. Pediatr. Scand.; 78:40-3.
- Bienvenu, J.; Sann, L.; Bienvenu, F.; Lahet, C.; Divry, P.; Cotte, J.;
Bethenod, M.; (1981)**
Laser nephelometry of orosomucoid in serum of newborns: Reference
intervals and relation to bacterial infections.
Clin. Chem.; 27 (5): 721-6.
- Bourke, G.J.; Daly, L.E.; McGilvary, J.; (1985)**
Interpretation and uses of medical statistics.
3rd Edition; 13: 238-58.
Blackwell Scientific Publications.
- Boyle, R.J.; Chandler, B.D.; Stonestreet, B.S., Oh, W.; (1978)**
Early identification of sepsis in infants with respiratory distress.
Pediatr.; 62 (5): 744-50.
- Cairo, M.S.; Worcester, C.; Rucker, R.; Bennetts, G.A.; Amlie, R.;
Perkin, R.; Anas, N.; Hicks, D.; (1987)**
Role of circulating complement and polymorphonuclear leukocyte
transfusion in treatment and outcome in critically ill neonates with
sepsis.
J. Pediatr.; 110: 935-41.
- Cario, M.S.; Worcester, C.C.; Rucker, R.W.; Hanten, S.; Amlie,
R.N.; Sender, L.; Hicks, D.A.; (1992)**
Randomized trial of granulocyte transfusions versus intravenous immune
globulin therapy for neonatal neutropenia and sepsis.
J. Pediatr.; 120:281-5.
- Castaneda, M.R.; (1947)**
A practical method for routine blood cultures in brucellosis.
Proc. Soc. Exp. Biol. 64:114-15.

References

- Champagne, S.; Fussell, S.; Scheifele, D.; (1984)**
Evaluation of skin antiseptics prior to blood culture in neonates.
Infect. Control; 5:489-94.
- Chirico, G.; Pondini, G.; Plebani, A.; Chiara, A.; Massa, M.; Ugazio, A.G.; (1987)**
Intravenous gammaglobulin therapy for prophylaxis of infection in high-risk neonates.
J.Pediatr.; 110:437-42.
- Chiu, K.M.; Mortesen, R.F.; Osmand, A.P.; Gewurz, H.; (1977)**
Interactions of alpha-1- acid glycoprotein with the immune system.
Immunology; 32:997-1004.
- Clapp, D.W.; Kleigman, R.M.; Baley, J.E.; (1989)**
Use of intravenously administered immune globulin to prevent nosocomial sepsis in low birth weight infants: Report of a pilot study.
J. Pediatr.; 115: 973-8.
- Claus, D.R.; Osmand, A.P.; Gewurz, H.; (1976)**
Radioimmunoassay of human CRP and levels in normal sera.
J.Lab. Clin. Med.; 87 (1): 120.
- Christensen, R.D.; Rothstein, G.; (1980)**
Exhaustion of mature marrow neutrophils in neonates with sepsis.
J. Pediatr.; 96 (2): 316-18.
- Christensen, R.D.; Bradley, P.P.; Rothstein, G.; (1981)**
The leukocyte left shift in clinical and experimental neonatal sepsis.
J. Pediatr.; 98 (1): 101-5.
- Christensen, R.D.; Anstall, H.B.; Rothstein, G.; (1982)**
Review: Deficiencies in neutrophil system of newborn infants and the use of leukocyte transfusions in the treatment of neonatal sepsis.
J. Clin. Apheresis; 1:33.
- Dacie, J. V.; Lewis, S.M.; (1984)**
Basic haematological techniques. Miscellaneous tests. In: Practical Haematology.
Sixth Edition, P. 38, 39, 40, 43, 44, 420, 421.
Edinburgh; Churchill livingstone.
- Dagan, R.; Gorodischer, R.; (1984)**
Infections in hypothermic infants younger than 3 months old.
AJDC; 138:483-5.

References

Deodhor, S.D.; (1989)

CRP: The best laboratory indicator available for monitoring disease activity.

Cleve. Clin. J. Med.; 56 (2): 126-30.

Dobson, S.R.M.; Baker, C.J.; (1990)

Enterococcal sepsis in neonates: Features by age at onset and occurrence of focal infection.

Pediatr.; 85: 165-71.

Donowitz, L.G.; Marsik, F.J.; Fischer, K.A.; Wenzel, R.P.; (1981)

Contaminated breast milk: A source of klebsiella bacteremia in a newborn intensive care unit.

Rev. Infect. Dis.; 394: 716-20.

Edwards, M.S.; (1986)

Complement in neonatal infections: an overview.

Pediatr. Infect. Dis.; 5 (3): S168-S170.

El-Diwany, K.M.; El-Tayeb, S.H.; Nour, N.M.; Eissa, A.; El-Feel, M.S.; Soliman, A.; El-Ramly, Z.; El-Tohamy, F.; El-Saway, A. G.M.; (1987)

Bacteriological and clinical aspect of neonatal bacterial infection.

The Gaz. Egypt. Paed. Ass.; 35 (1-2): 37-46.

El- Hefny, A.; Nour, S.; Helal, S.; Abu- Alam, S.; (1990)

Some Immunological aspects of neonatal septicemia.

M.D. Thesis, Pediatr.; Cario University.

El- Maraghi, S.; El-Shemi, N.; El-Badawi, A.; Samra, N.S.; (1991)

Screening early septicemia in high risk newborns.

M.Sc. Thesis, Pediatr.; Cario University.

Faden, H.S.; (1976)

Early diagnosis of neonatal bacteremia by buffy-coat examination.

J. Pediatr.; 88 (6): 1032-4.

Feigin, R.D.; Abramson, S.L.; Edwards; M.S.; (1987)

Infections in the foetus and neonate. In: Fanaroff, A.A.; Martin, R.J. (Eds.); Neonatal perinatal medicine diseases of the foetus and infant.

4 th Edition; 31: 763-8.

CV Mosby Company.

References

- Fischer, G.W.; (1988)**
Therapeutic gammaglobulin for pediatric infections: Prevention or treatment of neonatal sepsis.
The Pediatric Clinics of North America: 521-4.
- Frank, M.M.; (1992)**
Detection of complement in relation to disease.
J. Allergy Clin. Immunol.; 89 (3): 641-8.
- Gahmberg, C.G.; Andersson, L.C.; (1978)**
Leukocyte surface origin of human Alpha-1-acid glycoprotein (Orosomucoid).
J. Exp. Med.; 148: 507-21.
- Gerdes, J.S.; Polin, R. A.; (1987)**
Sepsis screen in neonates with evaluation of plasma fibronectin.
Pediatr. Infect. Dis. J.; 6: 443-6.
- Gerdes, J.S.; (1991)**
Clinicopathologic approach to the diagnosis of neonatal sepsis.
Clinics in Perinatology; 18 (2): 361-81.
- Gotoff, S.P.; (1992)**
The fetus and The Neonatal Infant. In: Behrman, R.E. & Kliegman, R.M.; Nelson Textbook of Pediatrics.
14 th Edition: 501-4.
Philadelphia; W.B. Saunders Comp.
- Gotoh, H.; Ishikawa, N.; Shioiri, T.; Hattori, Y.; Nomura, H.; Ogawa, J.; (1973)**
Diagnostic significance of serum orosomucoid level in bacterial infections during neonatal period.
Acta. Pediatr. Scand.; 62: 629-32.
- Gawenlock, A.H.; McMurray, J.R.; McLauchlan, D.M.; (1988)**
Acute phase reactants. In: Varley, S.L. (Ed.); Varley's. Practical Clinical Biochemistry.
6 th Edition; 422-36.
Heinemann Medical Books, London.
- Graves, G.R.; Rhodes, P.G.; (1984)**
Tachycardia as a sign of early onset neonatal sepsis.
Pediatr. Infect. Dis. J.; 3 :404.

References

- Gruskay, J.; Harris, M.C.; Costarino, A.T.; Polin, R.A.; Baumgart, S.; (1989)**
Neonatal staphylococcal epidermis meningitis with unremarkable CSF examination results.
Am.J. Dis. Child.; 143:580-82.
- Guerina, N.G.; (1991)**
Bacterial and Fungal infections. In: Cloherty, J.P. & Stark, A.R. (Eds.); Manual of Neonatal Care.
3 rd Edition; 146-67.
Boston; Little, Brown and Company.
- Guillois, B.; Berthou, C.; Awad, H.; Bendaoud, B.; Guillemin, M.G.; Alix, D.; Youinou, P.; (1989)**
The importance of C3d estimation in the diagnosis of generalized bacterial infections in newborn infants.
Acta. Pediatr. Scand.; 78: 369-72.
- Gussetti, N.; Ruga, E.; Giaquinto, C.; Memo, L.; Casellato, R.; D'Elia, R.; (1989)**
Is C-reactive protein a reliable indicator of bacterial infection in the newborn.
J.Inf. Dis.; 160 (5): 909-10.
- Gutteberg, T.J.; Askvik, K.; Jorgensen, T.; (1986)**
Serum lactoferrin and CRP in the mother and newborn after preterm rupture of membranes.
Acta Obstet. Gynecol. Scand.; 65 (3): 203.
- Hall, R. T.; Shigeoka, A.O.; Hill, R.; (1983)**
Serum opsonic activity and peripheral neutrophil counts before and after exchange transfusion in infants with early onset group B streptococcal septicemia.
Pediatr. Infect. Dis.; 2 (5): 356-8.
- Hammad, S.; Badr-El-Din, A.A.; Abdel-Latif, M.T.; Abdel-Aal, A.A.; Massoud, B.Z.; (1986)**
Early diagnosis of neonatal septicemia.
The Gaz. Egypt. Paed. Ass.; 34 (1-4): 7-13.
- Haruta, T.; Okura, K.; Yamamoto, H.; Kuroki, S.; Yamaoko, K.; Kobayashi, Y.; (1989)**
Clinical evaluation of cefmenoxime in infections of neonates.
Jpn. J. Antibiot.; 42 (12): 2661-71.

- Hashira, S.; Nonaka, C.; Fujii, R.; (1989)**
A preclinical and clinical study of cefmenoxime in newborns.
Jpn. J. Antibiot.; 42 (12): 2607-16.
- Hawrylyshyn, P.; Bernstein, P.; Milligan, J.E.; Soldin, S.; Pollard, A.; Papsin, F.R.; (1983)**
Premature rupture of membranes: The role of CRP in the prediction of chorioamnionitis.
Am. J. Obst. Gyn.; 147 (3): 240.
- Hill, H.R.; Shigeoka, A.O.; Augustine, N.H.; (1984)**
Fibronectin enhances the opsonic and protective activity of monoclonal and polyclonal antibody against group B streptococci.
J. Exp. Med.; 159: 1618.
- Hindocha, P.; Compnell, C.A.; Gould, J.D.M.; Wojciechowski, A.; Wood, C.B.S.; (1984)**
Serial study of C-reactive protein in neonatal septicemia.
Arch. Dis. Child.; 59: 435-8.
- Isaacman, S.H.; Heroman, W.M.; Lightsey, A.L.; (1984)**
Purpura fulminans following late onset group B beta haemolytic streptococcal sepsis.
Am. J. Dis. Child.; 138:915.
- Kelsy, M.C.; Lipscomb, A.P.; Mowles, J.M.; (1982)**
Limulus amoebocyte lysate endotoxin test: An aid to the diagnosis in the septic neonate.
J.Infect.; 4: 69.
- Khalifa, A.M.; Allam, M.; Nawar, F.; Sabry, S.I.; Hemeds, M.S.; (1989)**
Quantitative and qualitative changes in neonatal bacterial infection.
The Gaz. Egypy. Paed. Ass.; 37 (1-2): 1-5.
- Khashaba, A.A.; El Saadany, A.M.; Aly, A.I.; Abdel Meguid, A.R.; (1988)**
Study of some acute phase reactants in the newborn infant.
M.D. Thesis, Pediatr.; Benha University.
- Kim, K.S.; (1989)**
Use of intravenous immunoglobulin in the treatment of neonatal sepsis.
AJDC; 143: 1257-8.

References

- Kite, P.; Millar, M.R.; Gorham, P.; Congdon, P.; (1988)**
Comparison of five tests used in diagnosis of neonatal bacterpemia.
Arch. Dis. Child.; 63:639-43.
- Kite, P.; Langdal, V.; Todd, N.; Millar, M.P.; Mackay, P.; (1989)**
Direct isolation of coagulase negative staphylococci from neonatal blood samples.
J. Hosp. Infect.; 14: 135-40.
- Kite, P.; Todd, N.J.; Mackay, P.; Millar, M.; (1990)**
Plasma neutrophil elastase alpha-1-proteinase inhibitor levels in premature neonates with coagulase-negative staphylococcal bacteremia.
J. Pediatr.; 117 (4): 630-1.
- Kleiman, M.B.; Reynolds, J.K.; Schreiner, R.L.; Smith, T.W.; Allen, S.D.; (1984)**
Rapid diagnosis of neonatal bacteremia with acridine orange stained buffy coat smears.
J. Pediatr.; 105 (3): 419-21.
- Klein, J.O.; Dashafesky, B.; Norken, C.R.; Mayer; (1983)**
Selection of antimicrobial agents for treatment of neonatal sepsis.
Rev. Infect. Dis; 5: 55.
- Klerk, E.; Anderson, R.; (1982)**
Accurate laboratory measurement of serum CRP.
S.A. Med. J. (Abstact); 62: 1030-2.
- Laurenti, F.; Ferro, R.; Isacchi, G., Panero, A.; Savignoni, P.G.; Malagnino, F.; Palermo, D.; Mandelli, F.; Bucci, G.; (1981)**
Polymorphonuclear leukocyte transfusion for the treatment of sepsis in the newbron infant.
J. Pediatr.; 98 (1): 118-23.
- Liu, C.H.; Lehan, C.; Speer, M.E., Smith, E.B.; Gutgesell, M.E.; Fernbach, D.J.; Rudolph, A.J.; (1985)**
Early detection of bacteremia in an outpateint clinic.
Pediatr.; 75: 827-31.

References

- McCracken, G.H.; Shinefield, H.R.; (1966)**
Changes in the pattern of neonatal septicemia and meningitis.
Am. J. Dis. Child.; 112: 33-9.
- McCracken, G.H.; Freij, B.J.; (1987)**
Perinatal bacterial diseases. In: Feigin, R.D.; Cherry, J.D. (Eds.);
Textbook of Pediatric Infections Diseases.
2nd Edition; 25: 940-66.
W.B. Saunders Co.
- Mcintoch, K.; (1984)**
Bacterial infections of the newborn in Schaffer's diseases of the
newborn.
5th ed. P. 732, 734, 740, 751, 757.
Philadelphia, W.B. Saunders Company.
- Macleod, C.M.; Avery, O.T.; (1950)**
Detection of CRP in human serum.
J. Exp. Med.; 73:183.
- Manroe, B.L.; Rosenfeld, C.R.; Weinberg, A.G.; Browne, R.;
(1977)**
The differential leukocyte count in the assessment and outcome of
early-onset neonatal group B streptococcal disease.
J. Pediatr.; 91 (4): 632-7.
- Manroe, B.L.; Weinberg, A.G.; Rosenfeld, C.R.; Browne, R.;
(1979)**
The neonatal blood count in health and disease: Reference values for
neutrophilic cells.
J Pediatr.; 95 (1): 89-98.
- Martin, M.A.; Pfaller, M.A.; Massanori, R.M.; Wenzel, R.P.;
(1989)**
Use of cellular hydrophobicity, Slime production and species
identification markers for the clinical significance of coagulase.
negative staphylococcal isolates.
Am. J. Infect. Control.; 17: 130-5.
- Mathers, N.J.; Pohlandt, F.; (1987)**
Diagnosis audit of CRP in neonatal infections.
Eur. J. Pediatr.; 146 (2): 147.

References

- Melish, M.E.; Glasgow, L.A.; Turner, M.D.; (1972)**
The staphylococcal scalded - skin syndrome: Isolation and partial characterization of the exfoliative toxin.
J. Infect. Dis.; 125:129.
- Millar, M.R.; Tadd, N; Mackay, P.; (1990)**
Neonatal infections with coagulase negative staphylococci.
Arch. Dis. Child.; 63 (10): 1105-6.
- Moffet, H.L.; (1989)**
Perinatal syndromes, neonatal sepsis and meningitis. In: Pediatric infectious Diseases.
3 rd Edition: 496-501.
Philadelphia, J.B. Lippincott Company.
- Mohy El Din, O.; Badrawy, N.; Seound, E.; El Saify, A.; Choucri, C.S.; (1982)**
Laboratory evaluation of the indirect techniques in the diagnosis of neonatal infection.
M.Sc. Thesis. Clin. Pathol.; Cairo University.
- Oliveria, E.B.; Gostchlich, E.C.; Liu, T.Y.; (1977)**
Primary structure of human CRP.
Proc. Nat. Acad. Sci. USA; 74:3148.
- Payne, N.R.; Burke, B. A.; Day, D.L.; Christenson, P.D.; Thompson, T.R.; Ferrieri, P.; (1988)**
Correlation of clinical pathologic findings in early onset neonatal group B streptococcal infection with disease severity and prediction of outcome.
Pediatr. Infect. Dis. J.; 7 (12):L 836-47.
- Peakman, M.; Senaldi, G.; Liossis, G.; Gamsu, H.R.; Vergani, D.; (1992)**
Complement activation in neonatal infection.
Arch. Dis. Child.; 67: 802-7.
- Pelet, B.; (1980)**
C3, factor B, alpha-1-antitrypsin in neonatal septicemia with sclerema.
Arch. Dis. Child.; 55: 782-8.

- Peltola, H.O.; (1982)*
CRP for rapid monitoring of infections of the central nervous system.
Lancet; i: 980.
- Pepys, M.B.; (1981)*
C-Reactive Protein, Fifty Years On.
Lancet i: 653-6.
- Philip, A.G.S.; (1979)*
The protective effect of acute phase reactants in neonatal sepsis.
Acta Pediatr, Scand.; 68: 481-3.
- Philip, A.G.S.; Hewitt, J.R.; (1980)*
Early diagnosis of neonatal sepsis.
Pediatr.; 65 (5): 1036-41.
- Philip, A.G.S.; (1982)*
Detection of neonatal sepsis of late onset.
JAMA; 247 (4): 480-92.
- Philip, A.G.S.; (1984)*
Acute-Phase proteins in neonatal infection.
J.Pediatr.; 105 (6): 940-42.
- Philip, A.G.S.; (1985)*
Response of C- reactive protein in neonatal group B streptococcal infection.
Pediatr. Infect. Dis.; 4 : 145-8.
- Philip, A.G.S.; (1987)*
Neonatal infection. In: Neonatology, A practical guide.
3 rd Edition: 190-5.
Philadelphia, W.B. Saunders company.
- Pierce, J.R.; Merenstein, G.B.; Stocker, J.T.; (1984)*
Immediate post-partum cultures in an intensive care nursery.
Ped. Inf. Dis.; 3: 501-3.
- Pipkin, F.B.; (1984)*
Medical statistics made Easy.
First Edition: 112-121.
Edinburgh; Churchill Livingstone.

References

- Placzek, M.M.; Whitelaw, A.; (1983)**
Early and late neonatal septicemia.
Arch. Dis. Child.; 58: 728-31.
- Polin, R. A.; Geme, III, J.W.St.; (1992)**
Neonatal Sepsis. In: Aronoff; Hughes; Kohl; Speck; Wald; Advances
in Pediatric Infectious diseases.
Vol. 7: 25-61.
St. Louis; Mosby- Year Book. Inc.
- Pommier, C.G.; O'Shea, J.; Chused, T.; (1984)**
Studies on the fibronectin receptors of human peripheral blood
leukocytes.
J. Exp. Med.; 159: 137.
- Pourcyrus, M.; Bada, H.S.; Korones, S.B.; Barrett, F.F.;
Jennings, W.; Lockey, T.; (1991)**
Acute phase reactants in neonatal bacterial infection.
J. Perinatol. (Abstract); 11 (4): 319-25.
- Ralph, D.; Stuart, L.; Morven, S.; (1987)**
Infection in the fetus and neonate in neonatal perinatal medicine.
4th ed. P. 765-94.
C.V. Mosby Co. Louis,
- Reed, R.K.; Larter, W.E.; Sieber, D.F.; (1976)**
Peripheral nodular lesions in pseudomonas sepsis: The importance of
incision and drainage.
J. Pediatr.; 88:97.
- Rodwell, R.I.; Leslie, A.L.; Tudehope, D.I.; (1988)**
Early diagnosis of neonatal sepsis using a hematologic scoring
system.
J. Pediatr.; 112 (5): 761-7.
- Russell, G.A.B.; Smyth, A.; Cooke, R. W.I.; (1992)**
Receiver operating characteristic curves for comparison of serial
neutrophil band forms and C-reactive protein in neonates at risk of
infection.
Arch. Dis. Child.; 67: 808-12.

References

- Sann, L.; Bienvenu, F.; Bienvenu, J.; Bourgeois, J.; Bethenod, M.; (1984)**
Evaluation of serum prealbumin, C-reactive protein, and orosomucoid in neonates with bacterial infection.
J. Pediatr.; 105: 977-81.
- Saxtad, J.; Nilsson, L.A.; Hanson, L.A.; (1970)**
C-reactive protein in serum from infants as determined with immunodiffusion techniques.
Acta. Pediatr. Scand.; 59: 25-27.
- Scheifele, D.W.; Melton, P.; Whitchelo, V.; (1981)**
Evaluation of the limulus test for endotoxaemia in neonates with suspected sepsis.
J. Pediatr.; 98:899.
- Shatla, H.; Baghat, N.; Yossef, S.; Abo El Maged, L.; Sabry, W.; (1988)**
Diagnostic value of alpha-1-acid glycoprotein and fibrinogen in neonatal bacterial infections.
Egypt. J. Pediatr.; 5 (1-2): 131-8.
- Shigeoka, A.O.; Hall, R.T.; Hill, H.R.; (1978)**
Blood transfusion in group B streptococcal sepsis.
Lancet; 1: 635.
- Shine, B.; Gould, J.; Campbell, C.; Hindocha, P.; Wilmot, R.P.; Wood, C.B.; (1985)**
Serum CRP in normal and infected neonates.
Clin. Chim. Acta.; 148 (2): 97-103.
- Singer, J.M.; Platz, C.M.; Bader, E.; Elster, S.K.; (1957)**
The latex fixation test: Agglutination test for CRP and comparison with the capillary precipitin method.
Am. J. Clin. Pathol.; 28: 611-17.
- Spath, P.; Gabl, F.; (1976)**
Critical role of the conversion of the third complement component C3 for its immunochemical quantitation.
Clin. Chim. Acta; 73 (1): 171-5.
- Speer, Ch.; Bruns, A. ; Gahr, M.; (1983)**
Sequential determination of CRP, alpha-1-antitrypsin and haptoglobin in neonatal septicemia.
Acta. Pediatr. Scand.; 72: 679-83.

References

- Speer, C.P.; Ninjo, A.; Cari, M.; (1986)**
Elastase- alpha-1-proteinase inhibitor in early diagnosis of neonatal septicemia.
J. Pediatr.; 108:987.
- Tillet, W.S.; Francis T.Jr.; (1930)**
Serological reactions in pneumonia with a non protein fraction.
J.Exp. Med.; 52:561-71.
- Vain, N.E.; Mazlumian, J.R.; Swarner, O.W.; Cha, C.C.; (1980)**
Role of exchange transfusion in the treatment of sever septicemia.
Pediatr.; 66 (5): 693-7.
- Volkanakis, J.E.; Clements, W.; Schrohenloher, R.; (1978)**
CRP purification by affinity chromatography and physical and chemical characterization.
J. Immunol. Methods; 23:285-95.
- Varley, H.; (1980)**
The plasma proteins. In: Varley, Harold: Practical clinical chemistry.
5 th Edition; 590-91.
William Heinemann Medical Books, London.
- Vawter, G.F.; (1981)**
Perinatal listeriosis. In: Rosenberg, H.S.; Bern stein, J. (Eds.);
Prespectives in pediatric pathology.
2 nd Edition; 6: 153.
Masson Publishing, USA Inc.
- Visintine, A.M.; Oleske, J.M.; Nahmias, A.J.; (1977)**
Listeria monocytogenes infection in infants and children.
Am.J. Dis. Child.; 131: 393.
- Voora, S.; Srinivasan, G.; Lilien, L.D.; Yeh.; T.F.; Pildes, R.S.; (1982)**
Fever in full-term newborns in the first four days of life.
Pediatr.; 69:40-44.

References

- Weisman, L.E.; Stoll, B.J.; Kueser, T.J.; Rubio, T.T.; Frank, C.G.; Heiman, H.S.; Subramanian, K.N.; Hankins, C.T.; Anthory, B. F.; Cruses, D.F.; Hemming, V.G.; Fischer, G.W.; (1992)**
Intravenous immune globulin therapy for early-onset sepsis in premature neonates.
J. Pediatr.; 121 (3): 434-43.
- Whicher, J.T.; Martin, J.R.; Dieppe, P.A.; (1980)**
Prostaglandin stimulated increase in acute phase proteins in man and its failure in systemic sclerosis.
Lancet; ii: 1187.
- Whitelaw, A.; (1990)**
Treatment of sepsis with IgG in very low birthweight infants.
Arch. Dis. Child.; 65: 347-8.
- Yeyng, C.Y.; Lee, V.W.; Yeung, M.B.; (1973)**
Glucose disappearance rate in neonatal infection.
J. Pediatr.; 82:486.
- Yoder, M.C.; Douglas, S.D.; Gerdes, J.; (1983)**
Plasma fibronectin in healthy newborn infants, respiratory distress syndrome and perinatal asphyxia.
J.Pediatr.; 102 (5): 777-9.
- Yoder, M.C.; Polin, R.A.; (1986)**
Immunotherapy of neonatal septicemia. In: Polin, R.A.; Fox, W.W. (Eds.); The Pediatric Clinics of North America: The newborn II; 33 (3): 481-501.
Philadelphia: Saunders.
- Zipursky, A.; Jaber, H.; (1978)**
The haematology of bacterial infection in newborn infants.
Clin. Haematol.; 7:175.