

References

- Abdel - Aziz, D., (1977) :** Fevers and infectious diseases . El- Naser modern bookshop 2nd edition .
- Abramson, J.S. , Hampton, KD, Bahus Wasilukas, B.L. ,and Macron, M.J , (1985) :** The use of c.reactive protien of CSF for differentiation of miningitis from other CNS diseases . J. infect . Dis . 151 : 5 .
- Aggarwall, B.B , Hanzel, W.J., Moffat Kohr, W.J., and Harkins, R.N. (1985):** Primary structure of human lymphotoxin drived from 1788 lympho - blastiod cell line. J. Biol . Chem. vol, (260) p. 2334 - 2344.
- Aggarwall, B.B. , and Vilcek , eds (1992) :** Tumor necrosis factor, structure, function and mechanism of action Marcel Dekker press, New York, p. 1 - 600
- Anders Waage, petter Brandtzaeg, Alfred Halstensen, petter Kierulf and Terge Espevik (1989) :** Complex protein of cytokines.of patients with meningococcal septic shock. J.Exp. Med.. 169 jan 333 - 338 .
- Annual Reports of Scottish Mycobacterial Reference laboratory, Mearnskirk Hospital, Glesgow , 1970 - 1975 .**
- Arditi M, Kirk, R., Manogue Michael Caplan, and Ram Yagev. (1990) :** Cerbrospinal fluid tumor necrosis factor alpha and platlet activating factor concentrations and severity of bacterial meningitis in children. J of Infect Dis . Vol, (162) : 139 -147 .
- Balan, G., and Branza, M. (1985) :** Acute bacterial meningitis in children and adults . Med .Clin North Am; 219 - 231 .

Balk, R.A. (1987) : A controlled clinical trial of high dose of methylprednisolone in the treatment of severe sepsis and septic shock . New Eng J.Med. Vol, (317) : 653 - 658 .

Balzer, S. , Berant M. , Alon, U. (1983) : Effects of antibiotic treatment on cerebrospinal fluid, Am J. Clin . Path. Vol, (80) : 380 - 389 .

Band J.D, Fraser D.W, Ajella G, *etal* (1984) :Prevention of H.influenzae type b disease JAMA 251 : 2381 .

Bell, W.E. , (1982) : Bacterial infections of the nervous system in practice of pediatric neurology by swaiman and wright 2nd ed P. 659 . C.V Mosby company st. Lois. Toronto, London .

Beutler, B., Cerami, A. , Figorri, L.S., Palladine, M.A., and O'connor, J.V.(1986) : TNF- α is an endogenous pyrogen and induce production of IL-1- , J. Exp. Med. Vol, (163): P. 1433 - 1450 .

Beutler, B., Cerami, A. (1987) : Cachectin more than a tumor necrosis factor, New Eng J. Med. Vol , (316) : 379 - 385 .

Beutler, B., Cerami , A. (1988) : Cachectin, cachexia, and shock. AJNR, Vol,(39) : 75 - 83 .

Bihari, D.J. , Tinker, J. (1988) : The therapeutic value of vasodilator prostaglandin in multiple organ failure associated with sepsis. Intensive care Med, 15 : 3 - 7 .

Bjarne Moller ., Soren, C., Mogensen, *etal*; (1991) : Bioactive and inactive forms of TNF- α in spinal fluid from patients with meningitis. J.infect . Dis. Vol, (163) : 886 - 89 .

Blackwall, C.C. , Weir DM., James, V.S., Cartwright K.A.V., Stuart, J.M., Jones, D.M. (1989) : Secretor status and carriage of *Nisseria meningitidis* . *Epidem . infect.* Vol, (102) : 1 - 10 .

Bone , R.C, Fisher C.J, Clemmer T.P, *etal* ; (1987) :A controlled trial of high dose methylprednisolone in the treatment of septic shock . *New England J. Med* , 317 : 653 - 658 .

Brandtzaeg, P. , Kierulf , Gaustad, P. , Skulberg, A., Halvorsen, S. (1989): Plasma endotoxin as a predictor of multiple organ failure and death in septic meningitis . *J. infect Dis.* Vol, (159) : 195 - 204 .

Brandtzaeg, P. , Mollnes, T.E. , Keirulf, P. (1989) : Complement activation of endotoxin level in septic meningitis, *J. infect. Dis.* 160 : 58- 65.

Brandtzaeg, P., Brusletto B, Kierulf, P. (1990) : Plasminogen activator inhibitor 1 and 2 , Alpha - 2 - antipiasmin and endotoxin levels in systemic meningococcal disease *Thomb, Res*, 57 : 271 - 278 .

Brook, I. , Bricknell, K.S. , Overturt, G.D., Finegold, S.M., (1978) : Measurement of lactic acid in C.S.F of patients with infections of central nervous system. *J. infect. .Dis.* Vol , (137) : 384 - 90 .

Broome C.V, (1981) : Pneeummococcal polysaccharide vaccines .*Rev. Infect. Dis.*3 (Supl) : 582 .

Charles, A. Dinarello (1991) : The proinflammatory cytokines Interleukin - I - and Tumor necrosis factor and treatment of septic shock syndrome . *J. infect. Dis.* Vol, (163) : 1177 - 1184 .

Cherry, J.D. (1981) : Text book of pediatric infectious diseases
Philadelphia, W.B. , Saunders : page 1316 .

Chouaib, S., Marchiol, C., and Fandelizi, D. (1987) : Inhibition of human
Tumor necrosis alpha by adenylate cyclase activators . Immunobiol. Vol, (125)
no 1/2 P. 24 .

Corral, C.J. Peple, J.M., Moxon, E.R., and Hughes, W.T (1981) : C-
reactive protein in spinal fluid of children with meningitis. J. of pediatr. Vol ,
99 (3) : 365 - 369 .

Crokson, R.A., Payne, G.J., and Ratchiff, A.P. (1986) : Time sequence of
acute phase reactive proteins following surgical trauma. Clin.Chem.. Acta. Vol,
(14) : 435 - 441 .

Curtis, G.D.W., Slack, M.P.E.,and Rompkins, D.S. (1981) : Cerebrospinal
fluid lactate and the diagnosis of meningitis . J. Inf . Dis . Vol ,(3): 159 - 195 .

Diamond, R.D., and Bennett, J.E. (1974) : Ann .Intern.Med. Vol , (80) :
p. 176 .

Dinarello, C.A., Cannon, J.G., Walff, S.M., Bernheim, H.A., Beutler,B.,
Cerami, A., and Palladino, M.A. (1986) : Tumor necrosis factor is an
endogenous pyrogen and induces production of interleukin - I - J. Exp. Med.
Vol, (163) : 1433 - 1450 .

Dinarello, C.A. (1989) : Interleukin -I- and its biologically related cytokines.
Adv. Immunol . Vol, (44) : 153 - 205 .

El- Batawi, Y.A. (1981) : Manual of microbiol. infections of man . 4th
edition, part III. P.15 .

Feigin, R.D. and Dodge, P.R.(1970) : Pediatr. Clin. North Am..Vol, (23):
541

Feign, R.D (1983) : Acute bacterial meningitis beyond the neonatal period
In : Nelsson text book of pediatrics, Saundres company 12th edition. P. 619 -
624

Figueroa, J.E., Densen, P. (1991) : Infectious diseases associated with
complement deficiencies . clin. microbiol. Rev, (4) : 359 - 395 .

Fischers, C.L., and Gill, O.W. (1977) : Acute phase proteins, serum proein
abonormalities. Ed. by Ritzmannsf., Danniels, J.G., Boston little and Blown, P.
331 - 350 .

Fischl, M *etal*; (1987) : The efficacy of azidothymidine in the treatment of
patients with AIDS. N.Eng J. Med. Vol, (317) : p 185 .

Flagestad, T., Guttegerg, T., Kristiansen, B.E (1990) : Antibodies to
meningococci in children with meningococcal disease. Scand .J.infect . Dis. Vol,
(22) : 547 - 551 .

Fraser, W.E., darby., C.P ., and Koehler, R.E (1983) : Risk factors in
bacterial meningitis : charleston, country, south Coloformia . J.Infect . Dis. Vol
(127) : 271 - 373 .

Frash, C.E(1989) : Vaccines for prevention of meninococcal diseases . Clin.
microbiol. Rev, 2: 5134

Fryden, A., Link, H., and Norrby, E. (1978) : Cerbrospinal and serum
immunoglobulins and antibody titre in mumbs meningitis and aseptic
meningitis of other aetiology. Infection and Immunity. Vol ,(21) : 852 - 61 .

Gamble, J.R., Harlan, J.M., Klebanoff, S.J., Vadas, M.A.(1985) :
Stimulation of the adherence of neutrophils to umbilical vein endothelium by human recombinant tumor necrosis factor. Proc. Natl. Acad : USA. (82) : 8667 - 8671 .

Gerard, P., Moriau, M., Bachy, A., Malavaux, P., Demyer, R. (1983) :
Meningococcal purpura : Report of 19 patients treated with heparin. J. ped . Vol , (82) : 780 - 780 .

Glauser, M., Zanett, G., Bamgartner, J.D., Cohen, J.(1991) : Septic shock pathogenesis . Lancet 338 : 732 - 736 .

Gowenlock, A.H., Murray, J.R., and Mc . Lauchlan, D.M.(1988) : Normal cerebrospinal fluid in varleys practical clinical Biochemistry 6th ed pp, 443 - 51 .

Greene, G.R. (1978) : Meningitis due to H.influenzae other than type b: case report & review, pediatrics. Vol , (62) : 1021 .

Gurura J. V.J., Russo, R.M., Allen, J.E., *etal*; (1973) : What are the best indication for performing a lumbar puncture in an out patient child ? Clin . ped. Vol, (12) : 488 .

Guttberg, T.J., Flagestad, T., Jorgensen, T. (1986) : Lactoferrin, c-reactive protein, Alpha -1- antitrypsin and immunoglobulin -A- in cerebrospinal fluid in meningitis. Acta . Pediat scandinavica. Vol, (75) : 596 - 572 .

Hanshaw J.B, Dudgeon J.A, Marshall W.C (1985) : Viral diseases of fetus and newborn . Philadelphia , W.B. Saunders .

Harrison's *etal*;(1992) : Principle of internal medicine 12th edition . Vol,(1) :p. 493- 498 .

Heicappell, R., Saukurai, A., Haranaka, R., and Haranaka, K. (1987):
The mechanism of antitumor activity of TNF regarding factors influencing
antitumor activity and side effects immunobiol. Vol, (175) : No $\frac{1}{2}$ p. 31.

Hellman, S.(1990) : Principles of radiation therapy in cancer. Edited by
Devita, V.T., Hellman, S, et al ; 3rd edition . J.B. Lippincott company.
Philadelphia. chapter 15, p. 247 - 274 .

Heyderman, R.S., Klein, N. J., Shennan, J.I, and Levin, M. (1991) :
Deficiency of prostacyclin production in meningococcal shock. Arch. Dis
child. Vol, (66) : 1296 - 1299 .

Hoeprich , P.D(1983) : Infectious diseases . 3 rd edition . Harres and Raw
publishers . Philadelphia , p; 171 .

**Hoffman S.L., Punjabi, N.H., Kumala, S., Moechtar, M.A., Pulunish,
S.P., and Rivai, A.R.(1984) :** Reduction in chloramphenicol - treated severe
typhoid fever by high dose dexametasone. N.Eng J. Med.Vol, (310) : 82 - 88 .

Hudson, L., and Hay, F.C. (1989) : Immunoassay in practical immunology.
Edited by Hudson, L., and Hay, F.C., 3rd edition, Blackwell scientific
publication Oxford, London .

Jawetz, E., Melnick, L.J. and Adel berg A.E.(1982) : Review of medical
microbiology 17th edition Loss Altos, California. p, 230 .

Jones, D.M., Sutcliffe, E.M *etal* .,(1990) : meningococci with reduced
susceptibility to penicillin . Lancet 325 : 863 - 864 .

Kawakami, M., Ikeda, Y., Letrang, N., Vine, W., and Cerami, A. (1984) : Studies of conditions and agents that stimulate and inhibit the production of cachectin by macrophages . Proc. Int union pharma, 377 .

Keheri, J., Miller, A., and Fanc., A. (1987) : Effects of tumor necrosis factor - α on mitogen activated human .B. cells, J.Exp Med, Vol (166) : 786.

Klein, N.J., Heyderman, R.S., Levin, M. (1992) : Antibiotic choices for meningitis beyond the neonatal peroid. Arch Dis child. Vol, (67): 157 - 161 .

Knight, J.A. (1981) : Early chemical diagnosis of bacterial meningitis. C.S.F glucose, lactate, and lactate dehydrogenase . Clin . chemist. Vol, (27): 1431 - 1434 .

Kunknel, S.L., Larrick, J.W.,m ans Remick, D.G. (1987) : Prostaglandin E2 and prostacyclin regulate TNF production at the cellular and molecular levels . immunobiol. Vol ,(175) :No $1\frac{1}{2}$, p. 80

Landaas, S., etal., (1985) : Chemical analysis for early differential diagnosis between bacterial and viral meningitis .Scand cluin lab Invest.Vol, (45) : 525 - 529 .

Le . T., Weistein, D., Gubler, U., and Vilcek, J.(1987) : Induction of membrane associted IL-1 by TNF - α in human fibroblast . J. Immunol. Vol, (1381).P. 2137 - 2142 .

Leist, T.P., Feri, K., Kam Hansen, Zinkernagel, R.M. Fontana, A. (1988): Tumor necrosis factor alpha in CSF during bacterial meningitis. Evaluation in murine model infections and in patients . J. Exp. Medicine. Vol , (167) : 1443 - 1448 .

Lenette, E.H., Spaulding, H., and Traut, J.P.(1974) : Manual of clinical microbiology. 1st Ed. American society for microbiology.p 323 .

Levin M.J, (1994) : Treatment and prevention option of respiratory syncytial virus infection . J.pediatr. Vol,(124) : 5 part 2 : 522- 527 .

Lopez - Cortes, L.F., Cruz Ruiz , M., Gomez Mateos , J., Jimenes Hernandez D., palamino, J., and Jimenez , E. (1993) : Measurment of level of TNF - α and IL - 1 β in CSF of patients with meningitis of different etiologies. Clin Infect Dis.Vol, 16 (4) : 534 - 539 .

Masahiro Higuchi and Bharat B. Aggarwal (1994) :Cytokine research laboratory , Department of clinical immunology and biological therapy , University of Texas M.D. Anderson cancer center , Huston , Texas 77030 .

McCracken, GH, Jr., Mustafa, M.M., Ramilo, O.,and Risser, R.C. (1989): CSF TNF - α and IL- 1 β and outcome from neonatal Gram negative enteric bacillary meningitis. Ped infect Dis . J. Vol, (8): 155-159 .

Meager, A (1990) : Mediators of immunoresponse and clinical appilication of cytokines, Ed. by Meager , A. First published British library, Buckingham, p.18 .

Mercier, J.C., Beaufils, F., Hartman , J.F., Azema, D.(1988) : Hemodynamic patteren of meningococcal shock in children Crit care Med Vol, (16) : 27 - 33 .

Middle kamp, J.N. (1981) : In feigin , R.D., cherry, J.D. (eds) : text book of pediatric infectious diseases philladelphia, W.B. Saunders ,. 319 .

Moller, B., Mogensen, S.C., Wendelboe, P., Bendtzen, K. and petresen, C.M. (1991) : Bioactive and inactive forms of TNF - α in spinal fluid from patients with meningitis. j. Infect. Dis. Vol,(163) : 886 - 889 .

Monto, A.S (1994) :Using antiviral agents to control outbreaks of influenzae Q infection . Geriatrics .Vol,(49) : 12 : 30 : 34 .

Moore, P.S., Heirholzer, J., Dewitt, W., Gouant, K., Lippeveld, T., plikaytis, B., and Broome , C.V. (1990) : Recesspiratory viruses and mycoplasma as cofactors for epidemic group A meningococcal meningitis. JAMA, (264) : 1271- 1275 .

Morquet, R.L., De Bruin, R.W., Feirs, W., and Jeekel, J. (1987) : Indomethacin reduce TNF - α toxicity without interfering with its efficacy. Immunobiology . Vol. 175 , N $\frac{1}{2}$, p. 54 .

Movat, H.Z (1987) : TNF - α IL-1 role in acute inflammation and microvascular injury. J.Lab clin. Med. Vol, (110) : 668 - 681 .

Mustafa, M.M., Ramilo, O., Mertsola, J., Resser, R.C., Beuttler, B., Hansen, E.J., McCracken, GH, Jr. (1989) : Modulation of inflammation and cachectin activity in relation to treatment of experimental Haemophilus infleunzae type b. meningitis. J. infect . Dis . Vol, (107) : 818 - 825 .

Nawroth, P.P ., Bank, L., Handly, D., Cassimeris .J., Chess, L., and stern , D. (1986) : Tumor necrosis factor interaction with endothelial cell receptors to induce release of interleukin - I . J. Exp . Med. Vol , (163) : 1363 - 1375 .

Nawroth. (1986) : Interferons and other regulatory cytokines. Edited by Edward , M.D., Guignard, J., and curie, I.A. Interscience and sons. p. 290 - 310

Neale, M.L., Pilli, R. A., and Matthews, N. (1987) : Involvement of arachidonate metabolism in the cytologic effects of TNF - α on tumor cells in vitro Immunobiol . Vol , (175) , NO 1/2 , p.84 .

Norden, C.W., Michaels, R.H., and Melish, M. (1976) : J. Infect. Dis. Res . : 103 .

Nye, J.F. (1983) : The value of initial lab investigations in the management of meningitis. J. Infect Dis. Vol, (7) : 31 - 38 .

Odio, C.M., faingezicht, I., paris, M., Baltodano , A., Regers, J., Saez loren, x., Oslen K, and McCracken, J. (1991) : The beneficial effects of early dexamethazone therapy with bacterial meningitis New. Eng. J. Med Vol, (324) : 1531 - 1535 .

Ohga S, Tomnobi Aoki, Okada, Hideki Akeda, Katsuyoshi: Fujioka, Akio Ohashima, Toshinori; Mori, Ikuko Minamishima, and Kohjiueda. (1994) : Cerebrospinal fluid concentration of TNF - α and IL- 1 β and interferon γ in bacterial meningitis . Arch. Dis. child Vol , (70) : 123 - 125.

Okusawa, S., Gelfand, J.A., Ikjiama, T., Conolly, R.J., and Dinarello, C.A. (1988) : Interleukin -1- induce a shock like state in rabbits . Synergism with tumor necrosis factor and the effect of cyclooxygenase inhibition. J.clin. Invest . Vol, (81) : 1162 - 1172 .

Old, L.J. (1987) : Tumor necrosis factor : pleiotropic action and interactions. Immunobiol, Vol, (175) No 1/2 , p.24 .

Paton , J.C *etal* (1986) : Antibody response to pneumococcal vaccine in children aged from 5 - 15 years . Am. J. Dis. child .140, p.135

Pepple, J., Moxon, E.R., and Yolken, R.H.(1980) : J. Pediatr.Vol,(97)
p. 233

Ramilo, O., Saez- Lorenz, X., Mertsola, J., *etal* ; (1990) : Tumor necrosis factor alpha and interleukin - I- β initiate meningeal inflammation. J.Exp. Med. Vol, (172): 497 - 507 .

Satomi, N., saukurai, A., Haranaka, R., and Haranaka, K.(1987) : The mechanism of antitumor activity of TNF - α regarding factors influencing antitumor activity and side effects . Immunobiol. Vol, (175). No 1/2 p. 31 .

Schaad, V.B., Nelson, J.D., McCracken, GH. Jr. (1981) : Relapse in bacterial meningitis of childhood. Pediatrics , Vol (87) : 188 - 195 .

Schlech, W.F. III ., Ward , J.I., Band, J.D., Hight Ower A., and Broome, C.V. (1985) : Bacterial meningitis in USA through 1981 . The neonatal bacterial meningitis surveillance study JAMA, Vol,(253) p. 1749 - 1754.

Schwartz, B., Al. tobaig, A., Al- Ruwais, A., Fontaine , R.E., Ashi, J.A., Hightower, A.W., Broom, C.V., and Music, S.I (1988) : Comparative efficacy of ceftriaxone and rifampicin in eradicating pharyngeal carriage of group A. *Nisseria meningitidis*. Lancet i : 1239 - 1243 .

Selt, S.H.W., Merrill, R.E , and Deyne , E.D, (1972) : Pediatrics. vol , 49
P.206 .

Shackel Ford , P.G; Campell , J. and Feigin R.D, (1974) : J. pediatrics 85 : 478 .

Shanson, D.C.(1982): Microbiology in clinical practice. wright PSG. Bristol. London. 167.

Sipple, J.E., Hider, P.A., and Centroni, G.(1984) : J. Clinical Microbiol. 215 : 884 .

Smith, A.L. (1975) : Tuberculous meningitis in childhood Med. J. Aust 1 : 57 : 60 .

Smith, E.W.P, and Haynes, R.E.(1982): Changing incidence of haemophilus influenzae meningitis. Pediatrics. Vol, (50): 723 - 725 .

Spear, C.H., Bruns, A., and Gahrn. (1983) : Sequential determination of C- reactive protein, alpha -1- antitrypsin and haptoglobulins in neonatal septicaemia . Acta . ped . Scand. Vol,(72) : 679- 683 .

Spratt, B.G., Zhang, Q., Jones, D.M., Hutchison, A., Brannigan, J.A., and Dowson, G.G.(1989) : Recruitment of penicillin binding protein gene from *N. flavescens* during the emerging of penicillin resistance in *N. meningitidis*, Proc. Natl. Acad Sc: , USA, Vol,(86) : 9899- 9982 .

Sprigges, D.R.(1991) : TNF - α basic principles and pre- clinical studies . In basic therapy of cancer. Edited by Devita, D.T., Hellman, S., and Rossenberg, S.A. chapter 16 p . 354 - 377 , L.B., Lippincot company, Philadelphia, New York, London - Hagerstown .

Stephens D.S.,and Farley, M.M. (1991) : Pathogenic events during infection by *N. meningitidis* and *H. Influenzae* . Rev. Infect. Dis. Vol, (12) : 22- 33 .

Sun, X., Hsueh, W. (1988) : Bowel necrosis induced by tumor necrosis factor in rats is mediated by platelet activating factor. J.Clin Invest. Vol,(81) : 1328 : 1331 .

Swartz , M.N. (1984) : Bacterial meningitis more involved than just the meninges . N.Eng. J.Med . Vol , (311) 912 - 914 .

Syrogianopoulos, G.A., Hansen, E.J., Erwin, A.L., *et al*; (1988) : Haemophilus influenzae type b lipopolysaccharide induces meningeal inflammation. J.Infect . Dis. Vol, (157): 237 - 244 .

Talmadage, T.E., Tribble, R., Pennington, R.W. Phillip, H., and Wittrout, R. (1987) : Immunomodulatory and immunotherapeutic properties of r TNF - α and IF - γ in mice cancer. Resp.,Vol, (47) p. 2563 - 2570 .

Tauber, M.G. and Sande, M.A. (1984) : Pathogenesis of bacterial meningitis : Contributions by experimental models in rabbits. Infections . Vol, (12) : 3-10 .

Taverne J., Tavernier, J., Fiers, W; and Playfair, J.H.(1987) : TNF inhibits malaria parasites in vivo but not in vitro. . clin. Exp. Immunol. Vol, : (67) p. 1 - 4 .

Todd, S. (1983) : Clinical diagnosis and laboratory methods. Edited by Israel D.M.D., Jhon, B., and Henry , M.D., Saunders company, Philadelphia, p. 1001

Tracey, K.J., Vlassara, H., and Cerami, A. (1989) : cachectin tumor necrosis factor. Lancet. Vol, (44) : 1122- 1126 .

Tuomanen, E., Tomasz, A., Hengstler, B., and, Zac, O.(1985) : The relative role of bacterial cell wall and capsule in the induction of inflammation in pneumococcal meningitis. J. Infect. Dis. Vol, (151): 535- 540.

Tuomanen, E. (1988) : Partner drugs : A new outlook for bacterial meningitis. Ann. Intern. Med. Vol, (109) : 690 - 692 .

Vander, J.W., Enders, S., Ionneman, G., Okuswa, S., cannon, J., IkeJima, T., Gelfand and Dinarello, C. (1987) : A newly developed, highly sensitive radioimmunoassay for TNF comparison of immunoreactive TNF with IL-1 production from stimulated monocytes. *Immunobiol.* Vol, (175): No 1/2 p. 87 .

Waren , M.K., and Raph, P.(1986) : Macrophage growth factor stimulates human monocyte production of INF, TNF and colony stimulating factor. *J.Immunol.* Vol, (137): p.. 2281 .

Washio M, Hamada, T, Goda, H, *etal* ; (1993) : Acyclovir resistant herpes zoster encephalitis treated with vidarabine : a case report. *Fukuoka - Igaku - Zasshi ;* 84 (10) : 436 : 439 .

Weinstein, L. (1985): Bacterial meningitis. *Medical clinics of North America*, Vol,(69). No. 2 : 219 - 229 .

Werner, M. (1969) : Serum protein changes during the acute phase proteins. *Clin. Chim . Acta.* Vol, (25) : 299- 305 .

Wilcocks, C., and Manson, B. (1978) : In Manson's tropical diseases . Bailliere Tindall. London 7th ed p . 113 .

Wilfert, c.M., Buckley, R.H., and Mahanokunur, T.(1977) : *New Eng. J. Med.* 296 : 1485 .

Xavier saez liorens, MD, Octavio Ramile, MD, Mahmoud M.Mustafa, M.D, Jussi Mertsola, M.D, and George H. McCracken, Tr., MD (1990) : Molecular pathophysiology of bacterial meningitis . *J. pediatri.* Vol, (116): .No 5 : 671 - 684 .

Yogev, R (1985) : Advances in diagnosis and treatment of childhood meningitis. Ped. Inf. Dis. J. Vol,(4) : 321 - 25

Ziegler, E.J. (1988) : Tumor necrosis factor in humans . New Eng. J.Med. Vol, (318) : 1533 - 1535 .