

REFERENCES

- Akiyama S.K. and Yamada K.M. (1982):** Fibronectin in diseases in connective tissue disease. Wanger B.M., Fleischmajer R. and Kaufman N. (eds.). Baltimore, London: Williams and Wilkins.
- Alaniz C. (1989):** Management of ascites. *Clin. Pharm.* 8:645-654.
- Alexander S., Calona G., Yamada K. and Paston I. (1978):** Molecular properties of a major cell surface protein from chick embryoblasts. *J. Biol. Chem.* 253:7787.
- Alex-Raplan L., Szabo H. and Ophen K.E. (1988):** Clinical chemistry interpretation and technique and test of hepatic function. 3rd edition, P. 226.
- Alitalo K., Kovi T. and Vaheri A. (1980):** Fibronectin is produced by human macrophages. *J. Exp. Med.* 151:171.
- Alvarez F.I., Bernard O., Brunelle F. and Hadchaul P., Odieuer M. and Alagille D. (1983):** Portal obstruction in children, clinical investigation and haemorrhage risk. *J. Ped.* 103(5):696-707.
- Bar-Meirs L.E. and Conn H.O. (1977):** Analysis of ascitic fluid in cirrhosis. *Dig. Dis. Sci.* 24:136-144.

Bayron C.P. (1930): Quoted by Hyatt R.E. and Smith J.R. (1954): The mechanism of ascites. *Am. J. Med.* 16(37):434-448.

Becker W. (1990): Die standerdisierung immunochoesisches L-C partigen pamphlet, Germany.

Behrman R.E. and Speck W.T. (1992): Ascites peritonitis. In: Nelson Text of Pediatrics. 14th edition. Philadelphia: W.B. Saunders Co., pp. 846-851.

Bender M.D. (1985): Disease of peritoneum. In: Wyngarden J.B. and Smith L.H. (eds.). Cecil Textbook of Medicine. 17th edition. Philadelphia: W.B. Saunders Co., P. 786.

Bichet D.G., Van Patten V.J. and Schrier R.W. (1982): Potential role of increased sympathetic activity in impaired sodium and water excretion in cirrhosis. *N. Engl. J. Med.* 307:1552.

Blumberg P. and Robbins P. (1975): Effects of proteases on activation of resting chick embryofibroblasts and on cell surface proteins. *Cell* 6:137.

Bockisch A., Ruhlmann J., Robertz G.M. and Biersack H.J. (1987): Scontio-graphic proof of peritoneal pleural leakage. *Excerpta Medica Int. Med.* 62(5):2.

Bohnsack J., Ashea J., Texahashi T. and Brown E. (1985): Fibronectin enhanced phagocytosis of an alternative pathway activator by human culture-derived macrophage is mediated by C₄b/C₃b complement receptor (CR). *J. Immunol.* 135:2680.

- Borhanmasesh F., Kekmat K., Vaezzadhe K. and Rezaei H.R. (1979):**
Tuberculous peritonitis, prospective study of 32 cases in
Iran. *Ann. Int. Med.* 76(4):677.
- Bowen T. and Muller E. (1981):** Influence of sample preparation of
estimation of blood fibronectin. *J. Clin. Path.* 36:233.
- Boyer T.D., Khan A.N. and Reynolds T.B. (1978):** Diagnostic value of
ascitic fluid lactic dehydrogenase protein and WBC levels.
Arch. Int. Med. 138:1103-1105.
- Boyer T.D. and Goldman I.S. (1986):** Treatment of cirrhotic ascites.
Adv. Int. Med. 31:359-377.
- Brenner D.A. and Alcron J.M. (1990):** Therapy for hepatic fibrosis.
Semin. Liver Dis. 10:75-83.
- Brook D.R., Altman R.S., Loebman W.W. and Seef B. (1981):**
Measurement of lactate in ascitic fluid. *Dig. Dis.
Sci.* 26:1089-1094.
- Brown B.P., Abu Yousef M., Farner R. and Labreequne-Gingrich R.
(1990):** Doppler sonography (non invasive method for
evaluation of hepatic VOD. *Am. J. Radiology* 145:721.
- Brunward E. (1991):** Heart failure. *Harrison's Principle of Internal
Medicine.* 9th edition. New York: McGraw-Hill Co., P. 1039.
- Buchwald H., Guzman E. and Wigness B.D. (1989):** The Minnesota
shunt. *A.S.A.I.O. Trans.* 35:168-170.

Buffon R. (1987): Principles of immunochemical techniques. In: Fundamental of Clinical Chemistry. 3rd edition. Philadelphia: W.B. Saunders Co., P. 152.

Buhac I. and Jarmolych J. (1978): Histology of intestinal peritoneum in patient with cirrhosis of the liver and ascites. *Dig. Dis. Sci.* 23:417-422.

Cafortune M., Patriquin H. and Pamin G. (1990): Haemodynamic changes in portal circulation after portosystemic shunt, use of doppler sonography in 43 cases. *Am. J. Radiology* 149:701.

Carre I.J., McCare W.M. and Mowat A.P. (1984): Ascites. In: Forfar J.O. and Arniel G.C. (eds.). Textbook of Pediatrics. 3rd edition. New York: Churchill Livingstone. Vol. II, pp. 499-500.

Castaldo G., Oriani G, Cimino L., Topa M., Mostarda I., Castellano L. and Salvatore F. (1994): Total discrimination of peritoneal malignant ascites from cirrhotics and hepatocellular carcinoma-associated ascites by assays of ascitic cholesterol and lactate dehydrogenase. *Clin. Chem. (Abst.)* 40(3):478-483.

- Chen A., Amrani D. and Mosesson M. (1977):** Heterogeneity of cold insoluble globulin of human plasma. *Biochem. Biophys. Acta* 310:228.
- Choate M., Mosher M. and Jennifer J. (1983):** Fibronectin concentration in plasma of patients with breast cancer, colon cancer and acute leukaemia. *A.S.A.I.O. Trans.* 35:175-178.
- Clark J.H., Fitzgerald J.F. and Kleiman M.B. (1984):** Spontaneous bacterial peritonitis. *J. Ped.* 104:4.
- Cohn E.M. (1985):** Ascites Bokus Gastroenterology. 4th edition. Philadelphia: W.B. Saunders Co., Vol. I, P. 177.
- Collaredo-Mels G., Bellati G., Auriemma L., Perani G., Leandro G., Vacca N., Pinna N. and Ideo G. (1991):** Fibronectin, cholesterol and triglycerides ascitic fluid concentration in the prediction of malignancy. *Ital. J. Gastroenterol. (Abst.)* 23(4):179-186.
- Colli A., Buccino G., Cocciolo M., Parravicini R., Mariani F. and Scaltrini G. (1986):** Diagnostic accuracy of fibronectin in the differential diagnosis of ascites. *Cancer* 1:58(11):2489-2493.
- Conn H.O. (1989a):** Cirrhosis in diseases of liver. 5th edition. Schiff L. and Schiff E.R. (eds.): Philadelphia, Toronto, pp. 847.

- Conn H.O. (1989b):** Rational approach to the therapy of ascites. *A.S.A.I.O. Trans.* 35:163-165.
- Dawson A.D. (1960):** Historical notes on ascites. *Gastroenterol.* 30:790-791.
- Dejegard A., Anderson T. and Gluide G. (1986):** The influence of insulin on the raised plasma fibronectin concentration in human obesity. *Acta Med. Scand.* 220:269.
- Denniston J.C. and Barker L.E. (1976):** Abdominal fluid accumulation measurement using electrical impedance. *Med. Biol. Eng.* 74:623.
- Domula M., Bykowaska K., Weszryniewicz Z., Lopaciuk C. and Kopic M. (1985):** Plasma fibronectin concentration in health and septic infants. *J. Ped.* 144:49.
- Donowitz M., Kerstein M.D. and Spiro H.M. (1974):** Pancreatic ascites. *Med.* 53(3):183-193.
- Dudley-Hart F. (1983):** French index of differential diagnosis. 10th edition. Philadelphia: W.B. Saunders Co., pp. 8-11.
- Edell S.L. and Gefter W.B. (1989):** Ultrasonic differentiation of types of ascitic fluid. *Am. J. Radiology* 133:111-114.
- Ekind O., Marien M. and Wasserman D. (1984):** Plasma fibronectin concentration in burned patients and influence of sepsis. *J. Trauma* 24:214.

- El-Gayar M.A. (1985):** Ascites. M.D. Thesis, Mansoura University.
- El-Hassan H.H. (1984):** Abdominal tumours. Postgraduate Doctor Middle East. Dec., 718.
- Elias H. (1979):** Ascites in French index of differential diagnosis. 11th edition. Hart F.D. (ed.). Bristol: John Right, P. 78.
- El-Khodary M. (1985):** Analytical procedures and instrumentations. 2nd edition. Cairo: Dar El-Maaref, P. 194.
- Epstein M. (1982):** Renal functional abnormalities in cirrhosis pathophysiology and management. In: Zakin D. and Boyer T.B. (eds.). Hepatology. 1st edition. Philadelphia, London, Toronto: W.B. Saunders Co., P. 446.
- Epstein M. (ed.) (1988):** The kidney in liver disease. 3rd edition. Baltimore: William and Wilkins, P. 3.
- Eriksen H., Kern-Hanser P., Brocks V. and Jensen B. (1987):** Plasma fibronectin concentration in normal pregnancy and pre-eclampsia. *Acta Obstet. Gynecol. Scand.* 66:25.
- Faloon W.W., Eckhardt R.D., Cooper A.M. and Davidson C.S. (1949):** The effect of human serum albumin, mercurial diuretics and low sodium diet on sodium excretion in patients with cirrhotic liver. *J. Clin. Invest.* 28:595.

Felisart J., Rimola A. and Arroyo V. (1985): Cefotaxime is more effective than ampicillin-tobramycin in cirrhosis with infection. *Hepatol* 5:457.

Forkman B., Ganrot P., Gennser G. and Rannevik G. (1972): Plasma protein pattern in recurrent cholestasis of pregnancy. *Scand. J. Clin. Lab. Invest.* 29 (Suppl.):124-189.

Franken E.A. (1982): Ascites in infants and children. *Radiology* 1033:393.

Fretz E.J., Stark D.D. and Metz C.E. (1990): Detection of hepatic metastasis, comparison of contrast enhanced C.T., un-enhanced M.R. imaging and iron oxide enhanced M.R. imaging. *Am. J. Radiology* 155:763.

Fukuda M. and Hakomori S. (1979): Carbohydrate structure of glycoprotein A. *J. Biol. Chem.* 254:S451.

Fujimura T. and Torisu M. (1987): Neutrophil-mediated tumour cell destruction in cancer ascites. *Clin. Immunol. Immunopathol.* 43:174-184.

Gahamberg C., Kiehn D. and Hakomori S. (1979): Change in surface labelled galactoprotein and in glycolipid concentration in cells transformed by temperature-sensitive virus mutant. *Nature* 298:413.

Galen R.S. and Gambino S.R. (1975): Beyond normality - the predictive value of efficiency of medical diagnosis. New York: John Wiley and Sons.

Garg R., Sood A., Arora S., Bhatia K.L and Chawla L.S. (1993): Ascitic fluid cholesterol in differential diagnosis of ascites. *J. Assoc. Physicians. India* (Abst.) 41(10):644-646.

Garrison R.N., Lawrence D., Rebecca H.G. and Louis S.H. (1986): Malignant ascites, clinical and experimental observations. *Ann. Surg.* 203(6):644-651.

Gerbes A.L., Jungst D, Xie Y, Permanther W. and Paumgartner G. (1991): Ascitic fluid analysis for the differentiation of malignancy-related and non-malignant ascites. *Cancer* 15(68):1808-1814.

Gerbes A.L., Xie Y., Mezger G. and Jungst D. (1990): Ascitic fluid concentrations of fibronectin and cholesterol: Comparison of differential diagnostic value with the conventional protein determination. *Liver* 10:152-157.

Gerding D.N., Khan M.Y. and Ewing J.W. (1976): *Pasteurella multocida* peritonitis in hepatic cirrhosis with ascites. *Gastroenterol.* 70:413.

Ginés P., Tito L.I. and Arroyo V. (1988): Randomized comparative study of therapeutic paracentesis with and without intravenous albumin in cirrhosis. *Gastroenterol.* 94:1493-1502.

Gittlein V., Stauffer J.C. and Silverter R.C. (1982): The pH of ascitic fluid in diagnosis of spontaneous bacterial peritonitis in alcoholics. *Hepatol.* 2:408-411.

Glickman R.M. and Isselbacher K.J. (1979): Abdominal swelling and ascites. In: *Harrison's Principles of Internal Medicine*. New York: McGraw-Hill Co., 9th edition, P. 211.

Goldberg B.B., Cleear-Field H.R., Goodman G.A. and Morales I.Q. (1973): Ultrasonic determination of ascites. *Arch. Int. Med.* 131:465.

Gottschalk A., Hoffer P.B., Polchan E.J. and Berger H.J. (1988): Textbook of Diagnostic Nuclear Medicine. 2nd edition. Baltimore, Hong Kong, London: Williams and Wilkins. Vol. II, Ch. 34.

Grange J.D., Amiot X. and Grange V. (1990): Amoxycillin-clavulinic acid therapy of spontaneous bacterial peritonitis: a prospective study in cirrhotic patients. *Hepatol.* 11:360.

Graninger W., Tatra G., Pirich K. and Nasr F. (1985): Low anti-thrombin III and high plasma fibronectin in pre-eclampsia. *Am. J. Obstet. Gynecol.* 122:213.

Granot E. and Deckelbaum R.J. (1983): Pseudoascites as presenting physical sign of celiac disease. *Am. J. Gastroenterol.* 78(11):730-731.

Grinnel F. (1976): The serum dependence of total Hamster kidney cell attachment to substratum. *Exp. Cell Res.* 97:265.

Grinnel F., Feld M. and Miner D. (1980): Fibroblast adhesion to fibrinogen and fibrin substrata: A requirement for cold insoluble globulin (plasma fibronectin). *Cell* 517:519.

Hayashi Y. and Torisu M. (1990): New approach to management of malignant ascites with streptococcal preparation OK-432 III. *Surg* 107:74-84.

Hecker R. and Sherlock S. (1956): Electrolyte and circulatory changes in terminal liver failure. *Lancet* ii:1121.

Henderson M.J., Stein S.F., Kutner M., Asley J.D. and Rudmon D. (1980): Analysis of twenty three plasma proteins in ascites. *Ann. Surg.* 192:738-742.

Henery J.B. (1979): Diagnosis and management by laboratory methods. 16th edition. Philadelphia: W.B. Saunders Co., P. 679.

Henriksen J.H. Bendsten F. and Schutten H.G. (1990): Hepatic intestinal disposal of endogenous human alpha atrial natriuretic factor in cirrhotics. *Am. J. Gastroenterol.* 85:1155.

Henriksen J.H. and Ring Larsen H. (1990): Ascites formation in liver cirrhosis: The how and the why. *Dig. Dis.* 8:152.

Herlong H.F. (1988): Portal hypertension, ascites, subacute bacterial peritonitis and hepato-renal syndrome. The principle and practice of medicine. 22nd edition. Harvey A.M., Jones R.S., McKusick V.A. and Owens A.H. (eds.). Prentice-Hall International Inc., P. 848.

Hirshman J.V. (1978): Pericardial constriction. *Am. J. Heart* 96:110.

Hoefs J.C. (1981): Increase in ascites white blood cells and protein concentration during diuresis in patients with chronic liver disease. *Hepatology* 1:249-254.

Hoefs J.C., Canawati H.N. and Sapico F.L. (1990): Spontaneous bacterial peritonitis. *Hepatology* 2:399.

Hofeler H. and Klingemann H. (1984): Fibronectin and factor VIII, related antigen in liver cirrhosis and acute liver failure. *J. Clin. Biochem.* 22:15.

Hurwich D.B., Lindor K.D., Hay J.E. and Gross J.B. (1993): Prevalence of peritonitis and the ascitic fluid protein concentration among chronic liver disease patient. *Am. J. Gastroenterol.* 88(8):1254-1257.

Hutchinson G.H. (1987): Diseases of peritoneum, diseases of the gut and pancreas. Oxford: Blackwell Scientific Publication, pp. 1056-1070.

Hynes R. (1986): Fibronectin. *Am. J. Sci.* 254:42.

Hynes R., Ali I. and Desiree A. (1978): A large glycoprotein lost from the surface of transformed cells. *Ann. Acad. Sci.* 312:317.

Jenkins P.F. and Ward M.J. (1980): Role of peritoneal biopsy in the diagnosis of ascites. *Postgraduate Med. J.* 56:102.

John H., Peters K., Richard J., Maunder A., Anthony D. and Charles G. (1989): Elevated plasma level of ED⁺ (cellular fibronectin in patients with vascular injury). *J. Lab. Clin. Med.* 113:586-597.

Jolles H. and Coculam C. (1980): C.T. of ascites differential diagnosis. *Am. J. Radiology* 135:3154-322.

Jungst D., Gerbes A.L., Martin R. and Paumgartner G. (1986): Value of ascitic lipids in the differentiation between cirrhotic and malignant ascites. *Hepatol. (Abst.)* 6(2):239-43.

Kao H.W., Rakov N.E., Savage E. and Reynolds T.B. (1983): The effect of large volume paracentesis on plasma volume. *Hepatol.* 5:403-407.

Katano M. and Torisu M. (1983): New approach to management of malignant ascites with streptococcal preparation OK-432 II. *Surg.* 93:365-373.

Kauma S., Clark M., White C. and Halne J. (1989): Production of fibronectin by peritoneal macrophages and concentration of fibronectin in peritoneal fluid from patients with or without endometriosis. *Obstet. Gynecol.* 13:72.

Kew M.C., Vema R.R. and Williams D. (1971): Renal and intrarenal blood flow in cirrhosis of the liver. *Lancet* 2:504.

Kleckner M.S. (1960): Cirrhosis of the liver. Springfield: Charles C. Thomas, P. 544.

Kline M.M., McCalum R. and Gath P.H. (1979): The clinical value of ascitic fluid culture and leucocytes studies in alcoholic cirrhosis. *Gastroenterol.* 70:408-412.

Kurt J., Isselbacher T., Jack R. and Wands H. (1991): Neoplasm of liver. In: Harrison's Principle of Internal Medicine. 12th edition. Vol. II, pp. 1348-1350.

Kusmo J.W. (1984): Basic Statistics for Health Science. 1st edition. Mayfield Publishing Co., California.

Lach J., Rimola A. and Navasa M. (1992): Incidence and predictive factors of first episode of spontaneous bacterial peritonitis in cirrhosis with ascites: relevance of ascitic fluid protein concentration. *Hepatology* 16(3):724-727.

Laffi G., La-Villa G., Pinzani M., Carloni V., Cassini-Raggi V. and Gentilini A. (1993): Refractory ascites. *Ann. Ital. Med.* (Abst.) 8(1):38-46.

Lai K.N., Leung J.W. and Lake J. (1987): Ultrafiltration by hemofilter - a new therapeutic measure in intractable ascites. *Int. J. Artif. Organs* 10:109-114.

Lanser M., Brown J., Mora R. and Seigel J. (1986): Neutrophil chemiluminescence and opsonic fibronectin level following blunt trauma. *J. Surg. Res.* 41:269.

Leedlt D.H., Lim J.H. and Koyt Y. (1990): Sonographic detection of pneumoperitoneum in patient with acute abdomen. *Y.D.L.* 154:107.

Le Veen E.G. (1989): The place of peritoneo-venous shunt in the treatment of ascites. *A.S.A.I.O. Trans.* 35:165-168.

Light R.W., Mac Gregor I., Luchsinger P.C. and Ball W.C. (1972): Pleural effusion, the diagnostic separation of transudate and exudate. *Ann. Int. Med.* 77:507-511.

Lindsay K.L., Reynolds T.B. and Hoefs J.C. (1981): Ascites (Clinical conference). *West J. Med.* 134:414-425.

Longmire-Cook S.J. (1993): Pathophysiologic factors and management of ascites. *Surg. Gynecol. Obstet.* 176(2):191-202.

Lund R.H. and Newkirk J.B. (1979): Peritoneovenous shunting system for surgical management of ascites. *Surg.* 14:31-45.

Mal F., Nizard G., Labadie H., Trinchet J.C., Carnier M. and Beaugrand M. (1987): Fibronectin in ascitic fluid: its diagnostic significance. *Gastroenterol. Clin. Biol.* 11(10):369-642.

Mann S.K. and Mann. N. (1986): Pancreatic ascites. *Am. J. Gastroenterol.* 71:2.

Marc E., Lanser M. and Thomas M. (1983): Correction of serum opsonic defects after burn and sepsis by opsonic fibronectin administration. *Arch. Surg.* 118:338.

Marsh M.N. (1984): Chronic liver disease. In: *Clinical Gastroenterology*. 1st edition. Oxford: Blackwell Scientific publication, P. 481.

Masenda M., Yamanaka K. and Mastuda A. (1982): Distribution of fibronectin in plasma and liver diseases. *Clin. Chem. Acta* 118:141.

McDermott W.V., Stone M.O., Bothe A. and Trey C. (1984): Budd-Chiari syndrome: historical and clinical review with analysis of surgical corrective procedures. *Am. J. Surg.* 147:463-467.

Meinke A.H., Ester N.C. and Ernst C.B. (1979): Chylous ascites following abdominal aortic aneurysm repair. *Ann. Surg.* 190:631-633.

Menzel E. and Dunky A. (1986): Plasma fibronectin in the aged. *J. Gastroenterol.* 19:152.

Mezger G., Perametter W., Gerbes A.L., Wilmanns W. and Lamerz R. (1988): Tumour associated antigens in diagnosis of serous effusions. *J. Clin. Pathol.* 41:633-643.

- Miller W.J., Federle M.P. and Comptell W.L. (1991):** Diagnosis and staging of hepatocellular carcinoma comparison of C.T. and sonography in 36 liver transplantation patient. *Am. J. Radiology* 157:303.
- Morgan T.R., Imada T. and Hollister A.S. (1988):** Plasma human atrial natriuretic factor in cirrhosis and ascites with and without functional renal failure. *Gastroenterol.* 95:1641.
- Mortensen P.B., Kristensen S.D., Bloch A., Jacobsen B.A. and Rasmussen S.N. (1988):** Diagnostic value of ascitic fluid cholesterol level in the prediction of malignancy. *Scand. J. Gastroenterol.* 23(9):1085-1088.
- Mosesson M. and Amrani D. (1980):** The structure and biological activities of plasma fibronectin. *Blood* 56:145.
- Mosesson M., Chen A. and Husbey R. (1975):** The cold-insoluble globulin of human plasma. *Biochem. Biophys. Acta* 386:509.
- Mosher D. (1975):** Cross linking of cold-insoluble globulin by fibrin-stabilizing factor. *J. Biol. Chem.* 250:6614.
- Mowat A.P. (1987):** Liver Disorders in Childhood. Butterworth and Co., pp. 270-271.

- Nance F.C. (1985):** Disease of the peritoneum, mesentery and omentum. Bockus Gastroenterology. 4th edition. Philadelphia: W.B. Saunders Co., Ch. 7, pp. 4177-4190.
- Okumura H., Aramaki T. and Katsuta Y. (1989):** Pathophysiology and epidemiology of portal hypertension. *Drugs* 37:2-12.
- Olsson R. and Waldentrom T. (1979):** Gamma-glutamyl transferase activity in ascitic fluid in diagnosis of hepatocellular carcinoma. *Br. Med. J.* 11:830-831.
- Paré B., Talbot G. and Hoefs G.S. (1983):** Serum-ascites albumin concentration gradient: A physiologic approach to the differential diagnosis of ascites. *Gastroenterol.* 85:240-244.
- Parikh S.S., Amarapurkar D.N., Viswanath N., Desai H.G. and Kalro R.H. (1991):** Ascitic fluid protein and cellular changes during diuretic therapy in cirrhosis of liver. *J. Assoc. Physicians India* (Abst.) 39(3):263-264.
- Pearlstein E., Gold L. and Garcia P. (1981):** Fibronectin. A review of its structure and biological activity. *Mol. Cell. Biochem.* 29:103.
- Perez-Ayuso R.M.K., Arroyo V. and Planas R. (1983):** Randomized comparative study of efficacy of furosemide versus spironolactone in non-azotemic cirrhosis with ascites. *Gastroenterol.* 84:961-968.

- Perez-Ayuso R.M.K., Arroyo V. and Camps J. (1989):** Renal kallikrein excretion in cirrhotics with ascites: a relationship to renal hemodynamics. *Hepatol.* 4:247.
- Perri R., Kye N. and McCarthy J. (1982):** Fibronectin enhances in vitro monocyte macrophage mediated tumouricidal activity. *Blood* 60:430.
- Petress T.J. (1977):** Gamma-glutamyl transferase activity in patients with hepatoma. *Br. Med. J.* 1:1576.
- Postlemwaite A., Keski O., Babian G. and Kang A. (1981):** Introduction of fibroblast chemotaxis by fibronectin. *J. Exp. Med.* 46:47.
- Prieto M., Gomez-Lechon M.J., Hoyos M., Castell J.V., Carrasco D. and Berenguer J. (1988):** Diagnosis of malignant ascites. Comparison of ascitic fibronectin, cholesterol and serum-ascites albumin difference. *Dig. Dis. Sci* 33(7):833-838.
- Press O.W., Press N.O. and Kanfmen S.D. (1985):** Evaluation and management of chylous ascites. *Ann. Int. Med.* 96:358-364.
- Proctor R., Prendergast E. and Mosher D. (1982):** Fibronectin mediates attachment of *Staphylococcus aureus* to human neutrophils. *Blood* 59:681.

- Proctor R., Textor J., Van J. and Mosher D. (1985):** Role of fibronectin in human monocyte and macrophage bacterial activity. *Infect. Immun.* 47:629.
- Pussell B., Peake P., Brown M. and Charesworth J. (1985):** Human fibronectin metabolism. *J. Clin. Invest.* 143:76.
- Quintero E., Ginés P. and Arroy V. (1985):** Paracentesis versus diuretics in the treatment of ascites. *Lancet* 1:611-612.
- Rector W.G. and Reynolds T.B. (1984):** Propranolol in the treatment of ascites. *Arch. Int. Med.* 144:1761-1763.
- Rector W.G., Adair O. and Hossack K.F. (1990):** Atrial volume in cirrhosis: relationship to blood volume and plasma concentration of atrial natriuretic factor. *Gastroenterol.* 99:766.
- Reynolds T.B. and Peter R.L. (1982):** Budd-chiari syndrome. In: *Disease of Liver*. 5th edition. Schiff L. and Schiff E.R. (eds.). Philadelphia: J.B. Lippincott, P. 1622.
- Reynolds T.B. and Campra J.L. (1985):** Ascites in liver disease. *Bockus Gastroenterology*. 4th edition. Philadelphia: W.B. Saunders Co., P. 3121.
- Reynolds T.B. and Cowan R.E. (1979):** Peritoneoscopy, differential diagnosis of ascites, liver and biliary disease. Wright R. et al. (eds.) London: W.B. Saunders Co., P. 544.

Rocco V.K. and Ware A.J. (1986): Cirrhotic ascites; pathophysiology, diagnosis and management. *Ann. Intern. Med.* 105:573-585.

Roger F., Shealf D. and Nalan P. (1986): Fibronectin depletion and microaggregate clearance following trauma. *J. Trauma A.P.* 2:339.

Rouslahti E, Vuento M. and Engvall E. (1978): Interaction of fibronectin with antibodies and collagen in radioimmunoassay. *Biochem. Biophys. Acta* 534:210.

Runyon B.A. (1986): Elevated ascitic fluid fibronectin concentration. *Hepatol.* 3:219-222.

Runyon B.A. and Hoefs J.C. (1985): Ascitic fluid chemical analysis before, during and after spontaneous bacterial peritonitis. *Hepatol.* 5(2):257.

Runyon B.A., Morrissey R.L., Hoefs J.C. and Syle F.A. (1985): Opsonic activity of human ascitic fluid, a potential important protective mechanism against spontaneous bacterial peritonitis. *Hepatol.* 5:634-637.

Saba T. (1970): Physiology and pathophysiology of reticuloendothelial system. *Arch. Int. Med.* 126:1031.

Saba T., Blumenstock F., Sham D., Landabura R., Hrinda M. and Deno D. (1986a): Reversal of opsonic deficiency in surgical trauma and burned patients by infusion of human plasma fibronectin, correlation with experimental observation. *Am. J. Med.* 80:129.

- Saba T., Kiener J. and Holman J. (1986b):** Fibronectin and the critically ill patient. *Intens. Care Med.* 12:350.
- Salerno F., Badalamenti S. and Incerti P. (1987):** Repeated paracentesis and intravenous albumin infusion to treat tense ascites in cirrhotic patients. *J. Hepatol.* 5:102-108.
- Salerno F., Badalamenti S. and Moser O. (1990):** Effect of large volume paracentesis. *Gastroenterol.* 98:1063-1070.
- Scholmerich J., Volk B.A., Kottgen E., Ehlers S. and Gerok W. (1984):** Fibronectin concentration in ascites differentiates between malignant and non-malignant ascites. *Gastroenterol.* 87(95):1160-1164.
- Schrier R.W., Arroyo V. and Bernardi M. (1988):** Peripheral arterial vasodilatation hypothesis: a proposal for initiation of renal sodium and water retention in cirrhosis. *Hepatol.* 8:1151.
- Scott D.L., Farr M., Crockson A.P. and Walton K.W. (1982):** Synovial fluid and plasma fibronectin levels in rheumatoid arthritis. *Clin Sci* 62:71-76.
- Shear L. (1990):** Ascites in cirrhosis: A medical or surgical problem. *Hepatol.* 11:323-325.
- Sherlock S. and Dooly G. (1993):** Ascites. In: Disease of Liver and Biliary System. 9th edition. Philadelphia: W.B. Saunders Co., Ch. 9, pp. 114-131.

Sharp P.F., Gemmell H.G. and Smith F.W. (1989): Textbook of Nuclear Medicine. 1st edition. Oxford, New York, Tokyo: IRL Press, Ch. 2.

Sherman S., Rohwedder J.J., Ravikrishnank P. and Weg I.G. (1980): Tuberculous ascites and peritonitis, Reports of 36 general hospital cases. *Arch. Int. Med.* 740:506-508.

Siddiqui R.A., Kochhar R, Singh V, Rajwanchi A., Goenka M.K. (1992): Evaluation of fibronectin as a marker of malignant ascites. *J. Gastroenterol. Hepatol.* 7(2):161-164.

Soriano G., Guarner C. and Teixido M. (1991): Selective intestinal decontamination prevents spontaneous bacterial peritonitis. *Gastroenterol.* 100(2):477-481.

Stanley M.M. (1989): Pathogenesis of ascites: a unitary hypothesis. *A.S.A.I.O. Trans. (Abst.)* 35(2):181-183.

Stanley M.M., Greenlee H.B. and Chejfec T.J. (1988): International abstracts from peritoneal fibrosis in cirrhotic with ascites treated with peritoneovenous shunt. *Gastroenterol. (Abst.)* 92(6):2.

Slovic T.L., Clopp S.K. and Farooki Z.Q. (1984): Non-invasive evaluation of inferior vena cava: *A.J.D.C.* 138:277.

Stathakis N. and Mosesson M. (1977): Interaction among heparin, cold insoluble globulin and fibronectin in the formation of heparin insoluble precipitate fraction of plasma. *J. Clin Invest.* 60:885.

- Stathakis N., Fountas A. and Tsianose E. (1981):** Plasma fibronectin in normal subjects and in various disease states. *J. Clin. Pathol.* 34:504.
- Strassen W.N. and McCallough A.J. (1985):** Management of ascites. *Semin. Liver Dis.* 5:291-307.
- Strassen W.N., McCullough A.J. and Bacon B.R. (1986):** Immediate diagnostic criteria for bacterial infection of ascitic fluids. *Gastroenterol.* 90:1247.
- Such J., Guarner C. and Soriano G. (1990):** Selective intestinal decontamination increases serum and ascitic fluid C₃ level in cirrhosis. *Hepatol.* 12:1175.
- Summerskil W.H.J. and Banldus W.P. (1975):** Ascites in disease of the liver. 4th edition. Lean Schiff (ed.): Philadelphia, Toronto, P. 424.
- Svanberg L. and Astedt B. (1975):** Coagulative and fibrinolytic properties of ascitic fluid associated with ovarian tumour. *Cancer* 35:1382-1387.
- Targan S.R., Chow A.W. and Guze L.B. (1976):** Spontaneous peritonitis of cirrhosis due to *Campylobacter fetus*. *Gastroenterol.* 71:311.

- Tessler F.N., Gehring R.J. and Gomomos A.S. (1991):** Diagnosis of portal vein thrombosis: Value of colour Doppler imaging. *Am. J. Radiology* 157:293.
- Thomas L. (1992):** Laboratory and Diagnosis. 4th edition.
- Tito L., Rimola A. and Ginés P.(1988):** Recurrence of spontaneous bacterial peritonitis in cirrhosis: Frequency and predictive factors. *Hepatol.* 8:27.
- Torisu M., Katano M. and Kimura Y. (1983):** New approach to management of malignant ascites with streptococcal preparation OK-432 I. *Surg.* 93:357-364.
- Toy P.T. and Reid M.E. (1984):** Sample collection for determination of plasma fibronectin concentration. *J. Clin. Pathol.* 37:951.
- Trichopoulos D., Sizaret P. and Tabor E. (1980):** Alpha fetoprotein in liver cancer patients and controls in European population. *Cancer* 46(4):736.
- Waeber B. (1988):** Clinical applications of antimineralocorticoids. *J. Steroid. Biochem. Mol. Biol.* 31:739-744.
- Wagner D. and Hynes R. (1979):** Somain structure of fibronectin and its relation to function disulfide and sulp-hydryl groups. *J. Biol. Chem.* 254:6746.
- Wang S.S., Tsai Y.T., Lee S.D., Chen H.T. and Lu C.W. (1991):** Spontaneous bacterial peritonitis in patients with hepatitis B-related cirrhosis and hepatocellular carcinoma. *Gastroenterol.* 101(6):1656-1662.

Ward P.C.J. (1982): Interpretation of ascitic fluid data. *Postgrad. Med.*
71:171.

Welch H.F., Welch C.S. and Carter J.H. (1964): Prognosis after
surgical treatment of ascites: result of side to side shunt in
40 patients. *Surg.* 56:75.

Wolff L., Timple R., Pecker L. and Steffen C. (1967): A two
component system of human serum agglutination gelatin-
coated erythrocyte. *Vox. Sang.* 12:443.

Wyllie R.A., Rosu T.S. and Fitzgerald G.F. (1980): Ascites
pathophysiology and management. *J. Ped.* 97(2):167-176.

Yamada K. and Weston J. (1974): Isolation of major cell surface
glycoprotein from fibroblasts. *Proc. North Acad. Sci.*
71:3492.

Yeh H. and Wolf B.S. (1977): Ultrasonography in ascites. *Radiology*
124:683-789.