

REFERENCES

REFERENCES

Abdou, I. (1995):

Immunosuppressive and cytotoxic chemotherapy. Long term complication. Ann. Intern. Med., 83: 87.

Abo-El-Fetoh, A.; Lymona, N.; Abo El-Magd, M.; Aref, S. and Shoker, M. (1999):

Chromosomal breakage in systemic lupus erythematosus. Thesis Submitted for Partial Fulfillment of M.D. in Rheumatology and Immunology, Faculty of Medicine, Mansoura University, 83.

Ahearn, J.M.; Provost, T.; Dorsch, C.A. (1982):

The interrelationships of HLADR, MB and MT phenotypes, autoantibody expression and clinical features in SLE. Arthritis Rheum., 25: 1031.

Ahearn, J.M.; Provost, T.; Dorsch, C.A. (1992):

The interrelationships of HLA DR, MB, MT phenotypes, antibody expression and clinical features in SLE. Arthritis Rheum., 25: 1031.

Albani, S.; Keystone, E.; Nelson, J.L. et al. (1995):

Positive selection in autoimmunity: abnormal immune responses to a bacterial dnaJ antigenic determinant in patient with early rheumatoid arthritis. Nature Med., 1: 448-452.

Altman, H. (1977):

DNA and mutation in immunocomponent cells from patient with rheumatoid diseases and corresponding animal models. Smdia Biophysica, 61: 89-96.

Andrews, B.S.; Eisenberg, R.A.; Thefilopoulos, A.N. et al. (1978):

Spontaneous murine lupus-like syndromes. Clinical and immune pathological manifestation in several strains. J. Exp. Med., 148: 1198.

Arlett, C.F. and Lehmann, A.R. (1996):

Xeroderma pigmentosa, cockayne syndrome and trichothiodystrophy sun-sensitive to DNA repair defect in skin cancer. In: Eles RA< Pander BAJ, Easton DE et al. eds. Genetic pre-disposition to cancer landon: Chapman and Hall 185: 206.

- Arlinger, M.; Wintersberger, W.; Steiner, C.W. et al. (1994):**
High levels of bcl-2 protein in circulating T-lymphocytes but not B-lymphocytes of patient with SLE.
Arthritis Rheum., 37: 1423.
- Arnett, F.C.; Edworthy, S.M.; Bloch, et al. (1988):**
The classification of rheumatoid arthritis.
Arthritis Rheum., 31: 315-324.
- Arnett, F.C. and Reveille, J.D. (1992):**
Genetics of systemic lupus erythematosus. *Rheum. Dis. Clin. North Am.*, 18: 865-93.
- Auclair, M.; Medsger, T.A.; Gallway, S.M. et al. (1987):**
Frequencies of sister chromatid exchange in patients with diffuse systemic sclerosis and their relatives.
Rev. Rheum. Malosteoartic. Nov., 15(6): 584-9.
- Austin III, H.A.; Boumpas, D.T.; Vaughan, E.M. and Balow, J.E. (1994):**
Predicting renal outcomes in severe lupus nephritis: contributions of clinical and histologic data.
Kidney Int., 45: 544-550.
- Bannwarth, B.; Labat, L. and Moride, Y. (1994):**
Methotrexate in rheumatoid arthritis AN. Update. *Drugs*, 47: 25-50.
- Barbut, D.; Borer, J.S.; Coronna, J.; Wallerson, D.; Asherson, R.A. (1994):**
SLE clinical and immunologic patterns of disease expression in cohort of 1000 patients.
Medicine (Baltimore) 72: 113-114.
- Belmont, H.M.; Buyon, R. and Abramson, S. (1994):**
Up-regulation of endothelial cell adhesion molecules characterizes disease activity in SLE. The Schwartzman Phenomenon. *Arthritis Rheum.*, 37: 376-383.
- Berczi, J. (1993):**
Prolactin, pregnancy and autoimmune disease.
J. Rheumatol., 20: 1095.

Berrino, J.R. (1993):

Chemical action and pharmacology of methotrexate, azothioprine and cyclophosphamide in men.
Arthritis Rheum., 16: 79-83.

Bernstein, R. (1999):

Cytogenetics and malignancy: the value of chromosome studies in the management of patient with haematological disorders.
The Oncotech Reporter, Issue, 6, September.

Bloom, D. (1994):

Many fold increase in sister chromatid exchange.
Proc. Natl. Acad. Sci., USA, 71: 4508-4512.

Bodmer, J.G.; March, S.G.E.; Albert, E.D. et al. (1994):

Nomenclature for factors of the HLA system.
Tissue Antigen, 44:1.

Borg, S.; Moride, Y.; Ouellette, A.M. and Latt, S.A. (1990):

Sister chromatid exchanges in systemic lupus erythematosus.
Arthritis, Rheum., 58: 119-140.

Boumpas, D.T.; Fessler, B.J.; Austin, III, H.A. and Balow, J.E. (1995):

Systemic lupus erythematosus: emerging concepts. Part 2: Dermatologic and joint diseases.
Ann. Intern. Med., 123: 42-53.

Boumpas, D.T.; Austin, H.A.; Vaughan, E.M. et al. (1996):

Risk for sustained a menorrhea in patient with SLE receiving intermittent pulse cyclophosphamide.
Ann. Intern. Med., 119: 366.

Bowness, P.; Davies, K.A.; Norsworthy, P.J. et al. (1994):

Hereditary cell deficiency and systemic lupus erythematosus.
Q.J. Med., 87: 455-464.

Brown, J.H.; Jardetzky, T.S.; Gorga, J.C. et al. (1993):

Three dimensional structure of the human class II histocompatibility antigen HLA-DRI.
Nature, 354: 433.

Bruckmann, E.; Wojcik, A. and Obe, G. (1999):

X-irradiate of G. Chocell induces SCG which both true and Brdu-Substituted. Chromosome Research, 7: 277-288.

Bryant, S. (1998):

Cytogenetic effect of near ultraviolet radiation in normal and systemic lupus erythematosus lymphocytes. J. Clin. Invest., 643-71.

Caron, D.A.; Chen, P.P.; Kipps, T.J. (1991):

New roles for rheumatoid factor.
J. Clin. Invest., 87: 379-383.

Caporossi, D.; Sebastiani, G.; Masala, C. and Nicoletti, B. (1991):

Cytogenetic effects of near ultraviolet radiation in normal and systemic lupus erythematosus, lymphocytes. Mutat. Res. Mar., 229(1): 43-7.

Chen, P.P.; Silverman, G.S.; Liu, M.F. and Carson, D.A. (1993):

Idiotypic and molecular characterization of human rheumatoid factors. Chem. Immunol., 48: 63.

Chrousos, G.P.; Detera-Wadleigh, S.D. and Karl, M. (1993):

Syndromes of glucocorticoid resistance.
Ann. Intern. Med., 119: 113.

Claman, H.W.; Giorno, R.C.; Seibold, J.R. (1991):

Endothelial and fibroblast activation in scleroderma: the myth of the "uninvolved skin".
Arthritis Rheum., 34: 1495-1501.

Clement, P.J.; Furst, D.E.; Carapion, D.S.; Bohan, A. and Harris (1992):

Muscle disease in systemic sclerosis: Diagnostic and therapeutic considerations.
Arth. Rheum., 21: 62-71.

Clement, P.J.; Furst, D.E.; Carapion, D.S.; Bohan, A.; Harris (1994):

Diagnostic and therapeutic considerations.
Arth. Rheum., 21: 62-71.

Clements, P.J.; Furst, D.E. (1996):

Pathology: Systemic sclerosis. In: Clement PJ, Furst DE (eds). Baltimore, Williams and Wilkins, P. 170-175.

Croastein, B.N.; Kimmel, S.C.; Levin, R.I. et al. (1993):

A mechanism for the anti-inflammatory effects of corticosteroid. The glucocorticoid receptor regulates leukocyte adhesion to endothelial cells and expression of endothelial leukocyte adhesion molecule-1 and intracellular adhesion molecule-1.

Proc. Natl. Acad. Sci., USA, 89: 9991.

Davidson, R.L.; Kaufman, E.R.; Dougherty, C.P.; Ouellette, A.M.; Difolo, C.M. and Latt, S.A. (1980):

Induction of sister chromatid exchanges by Budr is largely independent of the BudR content of DNA.

Nature, 284: 74-6.

Davies, E.J.; Snowden, N.; Hillarby, M.C. et al. (1995):

Mannose-binding protein polymorphism in systemic lupus erythematosus. Arthritis, Rheum., 38: 110-14.

Dean, B.J. and Danford, N. (1984):

Assays for the detection of chemically induced chromosome damage in cultured mammalian cells. In: Venitts, Parry-J.M., eds. Mutagenicity testing, A practical approach. Oxford: IRL Press, 187-232.

Degiannis, D.; Seibold, J.R.; Czarneck, M. et al. (1990):

Soluble and cellular markers of immune activation in patients with systemic sclerosis.

Clin. Immunol. Immunopathol., 56: 259.

Denman, S. (1995):

Detection of breaking capacity factor in patients with rheumatoid arthritis. Ann. Rheum., 3: 45-9.

Dersimonian, H.; Sugita, M.; Glass, D.N.; Maier, A.L.; Weinblatt, M.C.; Reme, T. and Brenner, M.B. (1993):

Clonal V α 12.1 + T-cell expansions in the peripheral blood of rheumatoid arthritis patients. J. Exp. Med., 177: 1623-1623.

- Donadio, J.V.Jr.; Glassock, R.J. (1993):**
Immunosuppressive drug therapy in lupus nephritis.
Am. J. Kidney, Dis., 21: 239.
- Douvas, A. (1996):**
Pathogenesis: serologic correlates. In: Clement PJ, Furst DE (eds): Systemic sclerosis. Baltimore, Williams & Wilkins, P. 175-202.
- Dovazen, R. and Gigli, L. (1992):**
Expression and localization of proteins of the complement system in human skin.
J. Clin. Invest., 90: 200-12.
- Egaila, S.; Hammad, G. and El-Nashar, E. (2001):**
Chromosomal damage in lymphocytes of patients with systemic lupus erythematosus.
Benha MJ, May: 151-162.
- Emerit, I. (1976):**
Chromosomal breakage in systemic sclerosis and related disorders. Dermatologica, 153: 145-56.
- Emerit, I. and Marteau, R. (1980):**
Chromosomal studies in 14 patient with disseminated sclerosis. Hum. Genet., 13: 25-33.
- Endoh, M.; Yamashina, M.; Oh, H. et al. (1993):**
Immuno-histochemical demonstration of membrane cofactor protein (MCP) of complement in normal and diseased kidney tissues. Clin. Exp. Immunol., 94: 182-8.
- Gebhart, E. (1981):**
Sister chromatid exchanges and structural chromosome aberration in mutagenicity testing.
Hum. Genet., 58: 235-54.
- Gelehrter, T.D.; Collins, F.S. and Ginsburg, D. (1998):**
The principles of medical genetics. Second edition, Williams & Wilkins, A Waverly company, P. 335-55.
- Gerber, N.L. and Steinberg, A.D. (1979):**
Clinical use of immunosuppressive drug. Part. I Drugs, 11: 90.

- Goronzy, J.J.; Bartz-Bazzanella, P.; Hu, W.; Jendroy, M.C.; Walser-Kuntz, D.R. and Weyand, C.M. (1994):**
Dominant clonotypes in the repertoire of peripheral CD4+ T-cells in rheumatoid arthritis.
J. Clin. Invest., 94: 2068-2076.
- Goronzy, J.J. and Weyand, C.M. (1995):**
T-cells in rheumatoid arthritis: Paradigms and Facts.
Rheum. Dis. Clin. North Am., 21: 655-674.
- Griffing, W.L.; Moore, S. and Lithra, H.S. (1980):**
Associations of antibodies to native DNA with HLA-DR W3: A possible major histocompatibility linked human immune response gene.
J. Exp. Med., 152: 319.
- Grumet, F.C.; Coukell, A.; Bodmer, J.G. et al. (1991):**
Histocompatibility HLA antigens associated with system lupus erythematosus. A possible genetic predisposition to disease. N. Engl. Med., 285: 193.
- Falanga, V.; Soter, N.A.; Altman, R. et al. (1990):**
Elevated plasma histamine levels in systemic sclerosis (scleroderma). Arch. Dermatol., 126: 336.
- Feig, D.J. and Loeb, L.A. (1996):**
Endogenous mutagenesis. In: Eeles RA Ponder BAJ, Easton DF, et al. eds. Genetic pre-disposition to cancer London: Chapman and Hall, 175-185.
- Felsenfeld, G. (1992):**
Chromatin as an essential part of the transcriptional mechanism. Nature, 355: 219.
- Ferguson, M. (1996):**
Sister chromatid exchanges in human chromosomes including observations. Cancer Genet. Cytogenet., 1: 363-380.
- Fialkow, P.J. (1997):** Chromosomal breakage induced by extracts of human allogeneic lymphocytes. Science, 155: 1676-1677.

Firestein, G.S. (1991):

The immunopathogenesis of rheumatoid arthritis.
Curr. Opin. Rheumatol., 3: 398-406.

Fritzler, M.J. and Salazar, M. (1991):

Diversity and origin of rheumatologic autoantibodies. Clin.
Microbiol. Rev., 4: 256.

Fritzler, M.J.; Kinsella, T.D. and Garbutt, E. (1991):

The CREST syndrome: A distinct serologic entity with
anticentromere antibodies. Am. J. Med., 69: 520.

Fritzler, M.J. (1992):

Immunofluorescent antinuclear antibody test in Rose NR de
Macario EC, Fahey JL Friedman H, Penn GM (eds.):
Manual of clinical laboratory immunology Washington DC,
American Society for Microbiology, P: 724.

Fritzler, M. and Rubin, R. (1993):

Drug induced lupus in Wallace DJ Hahn BH (eds.): Dubois
Lupus Erythematosus, Philadelphia, Lea, Febige, P. 442.

Fox, D.A. and McCune, W.J. (1994):

Immunosuppressive drug therapy of systemic lupus
erythematosus. Rheum. Dis. Clin. North Am., 20: 265.

Furukawa, F.; Kashiwara-Sawami, M.; Lyons, M.B. et al. (1990):

Binding of antibodies to the extractable nuclear antigens SS-
A/Ro and SS-B/La is induced on the surface of human
Leukocytes by ultraviolet light (U.V.L.).
Arthritis Rheum., 26: 78-100.

Hahan, B.H. and Ebling, F.M. (1989):

Suppression of murine lupus nephritis by administration of
an anti-idiotypic antibody to anti-DNA T. Immunol., 132-
187.

Hahan, B.H. and Ebling, F.M. (1993):

Idiotypes and idotype networks. In: Wallco DJ, Hahn BH
(eds): Dubois, lupus Erythematosus. Philadelphia, Lea,
Febiger, P. 137.

Halla, J.T. and Hardin, J.G. (1994):

Underrecognized postdosing reactions to methotrexate in patients with rheumatoid arthritis. *J. Rheumatol.*, 21: 1224.

Hammad, G.; Yousef, A.; Farouk, S. and Bakr, A. (1997):

Sister chromatid exchange in rheumatoid arthritis patients. *Benha M.J. Vol.*, 14: 567-77.

Hardin, J.A. and Thomas, J.O. (1993):

Antibodies to histones in systemic lupus erythematosus localization of prominent autoantibodies on histones H1 and H2 B. *Proc. Natl. Acad. Sci., USA*, 80-4710.

Harris, E.D.Jr. (1990):

Rheumatoid arthritis. Pathophysiology and implication for therapy. *N. Engl. J. Med.*, 322: 1277-1289.

Hebert, L.A.; Cosio, F.G.; Birmingham, D.J. et al. (1992):

The role of the complement system in renal injury semin. *Nephrol.*, 12: 408-27.

Hellmann, D.B.; Petri, M.; Whiting-O'Keefe (1993):

Fetal infection in systemic lupus erythematosus. *Medicine*, 6: 341.

Hex, Goronzy, J.J.; Zhong, W.; Xie, C. and Weyand, C.M. (1995):

VH3-21B cells escape from a state of tolerance in rheumatoid arthritis and secrete rheumatoid factor. *Molecular Med.*, 768-780.

Higley, H.; Persichitte, K.; Chu, S.; Waegell, W. and Black, C. (1994):

Immunocytochemical localization and serologic detection of transforming growth factor B1. Association with type I procollagen and inflammatory cell markers in diffuse and limited systemic sclerosis, morphea and Raynaud's Phenomenon. *Arthritis Rheum.*, 37: 278-83.

Hoffee, P.A. (1999):

Genetics Quick Look Medicine, USMLE Step I, A visually oriented review, Fence Greek Publishing. Madison, Connecticut. P., 356-401.

- Hohan, C. and Datta, S.K. (1996):**
Lupus key pathogenic mechanisms and contributing factors.
Clin. Immunol. Immunopath., 77: 209.
- Hulsey, M.; Goldstein, R.; Scully, L. et al. (1995):**
Anti-ribosomal P antibodies in systemic lupus erythematosus
a case-control study correlating hepatic and renal disease.
Clin. Immunol. Immunopathol., 74: 252.
- Isern, R.A.; Yaneva, M.; Weiner, E. et al. (1992):**
Autoantibodies in patients with primary pulmonary
autoantibodies in hypertension. Association with anti-Ku
Am. J. Med., 93: 307.
- Jones, S. (1996):**
Lower frequency of sister chromatid exchanges among
individuals with sporadic dysplastic nevi. Int. J. Biol.
Marker, 12: 3, 118-124.
- Kamel, O.W.; Van de Rijn, M.; Weiss, L.M. et al. (1993):**
Reversible lymphomas associated with Epstein-Barr virus
occurring during methotrexate therapy for rheumatoid
arthritis and dermatomyositis. N. Eng. J. Med., 328: 1317.
- Kerstens, P.J.S.M.; Boerbooms, A.M.T.; Jeurissen, M.E.C. et al.
(1993):**
Accelerated nodulosis during low-dose methotrexate therapy
for rheumatoid arthritis. An analysis of ten cases.
J. Rheumatol., 19: 867.
- Kitani, A.; Hara, M.; Hirose, T. et al. (1991):**
Elevated levels of endogenous IL-6 in SLE: A putative role
in pathogenesis. J. Immunol., 147: 117.
- Klinman, D.M. (1993):**
B-cell abnormalities characteristic of SLE. In: Wallace DJ,
Hahn BH (eds): DuBois' Lupus erythematosus. Philadelphia,
Lea and (γ) Tebiger, P. 77.
- Kremer, J.M.; Alarcon, G.S.; Lightfoot, R.W.Jr. et al. (1994):**
Methotrexate for rheumatoid arthritis suggested guidelines
for monitoring liver toxicity. Arthritis. Rheum., 37: 316.

Kurosaka, M. and Ziff, M. (1993):

Immunoelectron microscopic study of the distribution of T-cell subsets in rheumatoid synovium. *J. Exp. Med.*, 158: 1191-1210.

Lahita, R.G.; Bradlow, L.; Fishman, J. et al. (1982):

Estrogen metabolism in systemic lupus erythematosus patients and family members. *Arthritis Rheum.*, 25: 843.

Lai, J.S.; Kuo, H.W.; Liao, F.C. and Lien, C.H. (1998):

Short communication sister chromatid exchange induced by chromium compounds in human lymphocytes. *International Archives of occupational and Environmental Health. Abstract Volume*, 71, Issue 8: 550-553.

Lapez-Longo, F.J.; Rodriguez-Mohou, M.; Escalona, M. et al. (1994):

Heterogeneity of the anti-Ro (SS-A) response in rheumatic diseases. *J. Rheumatol.*, 21: 1450.

Le loire, O.; Croquette, M.F. and Houvenagel, E. (1993):

Chromosome anomalies in rheumatoid polyarthritis breakage level and study of breaking factor. *Rev. Rheum. Mal. Osteoartic*, Nov., 53.

Liao, H.X.; Haynes, B.F. (1995):

Role of adhesion molecules in the pathogenesis of rheumatoid arthritis. *Rheum. Dis. Clin. N. Am.*, 21: 715.

Linker-Israeli, M.; Quismorio, F.P.Jr.; Horwitz, D.A. et al. (1990):

CD8+ lymphocytes from patients with systemic lupus erythematosus sustain, rather than suppress, spontaneous polyclonal IgG production and synergize with CD4 cells to support autoantibody synthesis. *Arthritis. Rheum.*, 33: 1216.

Liszewski, J.E.; Spronk, P.E.; Bootsma, H. et al. (1995):

Changes in antibodies to C1a predict renal relapses in systemic lupus erythematosus. *Am. J. Kidney Dis.*, 26: 595-601.

Madden, D.R.; Gorga, J.C.; Strominger, I.L. et al. (1991):

The structure of HLA, B27 reveals nonamer self-peptides bound in an extended conformation. *Nature*, 353-321.

- Manilow, M.R.; Bardana, E.J.Jr.; Pirofsky, B. et al. (1982):**
Systemic lupus erythematosus like syndrome in monkeys fed alfalfa sprouts: Rde of a non protein amino acid.
Science, 216: 415.
- Maricq, H.R.; Weinrich, M.C.; Keil, J.E.; Smith, E.A.; Harper, F.E.; Fosal, E.J. (1989):**
Prevalence of scleroderma spectrum disorder in the general population of South Carolina. Arthritis Rheum., 32: 998-1006.
- Maricq, H.R.; Carpentier, P.H.; Weinrich, M.C.; Keil, J.E.; Franco, A.; Dronet, P. and Proncoto, C.M. (1993):**
Geographic variation in the prevalence of Raynaud's phenomenon: Charleston, SC, USA, Tarentaise, Savole France. J. Rheumatol., 20: 70-76.
- Masi, A.; Rodnan, G.P. and Medsger, T.A. (1980):**
Primilminory criteria a for classification of systemic sclerosis (scleroderma).
Arthritis Rheum., 23: 581-90.
- Maul, G.G.; French, B.T.; Vanvenrool, J.W.J. et al. (1994):**
Topo-isomeras I identified by scleroderma 70 antisera: Enviroment of topoisomerase I at the centromere in mouse mitotic cells before anaphase. Proc. Natl. Acad. Sci. USA, 83: 5145.
- McGrath, H.Jr. (1994):**
Ultraviolet-AJ irradiation decreases clinical disease activity and auto-antibodies in patients with SLE.
Clin. Exper. Rheumatol., 12: 129.
- Medsger, T.A.Jr. (1994):**
Epidemiology of systemic sclerosis.
Clin. Dermatol., 12: 207-216.
- Meller, R.F. and Young, T.D. (2001):**
Sister chromatid exchange. In: Emery's elements of medical genetics. Eleventh edition, Churchill Livingston, London, New York, P. 153-159.

Menton, D.N.; Hess, R.A. (1991):

The ultrastructure of collagen in the dermis of tight skin (TSK) mouse fibroblasts. *J. Invest. Dermatol.*, 74: 139.

Mery, M. and Rimoin, S. (1990):

Chromosomal structure in human.
Arch. Dermatol., 136: 321.

Morgan, A.B. and Crossen, E. (1981):

Frequencies of sister-chromatid exchanges in patient with connective tissue disease.
Mutat. Res. Aug., 162: 131-136.

Morgan, A.B. (1993):

6- mercaptopurine (6-MP) in the treatment of rheumatoid arthritis and related conditions. *Ann. Rheum. Dis.*, 24: 179-180.

Morris, P.; Shaman, J.; Ataya, M. et al. (1994):

An essential role for HLA-DM in antigen presentation by class II major histocompatibility molecules. *Nature*, 368: 551.

Mosrin, C. and Thuriaux, P. (1990):

The genetic of RNA polymerases in yeasts. *Curr Genet.*, 17: 367-373.

Nakajima, T.; Aono, H.; Hasunuma, T. et al. (1999):

Overgrowth of human synovial cells driven by the human T-cell leukemia virus type I tax gene. *J. Clin. Invest.*, 92: 186-193.

Natali, P.G. and Tan, E.M. (1993):

Experimental skin lesions in mice resembling.
Arthritis Rheum., 16: 579.

Needleman, B.W.; Wigley, F.M.; Stair, R.W. (1992):

Interleukin-1, inteleukin-2, interleukin-4, interleukin-6, tumor necrosis factor and interferon levels in sera of patient with scleroderma. *Arthritis Rheum.*, 53: 67.

- Nepom, G.T. (1992):**
Prediction of susceptibility to rheumatoid arthritis by human leukocyte antigen genotyping. *Rheum. Dis. Clin. North Am.*, 18: 785-792.
- Notman, D.D.; Kurata, N. and Tan, E.M. (1990):**
Profiles of antinuclear antibodies in systemic rheumatic disease. *Ann. Intern. Med.*, 83: 464.
- Palmer, R.G.; Dore, C.J.; Henderson, L. and Denman, A.M. (1985):**
Sister-chromatid exchange frequencies in fibroblast and lymphocytes of patients with systemic lupus erythematosus. *Mutat. Res.*, 125: 132.
- Palmer, R.G.; Dore, C.J. and Denman, A.M. (1986):**
Sister-chromatid exchange frequencies in lymphocytes of controls and patients with connective tissue diseases. *Mutat. Res.*, 113-120.
- Palmer, R.G. (1987):**
Sister chromatid exchange assessment of chromosome damage induced by cytotoxic drugs in patients with connective tissue diseases. *Lancet*, 1: 246-9.
- Palmer, R.G.; Smith-Burchnell, C.A.; Dore, C.J. and Denman, A.M. (1987):**
Sensitivity of lymphocytes from patients with systemic lupus erythematosus to the induction of sister chromatid exchanges but alkylating agents and bromodeoxy uridine. *Ann. Rheum. Dis.*, 110-113.
- Peng, S.L. and Craft, J. (1996):**
Antibodies to spliceosomal Sn RNAs. In Shoenfeld Y, Peter JB (eds.): *Textbook of autoantibodies*, Netherlands, Elsevier (in press), P. 131-146.
- Perry, P. and Wolff, S. (1974):**
New giemsa method for differential staining of sister chromatids. *Nature*, 251: 156-8.
- Pisetsky, D.S. (1992):**
The antiphospholipid antibody syndrome, pregnancy and hormonal therapy in SLE. *Rheum. Dis. Clin. NA*, 18: 437-545.
- Powell, M. (1994):**
Chromosomal abnormalities in patients with scleroderma and their relative. *Arthritis Rheumatism*, 3: 466-475.

- Prescott, R.J.; Freemont, A.J.; Jones, C.J.P.; Hyland, J. (1992):**
Sequential dermal microvascular and perivascular changes in the development of scleroderma. *J. Pathol.*, 166: 255-263.
- Randen, I.; Thompson, K.M.; Pascual, V. et al. (1992):**
Rheumatoid factor V genes from patients with rheumatoid arthritis are diverse and show evidence of an antigen-driven response. *Immunol. Rev.*, 128: 49-71.
- Rantappa Dahlquist, S. and Nordenson, I. (1996):**
Chromosomal changes in rheumatoid arthritis patient. *Clinical Rheumatology*, 15 (6): 584-9.
- Reeves, W.H.; Satoh, M.; Iingso, W. et al. (1994):**
Antibodies to DNA, DNA binding proteins and histones. *Rheum. Dis. Clin. North Am.*, 20-1.
- Reidenberg, M.M.; Drayer, D.E. (1992):**
Genetic regulation of drug metabolism and SLE. In: Lahita RG (ed) *SLE*. New York, Churchill Livingstone, P. 893.
- Reininger, L.; Rada, S.Z.; Kiewica, T.; Kosco, M. et al. (1992):**
Development of autoimmune disease in SCLD mice populated with long term in vitro proliferating (NZB x NZW) F1 pre-B cells. *J. Exp. Med.*, 176: 1639.
- Reveille, J.D.; Owerbach, Goldstein, R.; Iserh, R.A.; Arnett, F.A. (1992):**
Association of polar amino at position 26 of the HLADQB1 first domin with the anticentromere autoantibody response in systemic sclerosis. *J. Clin. Invest.*, 89: 1209-12.
- Robin, D.L.; Skilling, I.; Benisek, W.F.; Wistar, R.Jr. (1994):**
Estimation of the IgM rheumatoid factor secreted by rheumatoid synovial cells for human IgG subclasses. *Arthritis Rheum.*, 29: 722.
- Rodman, G.P.; Lipinski, E. and Luksick, J. (1993):**
Skin thickness and collagen content on progressive systemic sclerosis and localized scleroderma. *Arthritis Rheum.*, 22: 130-140.

- Rooney, S. and Czepulkowski, A. (1992):**
Cytological detection of mutocarcinogen expose by sister chromatid exchange. *Nature*, 258: 121-5.
- Rothfield, N.F. (1992):**
Autoantibodies in scleroderma. *Rheum. Dis. Clin. North Am.*, 18: 483.
- Rothschild, B.M.; Turner, K.R. and Deluca, M.A. (1993):**
Symmetrical erosive peripheral polyarthritis. *Science*, 241: 1498-1502.
- Salmon, J.E. (1993):**
Abnormalities in immune complex clearance and Fc receptor function. In: Wallace DJ, Hahn BH (eds.): *Dubois, Lupus Erythematosus Philadelphia Lea, Febiger*, P. 108.
- Sanchez-Guerrero, J.; Karlson, E.W.; Colditz, G.A. et al. (1994):**
Post menopausal hormones (PMA) used and the risk of developing systemic lupus erythematosus.
- Satoh, M.; Ajmoni, A.K.; Ogasawara, T. et al. (1994):**
Autoantibodies to RNA polymerase II are common in systemic lupus erythematosus and overlap syndrome. *J. Clin. Invest.*, 94: 1981.
- Schnobel, A.; Csernok, E.; Jsenberg, D.A. et al. (1995):**
Antineutrophil cytoplasmic antibodies in systemic lupus erythematosus. *Arthritis Rheum.*, 38: 633.
- Schultz, D.R. and Arnold, P.J. (1993):**
Heat shock (stress proteins and autoimmunity in rheumatic diseases semin. *Arthritis Rheum.*, 22: 357.
- Scofield, R.H.; Frank, M.B.; Neas, B.R. et al. (1994):**
Cooperative association of T-cell beta receptor and HLA-DQ alleles in the production of anti-RO in systemic lupus erythematosus. *Clin. Immunol. Pathol.*, 72: 335.
- Seibold, J.R.; Medsger, T.A.; Winkelstein, A. et al. (1996):**
Immune complexes in progressive scleroderma. *Arthritis Rheum.*, 25: 1167.

- Sesso, R.; Monteiro, M.; Sato, E. et al. (1994):**
A controlled trial of pulse cyclophosphamide versus pulse methyl prednisolone in severe lupus nephritis. *Lupus*, 3: 107.
- Sherer, S.; Abe, S. and Sasaki, M. (1983):**
Chromosomal breakage in collagen diseases. *Z. Rheumatol.* Mar. Apr., 39(3-4): 84-90.
- Silman, A.J. (1993):**
Scleroderma: Diseases definition and criteria. In: Silman AJ, Hochberg MC (eds): *Epidemiology of rheumatic disease*. Oxford, Oxford University Press, P. 192-219.
- Silver, R.M. (1994):**
The eosinophilia-myalgia syndrome. *Clin. Dermatol.*, 12: 457-465.
- Singh, R.R.; Ebling, F.M.; Sercarz, E.E. et al. (1995):**
Immune tolerance to autoantibody-derived peptides delays development of autoimmunity in SLE. *J. Clin. Invest.*, 96: 2990.
- Solomon, C. and Bobrow, M. (1995):**
Sister chromatid exchanges a sensitive assay of agents damaging human chromosomes. *Mutat. Res.*, 30: 273-280.
- Sontheimer, R.D.; Gilliam, J.N. (1992):**
Systemic lupus erythematosus and the skin. In: Lahita RG (ed): *S.L.E.*, 2nd ed. New York, Churchill Livingstone, PP. 657-681.
- Snell, G.D.; Dausset, I.; Nathenson, S.G. et al. (1996):**
Histocompatibility New York. Academic Press.
- Snmez, S.; Kaya, M.; Aktas, A.; Jkbal, M. and Senel, K. (1998):**
High frequency of sister chromatid exchanges in lymphocytes of the patients with Behcet's disease. *Mutat. Res.*, 397: 2, 235-238.
- Snowden, N.; Hay, E.; Holt, et al. (1993):**
Clinical course of patients with anti-RNA antibodies. *J. Rheumatol.*, 20: 1256.

- Stastny, P. (1978):**
Association of the B-cell alloantigen DRW4 with rheumatoid arthritis. *N. Engl. J. Med.*, 298: 869-871.
- Strand, V.; Steinberg, A.D. and Loeb, L.A. (1999):**
New Research of RA. *Archives of Internal Medicine*, 159: 2542-2450.
- Svejstrup, J.Q.; Wang, Z.; Feaver, W.J. et al. (1995):**
Different forms of TFIIF for transcription in DNA repair cell. *Science*, 80: 21-8.
- Takeuchi, M.; Ouellette, A.M. and Albert, E.D. (1993):**
Study of sister chromatid exchange in patient with systemic sclerosis. *Nature Med.*, 4: 448-457.
- Tan, E.M.; Cohen, A.S. et al. (1982):**
The 1982 revised criteria for classification of SLE. *Arthritis. Rheum.*, 25: 1271-1277.
- Tan, E.M.; Rodnan, G.P.; Garcia, J. et al. (1990):**
Diversity of antinuclear antibodies in progressive systemic sclerosis. Anti-centromere antibody and its relationship to CREST syndrome. *Arthritis Rheum.*, 23: 617.
- Tohda, H.; Horaguchi, K.; Takahashi, K.; Oikawa, A. and Matsushima, T. (1980):**
Epstein-Barr virus transformed human lymphoblastoid cells for study of sister chromatid exchanges and their evaluation as a test system. *Cancer Res.*, 40: 4775-4780.
- Total, N.; Garry, R.F. and Schur, P.H. (1990):**
A conserved idotype and antibodies to retroviral proteins in SLE. *J. Clin. Invest.*, 85: 1866.
- Topfer, F.; Gordon, T.; McCluskey, J. et al. (1995):**
Intra and intermolecular spreading of autoimmunity involving the nuclear self-antigens La (SS-B) and Ro (SS-A). *Proc. Natl. Acad. Sci. USA*, 92: 875.
- Toth, R.; Allegretta, M.; Kim, Y.; Amento, A. (1995):**
T-cells derived from the skin of patient with systemic scleroderma preferentially utilize the T-cell receptor VB3. *Arthritis Rheum.*, 38: S252.

Tsai, D.E. and Koene, J.D. (1993):

In vitro selection of RNA epitopes using autoimmune patient serum. *J. Immunol.*, 150: 1137.

Tsao, B.P. and Hahn, B.H. (1992):

Autoimmunity and tolerance in Ig-transgenic mice: Murine SLE as a model to study B-cell activation. *Immunol. Today*, 13: 431.

Tsao, B.P. and Hahn, B.H. (1994):

Autoimmunity and tolerance in Ig-transgenic mice: Murine SLE as a model to study B-cell tolerance. *Int. Rev. Immunol.*, 11: 305.

Van Zeben, D.; Rook, G.A.W.; Hazes, J.M.W.; Zwinderman, A.H.; Zhang, Y. and Breedveld, F.C. (1994):

Early agalactosylation of IgG is associated with a more progressive disease course in patients with rheumatoid arthritis. Results of a follow-up study. *Br. J. Rheumatol.*, 33: 37.

Varonos, S. (1993):

Chromosomal breaks in connective tissue diseases. *Z. Rheumatol.*, AP; 39(3-4): 84-90.

Walser-Kuntz, D.R.; Weyand, C.M.; Weaver, A.J.; O'Fallon, W.M. and Goronzy, J.J. (1995):

Mechanisms underlying the formation of the T-cell receptor repertoire in rheumatoid arthritis. *Immunity*, 2: 597-605.

Weinblatt, M.E. (1985):

Toxicity of low dose methotrexate in rheumatoid arthritis. *J. Rheumatol.*, 12 (Suppl. 12): 35.

Weinblatt, M.E.; Kajian, H.; Germain, B.F. et al. (1990):

Low dose methotrexate compared with auranofin in adult rheumatoid arthritis. A thirty-six week, double-blind trial. *Arthritis. Rheum.*, 33: 330.

Weiner, E.S.; Hildebrandt, S.; Senecal, J.L. et al. (1993):

Prognostic significance of anticentromere antibodies and anti-topoisomerase I antibodies in Raynaud's disease. *Arthritis Rheum.*, 34: 68.

Weyand, C.M.; Hicok, K.C.; Conn, D. and Gornzy (1992):

The influence of HLA-DRB1 genes on disease severity in rheumatoid arthritis. *Ann. Intern. Med.*, 117: 801-806.

Weyand, C.M.; Hicok, K.C.; Conn, D. and Goronzy, J.J. (1995):

Correlation between disease phenotype and genetic heterogeneity in rheumatoid arthritis. *J. Clin. Invest.*, 95: 2120-2126.

White, B. (1996):

Pathogenesis: Immune aspect. In: Clements PJ Furst DE (eds): systemic sclerosis. Baltimore, Williams & Wilkins, PP. 229-250.

Wigley, F.M. (1995):

Is Raynaud's isolated phenomenon: *Contemporary Intern. Med.*, 7: 43-56.

Wilmer, J.L.; Colvin, O.M.; Bloom, S.E. (1992):

Cytogenetic mechanisms in the selective toxicity of cyclophosphamide analogs and metabolites towards avian embryonic B-lymphocytes in vivo. *Mutat. Res.*, 268: 115-130.

Winchester, R. (1994):

The molecular basis of susceptibility to rheumatoid arthritis. *Adv. Immunol.*, 56: 389-466.

Winkelstein, J.A. and Sullivan, K.E. (1995):

Complement deficiencies in Rich RR, ed. *Clinical Immunology. Principles and Practice* St. Louis, Mosby, 662-76.

Wolf, A.; Smolen, J.S. and Kovac, R. (1994):

SCE frequency in lymphocytes of systemic lupus erythematosus. *Mutat. Res. Sep.*, 128(1): 167-69.

Woods, C.G.; Leversa, J.; Rogers, J. et al. (1995):

Severe intrauterine growth retardation and mitomycin sensitivity; a new chromosome breakage syndrome. *J. Med. Genet.*, 34: 203-6.

Wooley, P.H.; Sud, S.; Langendorfer, A. et al. (1995):

T-cells infiltrating the skin of TSK-2 scleroderma-like mice exhibit T-cell receptor bias. *Arthritis Rheum.*, 38: S253.

Yamana, K.; Takasaki, Y.; Nishida, Y. et al. (1992):

Detection and Quantification of anti-Ki antibodies by enzyme linked immunosorbent assay using recombinant Ki antigen. *Arthritis. Rheum.*, 35: 667.

Yanagida, M. (2000):

Cell cycle mechanism of sister chromatid separation, role of cut-1 separin and cut-2 securin. *Genes Cells*, Jan., 5(1): 1-8
ISSN: 1356: 9597.

Youngblood, K.; Fruchter, L.; Ding, G.; Lope, Z. and Davidson, A. (1994):

Rheumatoid factors from the peripheral blood of two patients with rheumatoid arthritis are genetically heterogeneous and somatically mutated. *J. Clin. Invest.*, 93: 852.

Yurovsky, V.V.; Sutton, P.A.; Schulze, D.H.; Wise, R.A.; White, B. (1994):

Expansion of selected V81 + Y/8T cells in systemic sclerosis patient. *J. Immunol.*, 153: 881-91.

Yurovsky, V.V.; Wigley, F.M.; Wise, R.A.; White, B. (1995):

T-cell repertoire in the lungs of patients with scleroderma. *Arthritis. Rheum.*, 38: S253.