

References

- 20th Bethesda Conference: insurability and employability of the patient with ischemic heart disease. October 3–4, 1988, Bethesda, Maryland. *J Am Coll Cardiol*. 1989; 14: 1003–1044.
- Ades PA, Huang D, Weaver SO. Cardiac rehabilitation participation predicts lower rehospitalization costs. *Am Heart J*. 1992; 123: 916–921.
- Ades PA, Pashkow FJ, Nestor JR. Cost-effectiveness of cardiac rehabilitation after myocardial infarction. *J Cardiopulm Rehabil*. 1997; 17: 222–231.
- Ades PA, Coello CE: Effects of exercise and cardiac rehabilitation on cardiovascular outcomes. *Med Clin North Am* 2000; 84(1): 251-65.
- Ades PA. Cardiac rehabilitation and secondary prevention of coronary heart disease. *N Engl J Med*. 2001; 345: 892–902.
- Agency for Healthcare Research and Quality (AHRQ) Technology Assessment Program. Randomized trials of secondary prevention programs in coronary artery disease: a systematic review, Rockville, Maryland 2005.
- Agmon, Y, Khandheria, BK, Meissner, I, et al. Independent association of high blood pressure and aortic atherosclerosis: A population-based study. *Circulation* 2000; 102:2087.
- Al Suwaidi, J, Higano, ST, Holmes, DR Jr, et al. Obesity is independently associated with coronary endothelial dysfunction in patients with normal or mildly diseased coronary arteries. *J Am Coll Cardiol* 2001; 37:1523.
- Allen JK, Scott LB, Stewart KJ, Young DR. Disparities in women's referral to and enrollment in outpatient cardiac rehabilitation. *J Gen Intern Med*. 2004; 19: 747–753.
- Ambrosetti M, Salerno M, Ageno W, et al: Is physical training contraindicated in patients with deep vein thrombosis during cardiac rehabilitation? *Monaldi Arch Chest Dis* 2005 Mar; 64(1): 24-6.

References

- Ambrosetti M, Salerno M, Laporta A, Pedretti RF: Metabolic syndrome in patients with intermittent claudication referred to vascular rehabilitation. *Int Angiol* 2006 Mar; 25(1): 14-7.
- American Association of Cardiovascular and Pulmonary Rehabilitation. Guidelines for Cardiac Rehabilitation and Secondary Prevention Programs. 4th ed. Champaign, IL: Human Kinetics; 2004.
- American Heart Association: Cardiac rehabilitation programs. A statement for healthcare professionals from the American Heart Association. *Circulation* 1994 Sep; 90(3): 1602-10.
- Bader DS, Maguire TE, Spahn CM, et al: Clinical profile and outcomes of obese patients in cardiac rehabilitation stratified according to National Heart, Lung, and Blood Institute criteria. *J Cardiopulm Rehabil* 2001 Jul-Aug; 21(4): 210-7.
- Balady G, Ades PA, Comoss P, et al. Core components of cardiac rehabilitation/secondary prevention programs: a statement for healthcare professionals from the American Heart Association and the American Association of Cardiovascular and Pulmonary Rehabilitation Writing Group. *Circulation* 2000;102:1069-1073.
- Barker, DJ, Osmond, C, Simmonds, SJ, Wield, GA. The relation of small head circumference and thinness at birth to death from cardiovascular disease in adult life. *BMJ* 1993; 306:422.
- Benetos, A, Zureik, M, Morcet, J, et al. A decrease in diastolic blood pressure combined with an increase in systolic blood pressure is associated with a higher cardiovascular mortality in men. *J Am Coll Cardiol* 2000; 35:673.
- Benjamin, EJ, Levy, D. Why is left ventricular hypertrophy so predictive of morbidity and mortality?. *Am J Med Sci* 1999; 317:168.
- Berkhuisen MA, Nieuwland W, Buunk6 BP, Sanderma2 R, Viersma4 JR, Rispens P: Effect of high versus low frequency exercise training in multidisciplinary cardiac rehabilitation on health-related quality of life. *J Cardiopulmonary Rehabil* 1999; 19: 22-28.

References

- Berkman LF, Blumenthal J, Burg M, Carney RM, Catellier D, Cowan MJ, Czajkowski SM, DeBusk R, Hosking J, Jaffe A, Kaufmann PG, Mitchell P, Norman J, Powell LH, Raczynski JM, Schneiderman N; Enhancing Recovery in Coronary Heart Disease Patients Investigators (ENRICH). Effects of treating depression and low perceived social support on clinical events after myocardial infarction: the Enhancing Recovery in Coronary Heart Disease Patients (ENRICH) Randomized Trial. *JAMA*. 2003; 289: 3106–3116.
- Bethell HJ: Cardiac Rehabilitation: from Hellerstein to the millennium. *Int J Clin Prac* 2000; 54(3): 92-97.
- Blake, GJ, Dada, N, Fox, JC, et al. A prospective evaluation of lipoprotein-associated phospholipase A(2) levels and the risk of future cardiovascular events in women. *J Am Coll Cardiol* 2001; 38:1302.
- Blankenberg, S, Rupprecht, HJ, Bickel, C, et al. Circulating cell adhesion molecules and death in patients with coronary artery disease. *Circulation* 2001; 104:1336.
- Boffetta, P, Garfinkel, L. Alcohol drinking and mortality among men enrolled in an American Cancer Society Prospective Study. *Epidemiology* 1990; 1:342.
- Bolli R. The late phase of preconditioning. *Circ Res*. 2000; 87: 972–983.
- Boushey, CJ, Beresford, SAS, Omenn, GS, et al. A quantitative assessment of plasma homocysteine as a risk factor for vascular disease: Probable benefits of increasing folic acid intakes. *JAMA* 1995; 274:1049.
- Brochu M, Poehlman ET, Savage P, et al: Modest effects of exercise training alone on coronary risk factors and body composition in coronary patients. *J Cardiopulm Rehabil* 2000; 20(3): 180-188.
- Brown, DW, Giles, WH, Croft, JB. Left ventricular hypertrophy as a predictor of coronary heart disease mortality and the effect of hypertension. *Am Heart J* 2000; 140:848.

References

- Calle, EE, Thun, MJ, Petrelli, JM, et al. Body-mass index and mortality in a prospective cohort of U.S. adults [see comments]. *N Engl J Med* 1999; 341:1097.
- Camargo, CA, Stampfer, MJ, Glynn, RJ, et al. Prospective study of moderate alcohol consumption and risk of peripheral arterial disease in US male physicians. *Circulation* 1997; 95:577.
- Cannistra LB, Balady GJ, O'Malley CJ, et al: Comparison of the clinical profile and outcome of women and men in cardiac rehabilitation. *Am J Cardiol* 1992 May 15; 69(16): 1274-9.
- Cardiac rehabilitation programs: a statement for healthcare professionals from the American Heart Association. *Circulation*. 1994; 90: 1602–1610.
- Carlson JJ, Johnson JA, Franklin BA, VanderLaan RL: Program participation, exercise adherence, cardiovascular outcomes, and program cost of traditional versus modified cardiac rehabilitation. *Am J Cardiol* 2000; 86(1): 17-23.
- Carrel T, Mohaesi P: Optimal timing of rehabilitation after cardiac surgery: the surgeon's view. *Eur Heart J* 1998; 19 Suppl O: O38-O41.
- Church TS, Barlow CE, Earnest CP, Kampert JB, Priest EL, Blair SN. Association between cardiorespiratory fitness and C-reactive protein in men. *Arterioscler Thromb Vasc Biol*. 2002; 22: 1869–1876.
- Church TS, Lavie CJ, Milani RV, Kirby GS. Improvements in blood rheology after cardiac rehabilitation and exercise training in patients with coronary heart disease. *Am Heart J*. 2002; 143: 349–355.
- Clark, AM, Hartling, L, Vandermeer, B, McAlister, FA. Meta-analysis: secondary prevention programs for patients with coronary artery disease. *Ann Intern Med* 2005; 143:659.
- Collins, R, Peto, R, MacMahon, S, et al. Blood pressure, stroke, and coronary heart disease. part 2, short-term reductions in blood pressure: overview of randomized drug trials in their epidemiological context. *Lancet* 1990; 335:827.

References

- Cooper, R, Cutler, J, Desvigne-Nickens, P, et al. Trends and disparities in coronary heart disease, stroke, and other cardiovascular diseases in the United States: Findings of the National Conference on Cardiovascular Disease Prevention. *Circulation* 2000; 102:3137.
- Culleton, BF, Larson, MG, Kannel, WB, Levy, D. Serum uric acid and risk for cardiovascular disease and death: The Framingham Heart Study. *Ann Intern Med* 1999; 131:7.
- Danesh, J, Whincup, P, Walker, M, et al. Fibrin d-dimer and coronary heart disease: prospective study and meta-analysis. *Circulation* 2001; 103:2323.
- Davis, PH, Dawson, JD, Riley, WA, Lauer, RM. Carotid intimal-medial thickness is related to cardiovascular risk factors measured from childhood through middle age: the Muscatine study. *Circulation* 2001; 104:2815.
- De Bacquer, D, De Backer, G, Kornitzer, M, et al. Prognostic value of ischemic electrocardiographic findings for cardiovascular mortality in men and women. *J Am Coll Cardiol* 1998; 32:680.
- del Sol, AI, Moons, KG, Hollander, M, et al. Is carotid intima-media thickness useful in cardiovascular disease risk assessment? The Rotterdam study. *Stroke* 2001; 32:1532.
- Dennis C: Rehabilitation of patients with coronary artery disease. In: Braunwald E, ed. *Heart Disease: A Textbook of Cardiovascular Medicine*. Philadelphia, PA: WB Saunders Company; 1997: 1392-1403.
- Denollet J, Brutsaert DL. Enhancing emotional well-being by comprehensive rehabilitation in patients with coronary heart disease. *Eur Heart J* 1995;16:1070-1078.
- Diez Roux, AV, Merkin, SS, Arnett, D, et al. Neighborhood of residence and incidence of coronary heart disease. *N Engl J Med* 2001; 345:99.

References

- Dimmeler S, Zeiher AM. Exercise and cardiovascular health: get active to "AKTivate" your endothelial nitric oxide synthase. *Circulation*. 2003; 107: 3118–3120.
- Downs, JR, Clearfield, M, Weis, S, et al for the AFCAPS/TexCAPS Research Group. Primary prevention of acute coronary events with lovastatin in men and women with average cholesterol levels: Results of AFCAPS/TexCAPS. *JAMA* 1998; 279:1615.
- Duffy, SJ, Biegelsen, ES, Holbrook, M, et al. Iron chelation improves endothelial function in patients with coronary artery disease. *Circulation* 2001; 103:2799.
- Dugmore LD, Tipson RJ, Phillips MH, et al: Changes in cardiorespiratory fitness, psychological well-being, quality of life and vocational status following a twelve-month cardiac exercise rehabilitation program. *Heart* 1999; 81(4): 359-366.
- Effects of estrogen or estrogen/progestin regimens on heart disease risk factors in postmenopausal women. The Postmenopausal Estrogen/Progestin Interventions (PEPI) Trial. The Writing Group for the PEPI Trial. *JAMA* 1995; 273:199.
- El-Khairi, L, Ueland, PM, Refsum, H, et al. Plasma total cysteine as a risk factor for vascular disease: the European Concerted Action Project. *Circulation* 2001; 103:2544.
- Evenson KR, Rosamond WD, Luepker RV. Predictors of outpatient cardiac rehabilitation utilization: the Minnesota Heart Surgery Registry. *J Cardiopulm Rehabil* 1998;18:192-198.
- Executive Summary of the Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). *JAMA* 2001; 285:2486.
- Fagard RH. Exercise characteristics and the blood pressure response to dynamic physical training. *Med Sci Sport Exerc*. 2001; 33: S484–S492.

References

- Fang, J, Alderman, MH. Serum uric acid and cardiovascular mortality the NHANES I epidemiologic follow-up study, 1971-1992. National Health and Nutrition Examination Survey. JAMA 2000; 283:2404.
- Ferrara N, Corbi G, Bosimini E, et al: Cardiac rehabilitation in the elderly: patient selection and outcomes. Am J Geriatr Cardiol 2006 Jan-Feb; 15(1): 22-7.
- Flaker GC, Singh VN: Prevention of myocardial reinfarction. Recommendations based on results of drug trials. Postgrad Med 1993; 94(6): 94-104.
- Fletcher GF: Current status of cardiac rehabilitation. Am Fam Physician 1998; 58(8): 1778-1782.
- Franklin BA, Kahn JK, Gordon NF, Bonow RO. A cardioprotective "polypill"? Independent and additive benefits of lifestyle modification. Am J Cardiol. 2004; 94: 162–166.
- Franklin, SS, Khan, SA, Wong, ND, et al. Is pulse pressure useful in predicting risk for coronary heart disease? The Framingham Heart Study. Circulation 1999; 100:354.
- Franklin, SS, Larson, MG, Khan, SA, et al. Does the relation of blood pressure to coronary heart disease risk change with aging?: the Framingham Heart Study. Circulation 2001; 103:1245.
- Friedman DB, Williams AN, Levine BD. Compliance and efficacy of cardiac rehabilitation and risk factor modification in the medically indigent. Am J Cardiol 1997;79:281-285.
- Fuchs, CS, Stampfer, MJ, Colditz, GA, et al. Alcohol consumption and mortality among women. N Engl J Med 1995; 332:1245.
- Fuchs, CS, Stampfer, MJ, Colditz, GA, et al. Alcohol consumption and mortality among women. N Engl J Med 1995; 332:1245.
- Gardin, JM, McClelland, R, Kitzman, D, et al. M-mode echocardiographic predictors of six- to seven-year incidence of coronary heart disease, stroke, congestive heart failure, and mortality

- in an elderly cohort (the Cardiovascular Health Study). *Am J Cardiol* 2001; 87:1051.
- Genest, JJ, Martin-Munley, SS, McNamara, JR, et al. Familial lipoprotein disorders in patients with premature coronary artery disease. *Circulation* 1992; 85:2025.
 - Gerstein, HC, Pais, P, Pogue, J, et al. Relationship of glucose and insulin level to the risk of myocardial infarction: A case-control study. *J Am Coll Cardiol* 1999; 33:612.
 - Giannuzzi, P, Temporelli, L, Corra, U, et al, for the ELVD Study Group. Attenuation of unfavorable remodeling by exercise training in postinfarction patients with left ventricular dysfunction. Results of the Exercise in Left Ventricular Dysfunction (ELVD) Trial. *Circulation* 1997; 96:1790.
 - Gohlke H, Gohlke-Barwolf C: Cardiac rehabilitation. *Eur Heart J* 1998; 19(7): 1004-1010.
 - Gordon NF, Haskell WL: Comprehensive cardiovascular disease risk reduction in a cardiac rehabilitation setting. *Am J Cardiol* 1997; 80(8B): 69H-73H.
 - Greenland, P, Daviglus, ML, Dyer, AR, et al. Resting heart rate is a risk factor for cardiovascular and noncardiovascular mortality: the Chicago Heart Association Detection Project in Industry. *Am J Epidemiol* 1999; 149:853.
 - Hamalainen H, Luurila OJ, Kallio V, et al: Long-term reduction in sudden deaths after a multifactorial intervention programme in patients with myocardial infarction: 10-year results of a controlled investigation. *Eur Heart J* 1989 Jan; 10(1): 55-62.
 - Hambrecht R, Adams V, Erbs S, Linke A, Krankel N, Shu Y, Baither Y, Gielen S, Thiele H, Gummert JF, Mohr FW, Schuler G. Regular physical activity improves endothelial function in patients with coronary artery disease by increasing phosphorylation of endothelial nitric oxide synthase. *Circulation*. 2003; 107: 3152–3158.

- Harris, TB, Cook, EF, Kannel, WB, et al. Proportional hazards analysis of risk factors for coronary heart disease in individuals aged 65 or older. *J Am Geriatr Soc* 1988; 36:1023.
- Haskell WL, Alderman EL, Fair JM, Maron DJ, Mackey SF, Superko HR, Williams PT, Johnstone IM, Champagne MA, Krauss RM, et al. Effects of intensive multiple risk factor reduction on coronary atherosclerosis and clinical cardiac events in men and women with coronary artery disease. The Stanford Coronary Risk Intervention Prevention Project (SCRIP). *Circulation*. 1994; 89: 975–990.
- Hebert, PR, Moser, M, Mayer, J, et al. Recent evidence on drug therapy of mild to moderate hypertension and decreased incidence of coronary heart disease. *Arch Intern Med* 1993; 153:578.
- Hedbäck, J. Perk and P. Wodlin, Long-term reduction of cardiac mortality after myocardial infarction: 10-year results of a comprehensive rehabilitation program. *Eur Heart J* 14 (1993), pp. 831–835.
- Heitzer, T, Schlinzig, T, Krohn, K, et al. Endothelial dysfunction, oxidative stress, and risk of cardiovascular events in patients with coronary artery disease. *Circulation* 2001; 104:2673.
- Hubert, HB, Feinleib, M, McNamara, PM, et al. Obesity as an independent risk factor for cardiovascular disease: A 26-year follow-up of participants in the Framingham Heart Study. *Circulation* 1983; 67:968.
- Hubinette, A, Cnattingius, S, Ekbom, A, et al. Birthweight, early environment, and genetics: a study of twins discordant for acute myocardial infarction. *Lancet* 2001; 357:1997.
- Hung, J, Whitford, EG, Parsons, RW, et al. Association of sleep apnea with myocardial infarction in men. *Lancet* 1990; 336:261.
- Hunt, ME, O'Malley, PG, Vernalis, MN, et al. C-reactive protein is not associated with the presence or extent of calcified subclinical atherosclerosis. *Am Heart J* 2001; 141:206.

- Iellamo F, Legramante JM, Massaro MA, Raimondi G, Galante A. Effects of residential exercise training on baroreflex sensitivity and heart rate variability in patients with coronary artery disease: a randomized, controlled study. *Circulation*. 2000; 102: 2588–2592.
- Inoue, T, Uchida, T, Kamishirado, H, et al. Antibody against oxidized low density lipoprotein may predict progression or regression of atherosclerotic coronary artery disease. *J Am Coll Cardiol* 2001; 37:1871.
- Jakicic JM, Clark K, Coleman E, Donnelly JE, Foreyt J, Melanson E, Volek J, Volpe SL; American College of Sports Medicine. American College of Sports Medicine position stand: appropriate intervention strategies for weight loss and prevention of weight regain for adults. *Med Sci Sports Exerc*. 2001; 33: 2145–2156.
- Jarvisalo, MJ, Jartti, L, Nanto-Salonen, K, et al. Increased Aortic Intima-Media Thickness: A Marker of Preclinical Atherosclerosis in High-Risk Children. *Circulation* 2001; 104:2943.
- Jolliffe JA, Rees K, Taylor RS, Thompson D, Oldridge N, Ebrahim S. Exercise-based rehabilitation for coronary heart disease. *Cochrane Database Syst Rev*. 2001; CD001800.
- Jolliffe, JA, Rees, K, Taylor, RF, et al. Exercise-based rehabilitation for coronary heart disease. *Cochrane Database Syst Rev* 2005; 2.
- Jousilahti, P, Vartiainen, E, Tuomilehto, J, et al. Sex, age, cardiovascular risk factors, and coronary heart disease: A prospective follow-up study of 14,786 middle-aged men and women in Finland. *Circulation* 1999; 99:1165.
- Joyner MJ. Effect of exercise on arterial compliance. *Circulation*. 2000; 102: 1214–1215.
- Juan, SH, Lee, TS, Tseng, KW, et al. Adenovirus-mediated heme oxygenase-1 gene transfer inhibits the development of atherosclerosis in apolipoprotein e-deficient mice. *Circulation* 2001; 104:1519.

- Kallio V, Hamalainen H, Hakkila J, Luurila OJ: Reduction in sudden deaths by a multifactorial intervention programme after acute myocardial infarction. *Lancet* 1979 Nov 24; 2(8152): 1091-4.
- Kannel, WB, Gordon ,T, Schwartz, MJ. Systolic versus diastolic blood pressure and risk of coronary heart disease: The Framingham Study. *Am J Cardiol* 1971; 27:335.
- Kannel, WB, McGee, DL. Diabetes and cardiovascular risk factors: The Framingham Study. *Circulation* 1979; 59:8.
- Kannel, WB, McGee, DL. Diabetes and glucose tolerance as risk factors for cardiovascular disease: The Framingham Study. *Diabetes Care* 1979; 2:120.
- Kannel, WB. Left ventricular hypertrophy as a risk factor: the Framingham experience. *J Hypertens Suppl* 1991; 9:S3.
- Kobashigawa JA, Leaf DA, Lee N, et al: A controlled trial of exercise rehabilitation after heart transplantation. *N Engl J Med* 1999; 340(4): 272-277.
- Kobashigawa, D.A. Leaf, N. Lee et al., A controlled trial of exercise rehabilitation after heart transplantation. *N Engl J Med* 340 (1999), pp. 272–277.
- Krauss, RM, Winston, M. Obesity: Impact on cardiovascular disease. *Circulation* 1998; 98:1472.
- Kris-Etherton P, Eckel RH, Howard BV, St. Jeor S, Bazzarre TL; Nutrition Committee Population Science Committee and Clinical Science Committee of the American Heart Association. AHA Science Advisory: Lyon Diet Heart Study: benefits of a Mediterranean-style, National Cholesterol Education Program/American Heart Association Step I dietary pattern on cardiovascular disease. *Circulation*. 2001; 103: 1823–1825.
- Kris-Etherton P, Harris WS, Appel LJ. American Heart Association. Nutrition Committee. Fish consumption, fish oil, omega-3 fatty acids, and cardiovascular disease. *Circulation*. 2002; 106: 2747–2757.

References

- Kristal-Boneh, E, Silber, H, Harari, G, Froom, P. The association of resting heart rate with cardiovascular, cancer and all-cause mortality. Eight year follow-up of 3527 male israeli employees (the CORDIS study). *Eur Heart J* 2000; 21:116.
- Kutner NG, Zhang R, Huang Y, Herzog CA: Cardiac rehabilitation and survival of dialysis patients after coronary bypass. *J Am Soc Nephrol* 2006 Apr; 17(4): 1175-80.
- Lamarche, B, Moorjani, S, Lupien, PJ, et al. Apolipoprotein A-I and B levels and the risk of ischemic heart disease during a five-year follow-up of men in the Quebec cardiovascular study. *Circulation* 1996; 94:273.
- Lamm G, Denolin H, Dorossiev D, Pisa Z: Rehabilitation and secondary prevention of patients after acute myocardial infarction. WHO collaborative study. *Adv Cardiol* 1982; 31: 107-11.
- LaMonde MJ, Durstine JL, Yanowitz FG, Lim T, DuBose KD, Davis P, Ainsworth BE. Cardiorespiratory fitness and C-reactive protein among a tri-ethnic sample of women. *Circulation*. 2002; 106: 403–406.
- LaMonte, MJ, Eisenman, PA, Adams, TD, et al. Cardiorespiratory fitness and coronary heart disease risk factors: the LDS hospital fitness institute cohort. *Circulation* 2000; 102:1623.
- Lauer, MS, Okin, PM, Larson, MG, et al. Impaired heart rate response to graded exercise. Prognostic implications of chronotropic incompetence in the Framingham Heart Study. *Circulation* 1996; 93:1520.
- Laukkanen, JA, Lakka, TA, Rauramaa, R, et al. Cardiovascular fitness as a predictor of mortality in men. *Arch Intern Med* 2001; 161:825.
- Lavie CJ, Milani RV, Cassidy MM, Gilliland YE: Benefits of Cardiac Rehabilitation and Exercise Training in Older Persons. *Am J Geriatr Cardiol* 1995 Jul; 4(4): 42-48.

References

- Lavie CJ, Milani RV: Cardiac rehabilitation and preventive cardiology in the elderly. *Cardiol Clin* 1999; 17: 233-242.
- Lavie CJ, Milani RV: Cardiac rehabilitation, exercise training, and psychosocial risk factors. *J Am Coll Cardiol* 2006 Jan 3; 47(1): 212; author reply 212-3.
- Lavie CJ, Milani RV: Effects of cardiac rehabilitation programs on exercise capacity, coronary risk factors, behavioral characteristics, and quality of life in a large elderly cohort. *Am J Cardiol* 1995; 76(3): 177-9.
- Lekakis, JP, Papamichael, CM, Cimponeriu, AT, et al. Atherosclerotic changes of extracoronary arteries are associated with the extent of coronary atherosclerosis. *Am J Cardiol* 2000; 85:949.
- Limacher MC: Exercise and cardiac rehabilitation in women. *Cardiol Review* 1998; 6(4): 240-248.
- Linke A, Erbs S, Hambrecht R: Exercise and the coronary circulation-alterations and adaptations in coronary artery disease. *Prog Cardiovasc Dis* 2006 Jan-Feb; 48(4): 270-84.
- Lloyd-Jones, DM, Larson, MG, Beiser, A, et al. Lifetime risk of developing coronary heart disease. *Lancet* 1999; 353:89.
- Maier W, Meier B: Interventional cardiology in perspective: impact on cardiac rehabilitation. *Eur Heart J* 1998 Nov; 19 Suppl O: O24-8.
- Malik, I, Danesh, J, Whincup, P, et al. Soluble adhesion molecules and prediction of coronary heart disease: a prospective study and meta-analysis. *Lancet* 2001; 358:971.
- Manson, JE, Colditz, GA, Stampfer, MJ, et al. A prospective study of obesity and risk of coronary heart disease in women. *N Engl J Med* 1990; 322:882.
- Mark DB, Lauer MS. Exercise capacity: the prognostic variable that doesn't get enough respect. *Circulation*. 2003; 108: 1534–1536.

References

- Mark DB, Naylor CD, Hlatky MA, et al: Use of medical resources and quality of life after acute myocardial infarction in Canada and the United States. *N Engl J Med* 1994 Oct 27; 331(17): 1130-5.
- Maxwell, AJ, Bruinsma, KA. Uric acid is closely linked to vascular nitric oxide activity. Evidence for mechanism of association with cardiovascular disease. *J Am Coll Cardiol* 2001; 38:1850.
- Meyers, DG. The iron hypothesis: does iron play a role in atherosclerosis?. *Transfusion* 2000; 40:1023.
- Miura, K, Daviglius, ML, Dyer, AR, et al. Relationship of blood pressure to 25-year mortality due to coronary heart disease, cardiovascular diseases, and all causes in young adult men. The Chicago Heart Association detection project in industry.
- Mochari H, Lee JR, Kligfield P, Mosca L: Ethnic differences in barriers and referral to cardiac rehabilitation among women hospitalized with coronary heart disease. *Prev Cardiol* 2006; 9(1): 8-13.
- Moore SM, Charvat JM, Gordon NH, et al: Effects of a CHANGE Intervention to Increase Exercise Maintenance Following Cardiac Events. *Ann Behav Med* 2006 Feb; 31(1): 53-62.
- Morrison, JA, Jacobsen, DW, Sprecher, DL, et al. Serum glutathione in adolescent males predicts parental coronary heart disease. *Circulation* 1999; 100:2244.
- Mukherjee D, Fang J, Chetcuti S, Moscucci M, Kline-Rogers E, Eagle KA. Impact of combination evidence-based medical therapy on mortality in patients with acute coronary syndromes. *Circulation*. 2004; 109: 745–749. [[Abstract/Free Full Text](#)]
- Mulcahy R. Twenty years of cardiac rehabilitation in Europe: a reappraisal. *Eur Heart J* 1991;12:92-3.
- Mustafa, A, Nityanand, S, Berglund, L, et al. Circulating immune complexes in 50-year-Old men as a strong and independent risk factor for myocardial infarction. *Circulation* 2000; 102:2576.

References

- Nanette K and Wenger M, Efficacy of cardiac rehabilitation after myocardial infarction UpToDate, 2006.
- National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III) final report. *Circulation*. 2002; 106: 3143–3421.
- Njolstad, I, Arnesen, E, Lund-Larsen, PG. Smoking, serum lipids, blood pressure, and sex differences in myocardial infarction. A 12-year follow-up of the Finnmark Study. *Circulation* 1996; 93:450.
- O'Connor GT, Buring JE, Yusuf S, Goldhaber SZ, Olmstead EM, Paffenbarger RS Jr, Hennekens CH. An overview of randomized trials of rehabilitation with exercise after myocardial infarction. *Circulation*. 1989; 80: 234–244.[\[Abstract/Free Full Text\]](#)
- Oberman, G.F. Fletcher, J. Lee et al., Efficacy of high-intensity exercise training on left ventricular ejection fraction in men with coronary artery disease (the training level comparison study). *Am J Cardiol* 76 (1995), pp. 643–647.
- Oldridge N, Guyatt G, Crowe J, et al: Goal attainment in a randomized controlled trial of rehabilitation after myocardial infarction. *J Cardiopulm Rehabil* 1999 Jan-Feb; 19(1): 29-34.
- Oldridge NB, Guyatt GH, Fischer ME, Rimm AA. Cardiac rehabilitation after myocardial infarction: combined experience of randomized clinical trials. *JAMA*. 1988; 260: 945–950.[\[Abstract\]](#)
- Packard, CJ, O'Reilly, DS, Caslake, MJ, et al. Lipoprotein-associated phospholipase A2 as an independent predictor of coronary heart disease. West of Scotland Coronary Prevention Study Group. *N Engl J Med* 2000; 343:1148.

References

- Pescatello LS, Franklin BA, Fagard R, Farquhar WB, Kelley GA, Ray CA; American College of Sports Medicine. American College of Sports Medicine position stand: exercise and hypertension. *Med Sci Sports Exerc.* 2004; 36: 533–553.
- Philip A. Ades, M.D.: Cardiac Rehabilitation and Secondary Prevention of Coronary Heart Disease. *NEJM*, September 20, 2001, Volume 345:892-902,
- Pollock ML, Franklin BA, Balady GJ, Chaitman BL, Fleg JL, Fletcher B, Limacher M, Pina IL, Stein RA, Williams M, Bazzarre T. AHA Science Advisory. Resistance exercise in individuals with and without cardiovascular disease: benefits, rationale, safety, and prescription: an advisory from the Committee on Exercise, Rehabilitation, and Prevention, Council on Clinical Cardiology, American Heart Association; position paper endorsed by the American College of Sports Medicine. *Circulation.* 2000; 101: 828–833.
- Prescott, E, Hippe, M, Schnohr, P, et al. Smoking and the risk of myocardial infarction in women and men: Longitudinal population study. *BMJ* 1998; 316:1043.
- Prevention of cardiovascular events and death with pravastatin in patients with coronary heart disease and a broad range of initial cholesterol levels. The Long-Term Intervention with Pravastatin in Ischaemic Disease (LIPID) Study Group. *N Engl J Med.* 1998; 339: 1349–1357.
- Randomised trial of cholesterol lowering in 4444 patients with coronary heart disease: the Scandinavian Simvastatin Survival Study (4S). *Lancet.* 1994; 344: 1383–1389.
- Rauramaa R, Li G, Vaisanen SB. Dose-response and coagulation and hemostatic factors. *Med Sci Sports Exerc.* 2001; 33: S516–S520, S528–S529.
- Redberg, RF, Rifai, N, Gee, L, Ridker, PM. Lack of association of C-reactive protein and coronary calcium by electron beam computed tomography in postmenopausal women: implications for coronary artery disease screening. *J Am Coll Cardiol* 2000; 36:39.

References

- Rich-Edwards, JW, Manson, JE, Hennekens, CH, Buring, JE. The primary prevention of coronary heart disease in women. *N Engl J Med* 1995; 332:1758.
- Ridker, PM, Buring, JE, Rifai, N. Soluble P-Selectin and the Risk of Future Cardiovascular Events. *Circulation* 2001; 103:491.
- Ridker, PM, Buring, JE, Shih, J, et al. Prospective study of C-reactive protein and the risk of future cardiovascular events among apparently healthy women. *Circulation* 1998; 98:731.
- Ridker, PM, Glynn, RJ, Hennekens, CH. C-reactive protein adds to the predictive value of total and HDL cholesterol in determining risk of first myocardial infarction. *Circulation* 1998; 97:2007.
- Ridker, PM, Hennekens, CH, Buring, JE, Rifai, N. C-reactive protein and other markers of inflammation in the prediction of cardiovascular disease in women. *N Engl J Med* 2000; 342:836.
- Ridker, PM, Rifai, N, Stampfer, MJ, Hennekens, CH. Plasma concentration of interleukin-6 and the risk of future myocardial infarction among apparently healthy Men. *Circulation* 2000; 101:1767.
- Ridker, PM. Evaluating novel cardiovascular risk factors: Can we better predict heart attacks? *Ann Intern Med* 1999; 130:933.
- Rimm, EB, Giovannucci, EL, Willett, WC, et al. Prospective study of alcohol consumption and risk of coronary disease in men. *Lancet* 1991; 338:464.
- Rimm, EB, Stampfer, MJ, Ascherio, A, et al. Vitamin E consumption and the risk of coronary heart disease in men. *N Engl J Med* 1993; 328:1450.
- Robert J, Francisco T, Richard S, Kelly L, and Lorenzo T: Cardiac Rehabilitation. *EMedicine*, Internet publication, December 29, 2006.
- Roest, M, van der, Schouw YT, de Valk, B, et al. Heterozygosity for a Hereditary Hemochromatosis Gene Is Associated With Cardiovascular Death in Women. *Circulation* 1999; 100:1268.

References

- Roncaglioni, MC, Santoro, L, D'Avanzo, B, et al. Role of family history in patients with myocardial infarction: An Italian case-control study. GISSI-EFRIM Investigators. *Circulation* 1992; 85:2065.
- Rosenberg, L, Kaufman, DW, Helmrich, SP, Shapiro, S. The risk of myocardial infarction after quitting smoking in men under 55 years of age. *N Engl J Med* 1985; 313:1511.
- Rosengren, A, Wedel, H, Wilhelmsen, L. Body weight and weight gain during adult life in men in relation to coronary heart disease and mortality. A prospective population study. *Eur Heart J* 1999; 20:269.
- Rozanski A, Blumenthal JA, Kaplan J. Impact of psychological factors on the pathogenesis of cardiovascular disease and implications for therapy. *Circulation*. 1999; 99: 2192–2217.
- Sacks FM, Pfeffer MA, Moya LA, Rouleau JL, Rutherford JD, Cole TG, Brown L, Warnica JW, Arnold JM, Wun CC, Davis BR, Braunwald E. The effects of pravastatin on coronary events after myocardial infarction in patients with average cholesterol levels. Cholesterol and Recurrent Events Trial investigators. *N Engl J Med*. 1996; 335: 1001–1009.
- Sacks, FM, Pfeffer, MA, Moya, LA, et al. The effect of pravastatin on coronary events after myocardial infarction in patients with average cholesterol levels. Cholesterol and Recurrent Events Trial investigators. *N Engl J Med* 1996; 335:1001.
- Salomaa V, Niemela M, Miettinen H, et al. Relationship of socioeconomic status to the incidence and prehospital, 28-day, and 1-year mortality rates of acute coronary events in the FINMONICA myocardial infarction register study. *Circulation* 2000;101:1913-1918.
- Salomaa, V, Matei, C, Alesksic, N, et al. Soluble thrombomodulin as a predictor of incidence coronary heart disease and symptomless carotid artery atherosclerosis in the Atherosclerosis Risk in Communities (ARIC) study: A case-cohort study. *Lancet* 1999; 353:1729.

- Sanderson BK, Bittner V: Women in cardiac rehabilitation: outcomes and identifying risk for dropout. *Am Heart J* 2005 Nov; 150(5): 1052-8.
- Schachinger, V, Britten, MB, Zeiher, AM. Prognostic impact of coronary vasodilator dysfunction on adverse long-term outcome of coronary heart disease. *Circulation* 2000; 101:1899.
- Schwartz GG, Olsson AG, Ezekowitz MD, Ganz P, Oliver MF, Waters D, Zeiher A, Chaitman BR, Leslie S, Stern T; Myocardial Ischemia Reduction with Aggressive Cholesterol Lowering (MIRACL) Study Investigators. Effects of atorvastatin on early recurrent ischemic events in acute coronary syndromes: the MIRACL study: a randomized controlled trial. *JAMA*. 2001; 285: 1711–1718.
- Sdringola, S, Patel, D, Gould, KL. High Prevalence of Myocardial Perfusion Abnormalities on Positron Emission Tomography in Asymptomatic Persons With a Parent or Sibling With Coronary Artery Disease. *Circulation* 2001; 103:496.
- Selhub, J, Jacques, PF, Wilson, PWF, et al. Vitamin status and intake as primary determinants of homocysteinemia in the elderly. *JAMA* 1993; 270:2693.
- Sesso, HD, Lee, IM, Gaziano, JM, et al. Maternal and paternal history of myocardial infarction and risk of cardiovascular disease in men and women. *Circulation* 2001; 104:393.
- Sesso, HD, Stampfer, MJ, Rosner, B, et al. Two-year changes in blood pressure and subsequent risk of cardiovascular disease in Men. *Circulation* 2000; 102:307.
- Sharma RK, Singh VN, Flaker GC: Signal-averaged electrocardiogram: A non-invasive diagnostic tool for arrhythmia management. . *Develop Cardiol* 1994; 4(12): 175-92.
- Shepherd, J, Cobbe, SM, Ford, I, et al. Prevention of coronary heart disease with pravastatin in men with hypercholesterolemia. West of Scotland Coronary Prevention Study Group. *N Engl J Med* 1995; 333:1301.

- Singer, DE, Nathan, DM, Anderson, KM, et al. Association of HbA1c with prevalent cardiovascular disease in the original cohort of the Framingham Heart Study. *Diabetes* 1992; 41:202.
- Singh VN: The role of gas analysis with exercise testing. *Prim Care* 2001; 28(1): 159-79.
- Smart N, Marwick TH. Exercise training for patients with heart failure: a systematic review of factors that improve mortality and morbidity. *Am J Med.* 2004; 116: 693–706.
- Smith SC Jr, Blair SN, Bonow RO, Brass LM, Cerqueira MD, Dracup K, Fuster V, Gotto A, Grundy SM, Miller NH, Jacobs A, Jones D, Krauss RM, Mosca L, Ockene I, Pasternak RC, Pearson T, Pfeffer MA, Starke RD, Taubert KA. AHA/ACC Scientific Statement: AHA/ACC guidelines for preventing heart attack and death in patients with atherosclerotic cardiovascular disease: 2001 update: a statement for healthcare professionals from the American Heart Association and the American College of Cardiology. *Circulation.* 2001; 104: 1577–1579.
- Staessen, JA, Wang, JG, Thijs, L. Cardiovascular protection and blood pressure reduction: a meta-analysis. *Lancet* 2001; 358:1305.
- Stamler, J, Stamler, R, Neaton, JD. Blood pressure, systolic and diastolic, and cardiovascular risks: U.S. population data. *Arch Intern Med* 1993; 153:598.
- Stampfer, MJ, Hennekens, CH, Manson, JE, et al. Vitamin E consumption and risk of coronary heart disease in women. *N Engl J Med* 1993; 328:1444.
- Stampfer, MJ, Malinow, MR, Willett, WC, et al. A prospective study of plasma homocysteine and risk of myocardial infarction in US physicians. *JAMA* 1992; 268:877.
- Stec, JJ, Silbershatz, H, Ofler, GH, et al. Association of fibrinogen with cardiovascular risk factors and cardiovascular disease in the Framingham offspring population. *Circulation* 2000; 102:1634.

References

- Stone JA, Arthur HM: Canadian Guidelines for Cardiac Rehabilitation and Cardiovascular Disease Prevention, second edition, 2004: Executive summary. *Can J Cardiol* 2005 Oct; 21 Suppl D: 3D-19D.
- Streuber SD, Amsterdam EA, Stebbins CL: Heart rate recovery in heart failure patients after a 12-week cardiac rehabilitation program. *Am J Cardiol* 2006 Mar 1; 97(5): 694-8.
- Suwaidi, JA, Hamasaki, S, Higano, ST, et al. Long-term follow-up of patients with mild coronary artery disease and endothelial dysfunction. *Circulation* 2000; 101:948.
- Tanaka H, Dinunno FA, Monahan KD, Clevenger CM, DeSouza CA, Seals DR. Aging, habitual exercise, and dynamic arterial compliance. *Circulation*. 2000; 102: 1270–1275.
- Tanasescu M, Leitzmann MF, Rimm EB, Willett WC, Stampfer MJ, Hu FB. Exercise type and intensity in relation to coronary heart disease in men. *JAMA*. 2002; 288: 1994–2000.
- Taylor R, Kirby B: Cost implications of cardiac rehabilitation in older patients. *Coron Artery Dis* 1999; 10(1): 53-56.
- Taylor RS, Brown A, Ebrahim S, Jolliffe J, Noorani H, Rees K, Skidmore B, Stone JA, Thompson DR, Oldridge N. Exercise-based rehabilitation for patients with coronary heart disease: systematic review and meta-analysis of randomized trials. *Am J Med*. 2004; 116: 682–697.
- Taylor, RS, Brown, A, Ebrahim, S, et al. Exercise-based rehabilitation for patients with coronary heart disease: systematic review and meta-analysis of randomized controlled trials. *Am J Med* 2004; 116:682.
- The sixth report of the Joint National Committee on prevention, detection, evaluation, and treatment of high blood pressure. *Arch Intern Med* 1997; 157:2413.
- The sixth report of the Joint National Committee on prevention, detection, evaluation, and treatment of high blood pressure. *Arch Intern Med* 1997; 157:2413.

- Thompson PD, Buchner D, Piña IL, Balady GJ, Williams MA, Marcus BH, Berra K, Blair SN, Costa F, Franklin B, Fletcher GF, Gordon NF, Pate RR, Rodriguez BL, Yancey AK, Wenger NK; American Heart Association Council on Clinical Cardiology Subcommittee on Exercise, Rehabilitation, and Prevention; American Heart Association Council on Nutrition, Physical Activity, and Metabolism Subcommittee on Physical Activity. Exercise and physical activity in the prevention and treatment of atherosclerotic cardiovascular disease: a statement from the Council on Clinical Cardiology (Subcommittee on Exercise, Rehabilitation, and Prevention) and the Council on Nutrition, Physical Activity, and Metabolism (Subcommittee on Physical Activity). *Circulation*. 2003; 107: 3109–3116. [Free](#) [Full Text](#)
- Thompson PD. Exercise for patients with coronary artery and/or coronary heart disease. In: Thompson PD, ed. *Exercise and Sports Cardiology*. New York, NY: McGraw Hill; 2001: 354–370.
- Thun, MJ, Peto, R, Lopez, AD, et al. Alcohol consumption and mortality among middle-aged and elderly US adults. *N Engl J Med* 1997; 337:1705.
- Touboul, PJ, Elbaz, A, Koller, C, et al. Common carotid artery intima-media thickness and brain infarction: the etude du profil Genetique de l'Infarctus Cerebral (GENIC) case-control study. *Circulation* 2000; 102:313.
- Tuomainen, TP, Kontula, K, Nyys#nen, K, et al. Increased Risk of Acute Myocardial Infarction in Carriers of the Hemochromatosis Gene Cys282Tyr Mutation: A Prospective Cohort Study in Men in Eastern Finland. *Circulation* 1999; 100:1274.
- US Department of Health and Human Services, Public Health Service, AHCPR: *Cardiac Rehabilitation: Clinical Practice Guideline*. Rockville, Md: 1995; Oct: 1-202.
- Vaccarino, V, Berger, AK, Abramson, J, et al. Pulse pressure and risk of cardiovascular events in the systolic hypertension in the elderly program. *Am J Cardiol* 2001; 88:980.

References

- Vanharanta, M, Voutilainen, S, Lakka, TA, et al. Risk of acute coronary events according to serum concentrations of enterolactone: a prospective population-based case-control study. *Lancet* 1999; 354:2112.
- Vasan, RS, Larson, MG, Benjamin, EJ, et al. Left ventricular dilatation and the risk of congestive heart failure in people without myocardial infarction. *N Engl J Med* 1997; 336:1350.
- Waldius, G, Jungén, I, Holme, I, et al. High apolipoprotein B, low apolipoprotein A-I, and improvement in the prediction of fatal myocardial infarction (AMORIS study): a prospective study. *Lancet* 2001; 358:2026.
- Wallace, AM, McMahon, AD, Packard, CJ, et al. Plasma Leptin and the Risk of Cardiovascular Disease in the West of Scotland Coronary Prevention Study (WOSCOPS). *Circulation* 2001.
- Wannamethee, SG, Shaper, AG, Whincup, PH. Serum urate and the risk of major coronary heart disease events. *Heart* 1997; 78:147.
- Wei, M, Gibbons, LW, Mitchell, TL, et al. Low fasting plasma glucose level as a predictor of cardiovascular disease and all-cause mortality. *Circulation* 2000; 101:2047.
- Wenger NK, Froelicher ES, Smith LK, Philip A, Ades PA, Berra K, Blumenthal JA, Certo CM, Dattilo AM, Davis D, DeBusk RF, Drozda JP Jr, Fletcher BJ, Franklin BA, Gaston H, Greenland P, McBride PE, McGregor CG, Oldridge NB, Piscatella JC, Rogers FJ. Clinical Practice Guidelines No. 17: Cardiac Rehabilitation as Secondary Prevention. Rockville, Md: US Department of Health and Human Services, Public Health Service, Agency for Health Care Policy and Research, National Heart, Lung and Blood Institute; 1995. AHCPR Publication 96-0672.
- Wiedermann, CJ, Kiechl, S, Dunzendorfer, S, et al. Association of endotoxemia with carotid atherosclerosis and cardiovascular disease: prospective results from the Bruneck Study. *J Am Coll Cardiol* 1999; 34:1975.

References

- Wilhelmsson, C, Vedin, JA, Elmfeldt, D, et al. Smoking and myocardial infarction. *Lancet* 1975; 1:415.
- Wilking, SVB, Belanger, AJ, Kannel, WB, et al. Determinants of isolated systolic hypertension. *JAMA* 1988; 260:3451.
- Wilson, G. Rayos, T.K. Yeoh et al., Dissociation between exertional symptoms and circulatory function in patients with heart failure. *Circulation* 92 (1995), pp. 47–53.
- Wilson, PW, D'Agostino, RB, Levy, D, et al. Prediction of coronary heart disease using risk factor categories. *Circulation* 1998; 97:1837.
- Wilson, PW, Kannel, WB. Hypercholesterolemia and Coronary Risk in the Elderly: The Framingham Study. *Am J Geriatr Cardiol* 1993; 2:56.
- Wilson, PW, Kauppila, LI, O'Donnell, CJ, et al. Abdominal Aortic Calcific Deposits Are an Important Predictor of Vascular Morbidity and Mortality. *Circulation* 2001; 103:1529.
- Wilson, PW. Established risk factors and coronary artery disease: The Framingham Study. *Am J Hypertens* 1994; 7:7S.
- Wolk, A, Manson, JE, Stampfer, MJ, et al. Long-term intake of dietary fiber and decreased risk of coronary heart disease among women. *JAMA* 1999; 281:1998.
- Woods, A, Brull, DJ, Humphries, SE, Montgomery, HE. Genetics of inflammation and risk of coronary artery disease: the central role of interleukin-6. *Eur Heart J* 2000; 21:1574.
- Wu, KK, Aleksic, N, Ahn, C, et al. Thrombomodulin Ala455Val polymorphism and risk of coronary heart disease. *Circulation* 2001; 103:1386.
- Wybe N, Marike A, Dirk J, Johan Brügemann, Martin L. J., Eric van Sonderen, K. I. Lie, Harry J. G and Piet Rispens: Differential effects of high-frequency versus low-frequency exercise training in rehabilitation of patients with coronary artery disease. *Journal of the American College of Cardiology*; 36(1), 2000, 202-207.

References

- Wybe Nieuwland, Marike A. Berkhuisen et al., Differential effects of high-frequency versus low-frequency exercise training in rehabilitation of patients with coronary artery disease. Journal of the American College of Cardiology. Volume 36, Issue 1 , July 2000, Pages 202-207.
- Zhang, R, Brennan, ML, Fu, X, et al. Association between myeloperoxidase levels and risk of coronary artery disease. JAMA 2001; 286:2136.
- Zhu, J, Nieto, FJ, Horne, BD, et al. Prospective study of pathogen burden and risk of myocardial infarction or death. Circulation (Online) 2001; 103:45.
- Zhu, J, Quyyumi, AA, Norman, JE, et al. Effects of total pathogen burden on coronary artery disease risk and C-reactive protein levels. Am J Cardiol 2000; 85:140.