

References

Ackard DM, Neumark-Sztainer D and Story M (2003):

Overeating among adolescents: prevalence and associations with weight-related characteristics and psychological health. Pediatrics; 111: 67-74.

Alger S, Larson K and Boyce VL (1993):

Effect of phenylpropanolamine on energy expenditure and weight loss in overweight women. Am. J. Clin. Nutr.; 57: 120-126.

Allison DB, Fontaine KR and Heshka S (2001):

Alternative treatments for weight loss: A critical review. Crit. Rev. Food Sci. Nutr.; 41: 1-28.

Almond L and Newberry I (2000):

The importance of physical activity in weight management. Obesity in Practice; 2(2): 10-12.

Anderson JW, Vichitbandra S and Qian W (1999):

Long-term weight maintenance after an intensive weight loss program. J. Am. Coll. Nutr.; 18(6): 620-627.

Anderson JW and Konz EC (2001):

Obesity and disease management: Effects of weight loss on comorbid conditions. Obes. Res.; 9: 326-334.

Armstrong N, Balding J and Gentle P (1990):

Patterns of physical activity among 11-16-year-old British children. British Medical Journal; 301: 203-205.

Arrone L (1998):

Obesity.

Med Clin North America 1998; 82(1): 161-181.

Astrup A and Toubro S (1993):

Thermogenic, metabolic and cardiovascular responses to ephedrine and caffeine in man.

Int. J. Obes. Relat. Metal. Disord.; 17(1): 41-3.

Astrup A, Breum L and Toubro S (1995):

Pharmacological and clinical studies of ephedrine and other thermogenic agents.

Obes. Res.; 3(4): 537-40.

Astrup A, Bueniann B and Toubro S (1996):

Low resting metabolic rate in subjects predisposed to obesity: a role for thyroid status.

Am. J. Clin. Nutr.; 63: 879-883.

Atkinson R (1989):

Low and very low caloric diets.

Med. Clin. North. Am.; 73: 203-215.

Badmaev V, Majeed M and Conte AA (1999):

Garcinia cambogia for weight loss.

JAMA; 282: 233-4.

Bahadori ST, Chang K and Hu M (2003):

Acupuncture as an alternative methods for treatment of obesity.

Obesity Reviews 4: 56-61.

Ballinger A and Peikin SR (2002):

Orlistat: its current status as an anti-obesity drug.

Eur. J. Pharmacol.; 440: 109-117.

Ballor DL and Poehlman ET (1994):

Exercise training enhances fat-free mass preservation during diet-induced weight loss: a meta-analytical finding.

International Journal of Obesity; 18: 35-40.

Balsiger BM, Murr MM and Poggio JL (2000):

Bariatric surgery. Surgery for weight control in patients with morbid obesity.

Medical Clinics of North America; 84(2): 477-489.

Bei Y, Fang X and Yao Z (2004):

Sixty-two cases of simple obesity treated by acupuncture combined with massage.

J. Tradit. Chin. Med. 24(1): 36-39.

Bhatt AD, Dalal DG, Shah JJ, et al (1995):

Conceptual and methodologic challenges of assessing the short term efficacy of guggul in obesity: data emergent from a naturalistic clinical trial.

J. Post. Grad. Med.; 41: 5-7.

Birch S, Hesselink JK and Jonkman FA (2004):

Clinical research on acupuncture. Part 1. What have reviews of the efficacy and safety of acupuncture told us so far?

J. Altern. Complement Med.; 10(3): 468-80.

Birketwedt GS, Sundsfjerd J and Florholmen JR (2002):

Hypothalamic-pituitary-adrenal axis in the night eating syndrome.

Endocrinology and Metabolism; 282: E366-E369.

Blackburn G (1995):

Effect of degree of weight loss on health benefits.

Obesity Research; 3: 211-216.

Bravata DM, Sanders L and Huang J (2003):

Efficacy and safety of low-carbohydrate diets. A systematic review.

JAMA; 289: 1837-1850.

Bray GA (1989):

Obesity basic consideration and clinical approaches.

Dis. Mon.; 35: 449-537.

Bray GA, Blackburn GL and Ferguson JM (1999):

Sibutramine produces dose-related weight loss.

Obes. Res.; 7: 189-198.

Bray GA (2003):

Low-carbohydrate diets and realities of weight loss.

JAMA; 289: 1853-1855.

Brenner JS, Kelly CS and Wenger AD (2001):

Asthma and obesity in adolescents: is there an association?

Journal of Asthma 2001; 38(6): 509-515.

British Nutrition Foundation (BNF) Task Force Report (1999):

Obesity. Blackwell, London.

Cabyoglu MT, Ergene N and Tan U (2006):

The treatment of obesity by acupuncture.

Int. J. Neurosci.; 116(2): 165-75.

Capstick F, Brooks BA and Burns CM (1997):

Very low calorie diets (VLCD): a useful alternative in the treatment of the obese NIDDM patient.

Diabetes Research and Clinical Practice; 36(2): 105-111.

Chambers R, et al (2000):

Weight loss, weight maintenance and improved cardiac risk factors after 2 years of treatment with orlistat for obesity.

Obesity research; 8: 49-61.

Chang SL, Lin JG and Chi TC (1999):

An insulin-dependent hypoglycemia induced by electroacupuncture at the Zhongwan (CV12) acupoint in diabetic rats.

Diabetologia.; 42(2): 250-255.

Chen Z, Hendner J and Hedner T (1996):

Substance P induced respiratory excitation is blunted by delta-receptor specific opioids in the rat medulla oblongata.

Acta. Physiologica. Scandinavica.; 157(2): 165-173.

Cheng Y, Huang X and Jia W (editors) (1990):

Acupuncture technique. In: Chinese acupuncture and moxibustion. 2nd edition. Published by: Foreign Languages Press. Beijing, p: 318-339.

Chernoff R (2001):

Nutrition and health promotion in older adults.

J. Gerontol. A. Biol. Sci. Med. Sci.; 56: 47-53.

Choi HK, Atkinson K, Karlson EW, et al (2002):

Weight gain, obesity, and incident gout in a large, long-term, prospective cohort study.

Program and abstracts of the American College of Rheumatology 66th. Annual Scientific Meeting; 25-29.

Choy DS and Eidenschenk E (1998):

Effect of tragus clips on gastric peristalsis: a pilot study.

J. Altern. Complement Med.; 4: 399-403.

Cicuttini FM, Baker JR and Spector TD (1996):

The association of obesity with osteoarthritis of the hand and knee in women: a twin study.

Journal of Rheumatology; 23(7): 1221-1226.

Clement K, Vaisse C and Manning B (1995):

Genetic variation in the B3 adrenergic receptor and an increased capacity to gain weight in patients with morbid obesity.

N. Eng. J. Med.; 333: 352-354.

Cohen P and Friedman JM (2004):

Leptin and the control of metabolism: Role of Stearoyl-CoA Desaturase-1 (SCD-1).

J. Nutr.; 134: 2455-2463.

Colditz GA, Willett WC and Rotnitzkya A (1995):

Weight gain as a risk factor for clinical diabetes.

Annals of Internal Medicine; 51: 1100-1105.

Comuzzie AJ (2001):

The genetic contribution to human obesity: the dissection of a complex phenotype. In: Obesity, growth and development. 2nd ed. Edited by: Johnston FE, Foster GD. Published by: Smith-Gordon & Co., Ltd.: London; pp: 21-36.

Connolly HM, Crary JL and McGoon MD (1997):

Valvular heart disease associated with fenfluramine-phentermine.

N. Engl. J. Med.; 337: 581-588.

Considine RV, Sinha MK and Heiman ML (1996):

Serum immunoreactive leptin concentrations in normal weight and obese humans.

New England Journal of medicine; 334: 292-295.

Costain L (2003):

Diet trials.

BBC Publications, London, 2003; 204.

Crespo CH, Smit E and Troiano RP (2001):

Television watching, energy intake and obesity in US children: results from the third National Health and Nutrition Examination Survey 1988-1994.

Archives of Pediatric and Adolescent Medicine; 115: 360-365.

Crowley VE, et al (2002):

Obesity therapy: Altering the energy intake and expenditure balance sheet.

Nat. Rev. Drug Disc.; 1:276.

Dai S, Labarthe DR and Grunbaum JA (2002):

Longitudinal analysis of changes in indices of obesity from age 8 years to age 18 years: Project heartbeat!

Am. J. Epidemiol.; 156(8): 720-729.

Daniels SR, Khoury PR and Morrison JA (1997):

The utility of body mass index as a measure of body fatness in children and adolescents: Differences by race and gender.

Pediatrics; 99(6): 804-807.

Ditschuneit HH, Flechtner-Mors M and Johnson TD (1999):

Metabolic and weight loss effects of a long term dietary intervention in obese patients.

American Journal of Clinical Nutrition; 69(2): 198-204.

Ditschuneit HH, Frier HI and Flechtner-Mors M (2002):

Lipoprotein responses to weight loss and weight maintenance in high risk obese subjects.

European Journal of Clinical Nutrition; 56(3): 264-270.

Drent ML and Ven der Veen EA (1993):

Lipase inhibition: a novel concept in the treatment of obesity.

Int. J. Obes.; 17: 241-244.

Dung HC (1986):

Attempts to reduce body weight through auricular acupuncture.

Am. J. Acupuncture; 14: 117-122.

Dymek MP, le Grange D and Neven K (2002):

Quality of life after gastric bypass surgery: A cross-sectional study.

Obes. Res.; 10(11): 1135-1142.

Egan BM (2003):

Insulin resistance and the sympathetic nervous system.

Current Hypertension Reports; 5(3): 247-254.

Eichenberger A, Proietti S and Wicky S (2002):

Morbid obesity and postoperative pulmonary atelectasis: an underestimated problem.

Anesthesia and Analgesia; 95(6): 1788-1792.

Engeli S, Feldpausch M and Gorzelniak K (2003):

Association between adiponectin and mediators of inflammation in obese women.

Diabetes; 52: 942-947.

Epstein LH, Valoski AM and Vara LS (1995):

Effects of decreasing sedentary behaviour and increasing activity on weight changes in obese children.

Health Psychology; 14: 1-7.

Ernst E and White A (2001):

Prospective studies of the safety of acupuncture: a systematic review.

Am. J. Med.; 110: 481-5.

Evans M and Straus S (2001):

Ripe for study: complementary and alternative treatments for obesity.

Crit. Rev. Food Sci. Nutr.; 41: 35-38.

Farin HM, Abbasi F and Reaven GM (2006):

Body mass index and waist circumference both contribute to differences in insulin mediated glucose disposal in nondiabetic adults.

Am. J. Clinical Nutrition; 83(1): 47-51.

Felson DT, Anderson JJ and Naimark A (1988):

Obesity and knee osteoarthritis. The Framingham Study.

Annals of Internal Medicine, 109(1): 18-24.

Finer N (2002):

Sibutramine: its mode of action and efficacy.

Int. J. Obes. Relat. Metab. Disord.; 26(4): 29-33.

Firenzuoli F and Gori L (1999):

Garcinia cambogia for weight loss.

JAMA 1999; 282; 234-7.

Flegal KM, et al (2002):

Prevalence and trends in obesity among US adults. From 1999-2000.

JAMA; 288: 1723.

Flier JS and Maratos-Flier E (2002):

The stomach speaks-ghrelin and weight reduction.

N. Engl. J. Med.; 346: 1662.

Fontanine KR, et al (2003):

Years of life lost due to obesity.

JAMA; 289.

Foster GD (2001):

Non-dieting approaches. In: Eating disorders and obesity: a comprehensive handbook. Edited by: Brownell KD, Fairburn CG . published by: Guilford: New York, 55.

Foster GD and McGuckin BG (2002):

Non dieting approaches: Principles, practices and evidence. In: Obesity: Theory and Therapy. Edited by: Wadden TA, Stunkard AJ. Published by: Guilford: New York, 122.

Freedman MR, King J and Kennedy E (2001):

Popular diets: a scientific review.
Obes. Res.; 9: 1-40.

Fu H (2000):

What is the material base of acupuncture? The nerves!
Medical Hypotheses; 54(3): 358-359.

Galtier-Dereure F, Boegner C and Bringer J (2000):

Obesity and pregnancy: complications and cost.
American Journal of Clinical Nutrition; 71: 1242.

Ganong W, editor (1997):

Energy balance, metabolism and nutrition. In: Review of Medical Physiology, 18th ed. Published by Appleton & Lange. London, Sydney, Mexico: 291-295.

Garrow JS, Webster JD and Pearson M (1989):

Inpatient-outpatient randomized comparison of the Cambridge diet versus milk diet in 17 obese women over 24 weeks.
International Journal of Obesity; 13: 521-529.

Garrow JS and Summerbell CD (1995):

Meta-analysis: effect of exercise with or without dieting, on body composition of overweight subjects.

European Journal of Clinical Nutrition; 49: 1-10.

Garrow JS (1999):

Obesity. British Nutrition Foundation, London; 164.

Gately PJ and Cooke CB (2003):

A residential summer camp programme for the treatment of obese and overweight children.

Obesity in Practice; 5(1): 2-5.

Giller RM (1975):

Auricular acupuncture and weight reduction: a controlled study.

Am. J. Acupuncture; 3: 151-153.

Ginsberg-Fellner F and Knittle J (1989):

Weight reduction in young obese children. Effects on adipose tissue cellularity and metabolism.

Pediatr. Res. 15: 1381-1389.

Gortmaker SL, Must A and Sobol AM (1996):

Television viewing as a cause of increasing obesity among children in the United States 1986-90.

Archives of Pediatric and Adolescent Medicine; 150: 356-362.

Grundy SM, Cleeman JI and Daniels SR (2005):

Diagnosis and management of the metabolic syndrome: an American Heart Association / National Heart, Lung and Blood Institute Scientific Statement.

Circulation; 112(17): 2735-2752.

Guidance for Treatment of Adult Obesity (1996):

Bethesda, MD, shape up.

America and the American Obesity Association.

Gustafsson F, Kragelund CB and Torp Pedersen C (2005):

Effect of obesity and being overweight on long-term mortality in congestive heart failure: influence of left ventricular systolic function.

Eur. Heart J.; 26(1): 58-64.

Guyton AC and Hall JE (editors):

Control of appetite, pain controlling system. In: Textbook of medical physiology, 10th edition, published by Saunders, Philadelphia, London, New York, Son Louis, Sydney, Toronto, p. 106-155 and 683-685.

Haller CA and Benowitz NL (2000):

Adverse cardiovascular and central nervous system events associated with dietary supplements containing Ephedra alkaloids.

N. Engl. J. Med.; 343: 1833-8.

Halpern A and Mancini MC (2003):

Treatment of obesity: on update on anti-obesity medications.

Obesity Reviews; 4: 24-42.

Hartz A, Barboriak P and Wong A (1990):

The association of obesity with infertility and related menstrual abnormalities in women.

Int. J. Obes.; 3: 57-60.

Hauptman J, Lucas C and Boldrin MN (2000):

Orlistat in the long-term treatment of obesity in primary care settings.

Arch. Fam. Med.; 9: 160-167.

Havely and Solman (1965):

Determination of serum serotonin: In Harper's biochemistry (1988), 21st edition, edited by: Murrar RK et al., (edition). Published by Saunders, Burgh, London. P: 306.

Haynes WG, Sivitz WI and Morgan DA (1997):

Sympathetic and cardiorenal actions of leptin.

Hypertension; 30: 619-623.

Health Education Authority (1991):

Health and lifestyle survey.

HEA, London.

Hill JO and Wyatt H (2002):

Outpatient management of obesity: a primary care perspective.

Obes. Res. 10(2): 124-130.

Himms-Hagen J (1989):

Brown adipose tissue thermogenesis and obesity.

Prog. Lipid Res.; 28: 67-115.

Huang MH, Yang RC and Hu SH (1996):

Preliminary results of triple therapy for obesity.

Int. J. Obes. Relat. Metab. Disord.; 20(9): 830-836.

Ian W and Devid W (Editors) (2005):

What is obesity, causes of obesity, treatment of obesity. In: Obesity, your questions answered. 1st edition, published by: Churchill Livingstone, Edinburgh, London, New York, 1-103.

Ingelsson E, Sundstrom J, Arnlov J, et al. (2005):

Insulin resistance and risk of congestive heart failure.

JAMA; 294: 334-41.

Jakicic JM, Marcus BH, Gallagher KI, et al (2003):

Effect of exercise duration and intensity on weight loss in overweight, sedentary women: a randomized trial.

JAMA; 290: 1323-1330.

Jankovic B (1994):

Neuroimmunomodulation: The state of the art.

New York: the New York Academy of Sciences.

Jeffrey SF and Daniel WA (1998):

Eating disorders: Obesity. In: Williams Textbook of Endocrinology. 9th ed, edited by: Daniel JD., et al. Published by W.B. Saunders Company, Philadelphia, London, Toronto, Ch. 22, p: 1061-1075.

Jeffrey SF and Eleftheria MF (2005):

Obesity and eating disorders. In: Harrison's principles of internal medicine, 16th edition, edited by Dennis, L.K., et al., published by McGraw-Hill, Medical Publishing Division, New York, p. 422-429.

Jin HO, Zhou L, Lee KY, et al. (1996):

Inhibition of acid secretion by electrical acupuncture is mediated via beta endorphin and so-matostatin.

American Journal of Physiology; 271: 6524-6530.

Jordan KM, Dennison EM and Syddall H (2002):

Birthweight and spinal osteoarthritis: evidence for intrauterine programming.

Program and abstracts of the American College of Rheumatology 66th Annual Scientific Meeting; October 25-29. New Orleans, LA. Abstract 1238.

Jung R (1997):

Obesity as a disease.

British Medical Bulletin; 53: 307-321.

Kahn R, Buse J, Ferrannini E and Stern M (2005):

The metabolic syndrome: time for a critical appraisal: joint statement from the American Diabetes Association and the European Association for the Study of Diabetes.

Diabetes Care; 28: 2289-304.

Khansori DN, Murgo AJ and Golt PW (1990):

Effects of stress on the immune system.

Immunology Today; 11: 170-175.

Kho H, and Robertson EN (1997):

The mechanisms of acupuncture analgesia: Review and update.

American Journal of Acupuncture; 25: 261-629.

Kim SH, Lee YM and Jee SH (2003):

Use of metabolic markers to identify overweight individuals who are insulin resistant.

Ann Intern Med; 139(10): 802-809.

Kissebah A and Krakower G (1994):

Regional adiposity and morbidity.

Physiol. Rev.; 74: 761-811.

Klein S (2001):

Alternative therapies for obesity: benefit or rip-off.

Crit. Rev. Food Sci. Nutr.; 41: 33-34.

Klein S, Sheard NF, Pi-Sunyer X, et al (2004):

Weight management through lifestyle modification for the prevention and management of type 2 diabetes: rationale and strategies. A statement of the American Diabetes Association, the North American Association for the Study of Obesity, and the American Society for Clinical Nutrition.

Am. J. Clinical Nutrition; 80(2): 257-263.

Kolotkin RL, Crosby RD, Williams GR, et al (2001):

The relationship between health related quality of life and weight loss.

Obes. Res.; 9(9): 564-571.

Kumar P and Clark M (editors) (2005):

Obesity, Textbook of clinical medicine, 6th edition,
Published by Saunders. Burgh, London, Philadelphia,
Toronto, Sydney, Tokyo, p. 252-259.

Kushner R and Foster GD (2000):

Obesity and quality of life.
Nutrition; 16: 947-952.

Lacey JM, Tershakovec AM and Foster GD (2003):

Acupuncture for the treatment of obesity: a review of the
evidence.
Int. J. Obes. Relat. Metab. Disord.; 27(4): 419-27.

Lamertz CM, Jacobi C, Yassouridis A, et al (2002):

Are obese adolescents and young adults at higher risk for
mental disorders? A community survey.
Obes. Res.; 10(11): 1152-1160.

Lao L (1996):

Acupuncture technique and devices.
J. Altern. Complem. Med.; 2: 23-25.

Lawlor DA, Lean M and Sattar N (2007):

ABC of obesity. Obesity and vascular disease.
BMJ; 14(144): 124-127.

Lee CD, Jackson AS and Blair SN (1998):

US weight guidelines: is it also important to consider cardio-
respiratory fitness?
International Journal of Obesity; 22(20): S2-S7.

Leibel RL, Rosenbaum M and Hirsch J (1995):

Changes in energy expenditure resulting from altered body
weight.
N. Engl. J. Med.; 332: 621-628.

Lemay CA, Cashman S, Savageau J, et al (2003):

Underdiagnosis of obesity at a community health center.

J. Am. Board. Fam. Med.; 16(1): 14-21.

Lethbridge-Cejku M, Creamer P and Scott WW (2002):

Weight loss is associated with reduced risk of incident radiographic knee osteoarthritis in men but not in women: data from the Baltimore Longitudinal Study of Aging.

Program and abstracts of the American College of Rheumatology 66th Annual Scientific Meeting; October 25-29. New Orleans, LA. Abstract 1235.

Li SJ, Tang J and Han JS (1982):

The implication of central serotonin in electro-acupuncture tolerance in rat.

Scientia. Sinica.; 25(6): 620-629.

Li WH, Wang JD, Gu LM, et al (2004):

Treatment of simple obesity with electro-acupuncture and auricular acupoint pressing: a report of 177 cases.

Zhong Xi Yi Jie He Xue Bao.; 2(6): 449-458.

Li Y, Tougas G, Chiverton SG, et al (1992):

The effect of acupuncture on gastrointestinal function and disorder.

American Journal of Gastroenterology; 87(10): 1372-1381.

Li Z, Maglione M, Tu W, et al (2005):

Meta-analysis: pharmacologic treatment of obesity.

Ann. Intern. Med.; 142(7): 532-546.

Linde K, Jonas WB, Melchart D, et al (2001):

The methodological quality of randomized controlled trials of homeopathy, herbal medicines and acupuncture.

Int. J. Epidemiol.; 30: 526-631.

Liu Z, Sun F, Li J, et al (1992):

Prophylactic and therapeutic effects of acupuncture on simple obesity complicated by cardiovascular diseases.

J. Tradit. Chin. Med.; 12: 21-29.

Liu Z, Sun F, Su J, et al (2001):

Study on action of acupuncture on ventromedial nucleus of hypothalamus in obese rats.

J. Tradit. Chin. Med.; 21(3): 220-4.

Loffe E, Moon B, Connolly E et al (1998):

Abnormal regulation of the leptin gene in the pathogenesis of obesity.

Proc. Natl. Acad. Sci. USA; 95: 11852-11857.

Lohman T (1981):

Skinfolds and body density and their relation to body fatness: A Review.

Hum. Boil.; 53: 181-225.

Lukaski HC, Johnson PE and Bolonchuk WW (1987):

Assessment of fat-free mass using bioelectrical impedance measurements of human body.

Am. J. Clin. Nutr.; 41: 810-7.

Lyznicki JM, Young DC, Riggs JA, et al (2001):

Obesity: assessment and management in primary care.

Am. Fam. Physcian.; 63: 185-219.

Maciocia G (1989):

The foundations of Chinese medicine.

New York: Churchill Livingstone.

Maes HH, et al (1997):

Genetic and environmental factors in relative body weight and human adiposity.

Behavior Genetics; 27: 325-351.

Malizia E, Andreucci G, Paolucci D, et al (1979):

Electroacupuncture and peripheral beta endorphin and ACTH levels.

Lancet; 2(8141): 535-536.

Manson J, Willett W, Stampfer M, et al (1995):

Body weight and mortality among women.

N. Engl. J. Med.; 333: 677-685.

Mantzoros CS (1999):

The role of leptin in human obesity and disease: A review of current evidence.

Ann. Intern. Med.; 130: 671-680.

Mao-Liang Q, Shan-Chen Z, Zhong-Quan Y, et al (editors) (1993):

The 12 regular meridians. In: "Chinese Acupuncture and Moxibustion" 1st ed. Published by: Longman, Singapore publishers Ltd., Edinburgh, London, Madrid, Melbourne, New York and Tokyo. Ch. 3: 61-182.

Mason EE, Tang S, Renquiat KE, et al (1997):

Decade of change in obesity surgery.

Obesity Surgery; 7: 189-197.

Mazzoni R, Mannucci E, Rizzello SM, (1999):

Failure of acupuncture in the treatment of obesity: a pilot study.

Eat. Weight. Disord.; 4: 198-202.

McDuffie JR, Calis KA, Booth SL, et al (2002):

Effects of orlistat on fat-soluble vitamins in obese adolescents.

Pharmacotherapy; 22: 814-822.

McFarlane T, Polivy J and McCabe RE (1999):

Help, not harm: psychological foundation for a nondieting approach toward health.

J. Soc. Iss.; 55: 261-276.

McMahon FG, Weinstein SP, Rowe E, et al (2002):

Sibutramine is safe and effective for weight loss in obese patients whose hypertension is well controlled with angiotensin-converting enzyme inhibitors.

J. Hum. Hypertens.; 16: 5-11.

McMillan DC, Sattar N, Lean M, et al (2007):

ABC of obesity. Obesity and cancer.

BMJ; 14(144): 128-130.

Mikdad AH, Bowman BA, Ford ES, et al (2001):

The continuing epidemics of obesity and diabetes in the United States.

JAMA; 286(10): 1195-1200.

Millar DB, Hough CJ, Mazorow DL et al (1990):

Beta endorphin's modulation of lymphocyte is done, donor and time dependent.

Brain Behavior and Immunity; 4: 232-242.

Miller WC, Koceja DM and Hamilton EJ (1997):

A meta-analysis of the past 25 years of weight loss research using diet, exercise or diet plus exercise intervention.

Int. J. Obes. Relat. Metab. Disord. 21: 941-947.

Montague CT (1998):

Congenital leptin deficiency is associated with severe early onset obesity in humans.

Nature; 387: 903.

Morgan WP (1997):

Physical activity and mental health.

Taylor and Francis Washington DC; 93-106, 107-127.

Murphy NF, MacIntyre K, Stewart S, et al (2006):

Long-term cardiovascular consequences of obesity: 20-year follow-up of more than 15000 middle aged men and women (the Renfrew-Paisley study).

Eur. Heart J.; 27: 96-106.

National Audit Office (2002):

Tackling obesity in England. Report by the comptroller and auditor general.

HC 220, Session 2000-2001, para 2-13.

National Center for Health Statistics (2002):

Prevalence of overweight and obesity among adults:

United States; 1999.

National Institute of Nutrition (2003):

List of caloric content of different Egyptian food components.

National Task Force for Prevention and Treatment of Obesity (1999):

Very low-calorie diets. Objective: to provide an overview of the published scientific information on the safety and efficacy of very low-calorie diets (VLCDs) and to provide rational recommendations for their use.

Journal of the American Medical Association; 270: 967-974.

National Task Force for Prevention and Treatment of Obesity (2000):

Overweight, obesity and health risk.

Archives of Internal Medicine; 160(7): 898-904.

NICE (National Institute for Clinical Excellence) (2002):

Technical appraisal No 46. Surgery to aid weight reduction for people with morbid obesity.

NICE, London. Online. Available: www.nice.org.uk.

NIH Consensus Development Panel. National Institutes of Health Consensus Development Conference Statement (1991):

Gastrointestinal surgery for severe obesity.

Ann. Intern. Med.; 115: 956-961.

NIH Consensus Conference (1998):

Acupuncture.

JAMA; 280(17): 1518-1524.

Odsberg A, Schill U, and Haker E (2001):

Acupuncture treatment: side effects and complications reported by Swedish physiotherapists.

Complement Ther. Med.; 9: 17-20.

Oleson T (1998):

Differential application of auricular acupuncture for myofascial, autonomic and neuropathic pain.

Medical Acupuncture; 9: 23-28.

Onis M and Blossner M (2000):

Prevalence and trends of overweight among preschool children in developing countries.

Am. J. Clin. Nutr.; 72: 1032-1039.

Pan B, Castro-Lopes JM and Coimbra A (1996):

Activation of anterior lobe corticotrophs by electroacupuncture or noxious stimulations in the anasesthetized rat, as shown by colocalization of fos protein with ACTH and β -endorphin and increased hormone release. Brain Research Bulletin; 40(3): 175-182.

Paylou KN, Krey S and Steffee WP (1989):

Exercise as an adjunct to weight loss and maintenance in moderately obese subjects.

American Journal of Clinical Nutrition; 49: 1115-1123.

Peeters A, Barendregt JJ and Willekens F (2003):

Obesity in adulthood and its consequences for life expectancy: a life-table analysis.

Ann. Intern. Med.; 138: 24-32.

Pi-Sunyer FX (1996):

Obesity. In: Cecil textbook of medicine 20th ed. Edited by: Bennett J and Plum F. Published by: WB Saunders Company. Philadelphia, London, Toronto, Montreal, Sydney and Tokyo; 1167.

Pi-Sunyer FX (2002):

The obesity epidemic: Pathophysiology and consequences of obesity.

Obes. Res.; 10: 975-104.

Pi-Sunyer FX (2003):

A clinical view of the obesity problem.

Science; 299(5608): 859-860.

Plotnikoff NP, Murgo AJ, Miller GC, et al (1985):

Enkephalins: Immunomodulators.

Federation Proceedings; 44(1): 118-122.

Reaven GM (2003):

Importance of identifying the overweight patient who will benefit the most by losing weight.

Ann. Intern. Med.; 138(5): 420-423.

Reilly J, Kelly L, Montgomery C, et al (2007):

Physical activity to prevent obesity in young children:
cluster randomized controlled trial.

BMJ; 14(144): 70-72.

Richards D and Marley J (1998):

Stimulation of auricular acupuncture points in weight loss.

Australian Family Physician 1998; 27(2): 73-77.

Richter WO, Kerscher P and Schwandt P (1983):

Beta endorphin stimulates in vivo lipolysis in the rabbit.

Life Sciences; 33(1): 743-746.

Robbers RE, Kaplan GA, Shema SJ et al (2000):

Are the obese at greater risk for depression?

Am. J. Epidemiol.; 152(2): 163-170.

Roberts RE, Strawbridge WJ, Deleger S, et al (2002):

Are the fat more jolly?

Annals of Benhavioral Medicine; 24(3): 169-180.

Röschke J, Wolf Ch, Muller MJ, et al (2000):

The benefit from whole body acupuncture in major
depression.

J. Affect Dis.; 57: 73-81.

Roth A (2002):

Diabetes mellitus and obesity.

Prim. Care; 29(2): 279-95.

Roux L and Donaldson C (2004):

Economics and obesity: Costing the problem or evaluating
solutions?

Obes. Res.; 12(2): 173-179.

Sacks LL (1975):

Drug addiction, alcoholism, smoking, obesity treated by auricular staplepuncture.

Am. J. Acupuncture 1975; 3: 147-150.

Saris WH (2001):

Very-low-calorie diets and sustained weight loss.

Obesity Research; 9(4): 295S-301S.

Sarwer DB, Wadden TA, and Fabricatore AN (2005):

Psychosocial and behavioral aspects of bariatric surgery.

Obes. Res.; 13(4): 639-648.

Schwartz MW (2001):

Brain pathways controlling food intake and body weight.

Exp. Biol. Med.; 226: 978.

Seidell JC (1999):

Obesity: a growing problem.

Acta. Paediatr.; 88: 46-50.

Seidell JC (2000):

Obesity, insulin resistance and diabetes a worldwide epidemic.

Br. J. Nutr.; 83: 5-8.

Shafshak TS (1995):

Electroacupuncture and exercise in body weight reduction and their application in rehabilitating patients with knee osteoarthritis.

Am. J. Chin. Med. 23: 15-25.

Shang C (1989):

Singular point organizing center and acupuncture point.

American Journal of Chinese Medicine; 17(3-4): 119-127.

Shiffman ML, Sugerman HJ, Kellum JM et al (1992):

Changes in gallbladder bile composition following gallstone formation and weight reduction.

Gastroenterology; 103(1): 214-221.

Shiraishi T, Onoe M, Kojima T, et al (1995):

Effects of auricular stimulation of feeding related hypothalamic neuronal activity in normal and obese rats.

Brain Res. Bull; 36: 141-148.

Shiraishi T, Onoe M, Kojima T, et al (2003):

Effects of bilateral auricular acupuncture stimulation on body weight in healthy volunteers and mildly obese patients.

Exp. Biol. Med.; 228(10): 1201-7.

Skender ML, Goodrick GK, Del Junco DJ, et al (1996):

Comparison of 2-year weight loss trends in behavioral treatments of obesity: diet, exercise and combination interventions.

J. Am. Diet Assoc.; 96: 342-346.

Soong YS (1975):

The treatment of exogenous obesity employing auricular acupuncture.

Am. J. Chin. Med.; 3: 285-287.

Spiegelman BM, Flier JS and Kim SL (2001):

Obesity and the regulation of energy balance.

Cell; 104: 531.

Sramek JJ, Leibowitz MT, Weinstein SP, et al (2002):

Efficacy and safety of sibutramine for weight loss in obese patients with hypertension well controlled by beta-adrenergic blocking agents: a placebo-controlled, double-blind, randomized trial.

J. Hum. Hypertens.; 16: 13-19.

Stafford RS, Farhat JH, Misra B, and Schoenfeld DA (2000):

National patterns of physician activities related to obesity management.

Arch. Fam. Med.; 9(7): 631-638.

Steiner RP (1993):

Acupuncture: cultural perspectives, part 1.

Postgrad. Med. J.; 74: 60-67.

Steppan CM, et al (2001):

The hormone resistin links obesity to diabetes.

Nature 2001; 409(6818): 207-212.

Stux G and Pomeranz B (editors) (1987):

Channel and organs in acupuncture: textbook and atlas. 4th edition. Published by Springer Verlag: Berlin; 125-147.

Summerbell CD, Waters E, Edmunds LD, et al. (2006):

Interventions for preventing obesity in children.

Cochrane Library; (3): 1871.

Sun Q, and Xu Y (1993):

Simple obesity and obesity hyperlipemia treated with otoacupoint pellet pressure and body acupuncture.

J. Tradit. Chin. Med.; 13: 22-26.

Takeshige C, Nakamura A, Asamoto S et al (1992):

Positive feed-back action of pituitary beta endorphin on acupuncture analgesia afferent pathway.

Brain Research Bulletin; 27(1): 37-44.

Tang X (1993):

75 cases of simple obesity treated with auricular and body acupuncture.

J. Tradit. Chin. Med.; 13: 194-195.

Thomas F, Bean K, Pannier B, et al (2005):

Cardiovascular mortality in overweight subjects: the key role of associated risk factors.

Hypertension; 46(4): 654-659.

Toubro S, Astrup A, Breum L, et al (1993):

The acute and chronic effects of ephedrine/caffeine mixtures on energy expenditure and glucose metabolism in humans.

Int. J. Obes. Relat. Metabl. Disord.; 17(3): 73-7.

Town G and Bradley S (1991):

Maximal metabolic responses of deep and shallow water running in trained runners.

Med. and Sci. in Sports and Exercise; 23: 238.

Truswell A (1994):

Obesity.

British Medical Journal. 11: 54-60.

Ukkola O (2002):

Resistin a mediator of obesity associated insulin resistance or an innocent bystander?

European Journal of Endocrinology; 147(5): 571-574.

Ullet GA, Han S and Han JS (1998):

Electroacupuncture: Mechanism and clinical application.

Biological Psychiatry; 44(2): 129-138.

Van loan, M and Mayclin P (1992):

Body composition assessment: Dual Energy X-ray Absorbtiometry (DEXA) compared to reference methods.

Eur. J. Clin. Nutr.; 46: 125-130.

Van Saase JL, Vandenbroucke JP and van Romunde LK (1988):

Osteoarthritis and obesity in the general population. A relationship calling for an explanation.

Journal of Rheumatology, 15(7): 1152-1158.

Vastag B (2003):

Experimental drugs take aim at obesity.

JAMA; 289: 1763-1764.

Vettor R, Pagano C, Fabris R, et al (1993):

Lipolytic effect of beta-endorphin in human fat cells.

Life Sciences; 52(2): 657-661.

Vickers A and Zollman C (1999):

ABC of complementary medicine: acupuncture.

BMJ 1999; 319: 973-976.

Vickers A, Wilkson P and Kleijnen J (2002):

Acupuncture.

Qual. Health Care 2002; 11(1): 92-7.

Vigneri P, Frasca F, Sciacca L, et al (2006):

Obesity and cancer.

Nutr. Metab. Cardiovasc. Dis.; 16: 1-7.

Wadden TA (1993):

Treatment of obesity by moderate and severe caloric restriction: results of clinical research trials.

Annals of Internal Medicine 119: 688-693.

Wadden TA, Brownell KD and Foster GD (2002):

Obesity: responding to the global epidemic.

J. Consult Clin. Psychol.; 70: 510-525.

Waine C (2002):

Obesity and weight management in primary care.

Blackwell Science, Oxford; p. 36.

Walston J, Silver K, Bogardus C, et al (1995):

Time of onset of non-insulin dependent diabetes mellitus and genetic variation of the B₃-adrenergic receptor gene.

N. Engl. J. Med.; 333: 343-347.

Wang K and Liu J (1988):

Needling sensation receptor of an acupoint supplied by the median nerve studies of their electro-physiological characteristics.

American Journal of Chinese Medicine; 17(3-4): 145-155.

Wang JQ, Mao L and Han JS (1992):

Comparison of the antinociceptive effects induced by electroacupuncture and transcutaneous electrical nerve stimulation in the rat.

International Journal of Neuroscience; 65(1-4): 117-129.

Wang H (2002):

Observation on the therapeutic effects of acupuncture for 60 cases of simple obesity.

J. Tradit. Chin. Med.; 22(3): 187-9.

Whaley M (1992):

Predictors of over and under achievement of age predicted maximal heart rate.

J. Med. Science in Sports and Exercise; 24: 173.

White LM, Gardner SF, Gurley BJ, et al (1997):

Pharmacokinetics and cardiovascular effects of mahuang (*Ephedra sinica*) in normotensive adults.

J. Clin. Pharmacol.; 37: 116-22.

White A, Hayhoe S, Hart A, et al (2001):

Adverse events following acupuncture: prospective survey of 32.000 consultations with doctors and physiotherapists.

BMJ; 323: 485-6.

Widen E, Lehto M, Kanninen T, et al (1995):

Association of a polymorphism in the B₃-adrenergic receptor gene with features of the insulin resistance syndrome in Finns.

N. Eng. J. Med.; 333: 348-351.

Wild SH and Byrne CD (2007):

ABC of obesity. Risk factors for diabetes and coronary heart disease.

BMJ Middle East; 14(144): 121-123.

Wing RR (2001):

Behavioral treatment of obesity. In: Obesity: theory & therapy. Edited by: Wadden TA, Stunkard AJ. Published by: Guilford: New York, 162-175.

Wissler RW and Strong JP (1998):

Risk factors and progression of atherosclerosis in youth.

Am. J. Pathol.; 153(4): 1023-1033.

Wolf AM (2002):

Economic outcomes of the obese patient.

Obes. Res.; 10: 58S-62.

Wong AM, Su TY, Tang FT, et al (1999):

Clinical trial of electrical acupuncture on hemiplegic patients.

American Journal of Physical Medicine and Rehabilitation; 78: 117-122.

World Health Organization (1997):

Obesity: preventing and managing the global epidemic.

WHO Technical Report Series, No. 984. Geneva, World Health Organization. Online, Available:

www.who.int/nut/publications.htm.

Yaegashi M, Jean R, Zuriqat M, et al (2005):

Outcome of morbid obesity in the intensive care unit.

J. Intensive. Care. Med.; 20(3): 147-154.

Yamashita H, Tsukayama H, Tanno Y et al (1998):

Adverse events related to acupuncture.

JAMA 1998; 280: 1563-1564.

Yamashita H, Tsukayama H, Tanno Y, et al (1999):

Adverse events in acupuncture and moxibustion treatment: a six-year survey at a national clinic in Japan.

J. Altern. Complement Med.; 5: 229-36.

Yang R, Xu A, Pray J, et al (2003):

Cloning of omentin, a new adipocytokine from omental fat tissue in humans. Program and abstracts of the 63rd. Scientific Session of the American Diabetes Association; 13-17; New Orleans, Online. Available: www.medscape.com/viewarticle/458393.

Yu Y, Kasahara T, Sato T, et al (1997):

Enhancement of splenic interferon-gama, interleukin-2, NK cytotoxicity by S36 acupoint acupuncture in 344 rats.

Japanese Journal of Physiology; 47(2): 173-178.

Yu Y, Kasahara T, Sato T, et al (1998):

Role of endogenous interferon-gama on the enhancement of splenic NK cell activity by electroacupuncture stimulation in mice.

Journal of Neuroimmunology; 90(2): 176-186.

Zhao Y, Yang C and Liu Z (1992):

Effect of acupuncture on carbohydrate metabolism in patients with simple obesity.

J. Tradit. Chin. Med.; 12: 129-132.

Zhao M, Liu Z and Su J (2000):

The time effect relationship of central action in acupuncture treatment for weight reduction.

J. Tradit. Chin. Med.; 20: 26-29.

Zollman C and Vickers A (1999):

ABC of complementary medicine. Users and practitioners of complementary medicine.

BMJ; 319: 836-8.