

- **Hutchin, C.J., Ferreira, C. J., Norman, T. J. Q. (1985):** A comparison of local agents in the relief of discomfort after episiotomy. American Journal of Obstetric and Gynecology, 66, 45-46.
- **Abraham, S. (1990):** Recovery after childbirth: A preliminary prospective study. Medical Journal, 152 (1), 9-12.
- **Attia, A. A., Guindi, F. K., & Abed El-Razek, A. A. (1999):** The effect of practicing physical measures on relieving lower Back discomfort during perinatal period. The New Egyptian Journal of Medicine , 20 (4), 26-35.
- **Bruce, E. (2006):** Episiotomy versus intact perineal located at <http://www.femalepatient.com>
- **Kettle C, Johanson R. B (2000b):** Continuous versus interrupted sutures for perineal repair. Cochrane Review 3. the Cochrane Library, Oxford.
- **Lita, J.P. (1997):** Postpartum perineal pain. Midwifery Journal, 10, 256.
- **Synder RR, Hammond TL, Hankins GD (1996):** Episiotomy and body image. Mod Midwife, 31: 18-19.
- **Takano, A. (2006):** Benefits of honey located at <http://www.anwaar.net>.
- **Thacker SB, Banta HD. (1983):** Benefits and risks of episiotomy: an interpretative review of the English language literature 1860-1980. Obstet Gynecol ; 38: 322-338.
- **Viktrup L, Lose G, Rolff M, Barfoed K. (1992):** The symptom of stress incontinence caused by pregnancy or delivery in primiparas. Obsetet Gynecol; 79: 945-949.