## **REFERNCES**

- American Association of Occupational Health Nurses (AAOHN), (1999): Delivery of occupational and environmental health services. New York: Atlanta C, PP.3-5-11.
- **Aarås,A., Dainoff, M., and Thoresen, M. (2002):** Can a more neutral position of the forearm when operating a computer mouse reduce the pain level for VDU operators?, International Journal of Industrial Ergonomics 30, pp. 307–324.
- **Al-Wehedy, A. Hammad, S.M., and Al-Gilany, A. (2000):** Some health hazards among computer users.Benha Medical Journal, 16(3), PP. 601-615.
- **Allender, J. and Spradly,B. (2001):** Community health nursing concepts and practice, 5<sup>th</sup> ed., Lippincott, New York, PP.586-597.
- **Allender, J. and Spradly,B.** (2005): Community health nursing:Promoting and protecting the public's health, 6<sup>th</sup> ed., Lippincott, New York, PP.700-710.
- Andersen J.H., Thomsen, J.F., Overgaard, E., and Lassen C.F. (2003): Computer use and carpal tunnel syndrome: a 1-year follow-up study, Journal of the American Medical Association 289, pp. 2963–2969.
- **Aptel, M., Aublet-Cuvelier, A., and Cnockaert, J.C. (2002):** Work-related musculoskeletal disorders of the upper limb, Joint Bone Spine 69, pp. 546–555.

- Armstrong, T. J., Foulke, J. A., Bernard. j., Gerson, J., and Remple, D.M. (200): Investigation of applied forces in alphanumeric work. Am Ind Hyg Assoc, 55, PP. 30-35.
- **Basavanthappa, B.T. (1998):** Community health nursing, 1<sup>st</sup> ed., Jaypee Brothers Publishers, Bangalore, PP. 546-549.
- **Bergamasco, R., Girola, C., and Colombini, D.(1998):** Guidelines for designing jobs featuring repetitive tasks Ergonomics 41, pp. 1364–1383.
- **Blatter, B,M., Bongers, P.M.** (2002): Duration of computer and mouse use in relation to musculoskeletal disorders of neck or upper limb, International Journal of Industrial Ergonomics, 30(5), PP. 295-306.
- **Bohr,P.C.** (2000): Efficacy of office ergonomics education, Occupational Rehabilitation Journal 10, PP. 243-255.
- Brisson, S., Montreuil, L., and Punnett, L. (1999): Effects of an ergonomic training program on workers with video display units, Scand J Work Environ Health 25, PP. 255-263.
- **Butler, A., Selmond, S., and Pellino, T.(1998):** Orthopedic nursing, 2<sup>nd</sup> ed., W.B Saunders company, Philadelphia, PP. 65-66.
- Burgess-Limerick, R. Shemmell, J., R. Scadden, R., and Plooy, A. (1999): Wrist posture during computer pointing device use, Clinical Biomechanics 14, pp. 280–286.

- Carayon, P., Smith, M.J. and Haims, M. C. (1999): Work organization, job stress, and work related musculoskeletal disorders, Hum Factors, 41(1), PP644-663.
- Centers for Disease Control and Prevention (2000): Computer Workstation Ergonomics. Available online: http://www.cdc.gov/od/ohs/Ergonomics/compergo.htm.
- **Chris,A., (2006):** Guide to setting up an ergonomic computer station; http:// ergonomics.about.com /od/office/ss/computer-setup4,hTm.
- Cook, C., Burgess-Limerick, R., and Chang, S. (2000): The prevalence of neck and upper extremity musculoskeletal symptoms in computer mouse users, International Journal of Industrial Ergonomics 26, pp. 347–356.
- Coury, G., Kumar, S., Rodgher, S., and Narayan, Y.(1998):

  Measurement of shoulder adduction strength in different postures.

  International Journal of Industrial Ergonomics 22, pp. 195–206.
- **Croasmun J. (2002):** Easy on eyes= better on the job. Ergonomic today TM. Available at: http://www.ergoweb.com.
- **Division of Safety, National Institutes of Health (2004):** Ergonomics: Maintain Your Curves. Available online: http://www.nih.gov/od/ors/ds/ergonomics/index.html.

- Dowler, E., Kappes,B., Fenaughty,A., and Pemberton,G. (2001): Effects of neutral poture on muscle tension during computer use. International Journal of occupational safety and ergonomics,7 (1), PP. 61-78.
- **Duff,S.V.(1997):** Tendinitis, entrapment neuropathies, and related conditions. In M. J. Sanders(Ed.), Management of cumulative trauma disorders. Boston: Butterworth-Heinemann,PP41-103
- **Emara, AM.(2001):** Health hazards associated with work on video display terminals A WHO consideration. Egyptian Occup Med, 15 (1), PP. 105-108.
- **Fagarasanu,M., Kumar, S.** (2003): Carpal tunnel syndrome due to keyboarding and mouse tasks: International Journal of Industrial Ergonomics 31, PP. 119-136.
- **Hickey,J., Ouimette,R., and Venegoni, s. (2000):** Advanced practice nursing: changing roles and clinical application, 2<sup>nd</sup> ed., Lippincott, New York, P. 348.
- **Gerr, F. (2002):** A prospective study of computer users: I. Study design and incidence of musculoskeletal symptoms and disorders. American Journal of Industrial Medicine, 41(4): 221–235.
- Gerr, F., Marcus, M., and Monteilh, C. (2004): Epidemiology of musculoskeletal disorders among computer users: lesson learned from the role of posture and keyboard use. Electromyoger Kinesiol, 14(1), PP. 25-31.

- Greene, L. I., David, M., and Olejnik, S. (2005): Effects of an active ergonomics training program on risk exposure, worker beliefs, and symptoms in computer users, International Journal of Industrial Ergonomics, 24 (1), PP. 41-52.
- **Guetschow, J., (2000):** New ergonomic lines. Ergonomic today TM. Available at: http://www.ergoweb.com.
- Guidelines for Office Ergonomics. (2006): Computer and workstation equipment,(6), P. 6105.
- Hamza, E., Abd Elmohsen, M. (2004): Introduction of computer science, Zagazig University, PP.16-17.
- Hernandes, S., Tamez, G., Matinez, A., and Mendez,R. (2003):

  Computer use increases the risk of musculoskeletal disorders among newspaper office workers, Archives of Medical Research 34,PP. 331-342.
- Idowu Bayo, Ogunbodede Eyitope, Idowu Bimbo (2003): Information And Communication Technology In Nigeria: The Health Sector Experience Journal Of Information Technology Impact ,3 (2). 69-76. ISSN 1098-1
- **Jensen, C.** (2003): Development of neck and hand-wrist symptoms in relation to duration of computer use at work. Scand J Work Environ Health, 29(3), PP. 197-205.

- **Joan, M.** (1996): Nursing care in the community, 2<sup>nd</sup> ed., Mosby company., PP. 582-596.
- **Jones, C.S., and Orr, B. (1998):** Computer related musculoskeletal pain and discomfort among high school students. Am. J. Health study, 14 (3), P. 26.
- **Jorgen, R.** (2004): Upper limb neuropathy in computer operators. Aclinical case study of 21 patient. BNC, Musculoskeletal disord 5, P. 26.
- **Ladou, S. (1997):** Occupational and environmental medicine, 2<sup>nd</sup> ed., Schuster comp., New Jersey., PP41-63
- **Kaplan and Sadock's (2001):** Anxiety disorder. In Synopsis of Psychiatry: Behavioural sciences and Psychiatry, 8<sup>th</sup> ed , Giza. Mass publishing co: PP 583-584.
- Ketola, R., Toivonen, M., Hakkanen, R., and Takala, E. (2002):

  Effects of ergonomic intervention in work with video display units, Scand J Work Environ Health 28, PP. 18-24.
- **Khan, K. M. (2001):** Overuse tendinosis, not tendonitis Pare 1: Anew paradigm for a difficult clinical problem, The Physician and Sportsmedicine 28, PP53-60.
- **Knowles,M.** (1998): The adult learner. A neglected species, 5<sup>th</sup> ed., Gulf publishing comp Houston.
- **Ladou, S. (1997):** Occupational and environmental medicine, 2<sup>nd</sup> ed., Schuster comp., New Jersey., PP41-63.

- Lamiaa, A., Sahar, S., and Amel, a. (2006): Health complaints associated with prolonged computer use as perceived by computer usres, 5 (1), PP. 65-75.
- **Liao, M.H. and Drury, C.G. (2000):** Posture, discomfort and performance in a VDT task. Ergonomics 43, pp. 345–359.
- **Lim, S.Y, Carayon, P. (2001):** An integrated approach to cumulative trauma disorders in computerized offices: the role of psychosocial work factors, psychological stree, and ergonomic risk factore, Advances in human factors /ergonomics, 19, PP 387-400.
- Lincoin, A.E., Vernick, J.S., Ogaitis, S., Smith, G.S., Mitchell, C.S., and J. Agnew, J. (2000): Interventions for the primary prevention of work-related carpal tunnel syndrome. American Journal of Preventive Medicine 18, pp. 37–50.
- **Linton S.J.,and Tulder, M.W. (2001).** Preventive interventions for back and neck pain problems: What is the evidence? Spine, 26(7): 778–787.
- MANOJ, D. (2005): Movements International Yoga for computer related health problems, Manoj Dash; http://groups.yahoo.com/group/Yoga.
- Mani, L., and Gerr, F. (2000): Work-related upper extremity musculoskeletal disorders. Occupational and Environmental Medicine 27 (2000), pp. 845–864.

- Marklin, R.W. and Simoneau, G.G.(2001): Effect of setup configurations of split computer keyboard on wrist angle. Physical Therapy 81, pp. 1038–1048.
- **Marras, W.S.** (2000): Occupational low back disorder causation and control. Ergonomics, 43(7): 880–902.
- Mekhora, K., Listonc, S., and Coleb,H. (2000): The effect of ergonomic intervention on discomfort in computer users with tension neck syndrome, International Journal of Industrial Ergonomics, 26 (3), PP. 367-379.
- Moustafa,T., and Maged,z.(2005): Introduction of computer science, Zagazig University,P.9.
- Nelson, J.E, Treaster, D.E., and W.S. Marras, W.S. (2000): Finger motion, wrist motion and tendon travel as a function of keyboard angles. Clinical Biomechanics 15 (2000), pp. 489–498.
- Nevid, s., Rathus, A., and Rubenstein, H. (1998): Health in the new millennium, 1<sup>st</sup> ed., Worth publishers, New York, PP. 173-174.
- Nies, M. A., and McEwen, M.(2001): Community health nursing, 3<sup>rd</sup> ed., Saunders comp., U.S.A., PP. 731-735.
- Palmer, K.T, Cooper, C., Walker, K., Syddall. H, and Coggon, D. (2001): Use of Keyboards and Symptoms in the Neck and arm: Evidence from a national survey. Occup. Med (Lond). Sept. 5196): 392-5

- **Pascarellis, E.F., and Kella, J.J. (2001):** Soft tissue injuries related to use of the computer keyboard: a clinical study of 35 severly injured persons. J Occup Med 3, PP. 522-532.
- **Peddie,S.** (1997): The repetitive strain injury sourcebook. Los Angeles: Lowell House, PP. 45-70
- **Pillinger J and Rutherford (2003):** Computer and neck pain. Health News. 39: 12-14.
- **Reda,T.** (2005): Introduction of computer science, Zagazig University, PP.14-16.
- Rempel, D.M., Harrison, R.J., and Barnhart, S. (2001): Work related cumulative trauma disorders of the upper extremity. JAMA 267, PP. 838.
- **Rogers, B.** (1998): Ocupational health Nursing Expertise. AAOHN Journal 46 (10), PP. 447-483.
- **Rogers, B.** (2003): Occupational health nursing: concepts and practice, St Louis, Mo, Elsevier, 16(1), PP. 143-161.
- Rufus A. Adedoyin, B., Idowu, O., and Rotimi E. (2004):

  Musculoskeletal Pain Associated With The Use Of Computer

  Systems In Nigeria. The Internet Journal of Pain, Symptom

  Control and Palliative Care, 3(2), PP. 197-212
- **Salazar, K.** (2001): Core curriculum for occupational and environment health nursing, 2<sup>nd</sup> ed, W.B Saunders company, U.SA., PP. 138-145.

- **Sauter,S.** (2001): Predictors of strain in VDT users and traditional office workers In:Grandijean E. Ergonomics and health modern offices. London Taylor and Francis, PP. 129-135.
- **Schneider, F. (2001):** Four steps to an effective office ergonomics program. Occupational Health and Safety, 70(9): 140–144.
- **Schuchmanm, J. (2002):**Occupational rehabilitation, 2<sup>nd</sup> ed.,U.S.A,W.B. Saunders company, PP.985-619.
- Serina, E., Tal, R., and Rempel, D. (1999): Wrist and forearm postures and motions during typing. Ergonomics 42 ,pp. 938–951
- **Silverton, L., (1997):** The art and science of midwifery, 5<sup>th</sup> ed., prentice Hall international comp., London, PP. 174-181.
- **Simoneau, G.G., Marklin, R.W., and Monroe, J.F. (2001):** Wrist and forearm postures of users of conventional computer keyboards. Human Factors 41, pp. 413–424.
- **Smith, M. j., Bayehi, A.D. (2003):** Do ergonomics improvements increase workers' productivity: An intervention study in a call center., Ergonomics, 46 (3), PP.3-16.
- Smith, M.J., Cohen, B.G., and Stammerjohn, L.W. (2001): An investigation of health complaints and job stress in video display operations. Hum Factors, 23, PP. 387-400.
- **Stanhope, M., and Lancaster, J.(2004):** Community and public health nursing, 6<sup>th</sup> ed, U.S.A, Mosby company,pp1066-1091.

- **Stanhope, M., and Lancaster, J.(2002):** Foundations of community health nursing,1<sup>st</sup> ed, Mosby comp., U.S.A, pp. 528.
- **Tepper, M., Voollenbroek, H. J, and Hermens, S. (2003):** The effect of ergonomic computer device on muscle activity of the upper trapezius muscle during typing, Applied Ergonomics, 34 (2), PP. 125-130.
- **Travers, P.H And Stanton, B.A.** (2002): Office workers and video display terminals: physical, psychological and ergonomic factors. AAOHN j, 50(11), PP. 489-493.
- **Turner, P. J. (2000):** Visual requirement for VDT operators. Australian Journal of Optometry, 65, PP. 58-64.
- **Verhagen AP, et al. (2006):** Ergonomic and physiotherapeutic interventions for treating work-related complaints of the arm, neck or shoulder in adults. Cochrane Database of Systematic Reviews (4). Oxford: Update Software.
- **Visser, B. Korte, E., and Kuijer, P. (2000):** The effect of arm and wrist supports on the load of the upper extremity during VDU work, *Clinical Biomechanics* 15 ,PP. 34–38.
- Waheed, A., Fouad, M., Bassiouni, H., and Abdel-Aziz, M. (2000): Health hazards of visual display terminals among exposed workers in Suez Canal cities. Mansoura Med ,30 (2), PP. 15-40.

- **Wassell JT, et al. (2000):** A prospective study of back belts for prevention of back pain and injury. JAMA, 284(21): 2727–2732.
- Werner, R., Armstrong, C., Bir, M., And Aylard, K. (1997):
  Intracarpal canal pressures: the role of finger, hand, wrist and forearm position, Clinical Biomechanics 12, PP. 44-51.
- Wilkinson, W. (1990): A conceptual Model of Occupational Health Nursing, AAOHN Journal 38(2), PP. 71-7.
- World Health Organization (WHO), (2001):Video display Units and human health, Fact Sheet No. 201. Available at:fact201.html www.who.int/inf-fs/en/