

# CONTENTS

	Page
<b>CHAPTER 1</b>	
<b>INTRODUCTION AND AIM OF THE WORK</b>	
INTRODUCTION.....	1
AIM OF THE WORK .....	4
<b>CHAPTER 2</b>	
<b>LITERATURE REVIEW</b>	
1- General remarks about venerid clams.....	5
2- Morphological characters used in venerids taxonomy.....	8
3- Application of RAPD-PCR technique in taxonomy.....	9
4- Biomonitor/bioindicators.....	13
5- Interactions between dissolved metals and aquatic organisms.....	14
5-1- Metal speciation.....	14
5-2- Metal bioaccumulation.....	16
5-3- Metal uptake and depuration.....	17
5-4- Metal toxicity and detoxification.....	19
6- Oil pollution.....	22
7- The use of biomarkers in biomonitoring.....	23
7-1- Histopathological biomarker.....	27
7-2- Genetic ecotoxicological biomarker.....	29
<b>CHAPTER 3</b>	
<b>MATERIALS AND METHODS</b>	
1- Study areas.....	32
1-1- Mediterranean Sea stations.....	32
1-2- Suez Canal stations.....	33
1-3- Red Sea stations.....	33
2- Morphological investigation.....	33
3- Ecological investigation.....	33
3-1- Physico-chemical parameters determination.....	34
3-1-1- Water temperature (°C).....	34
3-1-2- Hydrogen ion concentration (pH).....	34
3-1-3- Salinity (‰).....	34
3-1-4- Dissolved oxygen (DO) mg/l.....	34
3-2- Heavy metals determination.....	34
3-2-1- Heavy metals in surface sea water.....	34
3-2-2- Heavy metals in sediment.....	35
3-2-3- Heavy metals in soft tissues.....	35
3-2-4- Heavy metals in shells.....	36
3-3- Total petroleum hydrocarbons (TPHs) determination.....	37
4- Histopathological study.....	37

## CONTENTS/CONT.

	Page
5- Genetic diversity.....	38
5-1- Different species from the same station.....	38
5-1-1- Genomic DNA extraction and purification.....	38
5-1-2- Polymerase Chain Reaction (PCR) amplification.....	39
5-1-3- Preparations for gel electrophoresis.....	41
A- Running buffers.....	41
B- Ethidium bromide.....	41
C- Sample loading dye (5x).....	41
D- 1.5% Agarose gel preparation.....	41
5-2- One species from different stations.....	42
6- Statistical analysis.....	42
<b>CHAPTER 4</b>	
<b>RESULTS</b>	
1- Morphological investigation.....	44
1-1- Identification of the collected samples.....	44
1-2- Common venerid ( <i>Tapes decussatus</i> ) collected from the different stations.....	53
2- Ecological investigation.....	59
2-1- Physico-chemical parameters of water.....	59
2-1-1- Temperature (°C).....	59
2-1-2- Hydrogen ion concentration (pH).....	59
2-1-3- Salinity (‰) .....	60
2-1-4- Dissolved oxygen (DO).....	60
2-2- Heavy metals in surface sea water.....	65
2-3- Heavy metals in sediment.....	77
2-4- Heavy metals in <i>Tapes decussatus</i> whole soft tissues.....	89
2-5- Heavy metals in <i>Tapes decussatus</i> shell.....	101
2-6- Comparison between heavy metals content in surface sea water, sediment and <i>Tapes decussatus</i> whole soft tissues and shell overall stations and seasons.....	113
2-7- Total petroleum hydrocarbons (TPHs) in surface sea water.	113
3- Histopathological studies.....	116
3-1- Parasitic infection.....	116
3-1-1 Description of digenean larval stages inside the clam ...	116
3-1-1-1- <i>Cercaria lata</i> Lespés, 1857 larvae.....	116
3-1-1-2- Unknown Digenean sp. Larvae.....	118
3-1-2- Prevalence of infection.....	119
3-2- Histopathological investigations.....	130
A- Gonads.....	130
B- Digestive gland.....	131

## CONTENTS/CONT.

	<b>Page</b>
4- Genetic diversity.....	143
4-1- RAPD profiles for the collected venerid clams from Ismailia.....	143
4-2- RAPD profile for <i>Tapes decussatus</i> collected from different stations.....	164
<b>CHAPTER 5</b>	
<b>DISCUSSION</b>	
DISCUSSION.....	176
<b>SUMMARY</b> .....	192
<b>REFERENCES</b> .....	198
<b>ARABIC SUMMARY</b> .....	\

## LIST OF TABLES

	Page
<b>Table (1):</b> List of arbitrary primers sequences and their G+C contents used in RAPD analysis of venerid clams.	40
<b>Table (2):</b> Venerid clams species and stations of collection.	54
<b>Table (3):</b> Seasonal variations in temperature (°C) and hydrogen ion concentration (pH), of surface sea water at different stations in 2006.	61
<b>Table (4):</b> Seasonal variations in salinity (‰) and dissolved oxygen (DO mg/l) of surface sea water at different stations in 2006.	62
<b>Table (5):</b> Seasonal variations in Cd and Co concentrations (µg/ml) in surface sea water at different stations in 2006.	69
<b>Table (6):</b> Seasonal variations in Cu and Fe concentrations (µg/ml) in surface sea water at different stations in 2006.	70
<b>Table (7):</b> Seasonal variations in Mn and Ni concentrations (µg/ml) in surface sea water at different stations in 2006.	71
<b>Table (8):</b> Seasonal variations in Pb and Zn concentrations (µg/ml) in surface sea water at different stations in 2006.	72
<b>Table (9):</b> Seasonal variations in Cd and Co concentrations (µg/ g dry weight) in sediment at different stations in 2006.	81
<b>Table (10):</b> Seasonal variations in Cu and Fe concentrations (µg/ g dry weight) in sediment at different stations in 2006.	82
<b>Table (11):</b> Seasonal variations in Mn, and Ni concentrations (µg/g dry weight) in sediment at different stations in 2006.	83
<b>Table (12):</b> Seasonal variations in Pb and Zn concentrations (µg/g dry weight) in sediment at different stations in 2006.	84
<b>Table (13):</b> Seasonal variations in Cd and Co concentrations (µg/ g dry weight) in whole soft tissues of <i>Tapes decussatus</i> collected from different stations in 2006.	93
<b>Table (14):</b> Seasonal variations in Cu and Fe concentrations (µg/ g dry weight) in whole soft tissues of <i>Tapes decussatus</i> collected from different stations in 2006.	94
<b>Table (15):</b> Seasonal variations in Mn and Ni concentrations (µg/g dry weight), in whole soft tissues of <i>Tapes decussatus</i> collected from different stations in 2006.	95
<b>Table (16):</b> Seasonal variations in Pb and Zn concentrations (µg/g dry weight) in whole soft tissues of <i>Tapes decussatus</i> collected from different stations in 2006.	96

## LIST OF TABLES/CONTINUE

	Page
<b>Table (17):</b> Seasonal variations in Cd and Co concentrations ( $\mu\text{g/g}$ dry weight) in <i>Tapes decussatus</i> shells collected from different stations in 2006.	105
<b>Table (18):</b> Seasonal variations in Cu and Fe concentrations ( $\mu\text{g/g}$ dry weight) in <i>Tapes decussatus</i> shells collected from different stations in 2006.	106
<b>Table (19):</b> Seasonal variations in Mn and Ni concentrations ( $\mu\text{g/g}$ dry weight) in shells of <i>Tapes decussatus</i> collected from different stations in 2006.	107
<b>Table (20):</b> Seasonal variations in Pb and Zn concentrations ( $\mu\text{g/g}$ dry weight) in shells of <i>Tapes decussatus</i> collected from different stations in 2006.	108
<b>Table (21):</b> Comparison between heavy metals content in surface sea water, sediment and <i>Tapes decussatus</i> whole soft tissues and shell overall stations and seasons.	114
<b>Table (22):</b> Monthly infection prevalence of digenean larvae during sampling period for <i>Tapes decussatus</i> populations collected from Ismailia and Suez.	127
<b>Table (23):</b> Monthly infection prevalence of digenean larvae during sampling period for <i>Venerupis pullastra</i> and <i>Venerupis aureus</i> collected from Ismailia.	128
<b>Table (24):</b> Number of amplified bands based on RAPD arbitrary primers for seven species of venerid clams collected from Ismailia.	160
<b>Table (25):</b> Number of monomorphic bands based on RAPD arbitrary primers for seven species of venerid clams collected from Ismailia.	160
<b>Table (26):</b> Number of polymorphic bands based on RAPD arbitrary primers for seven species of venerid clams collected from Ismailia.	161
<b>Table (27):</b> Number of unique bands based on RAPD arbitrary primers for seven species of venerid clams collected from Ismailia.	161
<b>Table (28):</b> Size range of bands (molecular weights in base pairs [bp]) based on RAPD arbitrary primers for seven species of venerid clams collected from Ismailia.	162
<b>Table (29):</b> Genetic similarity matrix index calculated by pairwise comparison of seven species of venerid clams collected from Ismailia.	163

## LIST OF TABLES/CONTINUE

	Page
<b>Table (30):</b> Number of amplified and monomorphic bands based on RAPD arbitrary primers for <i>Tapes decussatus</i> collected from different stations.	173
<b>Table (31):</b> Number of polymorphic and unique bands based on RAPD arbitrary primers for <i>Tapes decussatus</i> collected from different stations.	173
<b>Table (32):</b> Size range of bands (molecular weights in base pairs [pb]) based on RAPD arbitrary primers for <i>Tapes decussatus</i> collected from different stations.	174
<b>Table (33):</b> Genetic similarity matrix index values calculated by pairwise comparison of four populations of <i>Tapes decussates</i> collected from different stations.	175

## LIST OF FIGURES

	Page
<b>Fig. (1):</b> A map showing the study areas.	32
<b>Fig. (2):</b> Seasonal variations in temperature at the investigated stations.	63
<b>Fig. (3):</b> Seasonal variations in pH at the investigated stations.	63
<b>Fig. (4):</b> Seasonal variations in salinity at the investigated stations.	64
<b>Fig. (5):</b> Seasonal variations in DO at the investigated stations.	64
<b>Fig. (6):</b> Seasonal variations in Cd concentration of surface sea water at the investigated stations.	73
<b>Fig. (7):</b> Seasonal variations in Co concentration of surface sea water at the investigated stations.	73
<b>Fig. (8):</b> Seasonal variations in Cu concentration of surface sea water at the investigated stations.	74
<b>Fig. (9):</b> Seasonal variations in Fe concentration of surface sea water at the investigated stations.	74
<b>Fig. (10):</b> Seasonal variations in Mn concentration of surface sea water at the investigated stations.	75
<b>Fig. (11):</b> Seasonal variations in Ni concentration of surface sea water at the investigated stations.	75
<b>Fig. (12):</b> Seasonal variations in Pb concentration of surface sea water at the investigated stations.	76
<b>Fig. (13):</b> Seasonal variations in Zn concentration of surface sea water at the investigated stations.	76
<b>Fig. (14):</b> Seasonal variations in Cd concentration of sediment at the investigated stations.	85
<b>Fig. (15):</b> Seasonal variations in Co concentration of sediment at the investigated stations.	85
<b>Fig. (16):</b> Seasonal variations in Cu concentration of sediment at the investigated stations.	86
<b>Fig. (17):</b> Seasonal variations in Fe concentration of sediment at the investigated stations.	86
<b>Fig. (18):</b> Seasonal variations in Mn concentration of sediment at the investigated stations.	87
<b>Fig. (19):</b> Seasonal variations in Ni concentration of sediment at the investigated stations.	87
<b>Fig. (20):</b> Seasonal variations in Pb concentration of sediment at the investigated stations.	88
<b>Fig. (21):</b> Seasonal variations in Zn concentration of sediment at the investigated stations.	88

## LIST OF FIGURES /CONTINUE

	Page
<b>Fig. (22):</b> Seasonal variations in Cd concentration of <i>Tapes decussatus</i> whole soft tissues at the investigated stations.	97
<b>Fig. (23):</b> Seasonal variations in Co concentration of <i>Tapes decussatus</i> whole soft tissues at the investigated stations.	97
<b>Fig. (24):</b> Seasonal variations in Cu concentration of <i>Tapes decussatus</i> whole soft tissues at the investigated stations.	98
<b>Fig. (25):</b> Seasonal variations in Fe concentration of <i>Tapes decussatus</i> whole soft tissues at the investigated stations.	98
<b>Fig. (26):</b> Seasonal variations in Mn concentration of <i>Tapes decussatus</i> whole soft tissues at the investigated stations.	99
<b>Fig. (27):</b> Seasonal variations in Ni concentration of <i>Tapes decussatus</i> whole soft tissues at the investigated stations.	99
<b>Fig. (28):</b> Seasonal variations in Pb concentration of <i>Tapes decussatus</i> whole soft tissues at the investigated stations.	100
<b>Fig. (29):</b> Seasonal variations in Zn concentration of <i>Tapes decussatus</i> whole soft tissues at the investigated stations.	100
<b>Fig. (30):</b> Seasonal variations in Cd concentration of <i>Tapes decussatus</i> shell at the investigated stations.	109
<b>Fig. (31):</b> Seasonal variations in Co concentration of <i>Tapes decussatus</i> shell at the investigated stations.	109
<b>Fig. (32):</b> Seasonal variations in Cu concentration of <i>Tapes decussatus</i> shell at the investigated stations.	110
<b>Fig. (33):</b> Seasonal variations in Fe concentration of <i>Tapes decussatus</i> shell at the investigated stations.	110
<b>Fig. (34):</b> Seasonal variations in Mn concentration of <i>Tapes decussatus</i> shell at the investigated stations.	111
<b>Fig. (35):</b> Seasonal variations in Ni concentration of <i>Tapes decussatus</i> shell at the investigated stations.	111
<b>Fig. (36):</b> Seasonal variations in Pb concentration of <i>Tapes decussatus</i> shell at the investigated stations.	112
<b>Fig. (37):</b> Seasonal variations in Zn concentration of <i>Tapes decussatus</i> shell at the investigated stations.	112
<b>Fig. (38):</b> Comparison between heavy metals content in surface sea water, sediment and <i>Tapes decussatus</i> whole soft tissues and shell overall stations and seasons.	115
<b>Fig. (39):</b> Monthly infection prevalence of digenean larvae in (A) <i>Tapes decussatus</i> (Ismailia), (B) <i>Tapes decussatus</i> (Suez), (C) <i>Venerupis pullastra</i> and (D) <i>Venerupis aureus</i> .	129



## LIST OF FIGURES /CONTINUE

	Page
<b>Fig. (40):</b> Photograph of transverse section through the normal testis of <i>Tapes decussatus</i> (A) scale bar = 100 $\mu\text{m}$ and (B) scale bar = 25 $\mu\text{m}$ .	133
<b>Fig. (41):</b> Photograph of transverse section through the testis of <i>Tapes decussatus</i> (A) normal and (B) light infected by <i>Cercaria lata</i> larvae.	134
<b>Fig. (42):</b> Photograph of transverse section through the testis of <i>Tapes decussatus</i> infected by <i>Cercaria lata</i> larvae (A) light infection and (B) moderate infection.	135
<b>Fig. (43):</b> Photograph of transverse section through the testis of <i>Tapes decussatus</i> infected by <i>Cercaria lata</i> larvae (A) moderate infection, scale bar = 5 $\mu\text{m}$ and (B) heavy infection, scale bar = 400 $\mu\text{m}$ .	136
<b>Fig. (44):</b> Photograph of transverse section through the normal ovary of <i>Tapes decussatus</i> (A) scale bar = 125 $\mu\text{m}$ and (B) scale bar = 50 $\mu\text{m}$ .	137
<b>Fig. (45):</b> Photograph of transverse section through the ovary of <i>Tapes decussatus</i> (A) normal and (B) light infected by <i>Cercaria lata</i> larvae.	138
<b>Fig. (46):</b> Photograph of transverse section through the ovary of <i>Tapes decussatus</i> infected by <i>Cercaria lata</i> larvae (A) moderate infection and (B) heavy infection.	139
<b>Fig. (47):</b> Photograph of transverse section through the normal digestive gland of <i>Tapes decussatus</i> (A) scale bar = 100 $\mu\text{m}$ and (B) scale bar = 15 $\mu\text{m}$ .	140
<b>Fig. (48):</b> Photograph of transverse section through the digestive gland of <i>Tapes decussatus</i> collected from Suez (A) uninfected, scale bar = 15 $\mu\text{m}$ and (B) infected by <i>Cercaria lata</i> larvae , scale bar = 100 $\mu\text{m}$ .	141
<b>Fig. (49):</b> Photograph of transverse section through the of digestive gland of <i>Tapes decussatus</i> infected by <i>Cercaria lata</i> larvae (A) scale bar = 100 $\mu\text{m}$ and (B) scale bar = 10 $\mu\text{m}$ .	142
<b>Fig. (50):</b> Bands generated by PCR using RAPD primer (P1) for the seven species of venerid clams (A) amplified bands and (B) bands types.	147
<b>Fig. (51):</b> Bands generated by PCR using RAPD primer (P2) for the seven species of venerid clams (A) amplified bands and (B) bands types.	148
<b>Fig. (52):</b> Bands generated by PCR using RAPD primer (P3) for the seven species of venerid clams (A) amplified bands and (B) bands types.	149

## LIST OF FIGURES /CONTINUE

	Page
<b>Fig. (53):</b> Bands generated by PCR using RAPD primer (P5) for the seven species of venerid clams (A) amplified bands and (B) bands types.	150
<b>Fig. (54):</b> Bands generated by PCR using RAPD primer (P7) for the seven species of venerid clams (A) amplified bands and (B) bands types.	151
<b>Fig. (57):</b> Bands generated by PCR using RAPD primer (P12) for the seven species of venerid clams (A) amplified bands and (B) bands types.	154
<b>Fig. (58):</b> Bands generated by PCR using RAPD primer (P14) for the seven species of venerid clams (A) amplified bands and (B) bands types.	155
<b>Fig. (59):</b> Bands generated by PCR using RAPD primer (P17) for the seven species of venerid clams (A) amplified bands and (B) bands types.	156
<b>Fig. (60):</b> Bands generated by PCR using RAPD primer (P18) for the seven species of venerid clams (A) amplified bands and (B) bands types.	157
<b>Fig. (61):</b> Bands generated by PCR using RAPD primer (P19) for the seven species of venerid clams (A) amplified bands and (B) bands types.	158
<b>Fig. (62):</b> Bands generated by PCR using RAPD primer (P20) for the seven species of venerid clams (A) amplified bands and (B) bands types.	159
<b>Fig. (63):</b> Unweighted pair group method using arithmetic average (UPGMA) dendrogram based on genetic similarity index values calculated from data of 13 arbitrary primers for seven species of venerid clams collected from Ismailia.	163
<b>Fig. (64):</b> Bands generated by PCR using RAPD primers (P1 and P2) for <i>Tapes decussatus</i> collected from different stations (A) amplified bands and (B) bands types.	168
<b>Fig. (65):</b> Bands generated by PCR using RAPD primers (P3 and P5) for <i>Tapes decussatus</i> collected from different stations (A) amplified bands and (B) bands types.	169
<b>Fig. (66):</b> Bands generated by PCR using RAPD primers (P10 and P11) for <i>Tapes decussatus</i> collected from different stations (A) amplified bands and (B) bands types.	170

## LIST OF FIGURES /CONTINUE

	Page
<b>Fig. (67):</b> Bands generated by PCR using RAPD primers (P12 and P14) for <i>Tapes decussatus</i> collected from different stations (A) amplified bands and (B) bands types.	171
<b>Fig. (68):</b> Bands generated by PCR using RAPD primers (P17 and P18) for <i>Tapes decussatus</i> collected from different stations (A) amplified bands and (B) bands types.	172
<b>Fig. (69):</b> Unweighted pair group method using arithmetic average (UPGMA) dendrogram based on genetic similarity index values calculated from data of 10 arbitrary primers for <i>Tapes decussatus</i> individuals collected from different stations	175

## LIST OF PLATES

	Page
<b>Plate (1):</b> Exterior and interior views of common venerid ( <i>Tapes decussatus</i> ) shell showing the morphological characters of the family	55
<b>Plate (2):</b> Exterior and interior views of the venerid clams shells.	56
<b>Plate (3):</b> Exterior and interior views of the venerid clams shells.	57
<b>Plate (4):</b> Venerid clams shells show some abnormalities.	58
<b>Plate (5):</b> Sporocyst of <i>Cercaria lata</i> from <i>Tapes decussatus</i> (A) stained with carmine and (B) diagram.	121
<b>Plate (6):</b> Redia of <i>Cercaria lata</i> from <i>Tapes decussatus</i> (A) stained with carmine and (B) diagram.	122
<b>Plate (7):</b> <i>Cercaria lata</i> from <i>Tapes decussatus</i> (A) stained in carmine and (B) diagram.	123
<b>Plate (8):</b> Sporocyst of unknown Digenea from <i>Venerupis pullastra</i> (A) stained with carmine and (B) diagram.	124
<b>Plate (9):</b> Redia of unknown Digenea from <i>Venerupis pullastra</i> (A) stained with carmine and (B) diagram.	125
<b>Plate (10):</b> Cercaria of unknown Digenea from <i>Venerupis pullastra</i> (A) stained with carmine and (B) diagram.	126