

REFERENCES

- Abou- El-Sherbini, Kh. S. & Hamed, M.A. (2000):** Pollution assessment in the aquatic ecosystem along the western coast of the Suez Bay. *Egy. Aquat. Biol. Fish.*, 4 (4): 37 – 59.
- Adams, S.M. (2001):** Biomarker/bioindicator response profiles of organisms can help differentiate between sources of anthropogenic stressors in aquatic ecosystems. *Biomarkers*. 6 (1): 33– 44.
- Ai-hui; Zhao-xia, L.I. & Gong-neng, F. (2009):** Phylogenetic relationship of the genus *Meretrix* (Mollusca: Veneridae) based on mitochondrial COI. *Gene Sequences*, 30 (3): 233-239.
- Amler, M.R. (1999):** Synoptical classification of fossil and recent Bivalvia. *Geologica et Palacontologica*, 33: 237-248.
- Anderson, W. D.; Keith, W.J.; Mills, F. H.; Steinmeyer, M.E. & Bailey, J.L. (1978):** A Comprehensive survey of South Carolina's hard clam resources. SC Marine Resources Center Technical Rept. 32. South Carolina wildlife and Marine Resources Department, Charleston. 14 pp.
- Andre, C.; Lindegarth, M.; Jonsson, P.R. & Sundberg, P. (1999):** Species identification of bivalve larvae using random polymorphic DNA (RAPD). *J. Mar. Biol. Assoc. UK*, 79(3): 563-565.
- APHA (1992):** Standard Methods for the Examination of Water and Waste Water. Washington, 18th Ed. DC.
- Aranishi, F. (2006):** A novel mitochondrial intergenic spacer reflecting population structure of Pacific oyster. *J. Appl. Genet.*, 47: 119–123.
- Aranishi, F. & Okimoto, T. (2004):** Genetic relationship between cultured populations of Pacific oyster revealed by RAPD analysis. *J. Appl. Genet.*, 45 (4): 435-443.
- Aranishi, F. & Okimoto, T. (2005):** Sequence polymorphism in a novel noncoding region of Pacific oyster mitochondrial

- DNA. J. Appl. Genet., 46: 201–206.
- Aranishi, F. & Okimoto, T. (2006):** A simple and reliable method for DNA extraction from bivalve mantle. J. Appl. Genet., 47 (3): 251–254.
- Atienzar, A.F.; Conradi, M.; Evenden, A.J.; Jha, A.N. & Depledge, M.H. (1999):** Qualitative assessment of genotoxicity using random amplified polymorphic DNA: comparison of genomic template stability with key fitness parameters in *Daphnia magna* exposed to benzo [a]-pyrene. Environ. Toxicol. Chem., 18: 2275-2282.
- Atienzar, F.A.; Cordi, B.; Donkin, M.E.; Evenden, A.J.; Jha, A.N. & Depledge, M.H. (2000):** Comparison of ultraviolet- induced genotoxicity detected by random amplified polymorphic DNA with chlorophyll fluorescence and growth in a marine macroalgae, *Palmaria palmate*. Aquat. Toxicol. 50, 1-12.
- Atienzar, F.A.; Venier, P.; Jha, A.N. & Depledge, M.H. (2002):** Evaluation of random amplified polymorphic DNA (RAPD) assay for the detection of DNA damage and mutations, Mutat. Res. 521, 151–163.
- Atta, M.M. (1991):** Rocky shore fauna of Mediterranean coastline of Alexandria. Egypt. Ger. Soc. Zool. 5, 211-224.
- Awad, H.; Michel, P. & El-Shazly, A. (1983):** Actual situation of oil pollution in the ecosystem along the Egyptian Red Sea Coast. Bull. Inst. Oceanogr. & Fish., 9: 102-108.
- Bailey, N.T.J. (1981):** Statistical Methods in Biology. 2nd Ed. (Biological Science Texts): 216 pp.
- Baozhong, L.; Feng, Y.; Bo, D. & Jianhai, X. (2006):** The comparison between allozyme and RAPD markers for the population genetic structure analysis of scallop *Chlamys farreri*. Chinese J. Oceano. & Limnol. 24 (3): 295-299.
- Barash, A. & Danin, Z. (1973):** The Indo- Pacific species of mollusca in the Mediterranean and notes on a collection from the Suez Canal. Israel J. of Zool., 21: 301-374.

- Barber, B.J.; Ford, S.E. & Haskin, H.H. (1988):** Effect of the parasite MSX (*Haplosporidium nelsoni*) on oyster (*crassostrea virginica*) energy metabolism: I. Condition index and relative fecundity. J. Shellfish Res., 7: 25–31.
- Barman, H.K.; Barat, A.; Yadav, B.M.; Banerjee, S.; Mehar, P.K.; Reddy, P.V.G.K. & Jana, R.K. (2003) :** Genetic variation between four species of Indian major carps as revealed by random amplified polymorphic DNA assay. Aquaculture. 217, 115-123.
- Barron, M.G. (2003):** Bioaccumulation and bioconcentration in aquatic organisms. Handbook of ecotoxicology (2nd Eds), (Eds.) Hoffman, D.J.; Rattner, B.A., Burton, G.A. (Jr.) & Cairns, J. (Jr.) Lewis Publishers, Boca Ration: 877–888.
- Barsiene, J. (1994):** Chromosome set changes in molluscs from highly polluted habitats, in A.R. Beaumont (Ed.), Genetics and Evolution of Aquatic Organisms, Chapman & Hall, London, 434-447.
- Bartell, S.M.; Campbell, K.R.; Lovelock, C.M.; Nair, S.K. & Shaw, J.L. (2000):** Characterizing aquatic ecological risks from pesticides using diquat dibromide case study. Part III. Ecological process models. Environ. Toxicol. Chem. 19, 1441-1453.
- Baudrimont , M.; Andres, S.; Durrieu, G.& Boudou, A. (2003) :** The key role of metallothioneins in the bivalve, *Corbicula fluminea* during the depuration phase, after in situ exposure to Cd and Zn. Aquat, Toxicol., 63 : 89–102.
- Baudrimont, M.; Montaudouin, X. & Palvadeau, A. (2006):** Impact of digenean parasite infection on metallothionein synthesis by the cockle (*Cerastoderma edula*): a multivariate field monitoring. Mar. Pollut. Bull., 52: 494–502.
- Bavay, M. (1898):** Note sur les mollusques du Canal de Suez. Bull. Soc. Zool. France, 23: 161-165.
- Beasley, G. and Kneale, P. (2002):** Reviewing the impact of metals and PAH's on macro invertebrates in urban watercourses.

- Prog. Phys. Geogr. 26 (2), 236-270.
- Beaumont, A.R. & Pether, S.M.J. (1996):** Allozyme variation and gene flow between cockle *Cerastoderma edule* populations in southern United kingdom. Fish. Res., 28: 263-275.
- Belfiore, N.M. & Anderson, S.L. (2001):** Effects of contaminants on genetic patterns in aquatic organisms: a review, Mutat. Res. 289, 97-122.
- Ben Abdallah, L.G.; El Menif, N.T. & Maamouri, F. (2009):** The morphology and behaviour of *Cercaria lata* Lespés, 1857 (Digenea, Faustulidae) from Mediterranean clam *Tapes decussata* (L.). J. of Helminthology, 83: 69–76.
- Bentzen, P.; Taggart, C.T.; Ruzzante, D.E. & Cook, D. (1996):** Microsatellite polymorphism and the population structure of Atlantic cod (*Gadus morhua*) in the northwest Atlantic. Can. J. Fish. Aquat. Sci, 53: 2706-2721.
- Berthelin, C.; Kellner, K. & Mathieu, M. (2000):** Storage metabolism in the Pacific oyster (*Crassostrea gigas*) in relation to Summer mortalities and reproductive cycle (west coast of France). Comp. Biochem. Physiol, 125 (B): 359-369.
- Bickham, J.W., & Smolen, M.J. (1994):** Somatic and heritable effects of environmental genotoxins and the emergence of evolutionary toxicology. Environ. Health Perspect. 102, 25-28.
- Bielawski, J. P. & Pumo, D.E. (1997):** Randomly amplified polymorphic DNA (RAPD) analysis of Atlantic coast striped bass. Heredity, 78: 32-40.
- Bishop, J.A & Cook, L.M. (Eds.) (1981):** Genetic consequences of Man Made Change, Academic Press, London, 209 pp.
- Bitar, G. & Kouli- Bitar, S. (1999):** Inventaire des mollusques marins benthiques du Liban et remarques bigeographiques sur quelques especes nouvellement signalees. Mesogee, 56: 37-44.
- Black, M.C.; Ferrell, J.R.; Horning, R.C. & Martin, L.K. (1996):** DNA strand breakage in freshwater mussels *Anodonta*

- grandis* exposed to lead in the laboratory and field. Environ. Toxicol. Chem. 15 (5): 802-808.
- Blust, R.; Kockelbergh, E. & Baillieul, M. (1992):** Effect of salinity on the uptake of cadmium by the brine shrimp *Artemia franciscana*. Mar. Ecol. Prog. Ser. 84, 245-254.
- Bodoy, A.; Maitre- Allain, T. & Riva, A. (1981):** Croissance Comparee de la palourde europeenne *Ruditapes decussatus* et de la palourde japonaise *Ruditapes philippinarum* dans un ecosysteme artificiel mediterraneen. Vie Marine, 2: 39-51.
- Boehm, P.D. & Quinn, J.G. (1977):** The persistence of chronically accumulated hydrocarbons in the hard clam *Mercenaria mercenaria*. Mar. Biol., 44(3): 227-233.
- Bolognesi, C.; Rabboni, R. & Roggieri, P. (1996):** Genotoxicity biomarkers in *M. Galloprovincialis* as indicators of marine pollutants. Comp. Biochem. Physiol. 113(2), 319-323.
- Bourgoin, B.P. (1990):** *Mytilus edulis* shell as a bioindicators of lead pollution: consideration on bioavailability. Mar. Ecol. Prog. Ser., 61 (3): 253 – 262.
- Brezonik, P.L.; King, S.O. & Mach, C.E. (1991):** The influence of water chemistry on trace metal bioavailability and toxicity to aquatic organisms. In : Metal ecotoxicology; Concepts and applications, (Eds.) Newman, M.C. & McIntosh, A.W. Lewis Publishers, Chelsea : 1-4 pp.
- Bridges, J.W.; Garattini, S.; Agarwal, D.K.; Barltrop, D.; Breuton, M.J.; Chernoff, N.; Clark, J.; Dean, J.H.; Dixon, R.; Fliedner, T.; Saxen, L; Suskind, R.; Van Leeuwen, F.X.R.; Khera, K. & Tabakova, S. (1990):** Methods to detect toxic effects in specific mammalian organs and physiological systems. In : Short-term toxicity tests for non- genotoxic effects , (Eds.) Bourdeau, P., Somers, E., Richardson, G.M. & Hickman, J.R. John Wiley & Sons, New York: 11 pp.
- Brokordt, K.B.; Himmelman, J.H. & Guderley, H.E. (2000):** Effect of reproduction on escape responses and muscle metabolic capacities in the scallop *Chlamys islandica*

- Muller 1776. J. Exp. Mar. Biol. Ecol., 251: 205–225.
- Bryan, G.W. (1976):** Some aspects of heavy metal tolerance in aquatic organisms. In: Effects of pollutants on aquatic organisms, (Eds.) lockwood, A.P.M. Combridge University Press, Cambridge: 7-35.
- Cajaraville, M.P.; Marigomez, J.A.; Diez, G. & Angulo, E. (1992):** Comparative effects of the water accommodated fraction of three oils on mussels: 2. Quantitative alterations in the structure of the digestive tubules. Comp. Biochem. Physiol., 102: 113–123.
- Calabrese, A & Davis, H.C. (1966):** The pH tolerance of embryos and larvae of *Mercenaria mercenaria* and *Crassostrea virginica*. Biological Bulletin, 131 (3): 427-436.
- Campbell, P.G.C. & Tessier, A. (1996):** Ecotoxicology of metals in the aquatic environment: Geochemical aspects. In: Ecotoxicology: A hierarchical treatment (Eds.) Newman, M.C. & Jagoe, C.H. Lewis Publishers, Boca Raton: 21 pp.
- Canapa, A.; Marota, I.; Rollo, F. & Olmo, E. (1996):** Phylogenetic analysis of Veneridae (Bivalvia): Comparison of molecular and palaeontological data. J. Mol. Evol. 43: 517-522.
- Caquet, T. & Lagadic, L. (2000):** Consequences of individual- level alterations on population dynamics and community structure and function. In: Use of biomarkers for environmental quality assessment, (Eds.) Lagadic, L., Caquet, T., Amiard, J.-C. & Ramade, F.A.A. Balkema, Rotterdam: 291 pp.
- Carell, B.; Forberg, S.; Grundelius, E.; Henrikson, L.; Johnels, A.; Lindh, U.; Mutvel, H.; Olsson, M.; Savardstrom, K. & Westermarck, T. (1987):** Can mussel shells reveal environmental history ?. Ambio., 16 : 2-10.
- Carmichael, N.G. & Fowler, B.A. (1981):** Cadmium accumulation and toxicity in the kidney of the bay scallop, *Argopecten irradians*. Mar. Biol. 65: 35-43.
- Carmichael, N.G.; Squibb, K.S.; Engel, D.W. & Fowler, B.A. (1979):** Metals in the subcellular distribution of Cd, Mn

- and Zn by the clam, *Mercenaria mercenaria*. Comp. Biochem. Physiol., 65A: 203-206.
- Carriker, M.R.; Palmer, R.E.; Sick, L.V. & Jonhson, C.C. (1980):** Interaction of mineral elements in sea water and shell of the oyster (*Crassostrea virginica* "Gmelin") cultured in controlled and natural systems. J. Exp. Mar. Biol. Ecol., 46: 279 – 296.
- Chapman, D. (1992):** Water quality assessments. Chapman & Hall, London: 181-215.
- Chapman, P.M. (1997):** Is bioaccumulation useful for predicting impacts. Mar. Pollut. Bull, 34(5): 282 –283.
- Cheng, T. C. (1993):** Noninfectious diseases of marine molluscs. In: J. A. Couch and J.W. Fournier, Editors, Pathobiology of Marine and Estuarine Organisms, CRC Press, Boca Raton, pp: 289-318.
- Choi, K. S. & Park, K.I. (2001):** Histopathology and spatial distribution of the protozoan parasite, *Perkinsus* sp. found in the Manila clam *Ruditapes Philippinarum* distributed in Cheju, Korea. Korean J. Environ. Biol., 19: 79-86.
- Chow, S. & Ushiamo, H. (1995):** Global population structure of albacorc (*Thunnus alalunga*) inferred by RFLP analysis of the mitochondrial ATPase gene. Mar. Biol., 123: 39-45.
- Clark, A.D. & Lanigan, C.M.S (1993):** Prospects for estimating nucleotide divergence with RAPDs. Mol. Biol. Ecol., 10: 1096–1111.
- Clements, W.H. (2000):** Integrating effects of contaminants across levels of biological organization: An overview. J. Aquat, Eco. Stress. Recov. 7(2): 113-116.
- Clements, W.H.; Carlisle, D.M.; Lazochak, J.M. & Johnson, P.C. (2000):** Heavy metals structure benthic communities in Colorado mountain streams. Ecol. Appl. 10 (2): 626-638.
- Coen, L.D. (1995):** A review of the potential impacts of mechanical harvesting on subtidal and intertidal shellfish resources. SCDNR- MRRI, 46 pp.
- Connel, D.W.; LAM, P.; Richardson, B. & WU, R. (1999):**

- Introduction to ecotoxicology. Blackwell Sciences, Oxford: 121-134.
- Cormier, S.M. & Daniel, F.B. (1994):** Biomarkers: Taking the science forward. *Environ. Toxicol. Chem.*, 13(7): 1011-1012.
- Coughlan, B.M; Hartl, M.G.J.; O'Reilly, S.J.; Sheehan, D.; Morthersill, C; van Pelt, F.N.A.M.; O'Halloran, J. & O'Brien, N.M. (2002):** Detecting genotoxicity using the Comet assay following chronic exposure of Manila clam *Tapes semidecussatus* to polluted estuarine sediments. *Mar. Pollu. Bulle.* 44: 1359-1365.
- Cravo, A.; Bebianno, M.J. & Foster, P. (2004):** Partitioning of trace metals between soft tissues and shells of *Patella aspera*. *Env. Int.*, 30(1): 87- 98.
- Cravo, A.; Foster, P. & Bebianno, M.J. (2002):** Minor and trace elements in the shell of *Patella aspera* (Roding 1998). *Environ. Interna.* 28, 295-302.
- CSIR (1962):** The Wealth of India: Raw Materials Vol. VI: L-M. National Institute of Science Communication and Information Resources, CSIR, New Delhi, India.
- Dame, R. (1996):** Ecology of marine bivalves: an ecosystem approach. CRC Marine Science Series. Boca Raton, Florida. 254 pp.
- Dame; R.; Bushek, D. & Prins, T. (2001):** The role of suspension feeders as ecosystem transformers in shallow coastal environments. In K. Reise (Ed), *The Ecology of Sedimentary Coasts*. Springer. Verlag, Berlin, 11-37 pp.
- Dang, C.; de montaudouin, X.; Bald, J.; Jude, F.; Raymond, N.; Lanceleur, L.; Paul- Pont, I. & Caill – Milly; N. (2009):** Testing the enemy release hypothesis: trematode parasites in the non-indigenous Manila clam *Ruditapes Philippinarum*. *Hydrobiologia*, 630: 139–148.
- De Salle, R. & Birstein, V.J. (1996):** PCR identification of black caviar. *Nature*, 381: 197–198.
- De Wolf, H.; Blust, R. & Backelijau, T. (2004):** The population

- genetic structure of *Littorina littorea* (Mollusca: Gastropoda) along a pollution gradient in the Scheldt estuary (The Netherlands) using RAPD analysis. *Science of the total Environ.* 325 (1-3): 59–69.
- De Wolf, H.; Ulomi, S.A.; Backeljau, T.; Pratap, H.B. & Blust, R. (2001):** Heavy metal levels in the sediments of four Dartes Salaam mangroves. Accumulation and effect on the morphology of the periwinkle, *Littoraria scabra* (Mollusca: Gastropoda). *Environ. Int.*, 26 (4): 243-249.
- Depledge, M.H. (1994):** Genotypic toxicity: implications for individuals and populations. *Environ. Health Perspect.* 102 (12), 101-104.
- Depledge, M.H. (1998):** The ecotoxicological significance of genotoxicity in marine invertebrates. *Mutat. Res.* 399, 109–122.
- Depledge, M.H. & Fossi, M.C. (1994):** The role of biomarkers in environmental assessment (2) Invertabrates. *Ecotoxicology*, 3: 161–172.
- Depledge, M.H.; Aagaard, A. & Gyorkos, P. (1995):** Assessment of trace metal toxicity using molecular, physiological and behavioural biomarkers. *Mar. Pollut. Bull.*, 31: 19-27.
- Devaux, A.; Flammarion, P.; Bernardon, V.; Garric, J. & Monod, G. (1998):** Monitoring of the chemical pollution of the River Rhone through measurement of DNA damage and cytochrome P4501A induction in chub (*Leuciscus cephalus*). *Mar. Environ. Res.*, 46: 257–262.
- Downs, C.A; Dillon, R.T. (Jr.); Fauth, J.E. & Woodley, C.M. (2001):** A molecular biomarker system for assessing the health of gastropods (*Llyanassa obsoleta*) exposed to natural and anthropogenic stressors. *J. Exp. Mar. Biol. Ecol.*, 259: 189–241.
- Doyotte, A.; Cossu, C.; Jacquin, M.-C.; Babut, M. & Vasseur, P. (1997):** Antioxidant enzymes, glutathione and lipid peroxidation as relevant biomarkers of experimental or field exposure in the gills and the digestive gland of the

- freshwater bivalve, *Unio tumidus*. Aquat. Toxicol. 39: 93-110.
- Dybern, B. (1983):** Field sampling and preparation of subsamples of aquatic organisms for analysis of metals and organochlorines, FAO, Fisher Tech. Paper, 212: 1-13.
- Elder, J.F. & Collins, J.J. (1991):** Freshwater mollusks as indicators of bioavailability and toxicity of metals in surface- water systems. Rev. Environ. Contam. Toxicol. 122: 37–79.
- El-Gamal, M.M. (1988):** Studies on marine edible molluscs of Egypt. M. Sc. Thesis, Fac. Sci., Tanta Univ., Egypt.
- El-Gamal, M.M. & Sharshar, Kh. M. (2004):** Heavy metals partitioning between soft tissues and shells of some freshwater and marine mollusks as accumulation indicators for their monitoring in the environment. Egy. Ger. Soc. Zool., 45 (D): 141-161.
- Ellis, M.S.; Barber, R.D.; Hillman, R.E. & Powell, E.N. (1998):** Histopathology analysis. NOAA Tech. Memo. NOS ORCA, 130: 198–215.
- El-Nady, F.E. (1996):** Heavy metals exchange among the aquatic environment of the Mediterranean coast of Egypt. Indian J. Mar. Sci., 25(3): 225-233.
- Elo, K.; Ivanoff, S.; Vuorinen, J.A. & Piironen, J. (1997):** Inheritance of RAPD markers and detection of interspecific hybridization with brown trout and Atlantic salmon. Aquaculture, 152: 55-65.
- El-Samra, M.I.; Abdallah, A.M. & Hamed, M.A. (1995):** Seasonal variation of some metal in the aquatic system of the Suez Bay. Environ. Sci., 9: 81-90.
- El-Shenawy, N. S. (2004):** Heavy metal and microbial depuration of the clam *Ruditapes decussatus* and its effect on bivalve behavior and physiology. Environ Toxicol 19: 143-153.
- El-Shenawy, N.S.; Moawad, T.I.S; Mohallal, M.E.; Abdel-Nabi, I.M. & Taha, I.A. (2009):** Histopathologic biomarker response of clam, *Ruditapes decussates*, to Organophosphorous pesticides reldan and roundup: A

- laboratory study. J. Ocean Science, 44: 27–34.
- El-Tokhi, M.M. & Mostafa, Y.M. (2001):** Heavy metals and petroleum hydrocarbons contamination of bottom sediments of El- Sukhna area, Gulf of Suez, Egypt. Petro. Scie. Tech., 19: 481-494.
- Emara, A.M. & Belal, A.A. (2004):** Marine fouling in Suez Canal, Egypt. Egypt. J. Aqua. Res., 30(A): 189-206.
- Emara, A.M. & Said, T.O. (2003):** A marine bivalve as biomonitor for oil pollution along the coastal area of the Suez Gulf, Egypt. Egypt. J. Aquat. Biol. & Fish., 7(2): 119-133.
- Encomio, V. & Chu, F.L. (2000):** The effect of PCBs on glycogen reserves in the eastern oyster *Crassostrea virginica*. Mar. Environ. Res, 50: 45-49.
- Engel, D. W. & Fowler, B.A. (1979):** Factors influencing cadmium accumulation and its toxicity to marine organisms. Environ. Health Perspect. 28: 81-88.
- Epifanio, J.M.; Koppelman, J.B.; Nedbal, M.A. & Philipp, D.P. (1996):** Geographic variation of paddlefish allozymes and mitochondrial DNA. Trans. Am. Fish. Soc., 125: 546-561.
- Eversole, A.G. (1987):** Species profiles: life histories and environmental requirements of coastal fishes and invertebrates (South Atlantic) hard clam. U.S. Fish Wild. Serv. Biol. Rep. 82 (11.75). U.S. Army Corps of Engineers, TR El.82.4. 33 pp.
- Farag, E.A.; Dekinesh, S.I. & El- Odessy, H. (2000):** Trace elements accumulation in soft parts of six edible bivalve molluscs inhabiting Alexandria coast, Egypt. Egy. Ger. Soc. Zool., 31 (D): 27-36.
- Farag, E.A.; El-Sherief, S.S.; Mahmoud, A.D. & Abd-Ellah, S.M. (1999):** A study of rocky shore molluscs from Alexandria, Egypt. J. Egypt. Ger. Soc. Zool., 30 (D): 1-12.
- Fernandez, A.; Garcia, T.; Asensio, L.; Rodriguez, M.A.; Gonzalez, I. & Cespedes, A. (2000):** Identification of the clam species *Ruditapes decussates* (grooved carpet shell), *Venerupis pullastra* (pullet carpet shell), and *Ruditapes*

- philippinarum* (Japanese carpet shell) by PCR- RFLP. J. Agric. Food Chem., 48: 3336-3341.
- Ferrel, R.E.; Carville, T.E. & Martinez, J.D. (1973):** Trace metals in oyster's shells. Environ. Lett., 4: 311-316.
- Fischer-Piette, E. (1976):** Les Veneridae indeterminées de la collection de Calcutta (J). Rec. Zool. Surv. India, 70: 235-257.
- Fishelson, L. (2000):** Marine animal assemblages along the littoral of the Israeli Mediterranean seashore: The Red-Mediterranean Seas communities of species. Italian. J. of Zool., 67 (3): 393-415.
- Fishelson, L.; Bresler, V.; Manelis, R.; Zuk-Rimon, Z.; Dotan, A.; Hornung, H. et al., (1999):** Toxicological aspects associated with the ecology of *Donax trunculus* (Bivalvia, Mollusca) in a polluted environment. Sci. Total Environ. 226: 121-131.
- Florence, T.M.; Morrison, G.M. & Stauber, J.L. (1992):** Determination of trace element speciation and the role of speciation in aquatic toxicology. Sci. Total Environ. 125: 1-13.
- Forbes, V.E. & Forbes, T.L. (1994):** Ecotoxicology in theory and practice. Chapman & Hall, London: 69-71.
- Ford, S.E. (1985):** Chronic infections of *Haplosporidium nelsoni* (MSX) in the oyster *Crassostrea virginica*. J. Inverteb. Pathol. 45: 94-107.
- Ford, S.E. & Figueras, A.J. (1988):** Effects of sublethal infection by the parasite *Haplosporidium nelsoni* (MSX) on gametogenesis, spawning, and sex ratios of oysters in Delaware Bay, USA. Dis. Aquat. Org., 4: 121-133.
- Formi, A. (1994):** Comparison of chromosome aberrations and micronuclei in testing genotoxicity in humans. Toxicol. Lett. 72, 185-190.
- Forstner, V. & Prosi, F. (1979):** Heavy metal pollution in freshwater systems. In: Biological aspects of freshwater pollution. (Ed.) Ravera, O. Pergamon Press, Oxford: 129-

162.

- Fritsch, P. & Rieseberg, L.H. (1996):** The use of random amplified polymorphic DNA (RAPD) in conservation genetics, in: T.B. Smither, R.K. Wayne (Eds.), *Molecular Genetic Approaches in Conservation*. Oxford University Press, 55-73.
- Futoshi, A. & Takane, O. (2004):** Genetic relationship between cultured populations of Pacific oyster revealed by RAPD analysis. *J. Appl. Genet.*, 54(4): 435-445.
- Gainey, L.F. & Shumway, S.E. (1988):** A compendium of the responses of bivalve molluscs to toxic dinoflagellates. *J. Shellfish Research*, 7: 6223-6228.
- Gallardo, M. H. & Carrasco, J.I. (1996):** Genetic cohesiveness among populations of *Concholepas concholepas* (Gastropoda, Muricidae) in Southern Chile. *J. Exp. Mar. Biol. Ecol.*, 197: 237-249.
- Garcia de leon, F.J.; Chikhi, L. & Bonhomme, F. (1997):** Microsatellite polymorphism and population subdivision in natural populations of European sea bass *Dicentrarchus labrax* (Linnaeus, 1758). *Mol. Ecol.*, 6: 51-62.
- Gardner, G.R.; Pruell, R.J. & Malcolm, A.R. (1992):** Chemical induction of tumors in oysters by a mixture of aromatic and chlorinated hydrocarbons, amines and metals. *Mar. Environ. Res.* 34, 59-63.
- Ghobashy, A.A.; Mohamed, S.Z.; Gabr, H.R. & Brand, A.R. (1992):** Community structure and seasonal variation of Mollusca at Lake Timsah (Suez Canal). *J. Egypt. Ger. Soc. Zool.*, 7(B): 145-160.
- Giusti, L.; Williamson, A.C. & Mistry, A. (1999):** Biologically available trace metals in *Mytilus edulis* from the coast of northeast England. *Environ. Int.*, 25(8): 968-981.
- Gofas, S.; Le Renard, J. & Bouchet, P. (2001):** Mollusca in: Costello, M.J. *et al.* (Ed.) (2001). *European register of marine species: a check-list of the marine species in Europe and a bibliography of guides to their identification*.

- Collection Patrimoines Naturels, 50: 180-213 pp.
- Hadrys, H.; Balick, M. & Schierwater, B. (1992):** Applications of random amplified polymorphic DNA (RAPD) in molecular ecology. *Mol. Ecol.*, 1: 55-63.
- Hall, K.A.; Cribb, T.H. & Barker, S.C. (1999):** V4 region of small subunit rDNA indicates polyphyly of the Fellodistomidae (Digenea) which is supported by morphology and life-cycle data. *System. Parasito.* 43: 81-92.
- Hallegraeff, G.M. (1993):** A review of harmful algal blooms and their apparent global increase. *Phycologia*, 32: 79-99.
- Hamed, M.A. (1996):** Determination of some micro – elements in aquatic ecosystem and their relation to the efficiency of aquatic life. Ph. D. Thesis, Faculty of Science, Mansoura University. p 134.
- Hamed, M.A. (2005):** Assessment of heavy metals among suspended particulates and dissolved phases in Suez Canal water. *Chemis. & Ecol.*, 21 (5): 387–397.
- Hamed, M.A. & Emara, A.M. (2006):** Marine molluscs as biomonitors for heavy metal level in the Gulf of Suez, Red Sea. *Mar. Syst.*, 60: 220-234.
- Hamed, M.A.; Said, T.O. & Nasser, M.Z. (2003):** Effect of heavy metals and aromatic hydrocarbons on phytoplankton biomass as represented by chlorophyll – a in the northern part of the Gulf of Suez. *Egy. Acad. Soc. Environ. Dev.* 4(1): 67-85.
- Hamza, D.S. (2003):** Surveillance and biological studies on limpets inhabiting Egyptian water. M.Sc. Thesis, Faculty of science, Zagazig University (Benha).
- Hanafy, M.H.; Gab-Alla, A.A. & Hassanine, R.M.E. (1997):** Larval trematode (Digenea: Lepocreadiidae) infection in the gonads of the commercial bivalve *Venerupis decussata* from lake Timsah, Suez Canal. *J. Egypt. Ger. Soc. Zool.*, 24: 167–181.
- Handy, R.D.; Galloway, T.S. & Depledge, M.H (2003):** A proposal for the use of biomarkers for the assessment of chronic

- pollution and in regulatory toxicology. *Ecotoxicology*, 12: 231–343.
- Harding, G.C; Kenchington, E.L.; Bird, C.J.; Pezzach, D.S. & Landry, D.C. (1997):** Genetic relationships among subpopulations of the American lobster (*Homarus americanus*) as revealed by random amplified polymorphic DNA. *Can. J. Fish. Aquat. Sci.*, 54: 1762-1771.
- Hare, L. (1992):** Aquatic insects and trace metals: bioavailability, bioaccumulation, and toxicity. *Crit. Rev. Toxicol.* 22: 327-369.
- Harper, E.; Taylor, J. & Carme, J. (eds.) (2000):** The evolutionary biology of the Bivalvia. Geological Society, London, Special Publication. 177 p.
- Haskin, H.H. & Andrews, J.D. (1988):** Uncertainties and speculations about the life cycle of the eastern oyster pathogen *Haplosporidium nelsoni* (MSX). In: W.S.Fisher Editor Disease Processes in Marine Bivalve Molluscs, American Fisheries Society, Bethesda, MD, 5-22 pp.
- Hassanine, R. (2006):** The life cycle of *Diploproctodaeum arothroni* Bray and Nahhas, 1998 (Digenea: Lepocreadiidae), with a comment on the parasitic castration of its molluscan intermediate host. *J. Natural History*, 40: 19 – 20.
- Heasman, M.P.; O’connor, W.A. & Frazer, A.W.J. (1996):** Digenean (Bucephalidae) infections in commercial scallops, *Pecten fumatus* Reeve, and doughboy scallops *Chlamys (Mimachlamys) asperrima* (Lamarck), in Jervis Bay, New South Wales. *J. Fish Dis.*, 19: 333-339.
- Heipel, D.A; Bishop, J. D.D.; Brand, A.R. & Thorpe, J.P. (1998):** Population genetic differentiation of the great scallop *Pecten maximus* in western Britain investigated by randomly amplified polymorphic DNA. *Mar. Ecol. Prog. Ser.* 162: 163-171.
- Heist, E.J. & Gold, J.R. (1999):** Microsatellite DNA variation in sandbar sharks (*Carcharhinus plumbeus*) from the Gulf of

Mexico and mid- atlantic Bight. *Copeia* 1999: 182-186.

- Herricks, E.E.; Schaeffer, D.J. & Perry, J.A. (1989):** Biomonitoring: Closing the loop in the environmental sciences. In: *Ecotoxicology problems and approaches* (Eds.) Levine, S.A.; Harwell, M.A.; Kelly, J.R. & Kimball, K.D. Springer – Verlag, New York : 351-352.
- Hinton, D.E. (1992):** Histopathologic biomarkers. In: *Biomarkers-biochemical, physiological and histological markers of anthropogenic stress* (Hugget F.J.; kimerle, R.A.; Mehrle, P.M.Jr; Bergman, H.L. eds). SETAC Special Publication Series. Boca Raton, FL: Lewis Publishers, 155- 209.
- Hirschfeld, B.M.; Dhar, A.K.; Rask, K. & Alcivar-Warren, A. (1999):** Genetic diversity in the eastern oyster (*Crassostrea virginica*) from Massachusetts using the RAPD technique. *J. Shell fish Res.*, 18: 121-125.
- Huang, B.X.; Peakall, R. & Hanna, P.J. (2000):** Analysis of genetic structure of blacklip abalone (*Haliotis rubra*) populations using RAPD, minisatellite and microsatellite markers. *Mar. Biol.*, 136: 207–216.
- Hyne, R.V. & Maher, W.A. (2003):** Invertebrate biomarkers: links to toxicosis that predict population decline. *Ecotox, Environ. Saf.* 54(3) : 366-374.
- Hyotylainen, T.; Karles, A. & Oikari, A. (2002):** Assessment of bioavailability and effects of chemicals due to remediation actions with caging mussels (*Anodonta anatina*) at a creosote- contaminated lake sediment site. *Water Res.*, 36: 4497-4504.
- Ibrahim, A.M.; Aly, R.H.; Kenchington, E. & Ali, T.G. (2008):** Genetic polymorphism among five populations of *Pinctada radiata* from the Mediterranean coast in Egypt indicated by RAPD-PCR technique. *Egypt J. Zool.*, 50: 467-477.
- Ireland, M.P. & Wootton, R.J. (1977):** Distribution of lead, zinc, copper and manganese in marine gastropods, *Thais lapillus* and *littorina littorea*, around the coast of Wales. *Environ. Pollut.* 12: 27-41.
-

- Jamil, K. (2002):** Bioindicators and biomarkers of environmental pollution and risk assessment. Science Publishers, Enfield: 4, 15, 56.
- Kappner, I. & Bieler (2006):** Phylogeny of venus clams (Bivalvia: Venerinae) as inferred from nuclear and mitochondrial gene sequences. *Mol. Phylo. & Evol.*, 40(2): 317-331.
- Kargin, F. & Cogun, H.Y. (1999):** Metal interactions during accumulation and elimination of zinc and cadmium in tissues of the freshwater fish *Tilapia nilotica*. *Bull. Environ. Contam. Toxicol.* 63: 511-519.
- Keen, A.M. (1969):** Superfamily Veneracea, pp: 670-690. In: Leslie Reginald Cox et al., part N (Bivalvia), *Mollusca* 6, vols. 1 and 2xxxvii+952 pp. Part of Raymond C. Moore, ed., *Treatise on Invertebrate Paleontology*.
- Khamdan, S.A.A. (1998):** Occurrence of *Bucephalus* sp. trematode in the gonad of pear oyster, *Pinctada radiata*. *Environ. Interpret.* 24: 117 – 120.
- Klerks, P.L. & Weis, J.S. (1987):** Genetic adaptation to heavy metals in aquatic organisms: a review. *Environ. Pollut.* 45: 173-205.
- Klinbunga, S.; Ampayup, P.; Tassanakajon, A.; Jarayabhand, P. & Yoosukh, W. (2000):** Developemnt of species – specific markers of the tropical oyster (*Crassostrea belcheri*) in Thailand. *Mar Biotech.* 2: 476- 484.
- Klinbunga, S.; Ampayup P.; Tassanakajon, A.; Jarayabhand, P. & Yoosukh, W. (2001):** Genetic diversity and molecular markers of cupped oysters (genera *Crassostrea*, *Caccostrea* and *Striostrea*) in Thailand revealed by randomly amplified polymorphic DNA analysis. *Mar Biotech.* 3: 133–144.
- Koh, H.L.; Hallam, T.G. & Lee, H.L. (1997):** Combined effects of environmental and chemical stressors on a model *Daphnia* population. *Ecol. Model.* 103: 19-32.
- Kristensen, P. & Tyle, H. (1991):** The assessment of bioaccumulation. In : *Bioaccumulation in aquatic ecosystems*, (Eds.) Nagel, R. & Loskill, R. VCH, Weinheim: 189 pp.

- Kurelec, B. (1993):** The genotoxic disease syndrome. *Mar. Environ. Res.* 35: 341–348.
- Kurelec, B.; Garg, A.; Krca, S.; Chacko, M. & Gupta, R. (1989):** Natural environment surpasses polluted environment in inducing DNA damage in fish. *Carcinogenesis*. 10: 1337–1339.
- Lagadic, L.; Amiard J. C. & Caquet, T. (2000):** Biomarkers and evaluation of the ecotoxicological impact of pollutants. In: Use of biomarkers for environmental quality assessment, (Eds.) Lagadic, L., Caquet, T., Amiard, J.-C & Ramade, F. A.A. Balkema, Rotterdam: 303–311.
- Landrum, P.; Harkey, G.A. & Kukkonen, J. (1996):** Evaluation of organic contaminant exposure in aquatic organisms: The significance of bioconcentration and bioaccumulation. In: *Ecotoxicology: A hierarchical treatment*, (Eds.) Newman, M.C. & Jagoe, C.H. Lewis Publishers, Boca Raton: 85–132.
- Langston, W.J.; Bebianno, M.J. & Burt, G.R. (1998):** Metal handling in mollusks. In: Langston W., Bebianno, M.J. editors. *Metal metabolism in aquatic environments*. London: Chapman & Hall: 219–283.
- Lauckner, G. (1983):** Diseases of Mollusca: Bivalvia. In kinne, O. (Ed.), *Diseases of Marine Animals, Introduction, Bivalvia to Scaphopoda*, vol. 2. Biologische Anstalt Helgoland, Hamburg, Germany, 632–762 pp.
- Lee, M.K.; Cho, B.Y.; Lee, S.J.; Kang, J.Y.; Jeong, H.D.; Huh, S.H. & Huh, M. D. (2001):** Histopathological lesions of Manila clam, *Tapes philippinarum*, from Hadong and Namhae coastal areas of Korea. *Aquaculture*, 201: 199–209.
- Lespés, C. (1857):** Observations sur quelques cercaires parasites de Mollusques marins. *Annales des Sciences Naturelles Zoologie* (Ser 4), 7: 113–117.
- Li, G. & Hedgecock, D. (1996):** Mitochondrial DNA variation within and among larval cohorts of Pacific oyster,

- Crassostrea gigas*, detected by PCR- SSCP analysis. J. Shellfish. Res., 15: 511-512.
- Lin, M.C. & Liao, C.M. (1999):** ^{65}Zn (II) accumulation in the soft tissue and shell of abalone *Haliotis diversicolor supertexta* via the alga *Gracilaria tenuistipitata* var. *liui* and the ambient water. Aquaculture, 178: 89–101.
- Lindner, G. (1987):** Interessante Schneckenfunde and der sudturkischen Mittelmeerküste (Reisemitbringsel der familie Schmidt, Feldkirchen). Club Conchylia Informationen, 19: 32-43.
- Lingard, S.M.; Evans, R.D. & Bourgoin, B.P. (1992):** Methods for the estimation of organic-bound and crystal – bound metal concentrations in bivalves shells. Bull. Environ. Contam. Toxicol. 48: 179–184.
- Lison, D. (1999):** Importance of biotransformation pathways for interpreting biological monitoring of exposure. Toxicol. Lett. 108: 91-97.
- Livingstone, D.R. (1993):** Biotechnology and pollution monitoring: Use of molecular biomarkers in the aquatic environment. J. Chem. Tech. Biotechnol., 57: 195- 211.
- Liyan, Z.; Ying, H. & Guangxing, L. (2005):** Using DNA damage to monitor water environment. Chinese J. Oceanol. Limnol. 23 (3), 340-348.
- Lowe, D.M.; Moore, M.N. & Clarke, K.R. (1981):** Effects of oil in the digestive cells in mussels: quantitative alterations in cellular and lysosomal structure. Aquat. Toxicol. 1: 213–226.
- Lumb, S.M.; Bray, R.A. & Rollinson, D. (1993):** Partial small subunit (18S) rRNA gene sequences from fish parasites of the families Lepocreadiidae and Fellodistomidae (Digenea) and their use in phylogenetic analysis. System. Parasito. 26: 141– 149.
- Lynch, M. (1990):** The similarity index and DNA fingerprinting. Mol. Biol. Evol., 7: 478-484.
- Lynch, M. & Milligan, B.G. (1994):** Analysis of population genetic

- structure within RAPD markers. *Molec. Ecol.*, 3: 91- 99.
- Ma, X. L.; Cowles, D.L. & Carter, R.L. (2000):** Effect of pollution on genetic diversity in the bay mussel *Mytilus galloprovincialis* and the acorn barnacle *Balanus glandula*. *Mar. Environ. Res.* 50, 559- 563.
- Mackenzie, C.L.; Morrison, A.; Taylor, D.L.; Burrell, V.G.; Jr; Arnold, W.S. & Wakida Kusonoki, A.T. (2002) :** Quahogs in Eastern North America : Part I, Biology, Ecology and Historical Uses. *Marine Fisheries Review*, 64(2): 55 pp.
- Mance, G. (1987):** Pollution threat of heavy metals in aquatic environments. Elsevier Applied Science, London: 287–298.
- Marigomez, I.; Soto, M.; Cajaraville, M.P.; Angulo, E. & Giamberini, L. (2002):** Cellular and subcellular distribution of metals in molluscs. *Microsc. Res. Tech.*, 56(5): 358–392.
- Marin, S.A.; Haye, P.A.; Marchant, S. & Winkler, F.M. (2007):** Molecular markers used to analyze species-specific status in abalones with ambiguous morphology. *J. Shellfish Res.* 26(3): 833-837.
- Martin, L.K. & Black, M.C. (1998):** Biomarker assessment of the effects of coal strip-mines contamination on channel catfish. *Ecotox. Environ. Saf.* 41 (3): 307-320.
- Martinez, G.; Brokordt, K.; Aguilera, C.; Soto, V. & Guderley, H. (2000):** Effect of diet and temperature upon muscle metabolic capacities and biochemical composition of gonad and muscle in *Argopecten purpuratus* Lamark 1819. *J. Exp. Mar. Biol. Ecol.*, 247: 29-49.
- McCarthy, J.F. & Shugart, L.R. (1990):** Biological markers of environmental contamination. In: Biomarkers of environmental contamination (Eds.), McCarthy, J.F. & Shugart, L.R. Lewis Publishers, Boca Raton: 3-16.
- McCarty, L.S.; Power, M. & Munkittrick, K.R. (2002):** Bioindicators versus biomarkers in ecological risk assessment. *Hum. Ecol. Risk Assess.* 8(1): 159-164.
- Mienis, H.K. (1999):** On the presence of the Indo- Pacific bivalve

- Gafrarium pectinatum* (Linnaeus, 1758) along the Mediterranean coast of Israel. Club Conchyliia Informationen. 31: 37- 44.
- Mikkel, P.; Bieler, R.; Kappner, I. & Rawlings, T. (2006):** Phylogeny of Veneroidea (Mollusca: Bivalvia) based on morphology and molecules. Zoological J. Linnean Society, 148: 439-521.
- Mithelmore, C.L.; Birmelin, C.; Livingstone, D.R. & Chipman, J.K. (1998):** Detection of DNA strand breaks in Isolated Mussel (*Mytilus edulus* L) digestive gland cells using the "Comet assay". Ecotox. & Enviro. Safety. 41 (1), 51-58.
- Miura, O.; Kuris, A.M.; Torchin, M.E.; Hechinger, R.F.; Dunham, E.G. & Chiba, S. (2005):** Molecular- genetic analyses reveal cryptic species of trematodes in the intertidal gastropod *Batillaria cumingi* (Crosse). Int. J. Parasit. 35: 793-801.
- Moazzo, P.G. (1939):** Mollusques testaces marins du Canal de Suez. Memoires de L'Institut d'Egypt, 38: 1-283.
- Mohamed, S.Z.; Gabr, H.R.; Ghobashy, A.F.A & Brand, A.R. (1992):** Species composition and distribution of benthic mollusks in Lake Timsah(Suez Canal). J. Egypt. Ger. Soc. Zool. 7: 161-174.
- Mona, M.M. (2009):** The interactions between bathogenic bacteria and biological activity of marine edible bivalvian mollusca. M.Sc. Thesis, Faculty of Science, Tanta University, Egypt.
- Moore, M.N.; Lowe, D.M.; Livingstone, D.R. & Dixon, D.R. (1986):** Molecular and cellular indices of pollutant effects and their use in environmental impact assessment. Wat. Sci. Tech., 18: 223-232.
- Moraga, D.; Mdelgi- Lasram, E.; Romdhane, M.S.; El Abed, A. ; Boutet , I. ; Tanguy, A. & Auffret, M. (2002):** Genetic responses to metal contamination in two clams: *Ruditapes decussatus* and *Ruditapes philippinarum*. Mar. Environ. Res., 521-525.
- Morton, B. (1996):** The evolutionary history of the bivalvian.

- Oxford University Press, Oxford. 337-356 pp.
- Moursy, A. (1978):** Afield study of oil pollution of the Egyptian Coast of the Red Sea. National Research Center, Cairo, Egypt.
- MSTAT-C V. 2.10 (1988):** A microcomputer program for the Design, Management and Analysis of Agronomic. Res. Exp. Michigan State Univ. U.S.A.
- Mulcahy, D.L.; Cresti, M.; Linskens, H.F.; Intrieri, C.; Silverstoni, O.; Vignani, R. & Pancaldi, M. (1995):** DNA fingerprinting of Italian grape varieties : a test of reliability in RAPDs. *Advanced Horticultural Science*. 9: 185-187.
- Nacci, D. & Jackim, E. (1989):** Using the DNA alkaline unwinding assay to detect DNA damage in laboratory and environmentally exposed cells and tissues. *Mar. Environ. Res.* 28, 333-337.
- Nell, J.A. & Peterson, K.J. (1997):** Salinity studies on the clams *Katelysia rhytiphora* (Lamy) and *Tapes dorsatus* (Lamarck). *Aqua. Res.*, 28 (2): 115– 122.
- Newell, R.I.E. (2004):** Ecosystem influences of natural and cultivated populations of suspension- feeding bivalve mollusks: A Review. *J. Shellfish Res.*, 23: 51-61.
- Newman, M.C. (1995):** Quantitative methods in aquatic excotoxicology. Lewis Publishers, Boca Raton: 60–63.
- Ngo, T.T.T. & Choi, K.S (2004):** Seasonal changes of *Perkinsus* and *Cercaria* infections in the Manila clam *Ruditapes philippinarum* from Jeju, Korea. *Aquaculture*, 239: 57–68.
- Ngo, T.T.T.; Berthe, F.C.J. & Choi, K.S. (2003):** Prevalence and infection intensity of the ovarian parasite *Marteilioides chungmuensis* during an annual reproductive cycle of the oyster, *Crassostrea gigas*. *Dis. Aquat.Org.*, 56: 259-267.
- Nicholson, S. (2001):** Ecocytological and toxicological responses to copper in *Perna viridis* (L.) (Bivalvia: Mytilidae) haemocyte lysosomal membranes. *Chemosphere*. 45, 399-407.
- O'Reilly, P. & Wright, J.M. (1995):** The evolving technology of

- DNA fingerprinting and its application to fisheries and aquaculture. J. Fish. Biol., 47: 29-55.
- Officer, C.B.; Smayda, T.J. & Mann R. (1982):** Benthic filter feeding: a natural entrophication control. Mar. Ecol. Progr. Series. 8: 203–210.
- Palombi, A. (1933a):** *Bacciger bacciger* (Rud.) Nicoll, 1914, forma adulta di *Cercaria pectinata* Huet, 1891. Bolletino Dello Societa Del Naturalisti di Napoli, 44: 217-219.
- Palombi, A. (1933b):** *Cercaria pectinata* Huet *Bacciger bacciger* (Rud.). Rapporti geneticie biologia. Bolletino Dello Zoologia Napoli, 4: 1-11.
- Palombi, A. (1934):** *Bacciger bacciger* (Rud.) e trematode digenetico: Fam Steringophorida Odhner Anatomia, sistematica e biologia. Pubblicazione della Stazione Zoologica di Napoli, 13: 438–478.
- Pandey, S.; Parvez, S.; Sayeed, I.; Haque, R.; Bin-Hafeez, B. & Raisuddin, S. (2003):** Biomarkers of oxidative stress: a comparative study of river Yamuna fish *Wallago attu* (BL. & Schn.). Sci. Total Environ. 309: 105-115.
- Paraense, W.L.; Pointier, J.P.; Dealy, B.; Pernot, A.F.; Incani, R.N.; Balzan, C. & Chrosciechowski, P. (1992):** *Piomphalaria pronsa* (Gastropoda: Planorbidae): A morphological and biochemical study. Mem. Inst. Oswaldo. Cruz., 87: 171-179.
- Park, K.I. & Choi, K.S. (2001):** Spatial distribution of protozoan parasite *Perkinsus* sp. found in Manila clams, *Ruditapes philippinarum*, in Korea. Aquaculture, 203: 9-22.
- Park, K.; Park, J.K.; Lee, J. & Choi, K.S. (2005):** Use of molecular markers for species identification of Korean *Perkinsus* sp. Isolated from Manila clams *Ruditapes philippinarum*. Dis. Aquat. Org. 66: 255-263.
- Patwary, M.U.; Kenchington, E.L.; Bird, C.J. & Zouros, E. (1994):** The use of random amplified polymorphic DNA markers in genetic studies of the sea scallop, *Placopecten magellanicus* (Gmelin, 1971). J. Shell Fish Res., 13: 547–

553.

- Peijnenburg, W.J.G.M. & Jager, T. (2003):** Monitoring approaches to assess bioaccessibility and bioavailability of metals: Matrix issues. *Ecotox. Environ. Saf.* 56(1): 63-77.
- Pérez, E.C. (2004):** Induced spawning behavior and larval development of the hard clam *Mercenaria mercenaria* (Linnaeus, 1758) in Puerto Rico PhD. Thesis, University of Puerto Rico, Mayaguez Campus.
- Perkins, F.O. (1993):** Infectious diseases of molluscs. In: J.A. Couch and J.W. Fournie Editors, *Pathobiology of Marine and Estuarine Organisms*, CRC Press, Boca Raton, 255-287 pp.
- Peterson, C.H. (1986):** Quantitative allometry of gamete production by *Mercenaria mercenaria* into old age. *Mar. Ecol. Progr. Series*, 29: 93-97.
- Phillips, D.J.H. (1980):** Quantitative aquatic biological indicators: Their use to monitor trace metal and organochlorine pollution. *Appl. Sci. Publish. London*, 21: 326-339.
- Phillips, D.J.H. & Rainbow, P.S. (1994):** Biomonitoring of trace aquatic contaminants. Chapman & Hall, London: 86-97.
- Pipe, R.K. & Coles, J.A. (1995):** Environmental contaminants influencing immune function in marine bivalve molluscs. *Fish Shellfish Immunol.* 5: 581- 595.
- Plette, A.C.C.; NedeRlof, M.M.; Temminghoff, E.J.M. & Van Riemsdijk, W.H. (1999):** Bioavailability of heavy metals in terrestrial and aquatic systems. A quantitative approach *Enviro. Toxicol. Chem*, 18 (9): 1882– 1890.
- Prosi, F. (1981):** Heavy metals in aquatic organisms. In: *Metal pollution in the aquatic environment* (2nd edition), (Eds.) Forstner, U. & Wittmann, G.T.W. Springer – Verlag. New York: 271- 318.
- Pucheva, L.N.; Vaschenko, M.A. & Durkina, V.B. (2006):** Histopathology of the digestive gland of bivalve mollusk *Crenomytilus grayanus* (Dunker, 1853) from South Western Peter, The Great Bay, Sea of Japan. *Russian J. Mar. Biol.*, 32 (3): 166-172.

- Puente, X.; Villares, R.; Carral, E. & Carballeira, A. (1996):** Nacreous shell of *Mytilus galloprovincialis* as a biomonitor of heavy metal pollution in Galiza (NW Spain). *Sci. Total Environ.* 183 (3): 205- 211.
- Rainbow, P.S. (1993):** The significance of trace metal concentrations in marine invertebrates. In: *Ecotoxicology of metals in invertebrates* (Dallinger, R.; Rainbow, P.S. eds.). SETAC Special Publications Series. Boca Raton, FL: Lewis Publishers, 3-24.
- Rainbow, P.S. (2002):** Trace metal concentrations in aquatic invertebrates: why and so what? *Environ. Pollut.* 120: 497-507.
- Rainbow, P.S. & Dallinger, R. (1993):** Metal Uptake, regulation, and excretion in freshwater invertebrates. In : *Ecotoxicology of metals in invertebrates*, (Eds.) Dallinger, R. & Rainbow, P.S. Lewis Publishers, Boca Raton: 121 pp.
- Ramon, M. (1993):** Estudio de las poblaciones de *Chamelea gallina* (Linnaeus, 1758) *Donax trunculus* Linnaeus, 1758 (Mollusca: Bivalvia) en el golfo de valencia (Mediterraneo Occidental). Thesis, University of Barcelona.
- Ramon, M.; Gracenea, M. & Gonzalez-Moreno, O. (1999):** *Bacciger bacciger* (Trematoda, Fellodistomidae) infection in commercial clams *Donax trunculus* (Bivalvia, Donacidae) from the sandy beaches of the Western Mediterranean. *Dis. Aquat. Org.*, 35: 37–46.
- Ravera, O.; Cenct, R.; Beone, G.; Dantas, M. & Lodigiani, P. (2003):** Trace element concentrations in freshwater mussels as related to those in their environment. *J. Limnol.*, 62 (1): 61 –70.
- Rego I.; Martinez, A.; Gonzalez-Tizon, A.; Vieites, J.; Leira, F. & Mendez, J. (2002):** PCR technique for identification in the Pacific cod *Gadus macrocephalus* around Japan revealed by mtDNA and RAPD markers. *Fish Sci.*, 64: 673-679.
- Reid, T.M.; Dunn, J.F. & Loeb, L.A. (1994):** Mutagenesis by metal

- induced oxygen radicals. Environ. Health Perspect. 102, 57-61.
- Riedl, R. (1970):** Fauna Und Flora Der Adria. Verlag Paul Parey, Hamburg Und Berlin. P, 457 and Plate 153.
- Roberts, D.; Rittschof, D.; Gerhart, D. J.; Schmidt, A.R. & Hill, L.G. (1989):** Vertical migration of the clam *Mercenaria mercenaria* (L.) (Mollusca: Bivalvia): environmental correlates and ecological significance. J. Exper. Mar. Biol. Ecol., 126: 271-280.
- Robillard, S.; Beauchamp, G. & Laulier, M. (2003):** The role of abiotic factors and pesticide levels on enzymatic activity in the freshwater mussel *Anodonta cygnea* at three different exposure sites. Comp. Biochem. Physiol., 135C: 49-59.
- Roesijadi, G. & Robinson, W. E. (1994):** Metal regulation in aquatic animals: Mechanisms of uptake, accumulation, and release. In: Aquatic toxicology: Molecular, biochemical, and cellubr perspectives, (Eds.) Malins, D.C. & Ostrander, G.K. Lewis Publishers, Boca Raton: 387–420.
- Roman, M.; Gracenea, M. & Gonzalez-Moreno, O. (1999):** *Bacciger bacciger* (Trematoda, Fellodistomidae) infection in commercial clams *Donax trunculus* (Bivalvia, Donacidae) from the sandy beaches of the Western Mediterranean. Dis. Aquat. Org., 35: 37-46.
- Roux, D.J.; Van Vliet, H.R. & Van Veelen, M. (1993):** Towards integrated water quality monitoring: Assessment of ecosystem health. Water SA, 19 (4): 275-280.
- Said, T. O. (1996):** Analytical determination of dissolved petroleum components in sea water using coupling system (GC-MS). Ph.D. Thesis, Chem. Dept, Fac. Sci., Mansoura Univ., 100pp.
- Saitoh, K. (1998):** Genetic variation and local differentiation in the Pacific cod *Gadus macrocephalus* around Japan revealed by mt DNA and RAPD markers. Fish. Sci., 54: 673-679.
- Salanki, J.; Turpaev, T.M. & Nichaeva, M. (1991):** Mussel as a test animal for assessing environmental pollution and the

- sub-lethal effects of pollutants. In: Bioindicators and environmental management , (Eds.) Jeffrey, D.W. & Madden, B. Academic Press, London: 236 pp.
- Sato, M. & Nagashima, K. (2001):** Molecular characterization of a mitochondrial DNA segment from the Japanese scallop (*Patinopecten yessoensis*): Demonstration of a region showing sequence polymorphism in the population. Mar. Biotechnol., 3: 370–379.
- Schindler, D.W. (1996):** Ecosystems and ecotoxicology: A personal perspective. In: Ecotoxicology: A hierarchical treatment, (Eds.) Newman, M.C. & Jagoe, C.H. Lewis Publishers, Boca Raton: 375 pp.
- Segar, D.A.; Collins, J.D. & Riley, J.P. (1971):** The distribution of the major and some minor elements in marine animals. Mar. Biol. Ass. U.K., 51: 131–136.
- Seilacher, A. (1970):** Arbeitskonzept zur konstruktion morphologie. Lethaia, 3: 393-396.
- Sen Gupta, R. & Kureishy, T.W. (1981):** Present state of oil pollution in the northern Indian Ocean. Mar. Pollut. Bull., 12: 295- 301.
- Sen Gupta, R.; Naik, S. & Varadachari, V.V.R. (1989):** Environmental pollution in coastal areas of India. Ecotoxicology and Climate, Edi. By Bourdeau, P.; Haines, J.A.; Klein, W. & Krishna Murti, C.R. Published by John Wiley & Sons Ltd. 235-246.
- Sharabati, D. (1984):** “Red Sea Shells”. KPI Ltd., London.
- Shimamoto, M. (1986):** Shell microstructure of the Veneridae (Bivalvia) and its phylogenetic implications. Tohoku Univ., Sci. Rep., 2nd ser. (Geol.), 56 (1): 1-39.
- Shugart, L.R. (1996):** Molecular markers to toxic agents. In : Ecotoxicology: A hierarchical treatment (Eds.) Newman, M.C. & Jagoe, C.H. Lewis Publishers, Boca Raton: 139 pp.
- Shumway, S.E. (1990):** A review of the effects of the algal blooms on shellfish and aquaculture. J. World Aqu. Soc. 21 (2): 65–104.

- Shumway, S.E. & Cucci, T.L. (1987):** The effects of the toxic dinoflagellate *Protogonvaulax tamarensis* on the feeding and behaviour of bivalve molluscs. *Aquatic Toxicology*, 10: 9-27.
- Sinderman, J. (1990):** Epizootics and mass mortalities of shellfish In: C.J. Sinderman, Editor. *Principal Disease of Marine Fish and Shellfish*, Academic Press, San Diego (1990), 359-409.
- Sivaramakrishna, B. & Radhakrishnaiah, K. (2000):** Mercury induced alterations in the energetics of hepatopancreas of two freshwater molluscs, *Pila globosa* and *Lamellidens marginalis*. In: *Trace elements: Their distribution and effects in the environment*, (Eds.) Markert, B. & Friese, K. Elsevier, Amsterdam. 389–409 pp.
- Sleem, S.H. & Ali, T.K. (2008):** Application of RAPD-PCR in taxonomy of certain freshwater bivalves of genus *Caelatura*. *Global J. Molecu. Scie.* 3 (1): 27-31.
- Smolders, R.; Bervoets, L.; Wepener, V. & Blust, R. (2003):** A conceptual framework for using mussels as biomonitors in whole effluent toxicity. *Hum. Ecol. Risk Assess.* 9(3): 741- 760.
- Sparks, A.K. (1993):** Invertebrate disease – an overview. In: J.A. Couch and J.W. Fournie, Editors, *Pathobiology of Marine and Estuarine Organisms*, CRC Press, Boca Raton, 245-253 pp.
- Spatz, L.; Vidigal, T.H.D.A.; Silva, M.C.A.; Cappa, S.M.G. & Carvalho, O.S. (2000):** Characterization of *Biomphalaria orbignyi*, *Biomphalaria peregrine* and *Biomphalaria oligosa* by polymerase chain reaction and restriction enzyme digestion of the internal transcribed spacer region of the RNA ribosomal gene. *Mem. Inst. Oswaldo. Cruz.*, 95(6): 807-814.
- Stanley, S.M. (1970):** Relation of shell form to life habits in the Bivalvia (Mollusca). *Geol. Soc. America, Inc., Mem.* 125: 296 pp.

- Stanley, J. G. & Dewitt, R. (1983):** Species profiles: life histories and environmental requirements of coastal fishes and invertebrates (North Atlantic) hard clam. U.S. Fish Wildl. Serv. FWS/OBS-82/11.18.U.S. Army Corps of Engineers, TR El-82-4. 19 pp.
- Stark, J.D. & Banken, J.A.O. (1999):** Importance of population structure at the time of toxicant exposure. *Ecotoxicol. Environ. Safety*. 42, 282-287.
- Stegeman, J.J.; Brouwer, M.; Di Giulio, R.T.; Forlin, L.; Fowler, B.A.; Sanders, B.M. & Van Veld, P.A. (1992):** Molecular responses to environmental contamination : Enzyme and protein systems as indicators of chemical exposure and effect. In : *Biomarkers: Biochemical, physiological and histological markers of anthropogenic stress*, (Eds.) Huggett, R.J.; Kimerle, R.A., Mehrle, P.M. (Jr.) & Bergman, H.L. Lewis Publishers, Boca Raton: 235- 355.
- Stein, J.E.; Reichert, W.L. & Varanasi, U. (1994):** Molecular epizootiology: assessment of exposure to genotoxic compounds in teleosts. *Environ. Health Perspect.* 102 (12), 19- 23.
- Steinert, S.A (1999):** DNA Damage as a bivalve biomarker. *Biomarkers*, 4(6) 492- 496.
- Stepien, C.A. (1995):** Population genetic divergence and geographic patterns from DNA sequences: examples from marine and freshwater fishes. *Am. Fish. Soc. Symp.*, 17: 263– 287.
- Stiles, S.; Choromanski, J.; Schweitzer, D. & Xue, Q.Z. (1996):** Preliminary investigations of genetics and breeding of the bay scallop, *Argopecten irradians*. *J. Shellfish Res.*, 15: 461- 470.
- Stureson, U. (1978):** Cadmium enrichment in shells of *Mytilus edulis*. *Ambio*. 7: 122– 125.
- Sunila, L. (1987):** Histopathology of mussels (*Mytilus edulis* L.) from the Tvarmine area, the Gulf of Finland (Baltic Sea). *Ann. Zool. Fenn.*, 24: 55– 69.
- Sures, B. (2008):** Host- parasite interactions in polluted
-

- environments. J. Fish Biology, 73: 2133– 2142.
- Szefer, P. & Szefer, K. (1985):** Occurrence of ten metals in *Mytilus edulis* L. and *Cardium glaucum* L., from Gdansk Bay. Mar. Pollut. Bull., 16 (11): 446- 450.
- Szefer, P. & Szefer, K. (1990):** Metals in molluscs and associated bottom sediments of the southern Baltic. Helgol Meeresunters, 44: 424- 441.
- Szefer, P.; Frelek, K.; Szefer, K.; Lee, Ch-B; Kim, B-S; Warzocha, J. et al. (2002):** Distribution and relationships of trace metals in soft tissue, byssus and shells of *Mytilus edulis trossulus* from the southern Baltic. Environ. Pollut. 120: 423- 444.
- Taggart, J.B. & Ferguson, A. (1990):** Minisatellite DNA fingerprints of salmonid fishes. Anim. Genet., 21: 377– 389.
- Taggart, J.B.; Verspoor, E.; Galvin, P. T.; Moran, P. & Ferguson, A. (1995):** A minisatellite DNA marker for discriminating between European and North – American Atlantic salmon (*Salmo salar*). Can. J. Fish. Aquat. Sci., 52: 2305-2311.
- Taskinen, J. (1998):** Influence of trematode parasitism on the growth of a bivalve host in the field. Int. J. Parasitol., 28: 599– 602.
- Taskinen, J. & Valtonen, E.T. (1995):** Age, size and specific infection of *Anodonta piscinalis* (Bivalvia: Unionidae) with *Rhipidocotyle fennica* (Digenea: Bucephalidae) and its influence on host reproduction. Can. J. Zool., 73: 887-897.
- Tay, K.L.; Teh, S.J.; Doe, K.; Lee, K. & Jackman, P. (2003):** Histopathologic and histochemical biomarker responses of Baltic clam, *Macoma balthica*, to contaminated Sydney Harbour sediment, Nova Scotia, Canada. Environ. Health. Perspectives, 111 (3): 273–280.
- Thompson, M.R.; Ratcliffe, N.A. & Bayne, B.L. (1974):** Effect of starvation on structure and function in the digestive gland of the mussel *Mytilus edulis*. Mar. Biol. Assoc. U.K., 54:

699–712.

- Tillier, L. & Bavay, A. (1905):** Les mollusques testaces du canal de Suez. Bulletin de la Societe Zoologique de France, 30: 170-181.
- Toll, R.B.; Prezant, R.S. & Rollins, H.B. (2003):** A novel method for locating tagged infaunal bivalves: Submersible pulse technology metal detectors. J. of Shellfish Research, 22: 501-503.
- Triebskorn, R. & Kohler, H.R. (1996):** The impact of heavy metals on the grey garden slug. *Deroceras reticulatum* (Muller): Metal storage, cellular effects and semi- quantitative evaluation of metal toxicity. Environ. Pollut. 93 (3): 327–343.
- Van der Oost, R.; Beyer, J. & Vermeulen, N. P. E. (2003):** Fish bioaccumulation and biomarkers in environmental risk assessment: a review. Environ. Toxicol. Chem., 13: 57-149.
- Van Gastel, C.A. & Van Brummelen, T.C. (1994):** Incorporation of the biomarker concept in ecotoxicology calls for a redefinition of terms. Ecotoxicology, 5: 217–225.
- Van Ginneken, L., Chowdhury, M.J. & Blust, R. (1999):** Bioavailability of cadmium and zinc to the common carp, *Cyprinus carpio*, in complexing environments: A test for the validity of the free ion activity model. Environ. Toxicol. Chem., 18 (10): 2295–2304.
- Venkataraman, K. & Wafer, M. (2005):** Coastal and marine biodiversity of India. Indian J. Mar. Sci., 34(1): 57-75.
- Volpe, J. P. & Ferguson, M.M. (1996):** Molecular genetic examination of the polymorphic Arctic charr *Salvelinus alpinus* of Thingvallavatn, Iceland. Mol. Ecol., 5: 763-772.
- Walker, C.H. (1992):** Biochemical responses as indicators of toxic effects of chemicals in ecosystems. Toxicol. Lett, 64-65: 527–533.
- Walker, C.H. (1998):** The use of biomarkers to measure the interactive effects of chemicals. **Ecotox. Environ. Saf.** 40:

65–70.

- Walsh, K.; Dunstan, R.H. & Murdoch, R.N. (1995):** Differential bioaccumulation of heavy metals and organopollutants in the soft tissue and shell of the marine gastropod, *Austrococklea constricta*. Arch. Environ. Contam. Toxicol. 28(1): 35-39.
- Wang, Y. & Evans, R.D. (1993):** Influence of calcium concentrations on cadmium uptake by the freshwater mussel *Elliptio complanata*. Can. J. Fish. Aquat. Sci., 50: 2591-2596.
- Ward, J.E. & Shumway, S.E. (2004):** Separating the grain from the chaff: particle selection in suspension – and deposit-feeding bivalves. J. Exp. Mar. Bio. & Eco. 300: 83-130.
- Ward, J.E.; Newell, R.I.E.; Thompson, R.J. & MacDonald, B.A. (1994):** *In vivo* studies of suspension – feeding processes in the eastern oyster, *Crassostrea virginica* (Gmelin). Biological Bulletin, 186: 221-240.
- Ward, R.D.; Elliott, N.G.; Innes, B.H.; Smolenski, A.J. & Grewe, P.M. (1997):** Global population structure of yellowfin tuna, *Thunnus albacares*, inferred from allozyme and mitochondrial DNA variation. Fish. Bull, 95: 566-575.
- Welsh, J. & McClelland, M. (1990):** Fingerprinting genomes using PCR with arbitrary primers. Nucleic Acids Res., 18: 7213-7218.
- Wepener, V. (1997):** Metal ecotoxicology of the Olifants River in the Kruger National Park and the effect thereof on fish haematology. Ph. D.-Thesis, Rand Afrikaans University, South Africa.
- Williams J.G.K.; Kubelik, A. R.; Livak, K.J.; Rafalski, J.A. & Tingey S.V. (1990):** DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. Nucleic Acids Res., 18: 6513-6535.
- Winstead, J.T. (1995):** Digestive tubule atrophy in eastern oysters, *Crassostrea virginica* (Gmelin, 1791) exposed to salinity and starvation stress. J. Shellfish Res., 14: 105-111.

-
- Winston, G.W. & Di Giulio, R.T. (1991):** Prooxidant and antioxidant mechanisms in aquatic organisms. *Aquat. Toxicol.* 19: 137– 161.
- Wittmann, G. (1981):** Toxic metals. In : Metal pollution in the aquatic environment (2nd edition), (Eds.) Forstner, U & Wittmann, G.T.W. Springer-Verlag, Berlin: 7- 9 pp.
- Wong, P.K. (1988):** Mutagenicity of heavy metals. *Bullet. Environ. Contami. Toxicol.* 40, 597- 603.
- Woods Hole Oceanographic Institute (WHOI) (2004):** The harmful Algae Page. Available: Accessed: January 2005.
- Wright, D.A. (1995):** Trace metal and major ion interactions in aquatic animals. *Mar. Pollut. Bull.*, 31: 8- 18.
- Yap, C.K.; Chua, B.H.; Teh, C.H.; Tan, S.G. & Ismail, A. (2007):** Patterns of RAPD markers and heavy metal concentrations in *Perna viridis* (L.), collected from metal- contaminated and uncontaminated coastal waters: Are they correlated with each other? *J. of Genetics*, 43, 544-550.
- Yevich, P.P. & Barszcz, C.A. (1983):** Histopathology as monitor of marine pollution. Results of histopathologic examination of the animals collected from the US 1976 Mussel Watch Program. *Rapp. P-V Reun. Cons. Int. Explor. Mer.*, 182: 96-102.
-
