

REFERENCES

- Abu Laimun B. (2004):** Symptomatic spontaneous pneumothorax in term newborn infants. *Pediatr Pulmonol.* May; 37(5):443-6.
- Abbassi V, Merchant K, Abramson D. (2000):** Postnatal triiodothyronine concentrations in healthy preterm infants and in infants with respiratory distress syndrome. *Pediatr Res*; 11:802-4.
- Ainsworth SB, Milligan DW. (2002):** Surfactant therapy for respiratory distress syndrome in premature neonates: a comparative review. *Am J Respir Med.* 1(6):417-33.
- Ainsworth SB. (2005):** Pathophysiology of neonatal respiratory distress syndrome: implications for early treatment strategies. *Treat Respir Med.* 4(6):423-37.
- Alfirevic Z. (2007):** Early amniocentesis versus transabdominal chorion villus sampling for prenatal diagnosis. *Cochrane Database Syst Rev.* ;(2):CD000077.
- Al Tawil K, Abu-Ekteish FM, Tamimi O, Al Hathlol K, Altman DG. (1991):** Relation between several variables. In: *Practical statistics for medical research.* Chapman and Hall, 325-64.
- American Academy of Pediatrics, (2004):** Medical Home Initiatives for Children with Special Needs Project Advisory Committee. The medical home. *Pediatrics.* ; 110:184–186.
- American Lung Association. (2006):** Respiratory distress syndrome of new born fact sheet. Accessed May 7.
- Antenatal Corticosteroids Revisited. (2004):** Repeat Courses. NIH Consensus Statement Online August 17-18; [cited year, month, day]; 17(2): 1-10.

- Anil N, Praveen K, Sourabh D and Rajesh K (2001):** Surfactant Therapy for Hyaline Membrane Disease Indian Pediatrics, 38:640-646.
- Attar MA, Donn SM. (2002):** Mechanisms of ventilator-induced lung injury in premature infants. Semin Neonatol.Oct; 7(5):353-60.
- Augusto LA, Li J, Synguelakis M, Johansson J & Chaby (2002):** Structural basis for interactions between lung surfactant protein C and bacterial lipopolysaccharide. J Biol Chem 277: 23484-23492.
- Ballard JL, Khoury JC & Wedig K. (1991):** New Ballard Score expanded to include extremely premature infants. J Pediatrics; 119:417-423.
- Barton L, Hodgman JE, Pavlova Z. (1999):** Causes of death in the extremely low birth weight infant. Pediatrics. Feb; 103(2):446-51.
- Bhakoo ,Narang, A., Karthikeyan, G. & Kumar, P (2000):** Spectrum of respiratory distress in very low birth weight neonates. Indian J Pediatr 67, 8034.
- Bhuta T, Henderson Smart DJ.(1997):** Elective high-frequency oscillatory ventilation versus conventional ventilation in preterm infants with pulmonary dysfunction: systematic review and meta-analyses. Pediatrics. Nov; 100(5):E6.
- Bhuta T, Henderson Smart DJ.(2000):** Rescue high frequency oscillatory ventilation versus conventional ventilation for pulmonary dysfunction in preterm infants. Cochrane Database Syst Rev.; (2):CD000438.
- Bhutta ZA, Yusuf K. (2001):** Neonatal respiratory distress syndrome in Karachi: some epidemiological considerations. Paediatr Perinat Epidemiol. Jan; 11(1):37-43.

- Bhutta ZA, Yusuf K. (2001):** Profile and outcome of the respiratory distress syndrome among newborns in Karachi: risk factors for mortality. *J Trop Pediatr.* Jun; 43(3):143-8.
- Bianco AC, Salvatore D, Gereben B, Berry MJ, Larsen PR (2002):** "Biochemistry, cellular and molecular biology, and physiological roles of the iodothyronine selenodeiodinases". *Endocr Rev* 23 (1): 38–89.
- Bober K, Swietliński J. (2006):** Diagnostic utility of ultrasonography for respiratory distress syndrome in neonates. *Med Sci Monit.* Oct; 12(10):CR440-6.
- Bongers, Schokking JJ, Koot HM, Wiersma D, Verkerk PH, and Keizer Schrama SM. (2000):** Influence of timing and dose of thyroid hormone replacement on development in infants with congenital hypothyroidism. *J Pediatr.* ; 136: 292 – 29.
- Bonanno C, Fuchs K, Wapner RJ. (2007):** Single versus repeat courses of antenatal steroids to improve neonatal outcomes: risks and benefits. *Obstet Gynecol Surv.* Apr; 62(4):261-71.
- Boo NY, Cheong KB, Cheong SK, Lye MS, Zulfiqar MA. (1997):** Usefulness of stable micro bubble test of tracheal aspirate for the diagnosis of neonatal respiratory distress syndrome. *J Paediatr Child Health.* Aug; 33(4):329-34.
- Boyd S. (2004):** Causes and treatment of neonatal respiratory distress syndrome. *Nurs Times.* Jul 27-Aug 2; 100(30):40-4.
- Boyle KM, Baker VL, Cassaday CJ. (2001):** Neonatal pulmonary disorders. In: Barnhart SL, Czervinske MP, eds. *Perinatal and Pediatric Respiratory Care.* 2nd ed. Philadelphia, Pa: WB Saunders Co; 445.

- Brem AS. (1992):** Electrolyte disorders associated with respiratory distress syndrome and bronchopulmonary dysplasia. Clin Perinatol. Mar; 19(1):223-32.
- Brion LP, Soll RF. (2000):** Diuretics for respiratory distress syndrome in preterm infants. Cochrane Database Syst Rev. ;(2):CD001454.
- Casulari LA, Mangieri P, Naves LA, Ferreira M, Domingues L (2004):** Non thyroidal illness syndrome in patients with subarachnoid hemorrhage due to intracranial aneurysm. Ar Qneuro-Psiquiat 62:26–32.
- Caukwell S & Murphy DJ (2002):** The effect of mode of delivery and gestational age on neonatal outcome of the non-cephalic-presenting second twin. Am J Obstet Gynecol 187:1356–1361.
- Center of disease control "CDC". (1990):** Low birth weight--United States, 1975-1987. MMWR; 39:148-51.
- Chambers CD, Hernandez-Diaz S, Van Marter LJ, Werler MM, Louik C, Jones KL, Mitchell AA.(2006):** Selective serotonin-reuptake inhibitors and Risk of persistent pulmonary hypertension of the newborn. N Engl J Med. Feb 9; 354(6):579-87.
- Chard T, Soe A, Costeloe K. (1997):** The risk of neonatal death and respiratory distress syndrome in relation to birth weight of preterm infants. Am J Perinatol. Oct; 14(9):523-6.
- Charles, L.; Anderson, J. and Stewart, I.E. (1998):** Retinopathy of prematurity: In Colherty J and Stark A (Eds), Manual of Neonatal Care 4th ed., Philadelphia, Lippincott-Raven, P: 643-648.

- Chatson, K., Fant, M.E., and Cloherty, J., (1998):** Temperature control: In Colherty J. and Stark A. (ed.) Manual of Neonatal Care. 4th. ed. Lippincott-Raven Philadelphia p-p 139.
- Chiang MF, Arons RR, Flynn JT, Starren JB. (2004):** Incidence of retinopathy of prematurity from 1996 to 2000: analysis of a comprehensive New York State patient database. *Ophthalmology*; 111:1317–1325.
- Chida S, Phelps DS, Cordle C, Soll R, Floros J, Taeusch HW. (1998):** Surfactant-associated proteins in tracheal aspirates of infants with respiratory distress syndrome after surfactant therapy. *Am Rev Respir Dis*. Apr; 137(4):943-7.
- Cho K, Matsuda T, Okajima S, Sagawa T& Kobayashi K (2000):** Prediction of respiratory distress syndrome by the level of pulmonary surfactant protein A in cord blood sera. *Biol Neonate* 77:83–87.
- Chauhan SP, Hendrix NW, Magann EF. (2005):** Neonatal organ dysfunction among newborns at gestational age 34 weeks and umbilical arterial pH<7.00. *J Matern Fetal Neonatal Med*. Apr; 17(4):261-8.
- Cleary GM, Wiswell TE. (1998):** Meconium stained amniotic fluid and the meconium aspiration syndrome. An update. *Pediatr Clin North Am*. Jun; 45(3):511-29.
- Cohen RN, Weintraub BD, Wondisford FE (2000):** Thyrotropin. In: Braverman LE, Utiger RD, eds. *Werner's and Ingbar's the thyroid*. Philadelphia: Lippincott- Raven; 202–219.
- Cole FS, Noguee LM, Hamvas A. (2006):** Defects in surfactant synthesis: clinical implications. *Pediatr Clin North Am*. Oct; 53(5):911-27.

- Courtney SE, Durand DJ, Asselin JM. (2002):** High frequency oscillatory ventilation versus conventional mechanical ventilation for very-low-birth weight infants. *N Eng J Med*; 347:643-652.
- Cosmi EV, La Torre R, Piazzè JJ, Maranghi GL, Lerro N, Bianco D, Anceschi MM.(1997):** Intra - amniotic surfactant for prevention of neonatal Respiratory distress syndrome (IRDS): rationale and personal experience. *Eur J Obstet Gynecol Reprod Biol.* Feb; 71(2):135.
- Cosmi EV. (1992):** Prenatal prevention of respiratory distress syndrome: new pharmacologic approaches. *Early Hum Dev.* Jun-Jul; 29(1-3):283-6.
- Cowett RM, Loughhead JL. (2002):** Neonatal glucose metabolism: differential diagnoses, evaluation, and treatment of hypoglycemia. *Neonatal Netw.* Jun; 21(4):9-19.
- Crouch E & Wright JR (2001):** Surfactant proteins a and d and pulmonary host defense. *Annu Rev Physiol* 63: 521-554.
- Crowley P. (2003):** Prophylactic corticosteroids for preterm birth. *Cochrane Database Syst Rev* ;(4):CD000065.
- Crowther CA, Alfirevic Z, Haslam RR. (2004):** Thyrotropin releasing hormone added to corticosteroids for women at risk of preterm birth for preventing neonatal respiratory disease. *Cochrane Database Syst Rev.* ;(2):CD000019.
- Crowther CA, Harding JE. (2007):** Repeat doses of prenatal corticosteroids for women at risk of preterm birth for preventing neonatal respiratory disease. *Cochrane Database Syst Rev.* Jul 18 ;(3):CD003935.

- Curstedt T, Robertson B. (1993):** Comparison of prophylaxis and rescue treatment with Curosurf in neonates less than 30 weeks' gestation: a randomized trial. *Pediatrics*. Dec; 92(6):768-74.
- Cuestas RA. (2000):** Thyroid function in healthy premature infants. *J Pediatr*; 92:963-7.
- Cuestas RA, Lindall A, Engel RR. (1999):** Low thyroid hormones and respiratory distress syndrome of the newborn. *N Eng J Med* ; 295:297-302.
- De Paoli A, Davis P, Faber B, Morley C. (2008):** Devices and pressure sources for administration of nasal continuous positive airway pressure (NCPAP) in preterm neonates. *Cochrane Database Syst Rev*. Jan 23 ;(1):CD002977.
- Dechant KL, Faulds D. (2002):** Colfosceril palmitate. A review of the therapeutic efficacy and clinical tolerability of a synthetic surfactant preparation (Exosurf Neonatal) in neonatal respiratory distress syndrome. *Drugs*. Nov; 42(5):877-94.
- Delahunty C, Richard K, Coughtrie M, et al. (2001):** Transient hypothyroxinemia in preterm infants. *Dev Med Child Neurol Suppl.*; 86:26–27.
- Delang F, Dalhem A, Bourdoux P, et al. (2001):** Increased risk of primary hypothyroidism in preterm infants. *J Pediatr*; 105:462-9.
- DeMello DE. (2004):** Pulmonary pathology. *Semin Neonatol*; 9: 311–329.
- Demissie K, Ananth CV, Martin J, Hanley ML, MacDorman MF(2002):** Fetal and neonatal mortality among twin gestations in The United States: the role of intrapair birth weight discordance. *Obstet Gynecol* 100:474–480.

- Donn SM. (2005):** Lucinactant: a novel synthetic surfactant for the treatment of respiratory distress syndrome. *Expert Opin Investig Drugs*. Mar; 14(3):329-34.
- Dyke MP, Kohan R, Evans S. (1995):** Morphine increases synchronous ventilation in preterm infants. *J Paediatr Child Health*. Jun; 31(3):176-9.
- Doyle NM, Lally KP. (2004):** The CDH study group and advances in the clinical care of the patient with congenital diaphragmatic hernia. *Semin Perinatol.* ; 28:174–184.
- Dubowitz LMS, Dubowitz V, Goldberg C. (1991):** Clinical assessment of gestational age in the newborn infant. *J Pediatr* ; 77:1-10.
- Edwards MS. (1997):** Antibacterial therapy in pregnancy and neonates. *Clin Perinatol*. Mar; 24(1):251-66.
- Egberts J, de Winter JP, Sedin G, de Kleine MJ, Broberger U, van Bel F, Egerman RS, Mercer BM, Doss JL, Sibai BM. (1998):** A randomized controlled trial of oral and intramuscular dexamethasone in the prevention of neonatal respiratory distress syndrome. *Am J Obstet Gynecol*. Nov; 179(5):1120-3.
- Egyptian National Neonatal Care Program [ENNCP]. (2001):** Gestational age assessment should be unhurried.
- Elias-Jones AC, Barrett DA, Rutter N, Shaw PN, Davis SS. (2002):** Diamorphine infusion in the preterm neonate. *Arch Dis Child*. Oct; 66(10 Spec No):1155-7.
- Emdin M, Passino C, Prontera C, Iervasi A, Ripoli A, Masini S, Zucchelli GC, Clerico A.(2004):** Cardiac natriuretic hormones , thyroid Neuro hormones and cytokines in normal subjects and patients with heart failure. *Clin Chem Lab Med* 42:627–636.

- Erickson SJ, Grauaug A, Gurrin L and Swaminathan M (2000):** Hypocarbia in the ventilated preterm infants and its effect on intraventricular hemorrhage and bronchopulmonary dysplasia. *J Paediatr Child Health*.38 (6):560-2.
- Filippi L, Cecchi A, Tronchin M, Dani C, Pezzati M, Seminara S, Gasperini S, Zammarchi E, Rubaltelli FF(2004):** Dopamine infusion and Hypothyroxinemia in very low birth weight preterm infants. *Eur J Pediatr* 163:7–13.
- Fisher DA. (1999):** Hypothyroxinemia in premature infants: is thyroxine treatment necessary? *Thyroid*; 9:715–720.
- Fiori HH, Linderholm B, Fiori RM & Robertson B. (2001):** Computerized image analysis of bubbles in gastric aspirate for prediction of respiratory distress syndrome. *Acta Paediatr* 90:1402–1404.
- Franklin R, O'Grady C.** Neonatal thyroid function: effects of nonthyroidal illness. *J Pediatr* 1985; 107:599-602.
- Friedman S, Shinwell ES. (2004):** Prenatal and postnatal steroid therapy and child neurodevelopment. *Clin Perinatol* 31:529–544.
- Fujiwara T, Maeta H, Chida S, Morita T, Watabe Y, Abe T.(1999):** Artificial surfactant therapy in hyaline-membrane disease. *Lancet*. Jan 12; 1(8159):55-9.
- Furman B, Shoham-Vardi I, Bashiri A& Mazor M. (2000):** Preterm premature rupture of membranes is not an independent risk factor for neonatal morbidity. *J Matern Fetal Med* 10:107–111.
- Gardner DS, Fowden AL, Giussani DA. (2002):** Adverse intrauterine conditions diminish the fetal defense against acute hypoxia by

increasing nitric oxide activity. *Circulation*. Oct 22; 106(17):2278-83.

Gerard P, Bachy A, Battisti O. (2002): Transient thyroid insufficiency in sick neonates. *Acta Paediatr Belg* ; 32:193-8.

Ghafoor T, Mahmud S, Ali S, Dogar SA. (2003): Incidence of respiratory distress syndrome. *J Coll Physicians Surg Pak*. May; 13 (5): 271-3.

Ghidini a, Poggi SH, Spong CY, Goodwin KM, Vink J, Pezzullo JC (2005): Role of lamellar body count for the prediction of neonatal Respiratory distress syndrome in non-diabetic pregnant women. *Arch Gynecol Obstet*. Apr; 271(4):325-8.

Gibson AT. (2002): Perinatal corticosteroids and the developing lung. *Paediatr. Respir. Rev*. Mar; 3(1):70-6.

Gill AB, Weindling AM. (1993): Echocardiographic assessment of cardiac function in shocked very low birth weight infants. *Arch Dis Child*. Jan; 68(1 Spec No):17-21.

Gilbert C, Fielder A, Gordillo L, et al. (2005): Characteristics of infants with severe retinopathy of prematurity in countries with low, moderate, and high levels of development: implications for screening programs. *Pediatrics*; 115(5).

Goldenberg RL, Hauth JC & Andrews WW (2000): Intrauterine infection and preterm delivery. *N Engl J Med* 342:1500–1507.

Goldstein RF, Thompson RJ, Oehler JM, Brazy JE. (2001): Influence of acidosis, hypoxemia, and hypotension on neurodevelopmental outcome in very low birth weight infants. *Pediatrics*. Feb; 95(2):238-43.

- Golombek SG, Truog WE. (2000):*** Effects of surfactant on gas exchange and clinical course in near-term newborns with RDS. *J Perinat Med.*; 28:436–442.
- Gomella TL. (2004):*** Neonatology: management, procedures, on-call problems, diseases and drugs. Gomella, TL, Cunningham, MD., Eyal, FG, Zenk, KE (Eds.). 5th edition. Medical Publishing Division, McGraw-Hill Companies.
- Greenough A, Donn SM. (2007):*** Matching ventilatory support strategies to respiratory pathophysiology. *Clin Perinatol.* Mar; 34(1):35-53.
- Greenough A, Milner AD and Dimtrou G. (2003):*** Synchronized mechanical ventilation for respiratory support in newborn infants (Cochrane Review). In: The Cochrane Library, Issue 4, Oxford: Update Software.
- Greenough A, Morley GJ and Robertson NRC (1996):*** Acute respiratory disease in the newborn. Respiratory distress syndrome in: *Textbook of Neonatology.* (Ed): Robertson NRC. 2nd edition, Churchill Livingstone, Edinburgh, 385-486.
- Greenough A (2000):*** Expanded use of surfactant replacement therapy. *Eur J Pediatr* 159,635-40.
- Grier DG and Halliday HL (2004):*** Effects of glucocorticoids on fetal and neonatal lung development. *Treat, Respir Med.* 3(5):295-306.
- Hack M, Wright LL, Shankaran S, Tyson JE, Horbar JD, Bauer CR, Younes N. (2000):*** Very low birth weight outcomes of the National Institute of Child Health and Human Development Neonatal Network, November 1989 to October 2000. *Am J Obstet Gynecol.* Feb; 172(2 Pt 1):457-64.

- Hacking D, Watkins A, Fraser S, Wolfe R, Nolan T. (2001):** Respiratory distress syndrome and birth order in premature twins. Arch Dis Child Fetal Neonatal Ed. Mar; 84(2):F117-21.
- Hallman M. (2002):** Is rapid surfactant analysis at birth feasible in the diagnostics of respiratory distress syndrome? Acta Paediatr 91:370–372.
- Hallman M, Glumoff V, Rämet M. (2002):** Surfactant in respiratory distress syndrome and lung injury. Comp Biochem Physiol A Mol Integr Physiol. May; 129(1):287-94.
- Hallman M, Haataja R, Marttila R. (2002):** Surfactant proteins and genetic predisposition to respiratory distress syndrome. Semin Perinatol. Dec; 26(6):450-60.
- Hallman M, Merritt TA, Kari A, Bry K. (2002):** Factors affecting surfactant responsiveness. Ann Med. Dec; 23(6):693-8.
- Hallman M, Teramo K. (2002):** Measurement of the lecithin/sphingomyelin ratio and phosphatidylglycerol in amniotic fluid: an accurate method for the assessment of fetal lung maturity. Br J Obstet Gynaecol. Aug; 88(8):806-13.
- Hallman M. (2002):** Lung surfactant in respiratory distress syndrome. Acta Anaesthesiol Scand Suppl.; 95:15-20.
- Hammoud MS, ElSORI HA, Hanafi EA, Shalabi AA, Fouda IA, Devarajan LV. (2003):** Incidence and risk factors associated with the patency of ductus arteriosus in preterm infants with respiratory distress syndrome in Kuwait. Saudi Med J. Sep; 24(9):982-5.
- Hameed B, Shyamanur K, Kotecha S, et al. (2004):** Trends in the incidence of severe retinopathy of prematurity in a

geographically defined population over a 10-year period.
Pediatrics. ; 113:1653–1657.

Hansen AK, Wisborg K, Uldbjerg N and Henriksen TB.(2007): Risk of respiratory morbidity in term infants delivered by elective caesarean section: cohort study. *BMJ.* Jan 12; 336(7635):85-7.
Epub .Dec 11.

Hansen AK, Wisborg K, Uldbjerg N, Henriksen TB. (2007): Elective caesarean section and respiratory morbidity in the term and near-term neonate. *Acta Obstet Gynecol Scand.*; 86(4):389-94.

Harte GJ, Gray PH, Lee TC, Steer PA, Charles BG. (1997): Haemodynamic responses and population pharmacokinetics of midazolam following administration to ventilated, preterm neonates. *J Paediatr Child Health.* Aug; 33(4):335-8.

Hein HA, Ely JW, Lofgren MA. (2000): Neonatal respiratory distress in the community hospital: when to transport, when to keep. *J Fam Pract.* Apr; 46(4):284-9.

Henderson-Smart DJ, Bhuta T, Cools F, Offringa M. (2007): Elective high frequency oscillatory ventilation versus conventional ventilation for acute pulmonary dysfunction in preterm infants. *Cochrane Database Syst Rev.* ;(2):CD000104.

Henderson-Smart DJ, Cools F, Bhuta T, Offringa M. (2007): Elective high frequency oscillatory ventilation versus conventional ventilation for acute pulmonary dysfunction in preterm infants. *Cochrane Database Syst Rev.* Jul 18 ;(3):CD000104.

Hermansen CL, Lorah KN. (2007): Respiratory distress in the newborn. *Am Fam Physician.* Oct 1; 76(7):987-94.

- Hjalmarson O & Sandberg K (2002):** Abnormal lung function in healthy preterm infants. *Am J Respir Crit Care Med* 165:83–87.
- Ho JJ, Henderson-Smart DJ, Davis PG. (2002):** Early versus delayed initiation of continuous distending pressure for respiratory distress syndrome in preterm infants. *Cochrane Database Syst Rev.* ;(2):CD002975.
- Ho JJ, Subramaniam P, Henderson-Smart DJ, Davis PG. (2002):** Continuous distending pressure for respiratory distress syndrome in preterm infants. *Cochrane Database Syst Rev.* ;(2):CD002271.
- Ho NK. (1993):** Factors affecting responses of infants with respiratory distress syndrome to exogenous surfactant therapy. *Singapore Med J.*Feb; 34(1):74-7.
- Horbar JD, Carpenter JH, Buzas J, et al. (2004):** Collaborative quality improvement to promote evidence based surfactant for preterm infants: a cluster randomized trial. *BMJ.* ; 329:1004.
- Howlett A, Ohlsson A. (2003):** Inositol for respiratory distress syndrome in preterm infants. *Cochrane Database Syst Rev.* ;(4): CD000366.
- Hume R, Simpson J, Delahunty C, van Toor H, Wu SY, Williams FLR, Visser TJ (2004):** Human fetal and cord serum thyroid hormones: Developmental trends and inter-relationships. *J Clin Endocrinol Metab* 89:4097–4103.
- Ibrahim M, Morreale de Escobar G, Visser TJ, Dura'n S, van Toor H, StrachanJ, Williams FLR, Hume R (2003):** Iodine deficiency associated With parenteral nutrition in extreme preterm infants. *Arch Dis Child* 88:F56–F57.

- Ikegami M & Jobe AH. (1998):** Surfactant protein-C in ventilated premature lamb lung. *Pediatr Res* 44: 860-864.
- Ingemarrson I. (2003).** Gender aspects of preterm birth. *BJOG* ;110(suppl 20):34-8.
- Jacohsen BB, Peitersen B, Iiummer L. (2001):** Serum concentrations of thyrotropin, thyroid hormones and thyroid hormone-binding proteins during acute and recovery stages of idiopathic respiratory distress syndrome. *Acta Paediatr Scand* ; 68:257-64.
- Jain L, Dudell GG. (2006):** Respiratory transition in infants delivered by cesarean section. *Semin Perinatol*. Oct; 30(5):296-304.
- Jain L, Eaton DC. (2006):** Physiology of fetal lung fluid clearance and the effect of labor. *Semin Perinatol*. Feb; 30(1):34-43.
- Jenkinson SG. (1993):** Oxygen toxicity. *New Horiz*. Nov; 1(4):504-11.
- Kala P, Ten Have T, Nielsen H, Dunn M, Floros J. (1998):** Association of pulmonary surfactant protein A (SP-A) gene and respiratory distress syndrome: interaction with SP-B. *Pediatr Res*. Feb; 43(2):169-77.
- Kamper J, Wulff K, Larsen C, Lindequist S. (1993):** Early treatment with nasal continuous positive airway pressure in very low-birth-weight infants. *Acta Paediatr*. Feb; 82(2):193-7.
- Kantor MJ, Leef KH, Bartoshesky L, Getchell J, Paul DA. (2003):** Admission thyroid evaluation in very-low-birth-weight infants: association with death and severe intraventricular hemorrhage. *Thyroid* 13:965–969.
- Kaplan LA, Chapman JF, Bock JL, Santa Maria E, Clejan S, Huddleston DJ, Reed RG, Bernstein LH, Gillen-Goldstein J. (2002) :** Prediction of respiratory distress syndrome using the

Abbott FLM-II amniotic fluid assay. Clin Chim Acta. Dec; 326 (1-2):61-8.

Kattwinkel JM. (2006): The new Textbook of Neonatal Resuscitation, 4th edition, Multimedia CD-ROM. American Academy of Pediatrics. (March 31, 2006).

Kester MH, Martinez de Mena R, Obregon MJ, Marinkovic D, Howatson A, Visser TJ, Hume R, Morreale de Escobar G (2004): "Iodothyronine levels in the human developing brain: major regulatory roles of iodothyronine deiodinases in different areas". J Clin Endocrinol Metab 89 (7): 3117–3128.

Keszler M, Durand DJ. (2001): Neonatal high-frequency ventilation. Past, present, and future. Clin Perinatol. Sep; 28(3):579-607.

Keszler M, Modanlou HD, Brudno DS, Clark FI, Cohen RS, Ryan RM, Kaneta MK, Davis JM. (1997): Multicenter controlled clinical trial of High-frequency jet ventilation in preterm infants with uncomplicated respiratory distress syndrome. Pediatrics. Oct; 100(4):593-9.

Kim A. Gerten, MD, a Dean V. Coonrod, MD, MPH, a R. Curtis Bay, PhD Linda R. & Chambliss, MD, MPH (2005): Cesarean delivery was an independent risk factor for RDS. The risk was reduced with labor before cesarean, but still elevated. This supports the importance of being certain of fetal lung maturity before cesarean delivery, particularly when done before labor. Pediatrics ; 63:303-8.

Kimya Y, Küçükkömürcü S, Ozan H, Uncu G. (1995): Antenatal ambroxol usage in the prevention of infant respiratory distress

syndrome. Beneficial and adverse effects. Clin Exp Obstet Gynecol.; 22(3):204-11.

Klein AH, Foley B. Kenny FM, Fisher DA. (2000): Thyroid hormone and thyrotropin responses to parturition in premature infants with and without the respiratory distress syndrome. Pediatrics; 63:380-5.

Klein AH, Foley B, Foley TP, MacDonald HM, Fisher DA. (2000): Thyroid function studies in cord blood from premature infants with and without RDS. J Pediatr ; 98:818-20.

Koivisto M, Marttila R, Saarela T, Pokela ML, Valkama AM, Hallman M. (2005): Wheezing illness and re-hospitalization in the first two years of Life after neonatal respiratory distress syndrome. J Pediatr. Oct; 147(4):486-92.

Kolås T, Saugstad OD, Daltveit AK, et al., (2006): Planned cesarean versus planned vaginal delivery at term: comparison of newborn infant outcomes. Am J Obstet Gynecol. Dec; 195(6):1538-43. Epub. Jul 17.

Kotecha S. (2001): growth and nutrition: Implications for the newborn infant. Arch Dis Child Fetal Neonatal Ed 82:F69–74.

Kok JH, Hart G. Endert E, Koppe JG, De Vijlder JJM. (1983): Normal ranges of T4 screening values in low birth weight infants. Arch Dis Child ; 58:19.

Krauss AN. (2003): New methods advance treatment for respiratory distress syndrome. Pediatr Ann. Sep; 32(9) : 585-91.

Kumar A, Bhat BV. (2005): Epidemiology of respiratory distress of newborns. Indian J Pediatr. Jan-Feb; 63(1) : 93-8.

- Kumar A, Bhatnagar V. (2005):** Respiratory distress in neonates. Indian J Pediatr. May; 72(5) : 425-8.
- Kumar P, Kiran PS. (2004):** Changing trends in the management of respiratory distress syndrome (RDS). Indian J Pediatr. Jan; 71(1):49-54.
- Kurl S, Heinonen KM, Kiekara O. (1999):** The first chest radiograph in neonates exhibiting respiratory distress at birth. Clin Pediatr (Phila). May; 36(5):285-9.
- Laubscher B, van Melle G, Fawer CL, Sekarski N, Calame A. (1996):** Haemodynamic changes during high frequency oscillation for respiratory distress syndrome. Arch Dis Child Fetal Neonatal Ed. May; 74(3):F172-6.
- Lemyre B, Davis PG, de Paoli AG. (2002):** Nasal intermittent positive pressure ventilation (NIPPV) versus nasal continuous positive airway pressure (NCPAP) for apnea of prematurity. Cochrane Database Syst Rev. 2002 ;(1):CD002272.
- Lewis V, Whitelaw A. (2002):** Furosemide for transient tachypnea of the newborn. Cochrane Database Syst Rev. ;(1):CD003064.
- Lumley J. (2003):** Method of delivery for the preterm infant. BJOG. Apr; 110 Suppl 20:88-92.
- Laubscher B, van Melle G, Fawer CL, Sekarski N, Calame A. (1996):** Hemodynamic changes during high frequency oscillation for respiratory distress syndrome. Arch Dis Child Fetal Neonatal Ed. May; 74(3):F172-6.
- Lemyre B, Davis PG, de Paoli AG. (2002):** Nasal intermittent positive pressure ventilation (NIPPV) versus nasal continuous positive

airway pressure (NCPAP) for apnea of prematurity. Cochrane Database Syst Rev. ;(1):CD002272.

Lewis V, Whitelaw A. (2002): Furosemide for transient tachypnea of the newborn. Cochrane Database Syst Rev. ;(1):CD003064.

Lin HC, Warg S, Hsy Y, Yeh T and Chang F. (2001): Risk for respiratory distress syndrome in preterm infants born to mothers complicated by placenta previa. Early Hum Dev; 60(3):215-24.

Lumley J. (2003): Method of delivery for the preterm infant. BJOG. Apr.; 110 Suppl 20:88-92.

Man DW, Hamdy MH, Hendry GM, Bisset WH, Forfar JO. (1983): Congenital lobar emphysema: problems in diagnosis and management. Arch Dis Child. Sep; 58(9):709-12.

Marino BS, Bird GL, Wernovsky G. (2001): Diagnosis and management of the newborn with suspected congenital heart disease. Clin Perinatol.; 28:91–136.

Mangieri P, Suzuki K, Ferreira M, Domingues L, Casulari LA (2003): Evaluation of pituitary and thyroid hormones in patients with subarachnoid hemorrhage due to ruptured intracranial aneurysm. Ar Qneuro-Psiquiat 61:14–19.

Marttila R, Haataja R, Ramet M, Lofgren J. & Hallman M. (2003): Surfactant protein B polymorphism and respiratory distress syndrome in premature twins. Hum Genet; 112:18-23.

Mazela J, Merritt TA, Finer NN.(2007): Aerosolized surfactants. Curr Opin Pediatr. Apr; 19(2):155-62.

Mendeloff EN.(2004): Sequestrations, congenital cystic adenomatoid malformations and congenital lobar emphysema. Semin Thorac Cardiovasc Surg.; 16:204–214.

- Mlay GS and Mauji KP. (2000):** Respiratory distress syndrome among neonates admitted at Matimbili Medical Center, Dar EL Salaam, Tanzania. *J. Trop. Pediatr*; Oct.46 (6):303-7.
- Murphy N, Hume R, van Toor H, Matthews TG, Ogston SA, Wu SY, Visser TJ, Williams FLR (2004):** The hypothalamic-pituitary-thyroid axis In preterm infants; responsiveness to birth over the first 24 hours of life. *J Clin Endocrinol Metab* 89:2824–2831.
- Milner AD. (1996):** Surfactant and respiratory distress syndrome. *Turk J Pediatr*. Jan-Mar; 38(1):37-43.
- Moen MD, Perry CM, Wellington K. (2005):** Lucinactant: in neonatal respiratory distress syndrome. *Treat Respir Med.* ; 4(2):139-45.
- Moriette G, Paris- Llado J, Walti H, Escande B, et al., (2001):** Prospective randomized multicenter comparison of high - frequency oscillatory ventilation and conventional ventilation in preterm infants of less than 30 weeks with respiratory distress syndrome. *Pediatrics*; 107(2): 363-372.
- Moutquin JM. (2003):** Classification and heterogeneity of preterm birth. *BJOG* 110 Suppl 20:30–33.
- Moss TJ. (2006):** Respiratory consequences of preterm birth. *Clin Exp Pharmacol Physiol*. Mar; 33(3):280-4.
- Moya FR, Montes HF, Thomas VL, Mouzinho AM, Smith JF, Rosenfeld CR. (2000):** Surfactant protein A and saturated phosphatidylcholine in respiratory distress syndrome. *Am J Respir Crit Care Med*. Dec; 150(6 Pt 1):1672-7.

- Mukhopadhyay K, Kumar P, Narang A. (1998):** Role of early postnatal dexamethasone in respiratory distress syndrome. *Indian Pediatr.* Feb; 35(2):117-22.
- Neil M, Peter H, Rosalind S. (1986):** Forfar & Arneil's textbook of pediatrics (6th Ed.).
- O'Broovich HM. (1996):** Immature epithelial Na⁺ channel expression is one of the pathogenetic mechanisms leading to human neonatal respiratory distress syndrome. *Proc Assoc Am Physicians.* Sep; 108(5):345-55.
- O'Grady CM, Franklin RC.(1984):** Thyroid hormone concentrations and free thyroxin status in neonatal venous and capillary serum pairs. *Ann Clin Biochem* 1984; 21:306-9.
- Ohlsson A, Aher SM.(2006):** Early erythropoietin for preventing red blood cell transfusion in preterm and/or low birth weight infants. *Cochrane Database Syst Rev.* Jul 19;3:CD004863.
- Onrust SV, Dooley M, Goa KL. (1999):** Calfactant: a review of its use in neonatal respiratory distress syndrome. *Paediatr Drugs.* Jul-Sep; 1(3):219-43.
- Osborn DA, Hunt RW. (2007):** Postnatal thyroid hormones for respiratory distress syndrome in preterm infants. *Cochrane Database Syst Rev.* Jan 24 ;(1):CD005946.
- Parvin CA, Kaplan LA, Chapman JF, McManamon TG, Gronowski AM. (2005):** Predicting respiratory distress syndrome using gestational age and fetal lung maturity by fluorescent polarization. *Am J Obstet Gynecol.* Jan; 192(1):199-207.
- Pfister RH, Soll RF, Wiswell T. (2007):** Protein containing synthetic surfactant versus animal derived surfactant extract for the

- prevention and treatment of respiratory distress syndrome. Cochrane Database Syst Rev. Oct 17 ;(4):CD006069.
- Paul DA, Leef KH, Stefano JL, Bartoshesky L (2000):** Thyroid function in very-low-birth-weight infants with intraventricular hemorrhage. Clin Pediatr 39:651–656.
- Ranjit MS. (2000):** Cardiac abnormalities in birth asphyxia. Indian J Pediatr.Mar; 67(3 Suppl):S26-9.
- Rennie, J.M. and Robertson, N.R.C. (2002):** Hypoxic Ischemic Encephalopathy. A Manual of neonatal Intensive Care, 4th edition, Arnold Publishers; pp. 315 – 320.
- Redding RA, Pereira C. (2003):** Thyroid function in respiratory distress syndrome (RDS) of the newborn. Pediatrics ; 54:423-8.
- Robertson NJ, McCarthy LS, Hamilton PA, Moss AL. (2000):** Nasal deformities resulting from flow driver continuous positive airway pressure. Arch Dis Child Fetal Neonatal Eds. Nov; 75(3):F209-12.
- Rodriguez RJ, Martin RJ, Fanaroff AA. (2003):** The respiratory distress syndrome and its management. In: Fanaroff AA, Martin RJ, editors. Neonatal-perinatal medicine, 7th ed. St Louis: Mosby-Year Book.
- Rudolph, N. (1998):**Low birth weigh: In Finberg, L. ed. Saunders manual of Pediatric Practice (1st ed) WB Saunders Philadelphia, PP: 64-69.
- Rodriguez RJ, Martin RJ. (2003):** Exogenous surfactant therapy in newborns. Respir Care Clin N Am. Dec; 5(4):595-616.
- Rodriguez RJ. (2003):** Management of respiratory distress syndrome: an update. Respir Care. Mar; 48(3):279-86; discussion 286-7.

- Sanders M, Allen M, Alexander, G.R. (2002):** Gestational age assessment in preterm neonates weighting less than 1500 grams. *Pediatrics.*, 88: 542-546.
- Sasidharan P. (2004):** An approach to diagnosis and management of cyanosis and tachypnea in term infants. *Pediatr Clin North Am.*; 54:999–1021.
- Scanlon MF, Toft AD. (2000):** Regulation of thyrotrophin secretion. In: Braverman LE, Utiger RD, eds. *Werner's and Ingbar's the thyroid*. Philadelphia: Lippincott-Raven; 234–253.
- Schmidt BK. (1994):** Antithrombin III deficiency in neonatal respiratory distress syndrome. *Blood Coagul Fibrinolysis*. Jan; 5 Suppl 1:S13-7; discussion S59-64.
- Schwartz RM, Luby AM, Scanlon JW, Kellogg RJ.(1994):** Effect of surfactant on morbidity, mortality, and resource use in newborn infants weighing 500 to 1500 g. *N Engl J Med*. May 26;330(21):1476-80.
- Schrag S, Gorwitz R, Fultz-Butts K, Schuchat A. (2002).** Prevention of perinatal group B streptococcal disease. Revised guidelines from CDC. *MMWR Recomm Rep*. Aug 16; 51(RR-11):1-22.
- Shah VA, Yeo CL, Ling YL, Ho LY. (2003):** Incidence, risk factors of retinopathy of prematurity among very low birth weight infants in Singapore. *Ann Acad Med Singapore*; 34:169–178.
- Shiao SY, Ou CN. (2007):** Validation of oxygen saturation monitoring in neonates. *Am J Crit Care*. Mar; 16(2):168-78.

Shinwell ES, Karplus M, Reich D, Weintraub Z, Blazer S, Bader D, Yurman S, Dolphin T, Kogan A, Dollberg S, Arbel E, Goldberg M, Gur I, Naor N, Sirota L, Mogilner S, Zaritsky A, Barak M, Gottfried E (2000):

Early postnatal dexamethasone treatment and increased incidence of cerebral palsy. Arch Dis Child 83:F177–F181.

Silverman WC, Anderson DH. (1956): Controlled clinical trial on effects of water mist on obstructive respiratory signs, death rate and necropsy findings among premature infants. Pediatrics ; 17: 1-4.

Simmons F; (2000): Goldstein, clinical trial of lactulose in hepatic encephalopathy. Gastro enterology; 59: 827-832.

Simpson J, Williams FLR, Delahunty C, Ogston SA, van Toor H, Wu SY, Visser TJ, Hume R,(2005): With collaboration from the Scottish Preterm Thyroid Group 2005 Serum thyroid hormones in preterm infants and relationships to indices of severity of intercurrent illness. J Clin Endocrinol Metab 90:1271–1279.

Sims EJ, Vermillion ST, Soper DE. (2002): Preterm premature rupture of the membranes is associated with a reduction in neonatal respiratory distress syndrome. Am J Obstet Gynecol.Aug; 187(2):268-72.

Sinha S, Moya F, Donn SM. (2007): Surfactant for respiratory distress syndrome: are there important clinical differences among preparations? Curr Opin Pediatr. Apr; 19(2):150-4.

Sinha SK, Donn SM. (2007): Volume-controlled ventilation. Variations on a theme. Clin Perinatol. Sep; 28(3):547-60.

So BH, Tamura M, Mishina J, Watanabe T, Kamoshita S. (1995): Application of nasal continuous positive airway pressure to

early extubation in very low birth weight infants. Arch Dis Child Fetal Neonatal Ed. May; 72(3):F191-3.

Soll RF, Blanco F. (2001): Natural surfactant extract versus synthetic surfactant for neonatal respiratory distress syndrome. Cochrane Database Syst Rev. ;(2):CD000144.

Soll RF, Morley CJ. (2001): Prophylactic versus selective use of surfactant in preventing morbidity and mortality in preterm infants. Cochrane Database Syst Rev. ;(2):CD000510.

Soll RF, Morley CJ. (2003): Prophylactic versus selective use of surfactant in preventing morbidity and mortality in preterm infants. Cochrane Database Syst Rev ;(4):CD000510.

Spaulding SW (2000): Biological actions of thyrotrophin. In: Braverman LE, Utiger RD, eds. Werner's and Ingbar's the thyroid. Philadelphia: Lippincott-Raven; 227–233. Seri I 1995 Cardiovascular, renal and endocrine actions of dopamine in neonates and children. J Pediatr 126:333–344.

Stephen Nussey and Saffron Whitehead (2001): The thyroid gland in Endocrinology: An Integrated Approach Published by BIOS Scientific Publishers Ltd. ISBN 1-85996-252-1.

Stoll, B.J. and Ira-Adams-Chapman. (2007): The fetus and the Neonatal infant. Noninfectious Disorders. Hypoxia-Ischemic. In Nelson Textbook of Pediatrics, Behrman R.E., Kleigman R.M. and Jenson H.B. (eds.) 18th ed., W.B. Saunders Co., London; pp. 566- 568.

Stutchfield P, Whitaker R, Russell I. (2005): Antenatal Steroids for Term Elective Caesarean Section (ASTECS) Research Team. Antenatal betamethasone and incidence of neonatal respiratory

distress after elective caesarean section: pragmatic randomised trial. *BMJ*. Sep 24; 331(7518):662. Epub Aug 22.

Subramanian P, Henderson-Smart DJ, Davis PG. (2000): Prophylactic nasal continuous positive airways pressure for preventing morbidity and mortality in very preterm infants. *Cochrane Database Syst Rev.* ;(2):CD001243.

Suresh GK, Soll RF. (2002): Lung surfactants for neonatal respiratory distress syndrome: animal-derived or synthetic agents? *Paediatr Drugs.* ; 4(8):485-92.

Sweet D, Bevilacqua G, Carnielli V, Greisen G, Plavka R, Didrik Saugstad O , Simeoni U , Speer CP , Valls-I-Soler A , Halliday H, (2007): Working Group on Prematurity of the World Association of Perinatal Medicine, European Association of Perinatal Medicine. European consensus guidelines on the management of neonatal respiratory distress syndrome. *J Perinat Med.*; 35(3):175-86.

Sweet DG, Halliday HL. (1999): Current perspectives on the drug treatment of neonatal respiratory distress syndrome. *Paediatr Drugs.* Jan- Mar; 1(1):19-30.

Synder JM, Mendelson CR. (2001): Insulin inhibits the accumulation of the major lung surfactant apoprotein in human fetal lung explants maintained in vitro. *Endocrinology.* Apr; 120(4):1250-7.

Tang W, Ridout D, Modi N. (1997): Influence of respiratory distress syndrome on body composition after preterm birth. *Arch Dis Child Fetal Neonatal Ed.* Jul; 77(1):F28-31.

- Teeratakulpisarn J, Taksaphan S, Pengsaa K, Wiangnon S, Kosuwon W. (2000):** Prediction of idiopathic respiratory distress syndrome by the stable micro bubble test on gastric aspirate. *Pediatr Pulmonol.* Jun; 25(6):383-9.
- Todd DA, Wright A, Smith J, NICUS Group. (2007):** Severe retinopathy of prematurity in infants _30 weeks' gestation in New South Wales and the Australian Capital Territory from 1992 to 2002. *Arch Dis Child Fetal Neonatal Ed.*; 92:F251–F254.
- Uhrmann S, Marks KH, Maisels MJ, et al. (2002):** Thyroid function in the preterm infant: a longitudinal assessment. *J Pediatr* ; 92:968-73.
- Vakrilova L, Kalaidzhieva M, Slŭncheva B, et al.(2004):** The significance of parameters of the acid-base status, blood gas and blood lactate level for the adequate resuscitation and prognosis in newborns with very low birth weight ; 43 (5):25-31.
- Van Wassenae AG, Kok JH, Brie't JM, Pijning AM, de VijlderJJM. (2004):** Thyroid function in very preterm infants, possible implications. *Thyroid.* ; 9:85–91.
- Van Marer LJ, Allced EN, Leviton A, Pagano M, Rarad R and Moore M (2001):** Antenatal glucocorticoid treatment does not reduce chronic lung Disease among surviving preterm infants. *Journal of Pediatrics*, 138(2): 198-204.
- Van Kaam AH, Rimensberger PC. (2007):** Lung-protective ventilation strategies in neonatology: what do we know--what do we need to know? *Crit Care Med.* Mar; 35(3):925-31.
- Van Marter LJ, Allred EN, Pagano M, Sanocka U, Parad R, Moore M, Susser M, Paneth N, Leviton A.(2000):** Do clinical markers of barotraumas and oxygen toxicity explain interhospital variation in rates of chronic lung disease? The Neonatology Committee

for the Developmental Network. *Pediatrics*. Jun; 105(6):1194-201.

Van Meurs K, Hintz S, Rhine W, Benitz W. (2006): The use of inhaled nitric oxide in the premature infant with respiratory distress syndrome. *Minerva Pediatr*. Oct; 58(5):403-22.

Vain NE, Szyld EG, Prudent LM, Wiswell TE, Augilar AM, Vivas N,(2004) : for the Meconium Study Network . Oropharyngeal and Nasopharyngeal suctioning of meconium-stained neonates before delivery of their shoulders: multicentre randomized controlled trial. *Lancet*. ; 364:597–602.

Verder H. (2007): Nasal CPAP has become an indispensable part of the primary treatment of newborns with respiratory distress syndrome. *Acta Paediatr*. Apr; 96(4):482-4.

Verma RP. (2000): Respiratory distress syndrome of the newborn infant. *Obstet Gynecol Surv*. Jul; 50(7):542-55.

Volpe, J.J. (2001a): Hypoxic-ischemic encephalopathy intrauterine assessment. In: *Neurology of the Newborn*. Volpe J.J. (ed.) 8th edition, WB Saunders, Philadelphia, pp. 227 –295.

Volpe J.J. (2001b): Hypoxic- ischemic encephalopathy: Clinical Aspects. In *Neurology of the Newborn*. Volpe J.J. (Ed.). 8th edition, WB Saunders, Philadelphia; pp. 331- 394.

Vulsma T, Kok JH. (1996): Prematurity associated neurodevelopmental abnormalities and neonatal thyroid hormone levels. *N Engl J Med*.: 334:857–858.

Walther FJ, Benders MJ, Leighton JO. (1993): Early changes in the neonatal circulatory transition. *J Pediatr*. Oct; 123(4):625-32.

- Weaver TE & Conkright JJ. (2001):** Function of surfactant proteins B and C. *Annu Rev Physiol* 63: 555-578.
- Walfisch, A., Hallak, M. & Mazor, and M. (2001):** Multiple courses of antenatal steroids: risks and benefits. *Obstet Gynecol* 98, 491-7.
- Wiswell TE, Knight GR, Finer NN, et al. (2002):** A multicenter, randomized, controlled trial comparing Surfaxin (lucinactant) lavage with standard care for treatment of meconium aspiration syndrome. *Pediatrics*; 109:1081–1087.
- Wiberg N, Källén K and Olofsson P.(2006):** Base deficit estimation in umbilical cord blood is influenced by gestational age, choice of fetal fluid compartment, and algorithm for calculation. *Am J Obstet Gynecol*. Dec; 195(6):1651-6.
- Wijnberger LD, Huisjes AJ, Voorbij HA, Franx A, Bruinse HW, Mol BW. (2001):** The accuracy of lamellar body count and lecithin/sphingomyelin ratio in the prediction of neonatal respiratory distress syndrome: a meta-analysis. *BJOG*. Jun; 108(6):583-8.
- Wiswell TE, Mendiola J. (1993):** Respiratory distress syndrome in the newborn: innovative therapies. *Am FAM Physician*. Feb 1; 47(2):407-14.
- Wiswell TE, Gannon CM, Jacob J, et al. (2000):** Delivery room management of the apparently vigorous meconium-stained neonate: results of the multicenter, international collaborative trial. *Pediatrics*. ; 105:1–7.
- Williams FLR, Simpson J, Delahunty C, Ogston S, Bongers C, van Toor H, Wu SY, Visser TJ, Hume R.(2005):** With collaboration from the Scottish Preterm Thyroid Group Developmental trends in

cord and postpartum serum thyroid hormones in preterm infants.
J Clin Endocrinol Metab 89:5314–5320.

Williams FLR, Mires GJ, Barnett C, Ogston SA, van Toor H, Visser TJ, Hume R (2005): Transient hypothyroxinemia in preterm infants: the role of cord sera thyroid hormone levels adjusted for prenatal and intrapartum factors. J Clin Endocrinol Metab 90:4599–4606.

Wood NS, Marlow N, Costeloe K, Gibson AT, Wilkinson AR, (2000): Neurologic and developmental disability after extremely preterm birth. N Engl J Med 343:378–384.

Wood DW, Downes' JJ, Locks HI. (2000): A clinical score for the diagnosis of respiratory failure. Amer J Dis Child; 123: 227-9.

Yu VY. (1998): Recent advances in assisted ventilation for neonatal respiratory distress syndrome. Indian Pediatr. Jul; 35(7):631-40.

Yeh TF, Lin YJ, Lin HC, Huang CC, Hsieh WS, Lin CH, Tsai CH, (2004): Outcomes at school age after postnatal dexamethasone therapy for Lung disease of prematurity. N Engl J Med 350:1304–1313.

Zimmermann LJ, Janssen DJ, Tibboel D, Hamvas A, Carnielli VP. (2005): Surfactant metabolism in the neonate. Biol Neonate; 87(4):296-307.