References

References

- **Abdalla H, Burton G, Kirkland A et al (1993)** Age, pregnancy and miscarriage: uterine versus ovarian factors. Hum Reprod 8:1512.
- **Abdalla H & Thum M (2004)** An elevated basal FSH reflects a quantitative rather than qualitative decline of the ovarian reserve. Hum Reprod 19, 893–898.
- **Adashi EY (1994)** The climacteric ovary as a functional gonadotropin-driven androgen-producing gland. Fertil Steril 62, 20–27.
- **Ahmed Ebbiary NA, Lenton, E A, Salt C et al (1994)** The significance of elevated basal follicle stimulating hormone in regularly menstruating infertile women. Hum Reprod 9, 245-252.
- Alexander SE, Aksel S, Hazelton JM, Yeoman RR& Gilmore SM(1990) The effect of aging on hypothalamic function in oopherctomized women. Am J Obstet Gynecol 162:446.
- Ammini AC, Pandey J, Vijyaraghavan M& Sabherwal U (1994) Human female phenotypic development: role of fetal ovaries, J Clin Endocrinol Metab 19:604.
- Baarends WM, Uilenbroek JT, Kramer P, Hoogerbrugge JW, van Leeuwen EC, et al (1995) Anti-Müllerian hormone and anti-Müllerian hormone type II receptor messenger ribonucleic acid expression in rat ovaries during postnatal development, the estrous cycle, and gonadotropin-induced follicle growth. Endocrinology, 136, 4951–4962.
- Baka S, Makrakis E, Tzanakaki D, Konidaris S, Hassiakos D, Moustakarias T, Creatsas G. (2006) Poor responders in IVF: cancellation of a first cycle is not predictive of a subsequent failure. Ann N Y Acad Sci., 1092:418-25.
- **Baker TG** (1963) A quantitative and cytological study of germ cells in human ovaries. Prot R Soc. Lond. Biol. J 158,417-433.
- **Balasch J, Creus M, Fábregues F et al (1996)** Inhibin, follicle-stimulating honnone and age as predictors of ovarian response in in vitro fertilization cycles stimulated with gonadotropin-releasing hormone agonist-gonadotropin treatment. Am J Obstet Gynecol 175, 1226-1230.
- Bancsi LFJM, Broekmans FJM & Te Velde ER (1997) Predictive value of serum inhibin B for ART outcome? Fertil Steril, 68, 947-948

- Bancsi LFJM, Huijs AV, Den Oude CT et al (2000) Basal follicle -stimulating hormone levels are of limited value in predicting ongoing pregnancy rate after in vitro fertilization Fertil Steril 73, 552-551.
- Bancsi LFJM, Broekmans FJM, Mol BWJ, Habbema JDF & te Velde ER (2003) Performance of basal follicle-stimulating hormone in the prediction of poor ovarian response and failure to become pregnant after in vitro fertilization: a meta-analysis. Fertil Steril 79,1091–1100.
- Bancsi LF, Brockmans FJ, Eijkemans MJ, de Jong FH, Habbema JD& te Velde ER (2002) predictors of poor ovarian response in in-vitro fertilization: a prospective study comparing basal markers of ovarian reserve, Fertil Steril 77,328–336.
- Barroso G, Oehninger S, Monzó A, Kolm P, Gibbons WE, Muasher SJ (2001) High FSH:LH ratio and low LH levels in basal cycle day 3: impact on follicular development and IVF outcome. J Assist Reprod Genet. 18:499-505.
- Bassil S, Wyns C, Toussaint-Demylle D, Nisolle M, Gordts S and Donnez J (1997) The relationship between ovarian vascularity and the duration of stimulation in in-vitro fertilization. Hum Reprod 12, 1240–1245.
- Bath LE, Wallace WH, Shaw MP, Fitzpatrick C & Anderson RA (2003) Depletion of ovarian reserve in young women after treatment for cancer in childhood: detection by anti-Mullerian hormone, inhibin B and ovarian ultrasound. Hum Reprod 18, 2368–2374.
- Battaglia DE, Goodrvin P, Krein NA& Soures MR (1996), Influence of maternal age on meiotic spindle assembly in oocytes from naturally cycling women, Hum Reprod 11:2217.
- Behre HM, Greb RR, Mempel A, Sonntag B et al (2005) Significance of a common single nucleotide polymorphism in exon 10 of the follicle stimulating hormone (FSH) receptor gene for the ovarian response to FSH: a pharmacogenetic approach to controlled ovarian hyperstimulation. Pharmacogenet Genomics 15,451–456.
- Behringer RR, Finegold MJ & Cate RL (1994) Müllerian inhibiting substance function during mammalian sexual development. Cell, 79, 415–425.
- **Block E (1952)** Quantitative morphological investigations of the follicular system in women, Variations at different ages. Acta Anat 14, 108-123.

- **Bowen S, Norian J, Santoro N, Pal L. (2007)** Simple tools for assessment of ovarian reserve (OR): individual ovarian dimensions are reliable predictors of OR. Fertil Steril,88:390-5
- Bridges NA, Cooke A, Healy MJ, Hindmarsh PC& Brook CG(1993) Standards for ovarian volume in childhood and puberty, Fertil Steril 60:456.
- Brodowska A, Starczewski A, Laszczyńska M, Szydlowski A (2005) Ovarian androgenesis in women after menopause. Pol Merkuriusz Lek,; 19: 90-93.
- Brodowska A, Laszczyńska M, Starczewski A (2007) Apoptosis in ovarian cells in postmenopausal women Folia Histochemica Et Cytobiologica., 45, 99-105
- Broekmans FJ, Kwee J ,Hendriks DJ ,Mol BW & Lambalk CB (2006)A systematic review of tests predicting ovarian reserve and IVF outcome Hum Reprod Update, 12, 685–718,
- Bukovsky A, Caudle MR, Svetlikova M, Upadhyaya NB (2004) Origin of germ cells and formation of new primary follicles in adult human ovaries. Reprod Biol Endocrinol; 28:1–20.
- **Bukulmez O, Arici A (2004)** Assessment of ovarian reserve. Curr Opin Obstet Gynecol; 16:231–7.
- Burger HG, Famada Y, Bangah ML, McCloud PI& Warne GL(1991), Serum gonadotropin. sex steroid, and immunoreactive inhibin levels in the first two years of life, J Clin Endocrinol Metab 72:682.
- Burger HG, Dudrey EC. Iropper IL, Groome N, Guthrie JR, Green A& Dennerstein L (1999) Prospectively measured levels of serum follicle stimulating hormone, estradiol and the dimeric inhibins during the menopausal transition in a population-based cohort of women, J Clin Endocrinol Metab 84:4025.
- **Buyalos RP, Daneshmand S & Brzechffa PR** (1997) Basal estradiol and folliclestimulating hormone predict fecundity in women of advanced reproductive age undergoing ovulation induction therapy Fertil Steril 68,272-277.
- **Cahill DJ, Prosser CJ, Wardle PG et al (1994)** Relative influence of serum follicle stimulating hormone, age and other factors on ovarian response to gonadotrophin stimulation Br. J Obstet Gynaecol 101,999-1007.
- Cate RL, Mattaliano RJ, Hession C, Tizard R et al (1986) Isolation of the bovine and human genes for Müllerian inhibiting substance and expression of the human gene in animal cells. Cell, 45, 685–698.

- **Chang MY, Chiang CH, Hsieh TT et al (1998)** Use of the antral follicle count to predict the outcome of assisted reproductive technologies. Fertil. Steril. 69,505-510.
- Centers for Disease Control and prevention, American Society for Reproductive medicine, Society for Assisted reproductive Technology, RESOLVE(2003), 2001 assisted reproductive technology success rates. Centers for Disease Control and Prevention. Atlanta, GA,.
- Clement PB (1991) Ovary. In Sternberg SS (ed.), Histology for Pathologists. Raven Press, New York, pp. 765-795.
- **Cohen HL, Eisenberg P, Mandel E& Haller JO(1992)**, Ovarian cysts are common in premenarchal girls: a sonographic study of 101 children 2-12 years old, Am J Radiol 159:89.
- Cohen HL, Shapiro MA, Mandel FS& Shapiro ML (1993), Normal ovaries in neonates and infants: a sonographic study of 77 patients 1 day to 24 months old, Am J Roentgenol 160:583.
- Cohen-Haguenauer O, Picard Mattei JY & Mattei MGet al (1987) Mapping of the gene for anti-Müllerian hormone to the short arm of human chromosome 19. Cytogenetics and Cell Genetics, 44, 2–6.
- **Cook CL, Siow Y, Taylor S& Fallat ME** (2000) Serum müllerian inhibiting substance levels during normal menstrual cycles. Fertility and Sterility, 73, 859–861.
- Cook CL, Siow Y, Brenner, AG & Fallat ME (2002) Relationship between serum müllerian-inhibiting substance and other reproductive hormones in untreated women with polycystic ovary syndrome and normal women. Fertil Steril, 77, 141–146.
- Corson SL, Gutman J, Batzet FR et al (1999) Inhibin-B as a test of ovarian reserve for infertile women. Hum Reprod 14, 2818-2821.
- **Cranrer DW, Xu H& Harlo BL(1995)**: Family history as a predictor of early menopause, Fertil Sreril 61:740,.
- Danforth DR, Arbogast LK, Mroueh J et al (1998) Dimeric inhibin: a direct marker of ovarian aging Fertil Steril. 70, 119-123.
- de Bruin JP, Bovenhuis H, van Noord PA, Pearson pL, van Arendonk JA, et al, (2001) The role of genetic factors in age at natural menopause, Hum Reprod 16:2014

- **de Koning CH, Popp-Snijders C, Schoemaker J, Lambalk CB(2000).** Elevated FSH concentrations in imminent ovarian failure are associated with higher FSH and LH pulse amplitude and response to GnRH. Hum Reprod;15:1452–6.
- de Koning CH, Benjamins T, Harms P, Homburg R, van Montfrans JM, Gromoll J, Simoni M and Lambalk CB (2006) The distribution of FSH receptor isoforms is related to basal FSH levels in subfertile women with normal menstrual cycles. Hum Reprod 21,443–446.
- de Vet A, Loven JS, de Jong FH, Themmen AP& Fauser BC (2002) Anti-Müllerian hormone serum levels: a putative marker for ovarian aging. Fertil Steril, 77, 357–362.
- **Deborah S. Waches, Mickey S. Coffler, Pamela J. Malcom and R& Jeffrey Chang (2006)** Serum anti-mullerain hormone concentrations are not altered by acute administration of follicle stimulating hormone in polycystic syndrome and normal women, J Clin Endocrinol metab 192, 1871-1874.
- den Tonkelaar I, te Veld ER& Looman CWN(1998) Menstrual cycle length preceding menopause in relation to age at menopause. Maturiras 29:115.
- **Dennefors BL, Janson PO, Hamberger L and Knutsson F (1982)** Hilus cells from human postmenopausal ovaries: gonadotrophin sensitivity, steroid and cyclic AMP production. Acta Obstet Gynecol Scand 61,413–416.
- di Clemente N, Goxe B, Rémy JJ, Cate RL, Josso N, Vigier B & Salesse R (1994) Inhibitory effect of AMH upon aromatase activity and LH receptors of granulosa cells of rat and porcine immature ovaries. Endocrine, 2, 553–558.
- **Donnez J& Jadoul B(2002)** What are the implications of myomas on fertility? A need for a debate? Hum Reprod 17:1424.
- **Dowsett M, Cantwell B, Lal A, Jeffcoate SL and Harris AL (1988)** Suppression of postmenopausal ovarian steroidogenesis with the luteinizing hormone-releasing hormone agonist goserelin. J Clin Endocrinol Metab 66,672–677.
- **Durlinger AL, Kramer P, Karels B, de Jong FH, Uilenbroek JT, Grootegoed JA** & Themmen AP (1999) Control of primordial follicle recruitment by anti-Müllerian hormone in the mouse ovary. Endocrinology, 140, 5789–5796.
- **Durlinger AL, Gruijters MJ, Kramer P et al (2001)** Anti-Müllerian hormone attenuates the effects of FSH on follicle development in the mouse ovary. Endocrinology, 142, 4891–4899.

- **Durlinger AL, Visser JA & Themmen AP (2002a)** Regulation of ovarian function: the role of anti-Mullerian hormone. Reproduction 124, 601–609.
- **Durlinger AL, Gruijters MJ, Kramer P, Karels B, et al (2002b)** Anti-Müllerian hormone inhibits initiation of primordial follicle growth in the mouse ovary. Endocrinology, 143, 1076–1084.
- **Ebbiary N, Lenton E and Cooke I.** (1994) Hypothalamic–pituitary ageing: progressive increase in FSH and LH concentrations throughout the reproductive life in regularly menstruating women. Clin. Endocrinol., 41, 199–206.
- Ebner T,Sommergruber M, Moser M, Shebl O, Schreier -Lechner E. and Tews G (2006)Basal level of anti-Müllerian hormone is associated with oocyte quality in stimulated cycles Human Reproduction Vol.21, No.8 pp. 2022–2026,
- **Ebrahim A, Rienhardt G, Monis S et al (1993)** Follicle-stimulating hormone levels on cycle day 3 predict ovulation stimulation response Assist. Reprod. Genet. 10, 130-136.
- Eldar-Geva, T, Ben-Chetrit A, Spitz, IM, Rabinowitz R, Markowitz, E, Mimoni, T, Gal M, Zylber-Haran E & Margalioth EJ (2005) Dynamic assays of inhibin B, anti-Müllerian hormone and estradiol following FSH stimulation and ovarian ultrasonography as predictors of IVF outcome. Hum Reprod, 20, 3178–3183.
- El Toukhy T, Khalaf Y, Hart R, Talor A, Baude P(2002) Young age does not protect against the adverse effects of reduced ovarian reserve—an eight year study. Hum Reprod;17:1519–24.
- Engmann L, Sladkevicius P, Agrawal R, Bekir JS, Campbell S and Tan SL (1999) Value of ovarian stromal blood flow velocity measurement after pituitary suppression in the prediction of ovarian responsiveness and outcome of in vitro fertilization treatment. Fertil Steril 71, 22–29.
- **Faber BM,Mercan R, Hamasher P, Musher SJ, Toner JP(1997)** The impact of an egg donor's age and her prior fertility on reciepient pregnancy outcome, Fertil Steril, 68:370
- **Fåbregues F, Balasch J, Creus M et al. (2000)** Ovarian reserve test with human menopausal gonadotropin as a predictor of in vitro fertilization outcome. J. Assist Reprod. Genet. 17, 13-19.
- Faddy MJ, Gosden RG, Gougeon A, Richardson SJ& Nelson JF (1992) Accelerated disappearance of ovarian follicles in midlife: implications for forecasting menopause, Hum Reprod 7:1342-1346

- **Fanchin R, de Ziegier D, Olivennes F et al (1994)** Exogenous follicle stimulating hormone ovarian reserve test (EFORT): a simple and reliable screening test for detecting poor responders' in in-vitro fertilization Hum Repro., 9, 1607-1611.
- Fanchin R, Schonauer LM, Righini C, Frydman N, Frydman R & Taieb J (2003a) Serum anti-Müllerian hormone dynamics during controlled ovarian hyperstimulation. Hum Reprod, 18, 328–332.
- Fanchin R, Schonauer LM, Righini C, Guibourdenche J, Frydman R & Taieb J (2003b) Serum anti-Müllerian hormone is more strongly related to ovarian follicular status than serum inhibin B, estradiol, FSH and LH on day 3. Hum Reprod, 18, 323–327.
- Fanchin R, Lozano MDH, Frydman N, Gougeon A, di Clemente N, Frydman R, and Taieb J (2007) Anti-Müllerian hormone concentrations in the follicular fluid of the preovulatory follicle are predictive of the implantation potential of the ensuing embryo obtained by in vitro fertilization. JCEM 92, 1796–1802
- **Farimani M, Amiri I, Hoseini S(2006)** Day 3 serum inhibin-B level is not predictive of ovarian assisted reproductive technologies outcome; Iranian Journal of Reproductive Medicine 4, 1-5.
- **Fiçicioğlu C, Kutlu T, Baglam E, and Bakacak Z (2006)** Early follicular antimüllerian hormone as an indicator of ovarian reserve Fertil Steril 85, 592-596.
- Focchi GR, Simoes MJ, Baracat EC, de Lima GR. (1995) Morphological and morphometrical features of the corpus albicans in the course of the postmenopausal period. Bull Assoc Anat,; 79:15-18
- Frattarelli JL, Lauria-Costab DF, Miller BT, Dergh PA, Scott RT(2000) Basal antral follicle number and mean ovarian diameter predict cycle cancellation and ovarian responsiveness in assisted reproductive technology cycles, Fertil Sreril 74:512-517.
- **Fujisawa M T, Yamasaki H, Okada& Kamidono S (2002)** The significance of anti-Mullerian hormone concentration in seminal plasma for spermatogenesis Hum Reprod, 17: 968 970.
- Galtier-Dereure F, Bouard V, de Picot MC et al (1996) Ovarian reserve test with the gonadotrophin-releasing hormone agonist busereline: a correlation with invitro fertilization outcome Hum Reprod, 11, 1393-1398.

- Gnoth C, Schuring AN, Friol K, Tigges J, Mallmann P and Godehardt E (2008)Relevance of anti-Mullerian hormone measurement in a routine IVF programHum Reprod 23:1359-1365.
- Gondos B, Bhiraleus P& Hobel C(1971), Ultrastructural observations on germ cells in human fetal ovaries, Am J Obstet Gynecol 110:644.
- Gondos B, Westcrgaard L& Byskov A (1986), Initiation of oogenesis in the human fetal ovary: ultrastructural and squash preparation study, AM J Obstet Gynecol 155:189.
- **Gosden RG (1985)** Maternal age: a major factor affecting the prospect and outcome of pregnancy, Ann N Y Acad Sci 442:45.
- **Gougeon A(1986)**, Dynamics of follicutar growth in the human: a model from preliminary results, Hum Reprod 1:81.
- **Gougeon A(1996)** Regulation of ovarian follicular development in primates: facts and hypotheses, Endocr Rev, 17:121.
- **Gougeon A, Echochard R& Thalabard JC(1994)**, Age-related changes of the population of human ovarian follicles: increase in the disappearance rate of non-growing and early-growing follicles in aging women, Biol Reprod 50:653.
- Greb RR, Grieshaber K, Gromoll J, Sonntag B, Nieschlag E, Kiesel L and Simoni M (2005) A common single nucleotide polymorphism in exon 10 of the human follicle stimulating hormone receptor is a major determinant of length and hormonal dynamics of the menstrual cycle. J Clin Endocrinol Metab 90,4866–4872.
- Gruijters MJ, Visser JA, Durlinger AL& Themmen AP (2003) Anti-Müllerian hormone and its role in ovarian function. Moll Cel Endocrinol, 211, 85–90.
- Guerriero S, Ajossa S, Lai MP, Risalvato A, Paoletti AM and Melis GB (1999) Clinical applications of colour Doppler energy imaging in the female reproductive tract and pregnancy. Hum Reprod Update 5, 515–529.
- Gülekli B, Bulbul Y, Onvural A et al. (1999) Accuracy of ovarian reserve tests. Hum Reprod, 14, 2822-2826.
- Haadsma ML, Bukman A, groen H, Roeloffzen EMA, Groenewoud ER et al(2007) The number of small antral follicles(2-6mm) determines the outcome of endocrine ovarian reserve tests in a subfertile population. Hum. Reprod. 22, 1925-1931

- Hall JE, Welt, CK & Cramer DW (1999) Inhibin A and inhibin B reflect ovarian function in assisted reproduction but are less useful at predicting outcome. Hum Reprod 14, 409-415.
- Hannoun A, Abu Musa A, Awwad J et al (1998) Clomiphene citrate challenge test: a cycle to cycle variability of cycle day 10 follicle stimulating hormone level. Clin Exp Obstet Gynecol 25, 155-156
- Hansen LM, Batzer FR, Gutman JN et al (1996) Evaluating ovarian reserve: follicle stimulating hormone and oestradiol vanability during cycle days 2-5. Hum Reprod, 11, 486-489.
- **Hassold T& Chiu D (1985)** Maternal age-specific rates of numerical chromosome abnormalities with special reference to trisomy, Hum Genet 70:11.
- Hazout A, Bouchard P, Seifer DB, Aussage P, Junca AM & Cohen-Bacrie P (2004) Serum antimüllerian hormone/müllerianinhibiting substance appears to be a more discriminatory marker of assisted reproductive technology outcome than follicle-stimulating hormone, inhibin B, or estradiol. Fertil Steril, 82, 1323–1329.
- Healy DL, Burger HG, Mamers P, Jobling T, Bangah M, Quinn M, Grant P, Day AJ, Rome R & Campbell JJ (1993) Elevated serum inhibin concentrations in postmenopausal women with ovarian tumors. New England Journal of Medicine, 329, 1539–1542
- Hehenkamp WJK, Looman CWN, Themmen APN, de Jong FH, te Velde ER and Broekmans FJM(2006) Anti-Müllerian hormone levels in the spontaneous menstrual cycle do not show substantial fluctuation. J Clin Endocrinol metab,91, 4057-4063
- Hendriks DJ, Broekmans FJ, Bancsi LF, Looman CW, de Jong FH, te Velde ER. (2005a) Single and repeated GnRH agonist stimulation tests compared with basal markers of ovarian reserve in the prediction of outcome in IVF. J Assist Reprod Genet; 22:65–73.
- Hendriks DJ, Mol BW, Bancsi LF, te Velde ER and Broekmans FJ (2005b) Antral follicle count in the prediction of poor ovarian response and pregnancy after in vitro fertilization: a meta-analysis and comparison with basal follicle-stimulating hormone level. Fertil Steril 83,291–301.
- **Hengster P& Menardi G (1992)**, Ovarian cysts in the newborn, Pediatr Surg Int 7:372.

- Hershlag A, Lesser M, Montefusco D et al (1992) Interinstitutional variability of follicle-stimulating hormone and estradiol level Fertil Steril. 58. 1123-1126.
- Himelstien-Braw R, Byskov AC, Peters H& Faber M(1976), Follicular atresia in the infant human ovary, J Reprod Fertil 46:55.
- Hirobe S, He WW, Gustafson ML, MacLaughlin DT & Donahoe PK (1994)

 Mullerian inhibiting substance gene expression in the cycling rat ovary correlates with recruited or graafian follicle selection. Biological Reproduction, 50, 1238–1243.
- Ho JY, Guu HF, Yi YC, Chen MJ, Ho ES(2005) . The serum follicle-stimulating hormone-to-luteinizing hormone ratio at the start of stimulation with gonadotropins after pituitary down-regulation is inversely correlated with a mature oocyte yield and can predict "low responders". Fertil Steril.;83:883-8.
- Hofmann GE, Sosnowski J, Scott RT et al (1996) Efficacy of selection criteria for ovarian reserve screening using the clomiphene citrate challenge test in a tertiary fertility center population. Fertil Steril 66, 49-53.
- **Hofmann GE, Danforth DR & Seifer DB** (1998) *Inhibin-B: the physiologic basis of the clomiphene citrate challenge test for ovarian reserve screening. Fertil Steril* 69, 474-477.
- Hudson PL, Dougas I, Donahoe PK, Cate RL, Epstein J, Pepinsky RB &MacLaughlin DT (1990) An immunoassay to detect human mullerian inhibiting substance in males and females during normal development. J Clin Endocrino Metab 70, 16–22.
- Hughes EG, Robertson DM, Handelsman DJ, Hayward S, Healy DL& de Kretser DM (1990) Inhibin and estradiol responses to ovarian hyperstimulation: effects of age and predictive value for in viro fertilization outcome, J Clin Endocrinol Metab 70:358.
- **Hughesdon PE (1982)** Morphology and morphogenesis of the Stein–Leventhal ovary and of so-called 'hyperthecosis'. Obstetrical and Gynecological Survey, 37, 59–77.
- Hull MG, Fleming CF, Hughes AO& McDermorr A (1996). The age-related decline in female fecundity: a quantitve controlled study of implanting capacity and survival of individual embryos after in vitro fertilization. Fertil Steril 65:783.

- **Hsieh YY, Chang CC and Tsai HD (2001)** Antral follicle counting in predicting the retrieved oocyte number after ovarian hyperstimulation. J Assist Reprod Genet 18,320–324.
- Imbeaud S, Carre-Eusebe D, Rey R, Belville C, Josso N & Picard JY (1994) Molecular genetics of the persistent müllerian duct syndrome: a study of 19 families. Human Molecular Genetics, 3, 125–131.
- **Imbeaud S, Faure E, Lamarre I et al (1995)** Insensitivity to anti-Müllerian hormone due to a spontaneous mutation in the human anti-Müllerian hormone receptor. Nature Genetics, 11, 382–388.
- Ingraham HA, Hirokawa Y, Roberts LM et al (2000) Autocrine and paracrine Müllerian inhibiting substance hormone signaling in reproduction. Recent Progress in Hormone Research, 55, 53–67.
- **Jacobs SL, Metzger DA, Dodson WC, Haney AE (1990)** Effect of age on response to human menopausal gonadotropin stimulation, J Clin Endocrinol Metab 11:1525.
- **Jain T, Soules MR, Collins JA(2004)** Comparison of basal follicle-stimulating hormone versus the clomiphene citrate challenge test for ovarian reserve screening. Fertil Steril;82:180-5
- Johnson J, Canning J, Kaneko T, Pru JK & Tilly JL (2004) Germline stem cells and follicular renewal in the postnatal mammalian ovary. Nature 428 145–150.
- Johnson J, Bagley J, Skaznik-Wikiel M, Lee HJ, Adams GB, Niikura Y, et al (2005) Oocyte generation in adult mammalian ovaries by putative germ cells in bone marrow and peripheral blood. Cell 122 303–315.
- Josso N, Lamarre I, Picard JY, Berta P, Davies N, Morichon N, Peschanski M, Jeny R(1993) Anti-Müllerian hormone in early human development. Early Hum Dev 33:91–99.
- Kim YK, Wasser SK, Fujimoto VY, Klein NA, Moore DE and Soules MR (1997) Utility of follicle stimulating hormone (FSH), luteinizing hormone (LH), oestradiol and FSH: LH ratio in predicting reproductive age in normal women Hum. Reprod., 12, 1152–1155,
- Kim SU, Ku SY, Suh CS(2002). Clinical significance of transvaginal color Doppler ultrasonography of the ovarian artery as a predictor of ovarian response in controlled ovarian hyperstimulation for in vitro fertilization and embryo transfer. J Assist Reprod Genet, 19, 103-112.

- Klein NA, Illingworth PJ, Groome N P et al (1996a) Decreased inhibin B secretion is associated with the monotropic FSH rise in older ovulatory women: a study of serum and follicular fluid levels of dimeric Inhibin A and B in spontaneous menstrual cycles. Clin Endocrinol Metab ,81, 2742-2745.
- Klein NA, Battaglia DE, Fujimoto VY et al (1996b) Reproductive aging: accelerated ovarian follicular development associated with a monotropic follicle-stimulating hormone rise in normal older women. J Clin Endocrinol Metab, 81, 1038-1045.
- Klein NA, Battagtia DE, Clifton DK, Bremnir WJ& Soules MR(1996c), The gonadotropin secretion pattern in normal women of advanced reproductive age in relation to the monotropic FSH rise, J Soc Gynecol Investig 3:27.
- Klein NA, Battaglia DE, Miller PB, Branigan EF, Giudice LC, Soules MR, (1996d)Ovarian follicular development and the follicular fluid hormones and growth factors in normal women of advanced reproductive age, J Clin Endocrinol Metab 81:1946,.
- Klein NA, Harper AJ, Houmard BS, Sluss PM& Soules MR(2002), Is the short follicular phase in older women secondary to advanced or accelerated dominant follicle development?. J Clin Endocrinol Metab 87:5746.
- Klinkert ER, Broekmans FJ Looman CW and te Velde ER (2004) A poor response in the first in vitro fertilization cycle is not necessarily related to a poor prognosis in subsequent cycles. Fertil Steril 81,1247–1253.
- Klinkert ER, Broekmans FJ, Looman CW, Habbema JD and te Velde ER (2005) The antral follicle count is a better marker than basal follicle stimulating hormone for the selection of older patients with acceptable pregnancy prospects after in vitro fertilization. Fertil Steril 83,811–814.
- Knebelmann B, Boussin L, Guerrier D, Legeai L, Kahn A, Josso N & Picard JY (1991) Anti-Müllerian hormone Bruxelles: a nonsense mutation associated with the persistent Müllerian duct syndrome. Proceedings of the National Academy of Sciences of the United States of America, 88, 3767–3771.
- **Kuliev A, Cieslak J, Ilkevitch Y, Verlinsky Y(2003)** Chromosomal abnormalities in a series of 6,733 human oocytes in preimplantation diagnosis for age-related aneuploidies. Reprod Biomed Online 6:54,
- **Kupesic S and Kurjak A (2002)** Predictors of IVF outcome by three-dimensional ultrasound. Hum Reprod 17, 950–955.

- **Kupesic S, Kurjak A, Bjelos D and Vujisic S** (2003) Three-dimensional ultrasonographic ovarian measurements and in vitro fertilization outcome are related to age. Fertil Steril 79,190–197.
- **Kutlešić R, Ljubić A, Milosavljević M, et al. (2006)** Color Doppler and color Doppler energy imaging and measurements of ovarian stromal blood flow in controlled ovarian hyperstimulation for in vitro fertilization. Medicine and Biology 13, 104 108
- La Marca A & Volpe A (2006) Anti-Müllerian hormone (AMH) in female reproduction: is measurement of circulating AMH a useful tool? Clinical Endocrinology 64, 603–610
- La Marca A & Volpe A (2007) The Anti-Mullerian hormone and ovarian cancer Hum Reprod Update, 13, 265-273
- La Marca A, Malmusi S, Giulini S, Tamaro LF, Orvieto R, Levratti P.& Volpe A (2004a) Anti-Müllerian hormone plasma levels in spontaneous menstrual cycle and during treatment with FSH to induce ovulation. Hum Reprod, 19, 2738–2741.
- La Marca A, Orvieto R, Giulini S, Jasonni VM, Volpe A & De Leo V (2004b) Müllerian-inhibiting substance in women with polycystic ovary syndrome: relationship with hormonal and metabolic characteristics. Fertil Steril, 82, 970–972.
- La Marca A, De Leo V, Giulini S, Orvieto R, Malmusi S, Giannella L & Volpe A (2005a) Anti-Müllerian hormone in premenopausal women and after spontaneous or surgically induced menopause. Journal of the Society for Gynecologic Investigation, 12, 545–548.
- La Marca, A, Giulini S, Orvieto R, De Leo V & Volpe A (2005b) Anti-Müllerian hormone concentrations in maternal serum during pregnancy. Hum Reprod, 20, 1569–1572.
- La Marca A, Pati M, Orvieto R, Stabile G, Carducci Artenisio A & Volpe A (2006) Serum anti-Müllerian hormone levels in women with secondary amenorrhea. Fertil Steril, 85, 1547 1549
- La Marca A, Giulini S, Tirelli A, Bertucci E., Marsella T., Xella S. and Volpe A. (2007) Anti-Müllerian hormone measurement on any day of the menstrual cycle strongly predicts ovarian response in assisted reproductive technology Hum. Reprod. 22,766-771.

- Lamb NE, Freeman SB, Savage-Austin A, et al,(1996) Susceptible chiasmate configurations of chromosome 2l predispose to non-disjunction in both maternal meiosis I and meiosis II. Nat Genet 14:400.
- Lamb NE, Feingold E, Savage A, et al, (1997) Characterization of susceptible chiasma configurations that increase the risk for maternal nondisjunction of chromosome 21, Hum Mol Genet 6:1391.
- **Lambalk CB & De Koning CH (1998)** Interpretation of elevated FSH in the regular menstrual cycle. Maturitas 30, 215-220.
- **Lambalk CB, de Koning CH, Flett A, van Kasteren Y, Gosden R and Homburg R (2004)** Assessment of ovarian reserve. Ovarian biopsy is not a valid method for the prediction of ovarian reserve. Hum Reprod 19, 1055–1059.
- Lane AH, Lee MM, Fuller AF Jr, Kehas DJ, Donahoe PK & MacLaughlin DT (1999) Diagnostic utility of Müllerian inhibiting substance determination in patients with primary and recurrent granulosa cell tumors. Gynecologic Oncology, 73, 51–55.
- Lappohn, R.E., Burger, H.G., Bouma, J., Bangah, M., Kransde, M. & Bruijn, H.W. (1989) Inhibin as a marker for granulosa-cell tumors. New England Journal of Medicine, 321, 790–793.
- **Lass A (2001)** Assessment of ovarian reserve is there a role for ovarian biopsy? Hum Reprod 16,1055–1057.
- **Lass A (2004)** Assessment of ovarian reserve: is there still a role for ovarian biopsy in the light of new data? Hum Reprod 19,467–469.
- **Lass A& Brinsden P (1999)** The role of ovarian volume in reproductive medicine. Hum Reprod Update 5, 256-266.
- Lass A, Skull J, McVeigh E. et al. (1997a) Measurement of ovarian volume by transvaginal sonography before ovulation induction with human menopausal gonadotrophin for in vitro fertilization can predict poor response. Hum Reprod, 12, 294-297
- Lass A, Silye R, Abrams DC et al (1997b) Follicular density in ovarian biopsy of infertile women: a novel method to assess ovarian reserve Hum Reprod, 12, 1028-1031.
- Lawson R, El-Toukhy T. Kassab, A, Tallor A, Braude P, parsons J, Seed P(2003), Poor response to ovulation induction is a stronger predictor of early menopause than elevated basal FSH: a life table analysis. Hum Reprod 18:527.

- Lee MM, Donahoe PK, Hasegawa T, Silverman B, et al (1996) Müllerian inhibiting substance in humans: normal levels from infancy to adulthood. J Clin Endocrinol Metab, 81, 571–576.
- Lee MM, Donahoe PK, Silverman BL, Hasegawa T, Hasegawa Y, Gustafson ML, Chang Y& MacLaughlin DT(1997) Measurements of Serum Mullerian Inhibiting Substance in the Evaluation of Children with Nonpalpable Gonads N. Engl. J. Med., May 22, ; 336(21): 1480 1486.
- **Lee S, Lenton, E, Sexton L et al (1988)** The effect of age on the cyclical patterns of plasma LH, FSH, oestradiol and progesterone in women with regular menstrual cycles. Hum. Reprod., 3, 851–855.
- **Licciardi FL, Liu HC & Rosenwaks Z(1995)** Day 3 estradiol serum concentrations as prognosticators of ovarian stimulation response and pregnancy outcome in patients undergoing in vitro fertilization Fertil Steril, 64,991-994.
- Long WQ, Ranchin V, Pautier P, Belville C, et al (2000) Detection of minimal levels of serum anti-Müllerian hormone during follow-up of patients with ovarian granulosa cell tumor by means of a highly sensitive enzyme-linked immunosorbent assay. J Clin Endocrinol Metab, 85, 540–544.
- **LoumayeE, Billion JM, Mine IM et al (1990)** Prediction of individual response to controlled ovarian hyperstimulation by means of a clomiphene citrate challenge test. Fertil Steril ,53, 295-301.
- **Magarelli PC, Peulstone AC & Buyalos RP (1996)** Discrimination between chronological and ovarian age in infertile women aged 35 years and older: predicting pregnancy using basal follicle stimulating hormone, age and number of ovulation induction/intra-uterine insemination cycles Hum Reprod,11, 1214-1219.
- Maheshwari A, Flower P & Bhattacharya S (2006) Assessement of ovarian reserve-should we perform tests of ovarian reserve routinely Hum Reprod 21:2729-2735
- **Maroulis GB(1991)** Effect of aging on fertility and pregnancy, Seminars Reprod Endocrinol 9:165.
- Martin JSB, Nisker JA, Tummon IS et al (1996) Future in vitro fertilization pregnancy potential of women with variable elevated day 3 follicle-stimulating hormone levels, Fertil Steril, 65, 1238-1240.

- **McGee EA & Hsueh AJ (2000)** Initial and cyclic recruitment of ovarian follicles. Endocrine Reviews 21 200–214.
- **Meldrum DR(1993)** female reproductive ageing-ovarian and uterine factors, Fertil Steril 60:314.
- Millar DM, Blake JM, Stringer DA, Hrra H& Bablak C (1993)Prepubertal ovarian cyst formalion: 5 years experience, Obstet Gytecol 81:434.
- Misra MDT, MacLaughlin PK, Donahoe& Lee MM (2002) Measurement of Mullerian Inhibiting Substance Facilitates Management of Boys with Microphallus and Cryptorchidism. J Clin Endocrinol Metab, 87: 3598 3602.
- Mol BW, Verhagen TEM, Hendriks DJ, Collins JA, Coomarasamy A,Opmeer BC and Broekmans FJ(2006) Value of ovarian reserve testing before IVF: a clinical decision analysis. Hum Reprod 21, 1816–1823.
- Motta PM & Makabe S (1986) Germ cells in the ovarian surface during fetal development in humans. A three-dimensional microanatomical study by scanning and transmission electron microscopy, J Submicrosc Cytol Pathol 18:271.
- Motta PM, Makabe S & Nottola SA (1997) The ultrastructure of human reproduction. I. The natural history of the female germ cell: origin, migration and differentiation inside the developing ovary. Hum Reprod Update 3:281.
- Mukherjee T, Copperman A, Lapiinski R et al. (1996) An elevated day three follicle-stimulating hormone: luteinizing hormone ratio (FSH:LH) in the presence of a normal day 3 FSH predicts a poor response to controlled ovarian hyperstimulation. Fertil. Steril., 65, 588–593.
- Muttukrishna S, Suharjono H, McGarrigle H & Sathanandan M (2004) Inhibin B and anti-Müllerian hormone: markers of ovarian response in IVF/ICSI patients? BJOG 111, 1248–1253.
- Muttukrishna S, McGarrigle H, Wakim R, Khadum I, Ranieri DM and Serhal P (2005) Antral follicle count, anti-mullerian hormone and inhibin B: predictors of ovarian response in assisted reproductive technology? BJOG, 112, 1384–1390.
- **Nader S& Berkowitz AS (1991)** *Use of the hormonal response to clomiphene citrate as an endocrinological indicator of ovarian aging. Hum Reprod, 6, 931-933.*
- Nahum R, Shifren JL, Chang Y, Leykin L, Isaacson K and Toth TL (2001) Antral follicle assessment as a tool for predicting outcome in IVF—is it a better predictor than age and FSH? J Assist Reprod Genet 18,151–155.

- Nakano R, Shima K, Yamoto M, Kobayashi M, Nishimori K and Hiraoka J (1989) Binding sites for gonadotropins in human postmenopausal ovaries. Obstet Gynecol 73,196–200.
- **Nasmyth K** (2001), Disseminating the genome: joining, resolving, and separating sister chromarids during mitosis and meiosis, Annu Rev Genet 35:673.
- Nasmyth K, peters JM & Uhlmann F (2000), Splitting the chromosome: cutting the ties that bind sister chromatids, Science 288:1379.
- National Collaborating Center for Women's and Children's Health. (2004) Fertility: Assessment and Treatment for People with Fertility Problems. RCOG press, UK.
- Navot D, Rosenwaks Z & Mergalioth E J (1987) Prognostic assessment of female fecundity. Lancet 332, 645-647.
- **Ng EH, Tang OS and Ho PC** (2000) The significance of the number of antral follicles prior to stimulation in predicting ovarian responses in an IVF programme. Hum Reprod 15,1937–1942.
- Ng EHY, Chan CCW, Tang OS, Yeung WSB and Ho PC (2004) Effect of pituitary downregulation on antral follicle count, ovarian volume and stromal blood flow measured by three-dimensional ultrasound with power Doppler prior to ovarian stimulation. Hum Reprod 19, 2132–2137.
- **Ng EHY, Fong DYT, Yeung WSB and Ho PC** (2005) Ovarian stromal blood flow in the prediction of ovarian response during in vitro fertilization treatment Hum Reprod, 20. 3147–3151.
- Ng EHY, Tang OS, Chan CC, Ho PC. (2006) Ovarian stromal vascularity is not predictive of ovarian response and pregnancy. Reprod Biomed Online.;12):43-9
- **Nikolaou D & Templeton A(2003)** Early ovarian ageing: a hypothesis: Detection and clinical relevance. Hum Reprod 18:1137.
- Noci I, Borri P, Chieffi O et al (1995) Aging of the human endometrium: a basic morphological and immunohistochemical study. Eur J Obstet Gynecol Reprod Biol, 63:181
- **Padilla SL, Bayati J & Garcia JE** (1990) Prognostic value of the early serum estradiol response to leuprolide acetate in in vitro fertilization. Fertil Steri1, 53. 288-294.

- **Pearlstone AC, Fournet N, Gambone JC et al (1992)** Ovulation induction in women of age 40 and older: the importance of basal follicle-stimulating hormone level and chronological age. Fertil Srevil. 58. 674-679.
- **Pellestor E, Andreo B, Arnal E, et al (2003)** Maternal aging and chromosomal abnormalities: new data drawn from in vitro unfertilized human oocytes, Hum Genet 112:195.
- Peluso JJ, Steger RW, Jaszczak S and Hafez ES (1976) Gonadotropin binding sites in human postmenopausal ovaries. Fertil Steril 27,789–795.
- Peňarrubia J, Fábregues F, Manau D, Creus M, Casals G, Casamitjana R, Carmona F, Vanrell JA & Balasch J (2005) Basal and stimulation day 5 anti-Mullerian hormone serum concentrations as predictors of ovarian response and pregnancy in assisted reproductive technology cycles stimulated with gonadotropin-releasing hormone agonistgonadotropin treatment. Hum Reprod 20, 915–922.
- Perez MM, Gromoll J, Behre HM, Gassner C, Nieschlag E and Simoni M (2000) Ovarian response to follicle-stimulating hormone (FSH) stimulation depends on the FSH receptor genotype. J Clin Endocrinol Metab 85,3365–3369.
- **Piette C, De Mouzon J, Bachalot A et al. (1990)** In-vitro fertilization: influence of women's age on pregnancy rates. Hum Reprod 5, 56-59.
- **Pigny P, Jonard S, Robert Y & Dewailly D (2006)** Serum anti-Müllerian hormone as a surrogate for antral follicle count for definition of the polycystic ovary syndrome. J Clin Endocrinol Metab, 91, 941–945.
- Piltonen T, Koivunen R, Ruokonen A& Tapanainen JS (2003) Ovarian age related responsiveness to human chorionic gonadotropin, J Clin Endoainol Metab 88:3327.
- **Piltonen T, Morin-Papunen L, Koivunen R, Perheentupa A, Ruokonen A & Tapanainen JS** (2005) Serum anti-Müllerian hormone levels remain high until late reproductive age and decrease during metformin therapy in women with polycystic ovary syndrome. Hum Reprod, 20, 1820–1826.
- **Pohl CR, de Ridder CM& Plant TM (1995)** Gonadal and nongonadal mechanisms contribute to the prepupertal hiatus in gonadotropin secretion in the female rhesus monkey (Macaca mulatta), J Clin Endocrinol Metab 80:2094.
- Popovic-Todorovic B, Loft A, Lindhard A, Bangsboll S, Andersson AM and Andersen AN (2003) A prospective study of predictive factors of ovarian

- response in 'standard' IVF/ICSI patients treated with recombinant FSH. A suggestion for a recombinant FSH dosage normogram. Hum Reprod 18, 781–787.
- **Pritts EA (2001)**, Fibroids and infertility: a systematic review of the evidence, Obstet Gynecol Survey 56:483.
- **Qu J, Godin PA, Nisolle M and Donnez J (2000)** Distribution and epidermal growth factor receptor expression of primordial follicles in human ovarian tissue before and after cryopreservation. Hum Reprod 15,302–310.
- **Rabinovici J& Jaffe RB** (1990), Development and regulation of growth and differentiated function of human and subhuman primate fetal gonads, Endocr Rev11:532.
- Rajpert-De Meyts E, Jorgensen N, Graem N, Müller J, Cate RL & Skakkebaek NE (1999) Expression of anti-Müllerian hormone during normal and pathological gonadal development: association with differentiation of Sertoli and granulosa cells. J Clin Endocrinol Metab, 84, 3836–3844.
- **Reame NE, Wyman TL, Phillips DJ et al (1998)** Net increase in stimulatory input resulting from a decrease in inhibin B and an increase in activin A may contribute in part to the rise in follicular phase follicle stimulating hormone of ageing cycling women. J Clin Endocrinol Metab, 83, 3302-3307.
- **Redmer D and Reynolds L (1996)** Angiogenesis in the ovary. Rev Reprod 1, 182–192.
- Rey R, Lordereau-Richard I, Carel JC, Barbet P, Cate RL, Roger M, Chaussain JL, Josso N (1993) Anti-Müllerian hormone and testosterone serum levels are inversely related during normal and precocious pubertal development. J Clin Endocrinol Metab 77: 1220–1226.
- **Rey RA, Belville, C, Nihoul-Fekete C, et al (1999)** Evaluation of gonadal function in 107 intersex patients by means of serum antimüllerian hormone measurement. *J Clin Endocrinol Metab*, 84, 627–631.
- Rey R, Sabourin JC, Venara M, Long, WQ, Jaubert F, Zeller, WP, Duvillard P, Chemes H & Bidart JM (2000) Anti-Müllerian hormone is a specific marker of sertoli- and granulosa-cell origin in gonadal tumors. Human Pathology, 31, 1202–1208.
- **Rey R, Lukas-Croisier C, Lasala C & Bedecarras P (2003)** *AMH/ MIS: what we know already about the gene, the protein and its regulation. Molecular and Cellular Endocrinology, 15, 21–31.*

- **Richardson SJ, Senikas V& Nelson JF** (1987) Follicular depletion during the menopausal transition-evidence for accelerated loss and ultimate exhaustion, J Clin Endocrinol Metab 65:1231.
- Rolaki A, Drakakis P, Millingos S, Loutradis D, Makriginnakis A.(2005) Novel trends in follicular development, atresia and corpus luteum regression: role for apoptosis. Reprod Biomed Online, 11: 93-103.
- **Rubin JM, Bude RO, Carson PL, Bree RL and Adler RS (1994)** Power Doppler US: a potentially useful alternative to mean frequency-based color Doppler US. Radiology 190, 853–856.
- **Santoro N, Brown JR, Adel T& Skurnick JH(1996)**, Characterization of reproductive hormonal dynamics in the perimenopause, J Clin Endocrinol Metab 81:1495-1501.
- **Santoro N, Adel, T & Skurnick JH (1999)** Decreased inhibin tone and increased activin A secretion characterize reproductive aging in women. Fertil Steril 71, 658-662.
- Scheffer GJ, Broekrnans FJM, Dorland M et al. (1999) Antral follicle counts by transvaginal ultrasonography are related to age in women with proven natural fertility. Fertil Steril, 72, 845-851.
- Scheffer GJ, Broekmans FJ, Bancsi LF, Habbema JD, Looman CW and te Velde ER (2002) Quantitative transvaginal two- and three-dimensional sonography of the ovaries: reproducibility of antral follicle counts. Ultrasound Obstet Gynecol 20, 270–275.
- Scheffer GJ, Broekmans FJ, Looman CW, Blankenstein M, Fauser BC, teJong FH and teVelde ER (2003) The number of antral follicles in normal women with proven fertility is the best reflection of reproductive age. Hum Reprod 18,700–706.
- Schipper I, De Jong FH & Fauser BCJM (1998) Lack of correlation between maximum early follicular phase serum follicle stimulating hormone concentrations and menstrual cycle characteristics in women under the age of 35 years. Hum Reprod 13, 1442-1448.
- Schmidt KL, Ernst E, Byskov AG, Nyboe AA & Yding AC (2003) Survival of primordial follicles following prolonged transportation of ovarian tissue prior to cryopreservation. Hum Reprod 18,2654–2659.

- **Schwartz D& Mayaux MJ** (1982) Female fecundity as a function of age: results of artificial insemination in 2193 nulliparous women with azoospermic husbands. Federation CECOS, New Engl J Med 306:404,
- **Scott RT & Hofmann GE (1995)** Prognostic assessment of ovarian reserve. Fertil. Steril, 63, 1-11.
- **Scott RT, Toner JP, Muasher SJ et al** (1989) Follicle-stimulating hormone levels on cycle day 3 are predictive of in-vitro fertilization outcome. Fertil Steril 51, 651-654.
- **Scott RT, Hofmann GE, Oehninger S et al (1990)** *Intercycle variability of day 3 follicle-stimulating hormone levels and its effect on stimulation quality in in vitro fertilization. Fertil Steril, 54,297-302.*
- **Scott RT, Leonardi MR, Hofmann GE et al (1993)** A prospective evaluation of clomiphene citrate challenge test screening in the general infertility population. *Obstet. Gynecol.*, 82, 539-545.
- Scott M. Nelson, Robin W. Yates and Richard Fleming (2007) Serum anti-Müllerian hormone and FSH: prediction of live birth and extremes of response in stimulated cycles—implications for individualization of therapy. Hum Reprod 22,2414-2421.
- Scott RT, Opsahl MS, Leonardi MR et al (1995) Life table analysis of pregnancy rates in a general infertility population relative to ovarian reserve and patient age. Hum Reprod10, 1706-1710.
- **Seifer DB, Gardiner AC, Ferreira KA& Peluso JJ** (1996) Apoptosis as a function of ovarian reserve in women undergoing in vitro fertilization, Fertil Steril 66:593.
- **Seifer DB, Lambert-Messerlian G, Hogan JW et al (1997)** Day 3 serum inhibin B is predictive of assisted reproductive technologies outcome. Fertil Steril 67, 110-114.
- **Seifer DB, Scott RT, Bergh PA et al. (1999)** Women with declining ovarian reserve may demonstrate a decrease in day 3 serum inhibin B before a rise in folliclestimulating hormone. Fertil Steril, 72, 63-65.
- Seifer DB, MacLaughlin DT, Christian BP, Feng B & Shelden RM (2002) Early follicular serum müllerian-inhibiting substance levels are associated with ovarian response during assisted reproductive technology cycles. Fertil Steril, 77, 468–471.

- Sharara FI & McClamrock HD (1999) The effect of aging on ovarian volume measurements in infertile women. Obstet. Gynecol. 94, 51-60.
- **Sharara FI & Scott RT (2004)** Assessment of ovarian reserve. Is there still a role for ovarian biopsy? First do no harm! Hum Reprod 19,470–471.
- Sharara FI, Scott RT & Seifer DB (1998) The detection of diminished ovarian reserve in infertile women. Am J Obstet Gynecol 179,804-812.
- **Sharif K, Elgendy M, Lashen H et al (1998)** Age and basal follicle stimulating homone as predictors of in vitro fertilization outcome. Br J Obstet Gynaecol, 105, 107-112.
- Shenfield F, Doyle P, Valentine A, Steele SJ, Tan S-L, (1993) Effects of age, gravidity and male infertility status on cumulative conception rates following artificial insemination with cryopreserved donor sperm: analysis of 2998 cycles of treatment in one centre over 10 years, Hum Reprod 8:60.
- **Sherman B, West J and Korenman S (1976)** The menopausal transition: analysis of LH, FSH, estradiol, and progesterone concentrations during menstrual cycles of older women. J. Clin. Endocrinol. Metab., 42, 629–636.
- **Sherman BM, Wallace RB and Treloar AE (1979)** The menopausal transition: endocrinological and epidemiological considerations. J. Biosoc. Sci. Suppl.6, 19-35.
- Shideler SE, DeVane GW, Kalra PS, Benirschke K, Lasley BL(1989), Ovarian pituitary hormone interactions during the perimenopause, Maturitas, 11:331.
- Shrim A, Elizur SE, Seidman DS, Rabinovici J, Wiser A, Dor J. (2006) Elevated day 3 FSH/LH ratio due to low LH concentrations predicts reduced ovarian response. Reprod Biomed Online. 12,418-22.
- **Silber SJ, Nagy Z, Devroey P et al (1997)** The effect of female age and ovarian reserve on pregnancy rate in male infertility: treatment of azoospermia with sperm retrieval and intracytoplasmic sperm injection Hum Reprod, 12. 2693-2700.
- Silberstein T, MacLaughlin DT, Shai I, Trimarchi JR, Lambert-Messerlian G, Seifer DB, Keefe DL and Blazar AS(2006). Müllerian inhibiting substance levels at the time of HCG administration in IVF cycles predict both ovarian reserve and embryo morphology Hum Reprod 21, 159–163,
- Sinclair AH, Berta P, Palmer MS, et al (1990), A gene from the human sexdetermining region encodes a protein with homology to a conserved DNA-binding motif, Nature 346:240.

- Smeenk JMJ, Sweep FCGJ, Zielhuis GA, Kremer JAM, Thomas CMG, Braat DDM, (2007) Antimüllerian hormone predicts ovarian responsiveness, but not embryo quality or pregnancy, after in vitro fertilization or intracyoplasmic sperm injection Fertil Steril;87:223–6.
- Smotrich DB, Widra EA, Gindorff PR et al. (1995) Prognostic value of day 3 estradiol on in vitro fertilization outcome. Fertil Steril, 64, 1136-1140.
- **Speed RM,** (1988) The possible role of meiotic pairing anomalies in the atresia of human fetal oocytes, Hum Genet 78:260.
- **Speert H, (1996)** Obstetric & Gynecologic Milestones Illustrated, The Parthenon Publishing Group, New York.
- **Sperrof L&Fritz MA(2005)***Clinical gynecologic endocrinology and infertility,* 7th *edition, p 100.*
- Staessen C, Camus M, Bollen N, Devroey P, Van Steirteghem AC (1992) The relationship between embryo quality and the occurrence of multiple pregnancies. Fertil Steril. 57,626-30.
- **Stien ZA(1985)** A woman's age: childbearing and childrearing. Am J Epidemiol 121:327.
- Syrop CH, Willhoite A & Van Voorhis BJ (1995) Ovarian volume: a novel outcome predictor for assisted reproduction. Fertil Steril, 64, 1167-1171.
- **Syrop CH, Dawson JD, Husman KJ et al (1999)** Ovarian volume may predict assisted reproductive outcomes better than follicle stimulating hormone concentration on day 3. Hum Reprod, 14, 1752-1756.
- **Tabarowski Z, Szoltys M, Bik M, Slomczyñska M (2005)** Atresia of large ovarian follicles of the rat. Folia Histochem Cytobiol,; 43: 43-50.
- **Tanbo T, Dale PO, Aby-holm T et al (1989)** Follicle-stimulating hormone as a prognostic indicator in clomiphene citrate/human menopausal gonadotrophin-stimulated cycles for in-vitro fertilization. Hum Reprod 6, 647-650.
- **Tanbo T, Dale PO, Ludne O et al (1992)** Prediction of response to controlled ovarian hypertimulation: a comparison of basal and clomiphene citratestimulated hormone level: Fertil steril 57, 819: 824.

- **Tarlatzis BC, Zepiridis L, Grimbizis G and Bontis J (2003)** Clinical management of low ovarian response to stimulation for IVF: a systematic review. Hum Reprod Update 9,61–76.
- te Velde ER, Pearson PL(2002) The variability of female reproductive ageing. Hum Reprod Update 8:141.
- **Teixeira J, Maheswaran S & Donahoe PK (2001)** Müllerian inhibiting substance: an instructive developmental hormone with diagnostic and possible therapeutic applications. Endocrine Reviews, 22, 657–674.
- **Templeton. A.. Morris. J.K. and Parslo W(1996)** factor that affect outcome of invitro fertilisation treatment. Lancet- 348, 1402-1406.
- **Thatcher SS and Naftoline F (1991)** The aging and aged ovary. Semin. Reprod. Endocrinol.9,189-199.
- Tibiletti MG, Testa G, Vegetti W, Alagna F, Taborelli M, Dalpra L, Bolis PF, Crosignani PG(1999), The idiopathic forms of premature menopause and early menopause show the same genetic pattern. Hum Reprod 14:2731.
- **Tietze. C.** (1957) Reproductive span and rate of reproduction among Hutterite women. Fertil Sreril 8:89.
- **Tilly JL, Kowalski KI, Schomberg DW, Hsueh AJ(1992)**, Apoptosis in atreric ovarian follicles is associated with selected decreases in messenger ribonucleic acid transcripts for gonadotropin receptors and cytochrome P450 aromatase, Endocrinology 131:1670.
- Tomás, C., Nuojua-Huftunen. S. and Martikainen, H. (1997) Pretreatment transvaginal ultrasound examination predicts ovarian responsiveness to gonadotrophins in in-vitro fertilization. Hum Reprod, 12, 220-223.
- **Toner, J.P., Philput, C.B., Jones. G.S. et al. (1991)** Basal follicle-stimulating hormone level is a better predictor of in-vitro fertilization performance than age. Fertil Steril, 55, 784-191.
- **Treloar AE, (1981)** *Mensrual cyclicity and the pre-menopause. Maturitas 3:249.*
- **Treloar AE, Boynton RE, Borghild GB& Brown BW (1967),** Variation of the human menstrual cycle through reproductive life, Int J Fertil 12:77.
- **Tremellen KP, Kolo M, Gilmore A, Lekamge DN** (2005) Anti-mullerian hormone as a marker of ovarian reserve. Aust N Z J Obstet Gynaecol ;45:20–24.

- **Ushiroyama T and Sugimoto O (1995)** *Endocrine function of the peri- and postmenopausal ovary. Horm Res* 44, 64–68.
- van Noord PAH, Dubas JS, Dorland M, Boersma H, te Velde E(1997): Age at early menopause. In a population based screening cohort, the role of menarche, fecundity, and life style factors, Fertil Steril 68:95.
- Van Noord-Zaadstra BM, Looman CWN Alsbach H et al (1991) Delaying child-bearing: effect of age on fecundity and outcome of pregnancy. Br. Med. J., 302, 1361-1365.
- van Rooij IA, Broekmans FJ, te Velde ER, et al(2002) Serum anti-Müllerian hormone levels: a novel measure of ovarian reserve. Hum Reprod 17, 3065–3071.
- van Rooij IAL, Bancsi LFJMM, Broekmans FJM, Looman CWN, Habberna JDF and te Velde ER (2003) Women older than 40 years of age and those with elevated follicle-stimulating hormone levels differ in poor response rate and embryo quality in in vitro fertilisation. Fertil Steril 79,482–488.
- van Rooij IAJ, de Jong E, Broekmans FJM, Looman CW, Habbema JDF and te Velde ER (2004a) High follicle-stimulating hormone levels should not necessarily lead to the exclusion of subfertile patients from treatment. Fertil Steril 81, 1478–1485.
- van Rooij IA, Tonkelaar I, Broekmans FJ, Looman CW, Schefferde GJ, Jong FH, Themmente AP & te Velde ER (2004b) Anti-Müllerian hormone is a promising predictor for the occurrence of the menopausal transition. Menopause, 11, 601–606.
- van Rooij IA, Broekmans FJ, Scheffer GJ, Looman CW, Habbema JD, de Jong FH, Fauser BJ, Themmente AP & Velde ER (2005) Serum antimüllerian hormone levels best reflect the reproductive decline with age in normal women with proven fertility: a longitudinal study. Fertil Steril, 83, 979–987.
- van Zonneveld P, Scheffer GJ, Broekmans FJ, Blankenstein MA, de Jong FH, Looman CW, Habbema JD, te Velde ER(2003)Do cycle disturbances explain the age-related decline of female fertility? Cycle characteristics of women aged over 40 years compared with a reference population of young women, Hum Reprod 18:495.
- Vanhoutte L, De Sutter P, Van der Elst J and Dhont M(2005) Clinical benefit of metaphase I oocytes. Reproductive Biology and Endocrinology, 3:71

- Varasteh NN, Neurwirth RS, Levin B& Keltz MD(1999) pregnancy rates after hysteroscopic polypectomy and myomectomy in infertile women, Obstet Gynecol 94:168.
- Vaskivuo TE, Anttonen M, Herva R, Billig H, Dortand M, te Vetde ER, Stenback F, Heikinheimo (2001) Survival of human ovarian follicle from fetal to adult life apoptosis, apoptosis related proteins and transcription factor GATA-4. J Clin Endocrinol Metab 86:3421.
- Vegtti W, Marozzi A, Manfredini E, Testa C et al (2000), premature ovarian failure, Mol Cell Endorinol 161:53.
- **Volarcik K, Sheean L, Goldfarb J, et al (1998)**, The meiotic competence of in-vitro matured human oocytes influenced by donor age: evidence that folliculogenesis is compromised in the reproductively aged ovary, Hum Reprod 13:154.
- **Vermeulen A (1976)** The hormonal activity of the postmenopausal ovary. J Clin Endocrinol Metab 42,247–253.
- **Virro MR & Shewchuk AB (1984)** Pregnancy outcome in 242 conceptions after artificial insemination with donor sperm and effect of maternal age on the prognosis of successful pregnancy. Am J Obstet Gynecol. 148. 518-524.
- **Visser JA, Themmen AP(2005)** *Anti-mullerian hormone and folliculogenesis. Mol Cell Endocrinol*; 234:81–6.
- **Vladimirov IK, Tacheva DM, Kalinov KB (2004)***Mean Ovarian Diameter (MOD)* as a Predictor of Poor Ovarian Response. Journal of Assisted Reproduction and Genetics, 21, 73-77.
- **Wallace WH and Kelsey TW (2004)** Ovarian reserve and reproductive age may be determined from measurement of ovarian volume by transvaginal sonography. Hum Reprod 19, 1612–1617.
- **Warburton D (1987)** Reproductive loss: how much is preventable? New, Engl J Med 316:158.
- **Warburton D, Kline J, Srein 2., Strobino B(1986)** Cytogenetic abnormalities in spontaneous abortions of recognized conceptions. In: Porter IH, ed. Perinatal Genetics: Diagnosis and Treatment, Academic press. New York, 133.
- Webber LJ, Stubbs S, Stark J, Trew GH, Margara R, Hardy K & Franks S (2003) Formation and early development of follicles in the polycystic ovary. Lancet, 362, 1017–1021.

- Weenen C, Laven JS, Von Bergh AR, Cranfield M, Groome NP, Visser JA, et al (2004) Anti-Müllerian hormone expression pattern in the human ovary: potential implications for initial and cyclic follicle recruitment. Molecular Human Reproduction, 10, 77–83.
- **Weghofer A and Feichtinger W** (2006)*The forgotten variable: impact of luteinizing hormone on the prediction of ovarian reserve. Fertil Steril.* 8, 259-61.
- Wilcox AJ, Weiberg CR, O'Connor JF, Baird DD, Schtatierer JP, Canfield RE, Armstrong EG& Nisula BC (1988) incidence of early loss of pregnancy, New Engl J Med 319:189.
- Wildt L, Hausler A, Marshall G, Hutchison JS, Plant TM, Belchetz PE& Knobil E(1981)Frequency and amplitude of gonadotropin-releasing hormone Stimulation and gonadotropin secretion in the rhesus monkey, Endocrinology 109;376.
- Wilson JD, Griffin JE, George FW& Leshin M (1981) The role of gonadal steroids in sexual differentiation, Recent Prog Hor Res 37: 1.
- Winslow K.L., Toner J.P., Brzyski R.G. et aI (1991) The gonadotropin releasing hormone agonist stimulation test a sensitive predictor of performance in the flare-up in vitro fertilization cycle. Fertil Steril, 56. 711 717.
- Wolff E.F, and Taylor H.S. (2004) Value of the day 3 follicle-stimulating hormone measurement Fertil Steril; 81: 1486-8
- Zaidi J, Campbell S, Pittrof R, Kyei-Mensah A, Adel Shaker, Howard SJacobs and Seang Lin Tan(1995) Ovarian stromal blood flow in women with polycystic ovaries—a possible new marker for diagnosis? Hum Reprod 10 1992-1996.
- **Zaidi J, Barber J, Kyei-mensah A, Bekir J, Campbell S and Tan SL (1996)**Relationship of ovarian stromal blood flow at the baseline ultrasound scan to subsequent follicular response in an in vitro fertilization program. Obstet Gynecol 88, 779–784.
- Ziebe S, Loft A, Petersen JH, Andrsen AG, Lindenberg S, Petersen K& Andersen AN, (2001) Embryo quality and developmental potential compromised by age, Acta Obstet Gyecol Scand 80:169.
- Zinaman MJ, Clegg ED, Brown CC, O.Connor J& Selevan SG(1996), Estimates of human fertility and pregnancy loss, Fertil Steril 65:501.