

REFERENCES

Adlercreutz, H. and Mazur, W. (1997):

Phytoestrogens and western diseases.

Ann. Med., 29: 95-120.

Albertazzi, P., Pansini, F. and Bonaccorsi, G. (1998):

The effect of dietary soy supplementation on hot flushes.

Obstet. Gynecol., 91: 9-11.

Alexandersen, P., Toussaint, A. and Christiansen, C. (2001):

Ipriflavone Multicenter European Fracture Study. Ipriflavone in the treatment of postmenopausal osteoporosis; a randomized controlled trial. Menopause, 285 : 1482-1488.

Amato, P., Christopher, S. and Mellon, P.L. (2002):

Estrogenic activity of herbs commonly used as remedies for menopausal symptoms. Menopause, 9: 145-150.

Amy, C.E. (2002):

Allopathic and complementary alternative to hormone replacement therapy. Expert opin. pharmaco. Ther., 3: 949-955.

Anderiano, K. (2000):

The prevention of recurrence of osteoporotic fractures study.

Am. J. Med., 109: 267-276.

Anderson, J. and Garner, S.C. (1999):

Phytoestrogens and bone.

Baillieres Clin. Endocrinol. Metmb., 12: 543-557.

Atkinson, C., Compston, J.E. and Ringhara, S.A. (2000):

The effects of isoflavones Phytoestrogens on bone: preliminary results from a large randomized controlled trial (Abstract).

Endocri. Soc. Ann. Meet. Program Abs., 82: 43.

Atkinson, C.; Warren, R.M.L. and Bingham, S.A. (2004):

Red clover-derived isoflavones and mammographic breast density: a double-blind, randomized, placebo-controlled trial (ISRCTN 43940165). Breast Cancer Res., 6: R170-R179. Doi: 10.1186/bcr773).

Azevedo, G.D., Franco, R.F. and Baggio, M.S. (2005):

Procoagulant state after Raloxifene therapy in postmenopausal women. Fert. Sterility J. Dec., 84 (6): 1680-1684.

Baeza, I., de Castro, N.M. and Alvarado, C. (2007):

Improvement of immune cell functions in aged mice treated for five weeks with soybean isoflavones. Ann. N. Y. Acad. Sci., Apr. 1100: 497-504.

Bagger, Y.Z., Tanko, L.B.; Alexandersen, P. and Karsdal, M.A. (2005):

Oral salmon Calcitonin induced suppression of urinary collagen type II degradation in postmenopausal women: a new potential treatment of osteoarthritis. Bone, 37 : 425-430.

Barton, D.P.H.D (2002):

Pathophysiology and treatment of hot flushes.

Mayo Clinic. Proc., 77: 1207-1218.

Black, D.M., Bilezikian, J.P. and Enstrud, K.E. (2005):

One year of Alendronate after one year of parathyroid hormone (1-84) for osteoporosis. N. Engl. J. Med., 353 : 555-565.

Black, D.M., Greenspan, S.L. and Enstrud, K.E. (2003):

The effects of parathyroid hormone and Alendronate alone or in combination in postmenopausal osteoporosis. N. Engl. J. Med., 349 : 1207-1212.

Blumenthal, M. (2004):

The use of black cohosh to treat symptoms of menopause.

Sex Reproduc. Menopause, 2: 27-34.

Bordeleau, L.; Pritchard, K.; Goodwin, P. and Loprinzi, C. (2007):

Therapeutic options for the management of hot flushes in breast cancer survivors: An evidence-based review. Clin. Ther., 29 : 230-241.

Breslau, N.A., Brinkely, L., Hill, K.B. and Pak, C.Y. (1998):

Relationship of animal protein-rich diet to kidney stone formation and calcium metabolism. J. Clin. Endocrinol. Metab., 66: 140-146.

Btzezinski, A., Adlercreutz, H. and Shaoul, R. (1997):

Short term effects of Phytoestrogens-rich diet on postmenopausal women. Menopause, 4: 89-94.

Burk, G.I., Vitolins, M.Z. and Bland, D. (2000):

Soybean isoflavones as an alternative to traditional hormone replacement therapy. J. Nutr., 130: 6648-6658.

Burr, M.L., Fehily, A.M. and Holliday, R.M. (1999):

Effects of changes in fat and fiber intakes on myocardial infarction. Lancet, 2: 757-767.

Cagnacci, A., Renzi, A. and Cannolatta, M. (2006):

Tibolone and estradiol plus norethisterone acetate similarly influence endothelium-dependent vasodilatation. Infertility and Sterility, J. Aug., 86 : 480-483.

Campisi, R., Camilletti, J. and Guiglioni, A. (2006):

Tibolone improves myocardial perfusion in postmenopausal women with ischemic heart disease. J. Am. College, Cardiology, Feb. 7: 47 : 559-564.

Carton, M. (2006):

Breast cancer and hormone-replacement therapy:
Million Women Study, 362: 1328-1331.

Cecchini, M.G., Fleisch, H. and Muhlbauer, R.C. (1997):

Ipriflavone inhibits bone resorption in intact and ovariectomized rats. Calcif. Tissue Int., 61: 9-11.

Chapurlat, R.D. and Delmas, P.D. (2006):

Bisphosphonates for postmenopausal osteoporosis. Nat. Clin. Pract. Endocrinl. Metab, Apr. 2 (4): 2L 1-9; Quiz following 238.

Cheng, G., Wilezek, B. and Landgren, B.M. (2007):

Isoflavones treatment for acute menopausal symptoms.
Menopause, 14 : 468-473.

Christen, W.G., Gazino, J.M. and Hennekens, C.H. (2000):

Design of physician's health study. Randomized trial.
Ann. Epidemiol., 125-134.

Christopher Naquin, M.D. (2002):

Louisiana state university health science center kenner.
American Family Physician, 66: 129-134.

Clarkson, T.B. (2000):

Soy phytoestrogens: what will be their role in postmenopausal hormone replacement therapy? Menopause, 7: 71-75.

Cobellis, L. (2007):

Raloxifene therapy interacts with serum osteoprotegerin in postmenopausal women. Maturitas J. 20; 65 : 38-44.

Conner, P. (2007):

Breast response to menopausal hormone therapy, aspects on proliferation, apoptosis and Mammographic density. Ann. J. Med., 39 : 28-34.

Cornwell, T.; Cohick, W. and Raskin, I. (2004):

Dietary Phytoestrogens and health.
Phytochem., 65: 995-1016.

Coward, L., Srnith, M. and Barnes, S. (1998):

Chemical modifications of Isoflavones in soy food during cooking and processing. Am. J. Clin. Nutri., 68: 1486S-1491S.

Cruz, M.N., Agewall, S., Schenck-Gustafsson, K. and Kublickiene, K. (2007):

Acute dilatation of Endothelium by Phytoestrogens and estrogen receptor subtypes expression in small arteries from women with coronary heart disease. J. Atherosclerosis, Mar 14 .

Davis, S.R, Murkies, A.L. and Wilcox, G. (1998):

Phytoestrogens in clinical practice.
Integr. Med., 1: 27-34.

Deane, A., Constancio, L., Fogelman, I. and Hampson, G. (2007):

The impact of vitamin D status on changes in bone mineral density during treatment with Bisphosphonates and after discontinuation following long-term use in post-menopausal osteoporosis. BMC Musculoskeletal-disorders, 8: 3.

Delbressine, L., de Gooyer, M. and Kloosterober, H.L. (2000):

Sulphated metabolites of Tibolone show Sulphatase inhibiting activity. Progr. Endo, 8: 1639: 396.

Delmas, P.D., Bjarnason, N.H., Mitlak, B.H., Rovoux, A.C ; Shah, A.S. and Huster, W.J. (2004):

Effects of Raloxifene on bone mineral density, serum cholesterol concentration and uterine endometrium in postmenopausal women. Engl. J. Med., 337: 1641-1647.

Denes, P., Larson, J.C. and Greenland, P. (2007):

Major and minor ECG abnormalities in asymptomatic women and risk of cardiovascular events and mortality. JAMA , 297 : 978-985.

Duncan, A.M., Underhill, K.E. and Kurzer, M.S. (2000):

Modest hormonal effects of soy isoflavones in postmenopausal women J. Clin. Endocrinol. Metab., 85: 448.

Ederveen, A.G.H. and Kloosterboer, H.I. (2001):

Tibolone exerts its protective effect on trabecular bone loss through the estrogen receptors. J. Bone Miner. Res., 16: 1651.

Elbers, J., Hgeluken, C. and Wadhanv, A. (2007):

Tibolone (Livial, enhances Warfarin induced anti-coagulation in postmenopausal women. Maturitas J., 56 : 94-100.

Erdman, J.W., Stillman, R.J. and Boileau, R.A. (2000):

Provocative relation between soy and bone maintenance.
Am. J. Clin. Nurt., 72: 679-680.

Filshie, J., Bolton, T. and Ashley, S. (2005):

Acupuncture and self acupuncture for long term treatment of vasomotor symptoms in cancer patient-audit and treatment algorithm. Acupuncture Med., 23 (4): 171-80.

Fitzpatrick, L.A. (2003):

Soy isoflavones: hope or hype.
Maturitasj, (Suppl. 1): S21-S29.

Gass, M.G. and Taylor, M.D. (2001):

Alternatives for women through menopause.

Am. J. Obst. Gynecol., 185 : S47-S56.

Glazier, M.G. and Bowman, M.A. (2001):

A review of the evidence for the use of phytoestrogens as a replacement for traditional estrogen replacement therapy. Arch.

Intern. Med., 461: 1161-1172.

Gonzalez, C.A. and Salas-Salvado, J. (2006):

The potential of nuts in the prevention of cancer.

Br. J. Nutri., 96, Suppl. 2: S87-94.

Gordon, N., Sobel, D. and Tarazona, E. (1998):

Use of and interest in alternative therapies among adult primary care clinicians and adult members in a large health maintenance organization. West. J. Med., 169: 153-161.

Grady, D.; Wenger, N.I.C and Herrington, D. (2000):

Postmenopausal hormone therapy increases the risk for venous thromboembolic diseases: The Heart and Estrogen/Progestin Replacement Study. Ann. Intern. Med., 132: 689-696.

Greenspan, S.L., Bone, H.G. and Ettinger, M.P. (2007):

Treatment of Osteoporosis with Parathyroid Hormone Study Group; Effect of recombinant human parathyroid hormone (1-84) on vertebral fracture and bone mineral density in postmenopausal women with osteoporosis: a randomized trial. J. Ann. Intern. Med., 146 : 326-339.

Guttuso, T., Kurlan, R. and Kkburtz, K. (2003):

Gabapentin effects on hot flushes in postmenopausal women; a randomized controlled trial. *Obstet. Gynecol.*, 101: 337.

Hatourech, G.E., Khosla, S. and Sanyal, A. (2003):

Role of RANK ligand in mediating increased bone resorption in early postmenopausal women. *J. Clin. Invest.*, 111: 1221-1230.

Head, K.A. (1999):

Ipriflavone; an important bone-building isoflavones.
Altern. Med. Rev., 4: 10-22.

Helena, J., Teede, Fabien, S., Dalais and Barry, P. McGrath (2004):

Dietary soy containing phytoestrogens does not have detectable estrogenic effects on hepatic protein synthesis in postmenopausal women. *Am. J. Clin. Nutr.* Vol. 79, 396-401.

Hendrix, S. (1997):

Nonestrogen management of menopausal symptoms.
Endocrinol. Metab. Clin. North. Am., 26: 379-390.

Hidalgo, L.A., Chedraui, P.A. and San Miguel, G. (2005):

The effect of red clover isoflavones on menopausal symptoms, lipids and vaginal cytology in menopausal women. *The Official J. of the Intern. Soc. Gynec. Endocrinol.*, Nov. 215: 257-264.

Hirata, J., Swiersz, L. and Ettinger, B. (1997):

Does dong quai have estrogenic effects in postmenopausal women? A double blind, placebo-controlled trial. *Fertil. Steril.*, 68: 981-986.

Hodgson, J.M., Puddey, I.B. and Croft, K.D. (1997):

Supplementation with isoflavonoid phytoestrogens does not alter serum lipid concentrations; a randomized controlled trial in human. J. Nutr., 128: 728-732.

Holick, M.F. (1996):

Vitamin D and bone health.
J. Nutr. (Suppl. 4): S1159-S1164.

Honore, E.K., Williams, J.K. and Anthony, M.S. (1997):

Soy isoflavones enhance vascular reactivity in atherosclerotic female macaques. Fertil. Steril., 67: 148-154.

Htzpatrick, L.A. (2003):

Soy isoflavones; hope or hype.
Maturitas, 14: 44 Suppl. 1: S21-S29.

Ingram, D., Sanders, K. and Lopez, D. (1997):

Case-control study of phytoestrogens and breast cancer.
Lancet, 350: 990-994.

Ivarsson, T., Spetz, A.C. and Hammer, I.V.L. (1998):

Physical exercise and vasomotor symptoms in postmenopausal women. Maturitas, 29: 139-146.

Iwamoto, J., Takeda, T. and Sato, Y. (2006):

Efficacy and safety of Alendronate and Risendronate for post-menopausal osteoporosis. Curr. Med. Res. Opin., May 22 (5); 919-928.

Jackson, R.D., LaCroix, A.Z. and Barad, D. (2006):

Women's Health Initiative Investigators. Calcium plus vitamin D supplementation and the risk of fractures. N. Engl. J. Med., Feb., 16; 354 (7): 669-83.

Jane, L. Limer and Valeric S. (2004):

Phytoestrogens and breast cancer chemoprevention.
Breast Cancer Res., 6 : 119-127.

Jian, W.U., Jun; O and Yoshiko Ishimi (2006):

Effects of Isoflavone and Exercise on BMD and Fat Mass in Post-menopausal Japanese Women: A 1-year Randomized Placebo Controlled Trial Bone and Miner Research, 21: 780-789 (doi: 10.1359/jbmr.060209).

Jordan, V.C. (2007):

Beyond Raloxifene for the prevention of osteoporosis and breast cancer. Br. J. Pharmacology, 150 : 3-4.

Karin, H., Humphries, M. and Gill. S. (2003):

Risks and benefits of HRT; the evidence speaks.
CAMJ, 4: 1001-1010.

Kashani, E., Lyritis, G.P. and Matsouka, K. (2005):

Effect of intermittent administration of 200 IU intranasal salmon Calcitonin and low doses of alpha (OH) vitamin D3 on bone mineral density of the lumbar spine and hip region and biochemical bone markers in women with postmenopausal osteoporosis, a pilot study. Clin. Rheumatol., 24 : 232-238, Epub, Jan 13.

Kass-Annese, B.R. (2000):

Alternative Therapies for Menopause.
Cl. Obstet. Gynecology., 431 : 162-183.

Keller, C.; Fullerton, J. and Fleury, J. (1998):

Primary and secondary prevention strategies among older postmenopausal women. J. Nurse Midwif, 43: 262-272.

Keller, C., Fullerton, J. and Mobley, C. (1999):

Supplemental and complementary alternatives to hormone replacement therapy. J. Am. Academy of Nurse Practitioners, 11 : 187-198.

Kennely, E.J., Baggett, S. and Nuntanakarn, P. (2002):

Analysis of thirteen population of black cohosh for formomnetin. Phytomedicine, 9: 461-467.

Kipersztok, S. (1997):

Postmenopausal osteoporosis key issues.
University of Florida Continuing Medical Education Program.

Kirk, E.A., Sultherland, P. and Wang, S.A. (1998):

Dietary isoflavones reduce plasma cholesterol and atherosclerosis in C57BL/6 mice but not LDL receptor-deficient mice. J. Nutr., 128: 954-959.

Kloosterboer, H.L. (2000):

Intracrinology; the secret of the tissue specificity of Tibolone. J. Br. Menopause Soc., 6: 23-27.

Kronenberg, F. and Fugh-Berman, A. (2002):

Complementary and alternative medicine for menopausal symptoms: A review of randomized controlled trials. Ann. Intern. Med., 137: 805.

Kuiper, G., Lemmen, J. and Carsson, B. (1998):

Interaction of estrogenic chemicals and phytoestrogens with oestrogen receptors beta. Endocrinology, 139: 4252-4263.

Kurtay, G., Berker, B. and Demirer, C. (2004):

Transvaginal ultrasonographic assessment of the endometrium in asymptomatic, postmenopausal women using different HRT regimens containing Tibolone or estrogen. J. Reprod. Med., 49 ; 893-898.

Lambrinoudaki, I., Christodoulakos, G. and Botsis, D. (2006):

Bisphosphonates. Annals-of-the-New-York-Academy-of Sciences,, 1092: 397-402.

Lampe, J.W. (2003):

Isoflavonoid and lignan as dietary ligand in mediating increased bone resorption in early postmenopausal women. J. Clin. Invest., 111: 1221-1230.

Lau, E.M. and Woo, J. (1998):

Nutrition and osteoporosis.
Curr. Opin. Rheurn., 10: 368-372.

Lajous M., Lazcano E. –P.I Romieul (2006):

Folate, vitamin B6, and vitamin B12 Intake and the Risk of Breast Cancer among Mexican Women. Cancer Epidemiol. Biornarkers Prev. 15 : 443-448.

Lim, P.S.; Oug, F.B. and Adeeb, N. (2005):

Bone health in urban midlife Malaysian women: risk factors and prevention. Osteoporosis Int. J., 16 ; 2069-2079, Epub. Oct.19.

Loprinzi, C.L. and Barton, D.L. (2002):

Pilot evaluation of gabapentin for treating hot flushes.
Mayo Clin. Prac., 77 (1): 159-163.

Machelle, M. S. (2003):

Treating hot flushes without hormone replacement therapy applied evidence. Family Practice, 52: 291-296.

Mahady, G.B. (2003):

Is black cohosh estrogenic? Nutr. Rev., 61: 183-186.

Malmros, B., Mortensen, L. and Charles, P. (1998):

Positive effects of physiotherapy on chronic pain and performance in osteoporosis. Osteoporosis Int. J., 8: 215-221.

Maskarinec, G., Williams, A.E. and Carlin, L. (2003):

Mammographic densities in a one-year isoflavone intervention. Eur. J. Cancer Prev., 12: 165-169.

McDonnell, D.P., Connor, C.E. and Morris, J.D. (2002):

Definition of the molecular and cellular mechanisms underlying the Tissue-selective Agonist Antagonist Activities of Selective Estrogen Receptor Modulators. Recent Progress in Hormone Research, 57: 295-316.

Messalli, E.M., Mamini, G. and Cobellis, A.L. (2007):

Raloxifene therapy interacts with serum osteoprotegerin in postmenopausal women. Maturitas, Jan. 20, 56 : 38-44.

Messina, M.J. (1999):

Legumes and soy beans, overview of their nutritional profiles and health effects. Am. J. Clin. Nut., 70: 449S-450S.

Michael, D.S., James, H.L. and Robert, W. (2000):

Dietary soy isoflavones favorably influence lipids and bone turnover in healthy postmenopausal women. J.Nutr., 130:666-679

Miller ER3rd., Pastor-Barriuso, R. and Guallar, E. (2005):

High dosage vitamin E supplementation may increase all cause mortality. Ann. Intern. Med., 142 : 37-46.

Moen, M.D. and Scott, L.J. (2006):

Recombinant full-length parathyroid hormone (1-84).
Drugs, 66 : 2371-2381, discussion 2382-2385.

Morito, K., Hirose, T. and Kinjo, T. (2001):

Interaction of phytoestrogens with receptors alpha and beta.
Biol. Pharmaceut. Bull., 24: 351-356.

Nagi, S.L., Vonsaal, F.F. and Welshons, W.V. (1998):

The effective free fraction of estradiol and xenoestrogens in human serum measured by whole cell uptake assays, physiology of delivery modifies estrogenic activity. Proc. Soc. Exp. Biol. Med., 217: 300-309.

Nedstrand, E., Wijma, K. and Bammar, M. (2005):

Vasomotor symptoms decrease in women with breast cancer randomized to treatment applied relaxation or electro-acupuncture; a preliminary study. Climacteric, 8 (3): 243-250.

Newton, K.M., Buist, D.S. and LaCroix, A.Z. (2002):

Use of alternative therapies for menopause symptoms: Results of a population-based survey. Obstet Gynecol., 100: 18-25.

Nguyen, N.D., Eisman, J.A. and Nguyen, T.V. (2006):

Anti-hip fracture efficacy of Bisphosphonates.

J. Bone. Miner. Res., Feb. 21 : 340-349.

Nieves, J.W., Komar, L. and Lindsay, R. (1998):

Calcium potentiates the effect of estrogen and Calcitonin on bone mass: review and meta-analysis. Am. J. Clin. Nutr., 67: 18-24.

Pan, Y., Anthony, M. and Clarkson, T.B. (1999):

Effect of estradiol and soy phytoestrogens on choline acetyltransferase and nerve growth factor mRNAs in the frontal cortex and hippocampus of female rats. Proc. Soc. Exp. Biol. Med., 221: 118-125.

Peeters, P.H., Kienan-Boker, L. and Grobbee, D.E. (2003):

Phytoestrogens and breast cancer risk. Review of the epidemiological evidence. Breast Cancer Res. Treat. 77:171-183

Philp, H.A. (2003):

Hot flushes: a review of the literature on alternative and complementary treatment approaches. Alter Med. Rev., 8:284-302.

Potter, S.M., Baum, J.A. and Teng, H. (2002):

Soy protein and Isoflavones women.

Alternative Raven, 7: 317-327.

Powel, T. (2004):

Isoflavones and women's health. Breast Cancer Res., 6 : 140-142

Epub, Apr. 6.

Prince, R.L.; Devine, A. and Dick, I.M. (2006):

Effects of calcium supplementation on clinical fracture and bone structure: results of a 5-year, double-blind, placebo-controlled trial in elderly women. Arch. Intern. Med., Apr. 24, 166 (8): 869-75. Nat. Clin. Pract. Rheumatol. J., 3 : 18-19.

Quartarolo, N. and Arya, V. (2004):

Eosinophilia-myalgia syndrome. medicine Textbook, Retrieved Dec. 13, [http:// www Emedicine. Com /derm/ topic 891. htm](http://www.Emedicine.Com/derm/topic891.htm).

Quella, S.K., Loprinzi, C.L. and Barton, D.L. (2000):

Evaluation of soy phytoestrogens for the treatment of hot flashes in breast cancer survivors. A North Central Cancer Treatment Group Trial. J. Clin. Oncol., 18: 1068.

Rad, M.; Humpel, M. and Schaefer, O. (2006):

Pharmacokinetics and systemic endocrine effects of the phytoestrogenic 8-prenylnaringenin after single oral dose to postmenopausal women. Br. J. Clin. Pharmacol.. 62 : 288-96.

Raus, K., Brucker, C. and Wuttke, W. (2006):

First-time proof of endometrial safety of the special black cohosh extract (Actaea or Cimicifuga Racemosa extract) CR BNO 1055. Menopause, Jul-Aug., 13 : 678-691.

Reid, M.D., Tim Cundy, M.D., Andrew, B. and Grey, M.D. (2006):

Addition of Monofluorophosphate to estrogen therapy in postmenopausal osteoporosis. J. Clin. Endocrine Metab., first published ahead of print April 17, as doi:10.1210/jc.2264.

Riggs, B.L. and Hartmann, L.C. (2003):

Selective Estrogen Receptor Modulators, Mechanisms of Action and Application to Clinical Practice. N. Engl. J. Med. Vol. 34: 618-629.

Rossi, M., Garavello, W. and Negri, E. (2007):

Flavonoids and risk of squamous cell esophageal cancer. Int. J. Cancer; 120 : 1560-1564.

Rossi, M., Negri, E. and La Vccchia, C. (2006):

Flavonoids and colorectal cancer in Italy. Cancer Epidemiol. Biomarkers Prev., Aug. 15 : 1555-1558.

Ruiz-Larrea, M.B., Mohan, A.R. and Pice-Evanf (1997):

Antioxidant activity of phytoestrogenic isoflavones. Free Medic. Res., 26: 63-710.

Seidl, M.M. and Stewart, D.E. (1998):

Alternative treatments for menopausal symptoms: systematic review of scientific and lay literature. Can Fam Physician, 44: 1299-1308.

Setchell, K.D. and Cassidy, A. (1998):

Dietary isoflavones: biological effects and relevance to human health. J. Nutr., 129: 758-767.

Shen, C.L., Williams, J.S. and Yeh, J.K. (2007):

Comparison of the effects of tai chi and resistance training on bone metabolism in the elderly: a feasibility study. *Am. J. Chin. Med.*, 35 : 369-381.

Siefker, K. and DiSilvestro, R.A. (2006):

Safety and antioxidant effects of a modest soy protein intervention in hemodialysis patients. *J. Med. Food*, Fall, 9 : 368-367.

Silverman, S.L., Watts, N.B. and Lindsay, R. (2007):

Effectiveness of Bisphosphonates on nonvertebral and hip fractures in the first year of therapy. *Osteoporosis*, 18 : 25-34.

Sirtori, C.R. (2001):

Risks and benefits of soy Phytoestrogens in cardiovascular diseases, cancer, climacteric symptoms and osteoporosis. *Drug Safety*, 24: 665-682.

Stark, A. and Madar, Z. (2002):

Phytoestrogens: a review of recent findings.
J. Pediatr. Endocrinol. Metab., 15 : 561-572.

Stearns, V., Beebe, K.L. and Dube, E. (2003):

Paroxetine controlled release in the treatment of menopausal hot flushes. *JAMA*, 289: 2827-2834.

Studd JWW (2005):

Benefits and side effects of HRT after the Women's Health initiative (WHI) and million women's study (MWS) reports. In : *Progress in obstetrics and gynaecology*, vol 16. Editor studd J.E., Elsevier, London, PP 411,412.

Stygar, D., Muravitskayan, N. and Sahlin, L. (2003):

Effect of SERM treatment on growth and proliferation in rat uterus.
Reproductive Biology and Endocrinology, 1: 40.

Tait, D., Debra, L.; Charles, L. and Adjel, M.D. (2000):

Pathophysiology and treatment of hot flushes.
Mayo. Clin. Prac., 77: 1207-1218.

Tang, G.W.K. (2004):

The climacteric of Chinese factory workers.
Maturities, 19: 1777-1782.

Tanko, L.B., Bagger, Y.Z. and Alexandersen, P. (2004):

Safety and efficacy of a novel salmon Calcitonin (sCT) technology
based oral formulation in healthy postmenopausal women.
J., 19 (9): 1531-1538.

Tavani, A., Spertini, L. and La Vecchia, C. (2006):

Intake of specific flavonoids and risk of acute myocardial
infarction in Italy. Public. Health Nutr., 9 : 369-74.

Taya, L., McMillan and Mark, M.D. (2004):

Complementary and Alternative Medicine and Physical Activity
for Menopausal Symptoms. Journal of the American Medical
Women's Association, 59: 270-277.

Tham, D.M.; Gardner, C.D. and Haskell, W.L. (1998):

Potential health benefits of dietary Phytoestrogens; review of the clinical, epidemiological, and mechanistic evidence. J. CLin. Endocr. Metab., 83: 2223-2234.

Theodoratou, E., Kyle, J. and Campbell, H. (2007):

Dietary flavonoids and the risk of colorectal cancer.
J. Cancer Epidemiol. Biomarkers Prev., 16 (4): 684-693.

Tice, J.A., Ettinger, B. and Ensrud, K. (2003):

Phytoestrogens supplements for the treatment of hot flushes. The Isoflavones Clover Extract (ICE) Study: a Randomized Controlled Trial. JAMA, 290: 207.

Touillaud, M.S., Thiebaut, A.C. and Clavel-Chapelon, F. (2007):

Dietary lignan intake and postmenopausal breast cancer risk by estrogen and progesterone receptor, status. J. Natl. Cancer Inst., Mar. 21; 99 : 475-486.

Tucker, Hannon, M.T. and Chen, H. (1999):

Magnesium, fruit and vegetables intake associated with bone mineral density in elderly women. Am. J. Clin. Nutr., 69: 727-736.

Upmalis, D.H.; Lobo, R. and Bradley, L. (2000):

Vasomotor symptoms relief by soy isoflavones extracts tablets in postmenopausal women: A multicenter, double blind, and randomized, placebo-control led study. *Menopause*, 7: 236.

Van Patten, C.L., Olivotto, L.A. and Chambers, G.K. (2002):

Effect of soy Phytoestrogens on hot flushes in postmenopausal women with breast cancer: A randomized, controlled clinical trial. *J. Clin. Oncol.*, 20: 1449.

Vergne, S., Titier, K.; Bernard, V. and Sauvant, P. (2007):

Bioavailability and urinary excretion of isoflavones in humans; effects of soy-based supplements formulation and Equol production. *J. Pharm. Biomed. Anal.*, 12, 43 : 148S-1494.

Moralli V., M.D. (2002):

Alternative therapies for traditional disease states, menopause. *American Family Physician*, 66: 129-134.

Wagner, J.D., Anthony, M.S. and Cline, J.M. (2001):

Soy Phytoestrogens: research on benefits and risks. *Clin. Obstet. Gynecol.*, 44: 843-852.

Wagner, J.D., Zhang, L. and Greaves, K.A. (2000):

Soy protein reduces the arterial low-density lipoprotein (LDL) concentration and delivery of LDL cholesterol to the arteries of diabetic and non-diabetic male cynomolgus monkeys. *Metabolism*, 49: 1185-1196.

Wang, T.T.Y., Sathyamoorthy, N. and Phang, J.M. (1999):

Molecular effects of Genistein on estrogen receptor mediated pathways. *Carcinogenesis*, 171: 271-275.

Wei, H. (1998):

Photoprotective action of isoflavones, Genistein models, mechanisms and relevance to clinical dermatology. *J. Am. Acad. Dermatol.*, 39,: 271-272.

Wendy Klein, M.D. (2000):

Alternative to traditional HRT.

Women's Health Gend Based Med., 9: 235-243.

Whiting, P.W., Clouston, A. and Kerlin, P. (2002):

Black cohosh and other herbal remedies associated with acute hepatitis. *Med. J. Australia*, 177 (8): 440-443.

Wijma, K., Melin, A. and Hammar, M. (1997):

Treatment of menopausal symptoms with applied relaxation: A pilot study. *J. Behav. Ther. Exp. Psychiatry*, 28: 251-261.

Wuttke, W., Jerry, H. and Becker, T. (2003):

Phytoestrogens: endocrine disrupters or replacement for hormone replacement therapy. *Maturitas*, 14: 44, Suppl.S9-S20.

Wyon, K., Wijma, K. and Hamrnar, M. (2004):

A comparison of acupuncture and oral estradiol treatment of vasomotor symptoms in postmenopausal women. *Climacteric*, 7 : 153-164.

Yaffe, K., Sawaya, G. and Grady, D. (1998):

Estrogen therapy in postmenopausal women: effects on cognitive function and dementia. JAMA, 279: 688-695.

Yates, A.A., Schlicker, S.A. and Suiror, C.W. (1998):

Dietary reference intakes: the new basis for recommendations for calcium and related nutrients, B vitamins, and choline. J.Am. Diet Association, 98: 699-706.

Zaborowska, E., Brynhildsen, J. and Hammar, M. (2007):

Effects of acupuncture, applied relaxation, estrogens and placebo on postmenopausal women. Climacteric the Journal of the International Menopause Society, 10 : 38-45.

Zava, D.T. and Duwe, G. (1998):

Estrogenic and antiproliferative properties and other flavonoids in human breast cancer cells in vivi. Nutr. Cancer, 27: 31-40.

Zhang, R.H.; Chen, K.J. and Ma, X.C. (2005):

A clinical study of Yigu capsule in treating postmenopausal osteoporosis. Chin. J. Integr. Med. 11 : 97-103.

