

## REFERENCES

**Alegria A, Martins E, Dias M, Cunha A, Cardoso ML and Maire I.** Glycogen storage disease type IV presenting as hydrops fetalis. *J Inherit Metab Dis* 1999; 22: 330-332

**Amalfitano A, Bengur AR, Morse RP, Majure JM, Case LE, Veerling DL, Mackey J, Kishnani P, Smith W, McVie-Wylie A, Sullivan JA, Hoganson GE, Phillips JA 3rd, Schaefer GB, Charrow J, Ware RE, Bossen EH and Chen YT.** Recombinant human acid alpha-glucosidase enzyme therapy for infantile glycogen storage disease type II: results of a phase I/II clinical trial. *Genet Med* 2001; 3: 132-138

**Amartino H, Paineira D, Pomponio RJ, Niizawa G, Sabio Paz V, Blanco M and Chamoles N.** Two clinical forms of glycogen storage disease type II in two generations of the same family. *Clin Genet* 2006; 69: 187-188

**Annabi B, Hiraiwa H, Mansfield BC, Lei KJ, Ubagai T, Polymeropoulos MH, Moses SW, Parvari R, HersHKovitz E, Mandel H, Fryman M and Chou JY.** The gene for glycogen-storage disease type 1b maps to chromosome 11q23. *Am J Hum Genet* 1998; 62: 400-405

**Arion WJ, Wallin BK, Lange AJ and Ballas LM.** On the involvement of a glucose 6-phosphate transport system in the function of microsomal glucose 6-phosphatase. *Mol Cell Biochem* 1975; 6: 75-83

**Ausems MG, Lochman P, van Diggelen OP, Ploos van Amstel HK, Reuser AJ and Wokke JH.** A diagnostic protocol for adult-onset glycogen storage disease type II. *Neurology* 1999; 52: 851-853

**Bachrach BE, Weinstein DA, Orho-Melander M, Burgess A and Wolfsdorf JI.** Glycogen synthase deficiency (glycogen storage disease type 0) presenting with hyperglycemia and glucosuria: report of three new mutations. *J Pediatr* 2002; 140: 781-783

**Bandsma RHJ, Smit GPA and Kuipers F.** Disturbed lipid metabolism in glycogen storage disease type 1. *Eur J Pediatr* 2002;161:S65-9.

**Bao Y, Yang BZ, Dawson TL Jr and Chen YT.** Isolation and nucleotide sequence of human liver glycogen debranching enzyme mRNA: identification of multiple tissue-specific isoforms. *Gene* 1997; 197: 389-398

**Barthel A and Schmoll D.** Novel concepts in insulin regulation of hepatic gluconeogenesis. *Am J Physiol Endocrinol Metab.* 2003 Oct;285(4):E685-92.

**Bernier AV, Sentner CP, Correia CE, Theriaque DW, Shuster JJ, Smit GP and Weinstein DA.** Hyperlipidemia in glycogen storage disease type III: effect of age and metabolic control. *J Inherit Metab Dis.* 2008 Dec;31(6):729-32.

**Bianchi L.** Glycogen storage disease I and hepatocellular tumours. *Eur J Pediatr* 1993; 152 Suppl 1: S63-S70

**Bodamer OA, Feillet F, Lane RE, Lee PJ, Dixon MA, Halliday D and Leonard JV.** Utilization of cornstarch in glycogen storage disease type Ia. *Eur J Gastroenterol Hepatol* 2002; 14: 1251-1256

**Brody LC, Abel KJ, Castilla LH, Couch FJ, McKinley DR, Yin G, Ho PP, Merajver S, Chandrasekharappa SC and Xu J.** Construction of a transcription map surrounding the BRCA1 locus of human chromosome 17. *Genomics* 1995; 25: 238-247

**Brown BI and Brown DH.** Lack of an alpha-1,4-glucan: alpha-1,4-glucan 6-glycosyl transferase in a case of type IV glycogenosis. *Proc Natl Acad Sci USA* 1966; 56: 725-729

**Bruno C, van Diggelen OP, Cassandrini D, Gimpelev M, Giuffre B, Donati MA, Introvini P, Alegria A, Assereto S, Morandi L, Mora M, Tonoli E, Mascelli S, Traverso M, Pasquini E, Bado M, Vilarinho L, van Noort G, Mosca F, DiMauro S, Zara F and**

**Minetti C.** Clinical and genetic heterogeneity of branching enzyme deficiency (glycogenosis type IV). *Neurology* 2004; 63: 1053-1058

**Burwinkel B, Bakker HD, Herschkovitz E, Moses SW, Shin YS and Kilimann MW.** Mutations in the liver glycogen phosphorylase gene (PYGL) underlying glycogenosis type VI. *Am J Hum Genet* 1998; 62: 785-791

**Burwinkel B, Maichele AJ, Aagenaes O, Bakker HD, Lerner A, Shin YS, Strachan JA and Kilimann MW.** Autosomal glycogenosis of liver and muscle due to phosphorylase kinase deficiency is caused by mutations in the phosphorylase kinase beta subunit (PHKB). *Hum Mol Genet* 1997; 6: 1109-1115

**Cabrera-Abreu J, Crabtree NJ, Elias E, Fraser W, Cramb R and Alger S.** Bone mineral density and markers of bone turnover in patients with glycogen storage disease types I, III and IX. *J Inherit Metab Dis* 2004; 27: 1-9

**Capristo E, Addolorato G, Mingrone G, Greco AV and Gasbarrini G.** Effect of disease localization on the anthropometric and metabolic features of Crohn's disease. *Am J Gastroenterol* 1998;93:2411-9.

**Chen YT, Coleman RA, Scheinman JI, Kolbeck PC and Sidbury JB.** Renal disease in type I glycogen storage disease. *N Engl J Med* 1988; 318: 7-11

**Chen YT. Glycogen storage disease In: Scriver CR, Beaudet AL, Valle D, Sly WS and Kinzler V.** The metabolic and molecular bases of inherited disease. 8th ed. New York: McGraw-Hill 2001; p. 1521-51.

**Chen YT.** Glycogen storage diseases and other inherited disorders of carbohydrate metabolism. In: Kasper DL, Braunwald E, Fauci AS, et al, Eds. *Harrison's Principles of Internal Medicine*. New York: McGraw-Hill. 2005: p. 16:2319-23.

**Cheung YY, Kim SY, Yiu WH, Pan CJ, Jun HS, Ruef RA, Lee EJ, Westphal H, Mansfield BC and Chou JY.** Impaired neutrophil activity and increased susceptibility to bacterial infection in mice lacking glucose-6-phosphatase-beta. *J Clin Invest.* 2007 Mar;117(3):784-93.

**Chou JY and Mansfield BC.** Mutations in the glucose-6-phosphatase-alpha (G6PC) gene that cause type Ia glycogen storage disease. *Hum Mutat.* 2008 Jul;29(7):921-30.

**Chou JY, Matern D, Mansfield BC and Chen YT.** Type I glycogen storage disease: disorders of the glucose-6-phosphatase complex. *Curr Mol Med* 2002;2:121-43.

**Cleary MA, Walter JH, Kerr BA and Wraith JE.** Facial appearance in glycogen storage disease type III. *Clin Dysmorphol* 2002; 11: 117-120

**Coleman RA, Winter HS, Wolf B and Chen YT.** Glycogen debranching enzyme deficiency: long-term study of serum enzyme activities and clinical features. *J Inherit Metab Dis* 1992; 15: 869-881

**Cori GT, and Cori CF.** Glucose-6-phosphatase of the liver in glycogen storage disease. *J Biol Chem* 1952; 199: 661-667

**Correia CE, Bhattacharya K, Lee PJ, Shuster JJ, Theriaque DW, Shankar MN, Smit GP and Weinstein DA.** Use of modified cornstarch therapy to extend fasting in glycogen storage disease types Ia and Ib. *Am J Clin Nutr.* 2008 Nov;88(5):1272-6.

**Davidson JJ, Ozcelik T, Hamacher C, Willems PJ, Francke U and Kilimann MW.** cDNA cloning of a liver isoform of the phosphorylase kinase alpha subunit and mapping of the gene to Xp22.2-p22.1, the region of human X-linked liver glycogenosis. *Proc Natl Acad Sci USA* 1992; 89: 2096-210

**Demo E, Frush D, Gottfried M, Koepke J, Boney A, Bali D, Chen YT and Kishnani PS.** Glycogen storage disease type IIIhepatocellular carcinoma a long-term complication? J Hepatol 2007; 46: 492-498

**Ding JH, de Barsy T, Brown BI, Coleman RA and Chen YT.** Immunoblot analyses of glycogen debranching enzyme in different subtypes of glycogen storage disease type III. J Pediatr 1990; 116: 95-100

**Duarte IF, Goodfellow BJ, Barros A, Jones JG, Barosa C, Diogo L, Garcia P and Gil AM.** Metabolic characterization of plasma in juveniles with glycogen storage disease type 1a (GSD1a) by high-resolution (1)H NMR spectroscopy. NMR Biomed 2006; 20: 401-412

**Dunger DB, Holder AT, Leonard JV, Okae J and Preece MA.** Growth and endocrine changes in the hepatic glycogenoses. Eur J Pediatr. 1982 May;138(3):226-30.

**Ekstein J, Rubin BY, Anderson SL, Weinstein DA, Bach G, Abeliovich D, Webb M and Risch N.** Mutation frequencies for glycogen storage disease Ia in the Ashkenazi Jewish population. Am J Med Genet A 2004; 129: 162-164

**Emmett M and Narins RG.** Renal transplantation in type 1 glycogenosis. Failure to improve glucose metabolism. JAMA 1978; 239: 1642-1644

**Fernandes J, Berger R and Smit GP.** Lactate as a cerebral metabolic fuel for glucose-6-phosphatase deficient children. Pediatr Res. 1984 Apr;18(4):335-9.

**Franco LM, Krishnamurthy V, Bali D, Weinstein DA, Arn P, Clary B, Boney A, Sullivan J, Frush DP, Chen YT and Kishnani PS.** Hepatocellular carcinoma in glycogen storage disease type Ia: a case series. J Inherit Metab Dis 2005; 28: 153-162

**Galli L, Orrico A, Marcolongo P, Fulceri R, Burchell A, Melis D, Parini R, Gatti R, Lam C, Benedetti A and Sorrentino V.** Mutations in the glucose-6-phosphate transporter (G6PT) gene in patients with glycogen storage diseases type 1b and 1c. FEBS Lett 1999; 459: 255-258

**Gianni ML, Bonvissuto M, Gallieni M, Testolin C, Racchi E and Riva E.** Ramipril treatment in a patient with glycogen storage disease I non-A. J Inherit Metab Dis 2002; 25: 515-516

**Giuffre B, Parinii R, Rizzuti T, Morandi L, van Diggelen OP, Bruno C, Giuffre M, Corsello G and Mosca F.** Severe neonatal onset of glycogenosis type IV: clinical and laboratory findings

leading to diagnosis in two siblings. *J Inherit Metab Dis* 2004; 27: 609-619

**Gogus S, Kocak N, Ciliv G, Karabulut E, Akcoren Z, Kale G and Caglar M.** Histologic features of the liver in type Ia glycogen storage disease: comparative study between different age groups and consecutive biopsies. *Pediatr Dev Pathol* 2002; 5: 299-304

**Goh BH, Khan A, Efendic S and Portwood N.** Expression of glucose-6-phosphatase system genes in murine cortex and hypothalamus. *Horm Metab Res* 2006;38:1-7.

**Greene HL, Brown BI, McClenathan DT, Agostini RM Jr and Taylor SR.** A new variant of type IV glycogenosis: deficiency of branching enzyme activity without apparent progressive liver disease. *Hepatology* 1988; 8: 302-306

**Haines JL, Ozelius LJ, McFarlane H, Menon A, Tzall S, Martiniuk F, Hirschhorn R and Gusella JF.** A genetic linkage map of chromosome 17. *Genomics* 1990; 8: 1-6

**Heller S, Worona L and Consuelo A.** Nutritional therapy for glycogen storage diseases. *J Pediatr Gastroenterol Nutr.* 2008 Aug;47 Suppl 1:S15-21.

**Hendrickx J, Coucke P, Hors-Cayla MC, Smit GP, Shin YS, Deutsch J, Smeitink J, Berger R, Lee P and Fernandes J.** Localization of a new type of X-linked liver glycogenosis to the chromosomal region Xp22 containing the liver alpha-subunit of phosphorylase kinase (PHKA2). *Genomics* 1994; 21: 620-625

**Hendrickx J and Willems PJ.** Genetic deficiencies of the glycogen phosphorylase system. *Hum Genet* 1996; 97: 551-556

**Hidaka F, Sawada H, Matsuyama M and Nuno H.** A novel mutation of the PHKA2 gene in a patient with X-linked liver glycogenosis type 1. *Pediatr Int* 2005; 47: 687-690

**Hiraiwa H, Pan CJ, Lin B, Moses SW and Chou JY.** Inactivation of the glucose 6-phosphate transporter causes glycogen storage disease type 1b. *J Biol Chem.* 1999 Feb 26;274(9):5532-6.

**Howard TD, Akots G and Bowden DW.** Physical and genetic mapping of the muscle phosphofructokinase gene (PFKM): reassignment to human chromosome 12q. *Genomics* 1996; 34: 122-127

**Hunter I, Greene SA, MacDonald TM and Morris AD.** Prevalence and aetiology of hypothyroidism in the young. *Arch Dis Child* 2000;83:207-10.

**Ihara K, Nomura A, Hikino S, Takada H and Hara T.** Quantitative analysis of glucose-6-phosphate translocase gene expression in various human tissues and haematopoietic progenitor cells. *J Inherit Metab Dis* 2000; 23: 583-592

**Koeberl DD, Kishnani PS, and Chen YT.** Glycogen storage disease types I and II: treatment updates. *J Inherit Metab Dis* 2007; 30: 159-164

**Kotb MA, Abdallah HK and Kotb A.** Liver glycogenoses: are they a possible cause of polyneuropathy? A cross-sectional study. *J Trop Pediatr.* 2004 Aug;50(4):196-202.

**Kovacevic A, Ehrlich R, Mayatepek E, Wendel U and Schwahn B.** Glycogen storage disease type Ib without hypoglycemia. *Mol Genet Metab* 2007; 90: 349-350

**Laberge AM, Mitchell GA, van de Werve G and Lambert M.** Long-term follow-up of a new case of liver glycogen synthase deficiency. *Am J Med Genet A* 2003; 120: 19-22

**Lebo RV, Anderson LA, DiMauro S, Lynch E, Hwang P and Fletterick R.** Rare McArdle disease locus polymorphic site on 11q13 contains CpG sequence. *Hum Genet* 1990; 86: 17-24

**Lee P, Burch M and Leonard JV.** Plasma creatine kinase and cardiomyopathy in glycogen storage disease type III. *J Inherit Metab Dis* 1995; 18: 751-752

**Lee PJ, Muiesan P and Heaton N.** Successful pregnancy after combined renal-hepatic transplantation in glycogen storage disease type Ia. *J Inherit Metab Dis.* 2004;27(4):537-8.

**Lee PJ, Van't Hoff WG and Leonard JV.** Catch-up growth in Fanconi-Bickel syndrome with uncooked cornstarch. *J Inherit Metab Dis* 1995; 18: 153-156

**Lei KJ, Pan CJ, Shelly LL, Liu JL and Chou JY.** Identification of mutations in the gene for glucose-6-phosphatase, the enzyme deficient in glycogen storage disease type 1a. *J Clin Invest* 1994; 93: 1994-1999

**Lei KJ, Chen H, Pan CJ, Ward JM, Mosinger B Jr, Lee EJ, Westphal H, Mansfield BC and Chou JY.** Glucose-6-phosphatase dependent substrate transport in the glycogen storage disease type-1a mouse. *Nat Genet.* 1996 Jun;13(2):203-9.

**Lei KJ, Shelly LL, Pan CJ, Sidbury JB and Chou JY.** Mutations in the glucose-6-phosphatase gene that cause glycogen storage disease type 1a. *Science.* 1993 Oct 22;262(5133):580-3.

**Leuzzi R, Banhegyi G, Kardon T, Marcolongo P, Capecchi PL, Burger HJ, Benedetti A and Fulceri R.** Inhibition of microsomal glucose-6-phosphate transport in human neutrophils results in apoptosis: a potential explanation for neutrophil dysfunction in glycogen storage disease type 1b. *Blood* 2003; 101: 2381-2387

**Lewis GM, Spencer-Peet J and Stewart KM.** Infantile hypoglycaemia due to inherited deficiency of glycogen synthetase in liver. *Arch Dis Child* 1963; 38: 40-48

**L'hermine-Coulomb A, Beuzen F, Bouvier R, Rolland MO, Froissart R, Menez F, Audibert F and Labrune P.** Fetal type IV glycogen storage disease: clinical, enzymatic, and genetic data of a pure muscular form with variable and early antenatal manifestations in the same family. *Am J Med Genet A* 2005; 139: 118-122

**Lin CC, Tsai JD, Lin SP and Lee HC.** Renal sonographic findings of type I glycogen storage disease in infancy and early childhood. *Pediatr Radiol* 2005; 35: 786-791

**Liu PP, de Villa VH, Chen YS, Wang CC, Wang SH, Chiang YC, Jawan B, Cheung HK, Cheng YF, Huang TL, Eng HL, Chuang FR and Chen CL.** Outcome of living donor liver transplantation for glycogen storage disease. *Transplant Proc* 2003; 35: 366-368

**Lossos A, Barash V, Soffer D, Argov Z, Gomori M, Ben-Nariah Z, Abramsky O and Steiner I.** Hereditary branching enzyme dysfunction in adult polyglucosan body disease: a possible metabolic cause in two patients. *Ann Neurol* 1991; 30: 655-662

**Lucchiari S, Pagliarani S, Salani S, Filocamo M, Di Rocco M, Melis D, Rodolico C, Musumeci O, Toscano A, Bresolin N and Comi GP.** Hepatic and neuromuscular forms of glycogenosis type III: nine mutations in AGL. *Hum Mutat* 2006; 27: 600-601

**Maichele AJ, Burwinkel B, Maire I, Sovik O and Kilimann MW.** Mutations in the testis/liver isoform of the phosphorylase kinase gamma subunit (PHKG2) cause autosomal liver glycogenosis in the *gsd* rat and in humans. *Nat Genet* 1996; 14: 337-340

**Manz F, Bickel H, Brodehl J, Feist D, Gellissen K, Gescholl-Bauer B, Gilli G, Harms E, Helwig H and Nutzenadel W.** Fanconi-Bickel syndrome. *Pediatr Nephrol* 1987; 1: 509-518

**Marfaing-Koka A, Wolf M, Boyer-Neumann C, Meyer D, Odievre M and Labrune P.** Increased levels of hemostatic proteins are independent of inflammation in glycogen storage disease type Ia. *J Pediatr Gastroenterol Nutr.* 2003 Nov;37(5):566-70.

**Martin AP, Bartels M, Schreiber S, Buehrdel P, Hauss J and Fangmann J.** Successful staged kidney and liver transplantation for

glycogen storage disease type Ib: A case report. *Transplant Proc* 2006; 38: 3615-3619

**Matern D, Starzl TE, Arnaout W, Barnard J, Bynon JS, Dhawan A, Emond J, Haagsma EB, Hug G, Lachaux A, Smit GP and Chen YT.** Liver transplantation for glycogen storage disease types I, III, and IV. *Eur J Pediatr* 1999; 158 Suppl 2: S43-S48

**Melis D, Fulceri R, Parenti G, Marcolongo P, Gatti R, Parini R, Riva E, Della Casa R, Zammarchi E, Andria G and Benedetti A.** Genotype/phenotype correlation in glycogen storage disease type 1b: a multicentre study and review of the literature. *Eur J Pediatr* 2005; 164: 501-508

**Melis D, Havelaar AC, Verbeek E, Smit GP, Benedetti A, Mancini GM and Verheijen F.** NPT4, a new microsomal phosphate transporter: mutation analysis in glycogen storage disease type Ic. *J Inherit Metab Dis* 2004; 27: 725-733

**Melis D, Parenti G, Della Casa R, Sibilio M, Berni Canani R and Terrin G.** Crohn's like ileo-colitis in patients affected by glycogen storage disease Ib: two years follow up of patients with a wide spectrum of gastrointestinal signs. *Acta Paediatr* 2003;92:1415-21.

**Melis D, Pivonello R, Parenti G, Della Casa R, Salerno M, Lombardi G, Sebastio G, Colao A and Andria G.** Increased

prevalence of thyroid autoimmunity and hypothyroidism in patients with glycogen storage disease type I. *J Pediatr* 2007; 150: 300-305, 305.e1

**Moses SW and Parvari R.** The variable presentations of glycogen storage disease type IV: a review of clinical, enzymatic and molecular studies. *Curr Mol Med* 2002; 2: 177-188

**Moses SW, Wanderman KL, Myroz A and Frydman M.** Cardiac involvement in glycogen storage disease type III. *Eur J Pediatr* 1989; 148: 764-766

**Muhlhausen C, Schneppenheim R, Budde U, Merkel M, Muschol N, Ullrich K and Santer R.** Decreased plasma concentration of von Willebrand factor antigen (VWF:Ag) in patients with glycogen storage disease type Ia. *J Inherit Metab Dis* 2005; 28: 945-950

**Muller D, Santer R, Krawinkel M, Christiansen B and Schaub J.** Fanconi-Bickel syndrome presenting in neonatal screening for galactosaemia. *J Inherit Metab Dis* 1997; 20: 607-608

**Mundy HR, Hindmarsh PC, Matthews DR, Leonard JV and Lee PJ.** The regulation of growth in glycogen storage disease type 1. *Clin Endocrinol (Oxf)*. 2003 Mar;58(3):332-9.

**Newgard CB, Fletterick RJ, Anderson LA and Lebo RV.** The polymorphic locus for glycogen storage disease VI (liver glycogen phosphorylase) maps to chromosome 14. *Am J Hum Genet* 1987; 40: 351-364

**Noto RA, Vijayaraghavan V, Timoshin A and Sansobrino D.** Improved growth with growth hormone therapy in a child with glycogen storage disease Ib. *Acta Paediatr* 2003;92:977-9.

**Nuoffer JM, Mullis PE and Wiesmann UN.** Treatment with low dose diazoxide in two growth-retarded prepubertal girls with glycogen storage disease type Ia resulted in catch-up growth. *J Inherit Metab Dis* 1997; 20: 790-798

**Okada S, Seino Y, Kodama H, Yutaka T, Inui K, Ishida M, Yabuuchi H and Seino Y.** Insulin and glucagon secretion in hepatic glycogenoses. *Acta Paediatr Scand.* 1979 Sep;68(5):735-8.

**Okasaki K, Funato M, Kashima M, Nakama K, Inoue T and Hiura M.** Twenty-six-week repeat-dose toxicity study of a recombinant human granulocyte colony-stimulating factor derivative (nartograstim) in cynomolgus monkeys. *Toxicol Sci* 2002;65:246-55.

**Orho M, Bosshard NU, Buist NR, Gitzelmann R, Aynsley-Green A, Blumel P, Gannon MC, Nuttall FQ and Groop LC.** Mutations in the liver glycogen synthase gene in children with hypoglycemia due to glycogen storage disease type 0. *J Clin Invest* 1998; 102: 507-515

**Ozen H.** Glycogen storage diseases: new perspectives. *World J Gastroenterol.* 2007 May 14;13(18):2541-53

**Pabuccuoglu A, Aydogdu S and Bas M.** Serum biotinidase activity in children with chronic liver disease and its clinical significance. *J Pediatr Gastroenterol Nutr* 2002; 34: 59-62

**Pan CJ, Lin B and Chou JY.** Transmembrane topology of human glucose 6-phosphate transporter. *J Biol Chem.* 1999 May 14;274(20):13865-9.

**Parvari R, Moses S, Shen J, HersHKovitz E, Lerner A and Chen YT.** A single-base deletion in the 3'-coding region of glycogen debranching enzyme is prevalent in glycogen storage disease type IIIA in a population of North African Jewish patients. *Eur J Hum Genet* 1997; 5: 266-270.

**Pasqualini T, Ferraris JR and Jasper H:** Differences in anthropometric parameters and the IFG-I-IGFBP3 axis between liver and renal transplant children. *Transplantation* 70:472, 2000.

**Pomorski L, Bartos M, Matejkowska M and Kuzdak K.** The influence of a single acetic acid and lactic acid injection on rat normal thyroid tissue. *Cesk Patol* 2002;38:107-12.

**Rake JP, Visser G, Labrune P, Leonard JV, Ullrich K and Smit GP.** Glycogen storage disease type I: diagnosis, management, clinical course and outcome. Results of the European Study on Glycogen Storage Disease Type I (ESGSD I). *Eur J Pediatr* 2002; 161 Suppl 1: S20-S34

**Reddy SK, Austin SL, Spencer-Manzon M, Koeberl DD, Clary BM, Desai DM, Smith AD and Kishnani PS.** Liver transplantation for glycogen storage disease type Ia. *J Hepatol.* 2009 Sep;51(3):483-90

**Reitsma-Bierens WC.** Renal complications in glycogen storage disease type I. *Eur J Pediatr* 1993; 152 Suppl 1: S60-S62

**Roach PJ.** Glycogen and its metabolism. *Curr Mol Med* 2002; 2: 101-120

**Roe TF, Coates TD, Thomas DW, Miller JH and Gilsanz V.** Brief report: treatment of chronic inflammatory bowel disease in glycogen storage disease type Ib with colony-stimulating factors. *N Engl J Med* 1992; 326: 1666-1669

**Ryan IP, Havel RJ and Laros RK Jr.** Three consecutive pregnancies in a patient with glycogen storage disease type IA (von Gierke's disease). *Am J Obstet Gynecol* 1994; 170: 1687-1690; discussion 1690-1691

**Safouh H, Elsayyed R, Mansour S, Taha G, Elshabrawy M and Elkaraksy H.** Renal involvement in children with glycogen-storage disease. *Pediatrics* 2008 121: S143-b-S144

**Saltik IN, Ozen H, Ciliv G, Kocak N, Yuce A, Gurakan F and Dinler G.** Glycogen storage disease type Ia: frequency and clinical course in Turkish children. *Indian J Pediatr* 2000; 67: 497-501

**Saltik-Temizel IN, Coskun T, Yuce A and Kocak N.** Fanconi-Bickel syndrome in three Turkish patients with different homozygous mutations. *Turk J Pediatr* 2005; 47: 167-169

**Santer R, Kinner M, Steuerwald U, Kjærgaard S, Skovby F, Simonsen H, Shaiu WL, Chen YT, Schneppenheim R and Schaub J.** Molecular genetic basis and prevalence of glycogen storage disease type IIIA in the Faroe Islands. *Eur J Hum Genet* 2001; 9: 388-391

**Santer R, Schneppenheim R, Dombrowski A, Gotze H, Steinmann B and Schaub J.** Mutations in GLUT2, the gene for the liver-type glucose transporter, in patients with Fanconi-Bickel syndrome. *Nat Genet* 1997; 17: 324-326

**Schippers HM, Smit GP, Rake JP and Visser G.** Characteristic growth pattern in male X-linked phosphorylase-b kinase deficiency (GSD IX). *J Inherit Metab Dis* 2003; 26: 43-47

**Schroder JM, May R, Shin YS, Sigmund M and Nase-Huppmeier S.** Juvenile hereditary polyglucosan body disease with complete branching enzyme deficiency (type IV glycogenosis). *Acta Neuropathol (Berl)* 1993; 85: 419-430

**Servidei S, Riepe RE, Langston C, Tani LY, Bricker JT, Crisp-Lindgren N, Travers H, Armstrong D and DiMauro S.** Severe cardiopathy in branching enzyme deficiency. *J Pediatr* 1987; 111: 51-56

**Shen J, Bao Y, Liu HM, Lee P, Leonard JV and Chen YT.** Mutations in exon 3 of the glycogen debranching enzyme gene are associated with glycogen storage disease type III that is differentially expressed in liver and muscle. *J Clin Invest* 1996; 98: 352-357

**Shen JJ and Chen YT.** Molecular characterization of glycogen storage disease type III. *Curr Mol Med* 2002; 2: 167-175

**Shin YS.** Glycogen storage disease: clinical, biochemical, and molecular heterogeneity. *Semin Pediatr Neurol.* 2006 Jun;13(2):115-20.

**Slonim AE, Bulone L, Ritz S, Goldberg T, Chen A and Martiniuk F.** Identification of two subtypes of infantile acid maltase deficiency. *J Pediatr* 2000; 137: 283-285

**Sloop KW, Cao JX, Siesky AM, Zhang HY, Bodenmiller DM, Cox AL, Jacobs SJ, Moyers JS, Owens RA, Showalter AD, Brenner MB, Raap A, Gromada J, Berridge BR, Monteith DK, Porksen N, McKay RA, Monia BP, Bhanot S, Watts LM and Michael MD.** Hepatic and glucagon-like peptide-1-mediated reversal of diabetes by glucagon receptor antisense oligonucleotide inhibitors. *J Clin Invest.* 2004 Jun;113(11):1571-81.

**Spiegel R, Mahamid J, Orho-Melander M, Miron D and Horovitz Y.** The variable clinical phenotype of liver glycogen synthase deficiency. *J Pediatr Endocrinol Metab.* 2007 Dec;20(12):1339-42.

**Starzl TE, Demetris AJ, Trucco M, Ricordi C, Ildstad S, Terasaki PI, Murase N, Kendall RS, Kocova M and Rudert WA.** Chimerism after liver transplantation for type IV glycogen storage disease and type 1 Gaucher's disease. *N Engl J Med* 1993; 328: 745-749

**Talente GM, Coleman RA, Alter C, Baker L, Brown BI, Cannon RA, Chen YT, Crigler JF Jr, Ferreira P, Haworth JC, Herman GE, Issenman RM, Keating JP, Linde R, Roe TF, Senior B and Wolfsdorf JI.** Glycogen storage disease in adults. *Ann Intern Med* 1994; 120: 218-226

**Tang NL, Hui J, Young E, Worthington V, To KF, Cheung KL, Li CK and Fok TF.** A novel mutation (G233D) in the glycogen phosphorylase gene in a patient with hepatic glycogen storage disease and residual enzyme activity. *Mol Genet Metab* 2003; 79: 142-145

**Terzioglu M, Emre S, Ozen H, Saltik IN, Kocak N, Ciliv G, Yuce A and Gurakan F.** Glucose-6-phosphatase gene mutations in Turkish patients with glycogen storage disease type Ia. *J Inherit Metab Dis* 2001; 24: 881-882

**Toscano A and Musumeci O.** Tarui disease and distal glycogenoses: clinical and genetic update. *Acta Myol.* 2007 Oct;26(2):105-7.

**Trioche P, Francoual J, Capel L, Odievre M, Lindenbaum A and Labrune P.** Apolipoprotein E polymorphism and serum concentrations in patients with glycogen storage disease type Ia. *J Inherit Metab Dis* 2000; 23: 107-112

**van Schaftingen E and Gerin I.** The glucose-6-phosphatase system. *Biochem J* 2002; 362: 513-532

**van Schaftingen E and Gerin I.** The glucose-6-phosphatase system. *Biochem J.* 2002 Mar 15;362(Pt 3):513-32.

**Veiga-da-Cunha M, Gerin I, Chen YT, Lee PJ, Leonard JV, Maire I, Wendel U, Vikkula M and Van Schaftingen E.** The putative glucose 6-phosphate translocase gene is mutated in essentially all cases of glycogen storage disease type I non-a. *Eur J Hum Genet* 1999; 7: 717-723

**Visser G, Rake JP, Fernandes J, Labrune P, Leonard JV, Moses S, Ullrich K and Smit GP.** Neutropenia, neutrophil dysfunction, and inflammatory bowel disease in glycogen storage disease type Ib: results of the European Study on Glycogen Storage Disease type I. *J Pediatr* 2000; 137: 187-191

**Visser G, Rake JP, Kokke FT, Nikkels PG, Sauer PJ and Smit GP.** Intestinal function in glycogen storage disease type I. *J Inher Metab Dis* 2002; 25: 261-267

**Weinstein DA, Correia CE, Saunders AC and Wolfsdorf JI.** Hepatic glycogen synthase deficiency: an infrequently recognized cause of ketotic hypoglycemia. *Mol Genet Metab* 2006; 87: 284-288

**Wolfsdorf JI, Holm IA, and Weinstein DA.** Glycogen storage diseases. Phenotypic, genetic, and biochemical characteristics, and therapy. *Endocrinol Metab Clin North Am* 1999;28:801-23.

**Wolfsdorf JI and Weinstein DA.** Glycogen storage diseases. *Rev Endocr Metab Disord.* 2003 Mar;4(1):95-102.

**Yiu WH, Pan CJ, Allamarvdasht M, Kim SY and Chou JY.** Glucose-6-phosphate transporter gene therapy corrects metabolic and myeloid abnormalities in glycogen storage disease type Ib mice. *Gene Ther* 2007; 14: 219-226

**Zhang H, Cook J, Nickel J, Yu R, Stecker K, Myers K and Dean NM.** Reduction of liver Fas expression by an antisense oligonucleotide protects mice from fulminant hepatitis. *Nat Biotechnol.* 2000 Aug;18(8):862-7.