

References

- **Aldrete JA (1994):**
Discharge Criteria, Bailliere's.
Clinical Anesthesiology; 8:763-77.
- **Allen JG and Sury MR (2006):**
Sedation of children undergoing magnetic resonance
Imaging.
BJA; 97: 898–9.
- **American Academy of Pediatrics Committee on Drugs (2006):**
Guidelines for maintaining and management of pediatric
patients during and after sedation for diagnostic and
therapeutic procedures.
Pediatrics; 118:2587–2602.
- **American Society of Anesthesiologists (2002):**
Practice guidelines for sedation and analgesia by non-
anesthesiologists: a report by the American Society of
Anesthesiologists Task Force on sedation and analgesia by
non-anesthesiologists.
Anesthesiology; 96:1004–1017.
- **Anker JN (1996):**
Pharmacokinetics and renal function in preterm infants.
Acta Pediatr; 85:1393.

- **Ansell G. (1968):**
National survey of radiological complication.
Clin. Radio. 19,175 191.
- **Ansell G. (1970):**
Adverse reaction to contrast agents .Scope of problem.
Invest.radiol.5, 374-384.
- **Beauve B and Dearlove O (2008):**
Sedation of children under 4 weeks of age examination.
Pediatric Anesthesia; 18: 891–913.
- **Bell Ch (2002):**
Anesthesia in MRI suites.
Anesth. analge; 74: 508-17.
- **Berman WJ, Fripp RR and Rubler M (1990):**
Hemodynamic effects of ketamine in children undergoing
cardiac catheterization.
Pediatric Cardiol 11:72,1990.
- **Bettmann MA.(2004):**
Frequently asked questions: Iodinated contrast agents.
Radiographics 2004; 24:S3-10.
- **Betts K. (1977):**
Correspondence –nitrous oxide encephalography.
Anesth.Analg. 56, (3), 469.
- **Bissonnette B (1992):**
Temperature monitoring in pediatric anesthesia.
Int Anesthesiol Clin; 30:63–76.

- **Blitman NM, Lee HK and Jain VR (2007):**
Pulmonary atelectasis in children anesthetized for
cardiothoracic MR: evaluation of risk factors.
J Comput Assist Tomogr; 31:789–794.
- **Boidin Mp, Wolff GR and Doleman C (2002a):**
The technique of MRI under general anesthesia.
Top magn reson imaging; 14(1):19-30.
- **Boidin MP, Wolff GR and Doelman C (2002b):**
Sedation versus general anesthesia in MRI.
BJA; 84(6): 743-748
- **Bordes M, Semjen F and Sautereau A (2007):**
Which anesthesia for children undergoing MRI?
Ann Fr Anesth Reanim(English abstract); 26:287–291.
- **Cherny KW, Watson ML and Field SG. (2007):**
Aspiration pneumonitis requiring intubation after procedural
sedation and analgesia: a case report.
Ann Emerg Med; 49: 462–464.
- **Chinoy MR (2003):**
Lung growth and development.
Front Biosci; 8:D392–D415.
- **Chung SM. (2002):**
Safety issues in magnetic resonance imaging.
J Neuroophthalmol; 22:35-39.

- **Connolly L.A.(1996):**
Anesthesia for transjugular intrahepatic portosystemic shunt placement.
Anesthesiology 85. (4):946-947 1996.
- **Cote´ CJ (2005):**
Pediatric Anesthesia from. Miller RD., (2005).
Anesthesia. 6th edition. Churchill Livingstone; ch 60:2370-2376.
- **Cote´ CJ (2008):**
Round and round we go: sedation – what is it, who does it, and have we made things safer for children?
Pediatr Anesth; 18:3–8.
- **Coté CJ and Wilson S (2006):**
Guidelines for Monitoring and Management of Pediatric Patients During and After Sedation for Diagnostic and Therapeutic Procedures.
An Update Pediatrics; 118 (6): 2587-2602.
- **Cravero J P and Blike G T (2004):**
Review of Pediatric Sedation.
Anesth Analg; 99:1355-1364.
- **Cravero JP, Blike GT, Beach M, Gallagher SM, Hertzog JH, Havidich JE, Gelman B, and the Pediatric Sedation Research Consortium (2006):**
Pediatric Sedation Research Consortium. Incidence and nature of adverse events during pediatric

- sedation/anesthesia for procedures outside the operating room:
report from the Pediatric Sedation Research Consortium.
Pediatrics; 118(3):1087-1096.
- **Dalal P G, Murray D, Thomas Cox, McAllister J, and Snider R (2006):**
Sedation and Anesthesia Protocols Used for Magnetic Resonance Imaging Studies in Infants.
Provider and Pharmacologic Considerations. Anesth Analg; 103:863-868.
 - **David Gozala and Yaacov Gozal (2008):**
Pediatric sedation/anesthesia outside the operating room.
Current opinion in anesthesiology; 21:494-498.
 - **Dearlove O and Corcoran JP (2007):**
Sedation of children undergoing magnetic resonance imaging.
BJA; 98: 548–9.
 - **Dixon RL and Ekstrand KE(1999):**
Review basic spin physics.
Magn Reson Imaging Clin N Am; 7(4):607-27.
 - **D`Orsogna L,Sandor GGS,and Patterson MWH.(1989):**
Influence of echocardiography in preoperative cardiac catheterization in congenital heart disease.
Int J Cardio24:19,1989.

- **Dukes MNG. (1996):**
Radiological contrast media and radiopharmaceuticals . In
Dukes MNG (ed):
Meyler side effects of drugs.vol 13.New York
,Elsevier,1996,p 1382
- **Edelman R and Warach S (1993):**
Magnetic resonance imaging.
N Engle J Med; 328:708-16.
- **Feldman J M and Kalli I (2006):**
Equipment and environmental issues for none operating
room anesthesia.
Current Opinion in Anesthesiology; 19:450–452.
- **Fitzgerald M. (1991):**
The developmental neurobiology of pain . In :bond
MR,Charlton JE,Woolf CJ eds.Proceedings of the Vth World
Congress on Pain (pain research and clinical manegment
,vol. 4); 1990 Apr 1-6;Adelaide South Australia .
Amsterdam : Elsevier 1991: 253-61.
- **Forestner JE (1984):**
anesthesia for radiologic procedures .In Murphy C, Murphy
M (eds) Radiology for anesthesia and critical care.
New YORK Churchill Livingstone ,1987,p 302.
- **Gangarosa RE, Minnis JE and Nobbe J. (1997):**
operational safety issues in MRI Magnetic resonance

imaging; 15:287-292.

- **Goldberg M.(1984):**
Systemic reaction to intravascular contrast media.
Anesthesiology 60:46, 1984.
- **Gold farb S,Spinler S and Berns JS.(1993):**
low-osmolality contrast media and the risk of contrast
associated nephrotoxicity:
Invest Radiol 28:S7.1993.
- **Gow PJ, Ghabrial H and Smallwood RA (2001):**
Neonatal hepatic drug elimination.
Pharmacol Toxicol; 88:3–15.
- **Gozal D and Gozal Y (2008):**
Pediatric sedation/anesthesia outside the operating room.
Current Opinion in Anesthesiology; 21:494–498.
- **Green SM, Roback MG and Miner JR. (2007):**
Fasting and emergency department procedural sedation and
analgesia: a consensus-based clinical practice advisory.
Ann Emerg Med; 49:454–461.
- **Gregory GA (2005):**
Resuscitation of the Newborn from: Miller RD., (2005):
Anesthesia. 6th edition.
Churchill Livingstone; ch 59:2348-2353.
- **Heard CM, Joshi P and Johnson K (2007):**
Dexmedetomidine for pediatric MRI sedation: a review of a

series of cases.

Pediatr Anesth; 17: 888–892.

- **Holzman RS, Cullen DJ Eichhorn JH (1994):**
Guidelines for sedation by anesthesiologists during
diagnostic and therapeutic procedures .
JClin Anesth 6:265, 1994.
- **Hopkins LN.(1993):**
Endovascular approach:Role of neurosurgery.
Semin Intervent Radiol 10:189.1993.
- **Hornak J P (2008):**
the Basics of MRI.
4th edition ch.3:35-39,<http://www.cis.rit.edu/htbook/mri/>
visited in 2008.
- **Huang H K. (1987):**
Elements of digital radiology.A professional handbook and
guide.
NJ: Prentice-Hall.
- **Hubbard AM and Fellows KE.(1993):**
Pediatric interventional radiology:current practice and
innovations.
Cardiovascular Intervent Radiol 16.267-274 .1993.
- **ICRP.(1990):**
International Commission on Radiological Protection. 1990
Recommendations of the International Commission on
Radiological Protection.1991.

- Annals of the ICRP, Publication 60, Oxford: Pergamon.
- **Itai Shavit, Llan Keidan, Arie Augarten (2006):**
The practice of pediatric procedural sedation and analgesia in the emergency department.
European journal of emergency medicine; 13;270-275.
 - **Joint Commission on Accreditation of Healthcare Organization (2001):**
Anesthesia and Sedation.
Joint Commission Resources, Oakbrook Terrace,
<http://www.Jointcomission.org/> retrieved August 18, 2008.
 - **Jones DG, Shrimpton P C. (1991):**
Survey of CT practice in the UK Part 3: Normalised organ doses calculated using Monte Carlo techniques.
National Radiological Protection Board, Chilton, United Kingdom; NRPB-R250.
 - **Jones M, Leslie K and Mitchell P. (2004):**
Anesthesia for endovascular treatment of cerebral aneurysms.
Clin Neurosci 11. (5):468-470.2004.
 - **Jose PA, Fildes RD and Gomez RA (1994):**
Neonatal renal function and physiology.
Curr Opin Pediatr; 6:172
 - **Kanal E, Borgstede JP and Barkovich AJ. (2004):**
American College of Radiology white paper on MR safety: 2004 update and revisions.

Am J Roentgenol; 182:1111–1114.

- **Karian VE, Burrows PE and Zurakowski D. (2002):**
The development of a pediatric radiology sedation program.
Pediatr Radiol; 32: 348–53.
- **Keeter S., Benator RM, Weinberg SM. (1990):**
Sedation in pediatric CT: National survey of current practice
Radiology 175:745, 1990.
- **Kemper J, Klocke A, Kahl-Nieke B and Adam G (2005):**
Orthodontic brackets in high field MR imaging:
experimental evaluation of magnetic field interactions Tesla.
Fortschr Geb Rontgenstr Nuklearmed (English abstract);
177:1691–1698.
- **Kim G, Green SM and Denmark TK. (2003):**
Ventilatory response during dissociative sedation in children:
a pilot study.
Acad Emerg Med; 10: 140–5.
- **Kirberg A, Sagredo R and Montalva G. (2005):**
Ketamine for pediatric endoscopic procedures and as a
sedation complement for adult patients.
Gastrointest Endosc; 61:501–502.
- **Koroglu A, Demirbilek S, Teksan H, Sagir O, But AK, Ersoy OM (2005):**
Sedative, haemodynamic and respiratory effects of

dexmedetomidine in children undergoing magnetic resonance imaging examination: Preliminary results.
BJA; 94: 821–4.

- **Koroglu A, Teskan H, Sagir O, Yucel A, Toprak HI, and Ersoy OM (2006):**

A comparison of the sedative, hemodynamic and respiratory effects of dexmedetomidine and propofol in children undergoing magnetic resonance imaging.

Anesth Analg; 103: 63-7.

- **Krasin MJ,Xiong A, and Wu S.(2005):**

The effects of external beam irradiation on the growth of flat bones in children :modeling a dose-volume effects .

Int J. Radiation Oncology Biol Phys . 2005; 62:1458-1463.

- **Krauss B and Green SM (2006):**

Procedural sedation and analgesia in children.

Lancet; 367: 766–80.

- **Krauss B and Green SM (2008):**

Training and credentialing in procedural sedation and analgesia in children: lessons from the United States model.

Pediatr Anesth; 18:30–35.

- **Kuchta KF. (2003):**

Endovascular abdominal aortic aneurysm repair .

In J Cardiothorac Vasc Anesth 7(2): 205-211.2003.

- **Lucinda I. Everett (2001):**
anesthesia for diagnostic procedures.
Current opinion in anesthesiology; 14:629-633.
- **Lutterbey G, Wattjes MP and Doerr D. (2007):**
Atelectasis in children undergoing either propofol infusion
or positive pressure ventilation anesthesia for magnetic
resonance imaging.
Pediatr. Anesth; 17:121–125.
- **Machata AM, Willschke H, Kabon B, Kettner SC and
Marhofer P (2008):**
Propofol-based sedation regimen for infants and children
undergoing ambulatory magnetic resonance imaging.
BJA; 101 (2): 239–43.
- **Malviya S, Lewis T V, Eldevik O P Rockwell DT, Wong
JH, Tait AR. (2000):**
sedation and general anesthesia in children undergoing MRI
and CT: adverse events and outcomes.
BJA; 84(6): 743-8.
- **Mandell GA, Majd M, Shalaby-Rana EI and Gordon I
(2003):**
Procedure guideline for pediatric sedation in nuclear
medicine, 3.0. Reston (VA).
Society of Nuclear Medicine; 25: 5 p.
- **Manner JM and Codman VA.(1969):**
General anesthesia for cardiac catheterization in children:the

effect of spontaneous or controlled ventilation on the evaluation of congenital abnormalities.

Anesthesia 24:541,1969.

- **Mason K P (2008):**

The pediatric sedation service: who is appropriate to sedate? which medications I should use, who should prescribe the drugs, and how do I bill.

Pediatr Radiol ; 38 (2): 218–224.

- **Mason KP, Michna E and Dinardo JA. (2002):**

Evolution of a protocol for Ketamine-induced sedation as an alternative to general anesthesia for interventional radiological procedures in pediatric patients.

Radiology; 225:457–465.

- **Mason KP, Sanborn P and Zurakowski D. (2004a):**
Superiority of pentobarbital versus chloral hydrate for sedation in infants during imaging.

Radiology; 230:537–542.

- **Mason KP, Zurakowski D and Connor L. (2004b):**

Infant sedation for MR imaging and CT: oral versus intravenous pentobarbital.

Radiology; 233:723–728.

- **Mason K P, Zurakowski D, Zgleszewski S E, Robson CD, Carrier M, Hickey PR and Dinadro JA (2008):**

High dose dexmedetomidine as the sole sedative for

pediatric MRI.

Pediatric Anesthesia; 18: 403–411.

- **Merten GJ Yucel A and Toprak HI. (2004):**
prevention of contrast-induced nephrology with sodium carbonat :A ranomised ,controlled trial.
JAMA 2004; 291:2328-34.
- **Meschi M , Dinadro JA and Lewis T V.(2006):**
Facts and fallacies concerning the prevention of contrast medium- induced nephropathy.
Crit Care Med 2006; 34:2060-8.
- **Morgan GE, Mikhail MS and Murray MJ (2005):**
Clinical Anesthesiology.
Fourth edition. Lange McGraw-Hill.p:850.
- **Morris C G, McBrien M E, and Farling P A (2004):**
Anticipation and planning for anaesthesia provision within MR units in the UK Br.
J. Radiol; 77(915): 216 – 218.
- **Mueller PR,Wittenberg KH ,Kaufman JA (1997):**
Pattern of anesthesia and nursing care for interventional radiology procedures A national survey of physician practices and preferences .
Radiology 202:339, 1997.
- **Mulhern RK ,Palmer SL and Merchant TE.(2005):**
Neurocognitive consequences of risk-adapted therapy for

childhood medullblastoma .

J Clin Oncol.2005; 23:5511-5519..

- **Nixon C, Hirsch NP, Ormerod IEC and Johnson G (1986):**

Nuclear magnetic resonance.

Anesthesia; 41:131-137.

- **Osborn I P (2002):**

Magnetic resonance imaging anesthesia: new challenges and techniques.

Curr. Opin Anaesthesiol; 15:443-448.

- **Pearson G. A. (2002):**

Hand book of pediatric intensive care.

1st edition; ch 12:169-170.

- **Peden CJ, Menon DK and Hall AS. (1992):**

Magnetic resonance for the anaesthetist. Part II: anaesthesia and monitoring in MR units.

Anaesthesia; 47:508-517.

- **Pershad J, Wan J and Anghelescu DL (2007):**

Comparison of Propofol with

Pentobarbital/Midazolam/Fentanyl Sedation for Magnetic Resonance Imaging of the Brain in Children.

Pediatric; 120(3): 629-636.

- **Phan H and Nahata M C (2008):**

Clinical Uses of Dexmedetomidine in Pediatric Patients.

Pediatr Drugs; 10 (1): 49-69.

- **Ramsay MAE, Savage TM, Simpson BRJ. (1974):**
Controlled sedation with alphaxolone-alphadolone.
BMJ; 2:256.
- **Ronchera-Oms CL, Casillas C and Marti-Bonmati L. (2004):**
Oral chloral hydrate provides effective and safe sedation in
pediatric magnetic resonance imaging.
J Clin Pharm Ther; 19:239–243.
- **Rudolph AM (1970):**
The changes in the circulation after birth: Their importance
in congenital heart disease.
Circulation; 41:343- 359.
- **Rudolph AM and Heymann MA (1974):**
Fetal and neonatal circulation and respiration.
Annu Rev Physiol; 36:187.
- **Sanborn PA, Michna E and Zurakowski D. (2005):**
Adverse cardiovascular and respiratory events during
sedation of pediatric patients for imaging examinations.
Radiology; 237:288–94.
- **Serafini G and Zadra N (2008):**
Anesthesia for MRI in the pediatric patient.
Current Opinion in Anesthesiology; 21:499–503.
- **Sesay M, Tauzin-Fin P, Verdonck O, Dousset V and**

Maurette P (2008):

A wireless remote controlled infusion pump for anesthesia during magnetic resonance imaging.

BJA; 100(6):862-863.

- **Shankar W and Deshpande JK (2005):**

Procedural sedation in the pediatric patient.

Anesthesiol Clin North Am; 23:635–654.

- **Shavit I, Keidan I, Hoffmann Y, Mishuk L, Rubin O, Ziv A, Steiner IP. (2007):**

Enhancing patient safety during pediatric sedation: the impact of simulation-based training of non anesthesiologists.

Arch Pediatr Adolesc Med; 161(8):740-3.

- **Shrimpton P C, Jones D G, Hillier M C, Wall B F, Le Heron J C, Faulkner K. (1991):**

Survey of CT practice in the UK Part 2: Dosimetric aspects.

National Radiological Protection Board, Chilton, United Kingdom 1991; NRPB-R250.

- **Shukry M and Ramadhyani U (2005):**

Dexmedetomidine as the primary sedative agent for brain radiation therapy in a 21 month old child.

Pediatr Anesth; 15: 241-2.

- **Slater JD, Rossi CJ and Yonemoto LT.(2004):**

Proton therapy for prostate cancer: the initial Linda University experience.

- Int J Radiat Oncol Biol Phys. 2004; 59:348-352.
- **Soto RG, Fu ES and Vila H. (2004):**
Capnography accurately detects apnea during monitored anesthesia care.
Anesth Analg; 99:379–382.
 - **Stanger P, Heymann MA and Tarnoff H. (1974):**
Complication of cardiac catheterization of neonates, infant and children : A three-year study.
Circulation 50:595, 1974.
 - **Steward DJ (2005):**
manual of pediatric anesthesia, 6th edition.
Churchill livingstone. P:115.
 - **Stonell CA, Leslie K and Absalom AR (2006):**
Effect-site targeted patient-controlled sedation with propofol: comparison with anesthetist administration for colonoscopy.
Anesthesia; 61:240–247.
 - **Sundaram R, Brown AG and Koteeswaran SK. (2006):**
Anesthetic implication of uterine artery embolisation in management of massive obstetric haemorrhage.
Anesthesia 61. (3):248-252. 2006.
 - **Sury MR, Harker H and Begent J. (2005):**
The management of infants and children for painless imaging.
Clin Radiol; 60:731–41.

- **Taghon TA, Bryan YF, Kurth CD. (2006):**
Pediatric radiology sedation and anesthesia.
Int Anesthesiol Clin. ; **44(1):65-79.**
- **Tepel M, Kurth CD and Begent J(2000):**
Prevention of radiographic-contrast-agent-induced reduction
in renal function by acetyl cysteine.
N Engl J Med 2000;343:180-4.
- **Tobias JD (2007):**
Dexmedetomidine: application in pediatric critical care and
pediatric anesthesiology.
Pediatr Crit Care Med; 8:115–131.
- **Usher AG, Kearney RA and Tsui BC (2005):**
Propofol total intravenous anesthesia for MRI in children.
Pediatr Anesth; 15: 23–8.
- **Wappler F (2006):**
The propofol infusion syndrome [in German].
Deutsches Arzteblatt; 11:A705-A710.
- **Watkinson A.F, Francis I.S,and Torrie P.(2002):**
Commentary: the role of anesthesia in interventional
radiology.
Br J Radiol 75. 105 -106 .2002;Citation
- **Westbrook C and Kaut C (1993):**
MRI in practice.
1st edition Blackwell Science Inc US; ch.2:13-18.

- **White PF (1998):**
Ketamine update: its clinical uses in anesthesia.
Semin Anesthesia; 7:113–126.
- **Yamamoto L G (2008):**
Initiating a Hospital-Wide Pediatric Sedation Service
Provided by Emergency Physicians.
Clinical Pediatrics; 47(1): 37-48.
- **Young WL and Pile –Spellman J(1994):**
Anesthetic consideration for interventional neurosurgery.
Anesthesiology 80:427, 1994.
- **Zgleszewski S E, Zurakowski D and Fontaine P J. (2008):**
Is Propofol a Safe Alternative to Pentobarbital for Sedation
during Pediatric Diagnostic CT?
Radiology; 247(2): 528 – 534.