

Agarwal SS, Kumar L, Chavali KH and Mestri SC (2009).

Fatal Venous Air Embolism Following Intravenous Infusion.

Journal of Forensic Sciences: Volume 54, Number 3, May , pp. 682-684(3).

Agnelli G (2004).

Prevention of venous thromboembolism in surgical patients.

Circulation; 110(24 Suppl 1):IV4-12.

Agnelli G, Becattini C, Kirschstein T (2002).

Thrombolysis vs heparin in the treatment of pulmonary embolism: a clinical outcome-based meta-analysis.

Arch Intern Med;162:2537–2541.

Aklog L, Williams CS, Byrne JG, Goldhaber SZ (2002).

Acute pulmonary embolectomy: a contemporary approach.

Circulation;105:1416–1419.

Alikhan R, Peters F, Wilmott R, Cohen AT (2004).

Fatal pulmonary embolism in hospitalized patients: a necropsy review.

J Clin Pathol ;57:1254–1257.

Amaragiri SV and Lees T (2000).

Elastic compression stockings for prevention of deep vein thrombosis.

Cochrane Database of Systematic Reviews; Issue 1. Art. No.: CD001484. DOI: 10.1002/14651858.CD001484.

Andrew M, Michelson AD, Bovill E, Leaker M, Massicotte MP (1998).

Guidelines for antithrombotic therapy in pediatric patients.

J Pediatr;132:575-88.

Anne M, Jennifer A, Labiris R, James D et al. (2005).

Systematic Overview of Warfarin and Its Drug and Food Interactions
ARCH INTERN MED/VOL 165, MAY 23.

Ansell J, Hirsh J, Poller L, Bussey H, Jacobson A et al. (2004).

The Pharmacology and Management of the Vitamin K Antagonists.
Chest;126;204S-233S.

Ardissino D, Van de Werf F, Betriu A et al. (2003).

Management of acute myocardial infarction in patients presenting with ST-segment elevation: the Task Force on the Management of Acute Myocardial Infarction of the European Society of Cardiology.
Eur Heart J;24:28-66.

Arnold GJ, Azarnoff D, Barbutto JP, Branch RA, Campbell RK (2001).

A Comprehensive Reference for Generic and Brand Prescription Drugs. In: Schreffer J, ed. St. Louis, MO: Harcourt Health Sciences Company.

Augustinos P and Ouriel K (2004).

Invasive Approaches to Treatment of Venous Thromboembolism.
Circulation;110;I-27-I-34.

Bates S, Weitz J, Johnston M et al. (2001).

Use of a fixed activated partial thromboplastin time ratio to establish a therapeutic range for unfractionated heparin.
Arch Intern Med; 161:385–391

Bauer K, Turpie A, Eriksson B, Lassen M (2003).

Steering Committees of the Pentasaccharide Orthopedic Prophylaxis Studies.
Efficacy and safety of fondaparinux in major orthopedic surgery according to the timing of its first administration.
Thromb Haemost; 90:364–6

Bauer KA (2003).

New pentasaccharides for prophylaxis of deep vein thrombosis: pharmacology.
Chest, 124:364S–370S.

Bax J, Betriu A, Crea F, Falk V, Filippatos G et al. (2008).

Management of acute myocardial infarction in patients presenting with persistent ST-segment elevation.
European Heart Journal 29, 2909–2945.

Bousser MG, Bouthier J, Büller HR et al. (2008).

Comparison of idraparinux with vitamin K antagonists for prevention of thromboembolism in patients with atrial fibrillation.
Lancet 371 (9609): 315–21.

Brenner DJ and Hall EJ (2007).

Computed tomography:an increasing source of radiation exposure.
N Engl J Med;357:2277–2284.

Brotman DJ, Jaffer AK, Hurbanek JG, Morra N (2004).

Warfarin prophylaxis and venous thromboembolism in the first 5 days following hip and knee arthroplasty.
J Thromb Haemost; 92:1012.

Brunot S, Corneloup O, Latrabe V, Montaudon M, Laurent F (2005).

Reproducibility of multi-detector spiral computed tomography in detection of sub-segmental acute pulmonary embolism.
Eur Radiol;15:2057–2063.

Buller H, Agnelli G, Hull R, Hyers T, Prins M, Raskob G (2004).

Antithrombotic therapy for venous thromboembolic disease: the Seventh ACCP Conference on Antithrombotic and Thrombolytic Therapy.
Chest; 126: 401S–28S

Buller HR, Davidson BL, Decousus H et al. (2004).

Fondaparinux or enoxaparin for the initial treatment of symptomatic deep venous thrombosis: a randomized trial.

Ann Intern Med, 140:867–73.

Buller HR ,Van Belle A , Huisman MV, Huisman PM et al. (2006).

Effectiveness of managing suspected pulmonary embolism using an algorithm combining clinical probability, D-dimer testing, and computed tomography.

JAMA;295:172–179.

Casazza F, Bongarzone A, Capozzi A, Agostoni O (2005).

Regional right ventricular dysfunction in acute pulmonary embolism and right ventricular infarction.

Eur J Echocardiogr;6:11–14.

Cindy HL, Hankey GJ, Eikelboom JW (2005).

Venous thromboembolism: diagnosis and management of pulmonary embolism

MJA; 182: 569–574.

Chan YC, Valenti D, Mansfield AO, Stansby G (2000).

Warfarin induced skin necrosis.

Br J Surg 87 (3): 266–72.

Christiansen SC, Cannegieter SC, Koster T, Vandenbroucke JP (2005).

Thrombophilia, clinical factors and recurrent venous thrombotic events.

JAMA; 293:2352–61.

Cohen A, Heit J, Anderson FJ (2005).

Estimated annual number of incident and recurrent, non-fatal and fatal venous thromboembolism (VTE) events in the US.

Blood.;106:267A.

Cohen AT, Davidson BL, Gallus AS, Lassen MR et al. (2006).

Efficacy and safety of fondaparinux for the prevention of venous thromboembolism in older acute medical patients: randomised placebo controlled trial.

BMJ; 332:325–9.

Cohen AT, Tapson VF, Bergmann JF, Goldhaber SZ, Kakkar AK et al.(2008).

Venous thromboembolism risk and prophylaxis in the acute hospital care setting (ENDORSE study): a multinational cross-sectional study.

Lancet 371:387–394.

Crapo RO, Jensen RL, Wanger JS (2001).

Single-breath carbon monoxide diffusing capacity

Clin Chest Med.;22:637–649.

Cushman M, Albert W, Tsai, Richard H, White G et al. (2004).

Deep Vein Thrombosis and Pulmonary Embolism in Two Cohorts: The Longitudinal Investigation of Thromboembolism Etiology.

Am J Med.;117:19–25.

Cushman M, Glynn, RJ, Goldhaber, SZ et al. (2006).

Hormonal factors and risk of recurrent venous thrombosis: the prevention of recurrent venous thromboembolism trial.

J Thromb Haemost; 4:2199.

Delaney JA, Opatrny L, Brophy JM, Suissa S (2007).

Drug drug interactions between antithrombotic medications and the risk of gastrointestinal bleeding.

CMAJ 177 (4): 347–51

Di Nisio M, Sohne M, Kamphuisen PW, Buller HR (2005).

D-Dimer test in cancer patients with suspected acute pulmonary embolism.

J Thromb Haemost;3:1239–1242.

Di Nisio M, Squizzato A, Rutjes AW, Buller HR et al. (2007).

Diagnostic accuracy of D-dimer test for exclusion of venous thromboembolism:
a systematic review.

J Thromb Haemost;5:296–304.

Donat F, Duret JP, Santoni A et al. (2002).

The pharmacokinetics of fondaparinux sodium in healthy volunteers.

Clin Pharmacokinet, 41(Suppl 2):1–9.

Edwin JR, Harry RB , Matthijs O (2009).

Deep Vein Thrombosis and Pulmonary Embolism.

Thrombolysis for the Treatment of Pulmonary Embolism. pages 504-505

published by John Wiley & Sons, Ltd. ISBN: 978-0-470-51717-8

Elliott CG, Goldhaber SZ, Visani L, DeRosa M (2000).

Chest radiographs in acute pulmonary embolism. Results from the International
Cooperative Pulmonary Embolism Registry.

Chest;118:33–38

Eriksson BI, Borris LC, Friedman RJ et al. (2008)

Rivaroxaban versus enoxaparin for thromboprophylaxis after hip arthroplasty. N Engl
J Med.;358(26):2765-2775

Ferrari E, Benhamou M, Berthier F, Baudouy M (2005).

Mobile thrombi of the right heart in pulmonary embolism: delayed disappearance
after thrombolytic treatment.

Chest;127:1051–1053.

Finkelstein Y, Chitayat D, Schechter T, Keating S et al. (2005).

Warfarin embryopathy following low dose maternal exposure.

J Obstet Gynaecol Can;27(7):702-706.

Folsom AR (2007).

Update on factor V Leiden association with venous thromboembolism in the LITE Study.

Blood;109:1336–1337.

Francis CW (2007).

Prophylaxis for thromboembolism in hospitalized medical patients.

N Engl J Med ;356:1438-44. (Erratum, N Engl J Med 2007;357:203).

Geerts WH, Pineo GF, Heit JA, et al. (2004).

Prevention of venous thromboembolism: the Seventh ACCP Conference on Antithrombotic and Thrombolytic Therapy.

Chest; 126:338S.

Geibel A, Zehender M, Kasper W, Olschewski M (2005).

Prognostic value of the ECG on admission in patients with acute major pulmonary embolism.

Eur Respir J;25:843–848.

Gillet JL, Perrin, MR, Allaert FA (2007).

Short-term and mid-term outcome of isolated symptomatic muscular calf vein thrombosis.

J Vasc Surg; 46:513.

Ginsberg JS and McRae SJ (2005).

New anticoagulants for the prevention and treatment of venous thromboembolism.

Vascular Health and Risk Management:1(1) 41–53.

Goldhaber SZ (2002).

Echocardiography in the management of pulmonary embolism

Ann Intern Med. ;136:691–700.

Goldhaber SZ and Elliott CG (2003).

Acute Pulmonary Embolism: Part I: Epidemiology, Pathophysiology, and diagnosis.
Circulation.;108;2726-2729.

Goldhaber SZ and Elliott CG (2003).

Acute Pulmonary Embolism: Part I: Epidemiology, Pathophysiology, and diagnosis.
Circulation;108;2726-2729.

Goldhaber SZ and Tapson VF (2004).

A prospective registry of 5,451 patients with ultrasound-confirmed deep vein thrombosis.

Am J Cardiol; 93: 259–62

Goodman LR, Stein PD, Matta F, Sostman HD, Wakefield TW et al. (2007).

CT venography and compression sonography are diagnostically equivalent:
data from PIOPED II.

AJR Am J Roentgenol;189:1071–1076.

Gourlay T and LiEl-Ali KM (2006).

Assessment of the risk of systemic fat mobilization and fat embolism as a
consequence of liposuction: ex vivo study.

Plast Reconstr Surg , 117(7):2269-76.

Grassi CJ, Swan TL, Cardella JF et al. (2003).

Quality improvement guidelines for percutaneous permanent inferior vena cava filter
placement for the prevention of pulmonary embolism.

J Vasc Interv Radiol;14:S271-S275.

Green B, Greenwood M, Saltissi D, Westhuyzen J et al. (2004).

Dosing strategy for enoxaparin in patients with renal impairment presenting with
acute coronary syndromes.

Br J Clin Pharmacol; 59: 281-90.

Heit J, Petterson T, Farmer S, Bailey K, Melton L (2006).

Trends in Indicence of deep vein thrombosis and pulmonary embolism: a 35-year population-based study.

Blood.;108:430a.

Heit JA, Cohen AT, Anderson FA (2005).

VTE Impact Assessment Group. Estimated annual number of incident and recurrent, non-fatal and fatal venous thromboembolism (VTE) events in the U. S.

Blood ; 106:267.

Heit JA, O’FallonWM, Petterson TM, Lohse CM, Silverstein MD (2002).

Relative impact of risk factors for deep vein thrombosis and pulmonary embolism: a population-based study.

Arch Intern Med ;162:1245–124.

Hirsh J, Fuster V, Ansell J, Halperin J (2003).

American Heart Association/American College of Cardiology Foundation Guide to Warfarin Therapy.

Circulation.;107;1692-1711

Hirsh J, Kenneth A , Maria B, Donati , Gould M, Meyer M et al. (2008).

Parenteral Anticoagulants: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines (8th Edition)

Chest;133;141S-159S

Hoppe H (2009).

Optional Vena Cava Filters:Indications, Management, and Results.

Dtsch Arztebl Int; 106(24): 395–402

Horstkotte D, Follath F, Gutschik E, Lengyel M, Oto A, Pavie A et al. (2004).

Guidelineson prevention, diagnosis and treatment of infective endocarditis executive summary.

Eur Heart J ;25:267–276.

Hovens MM, Snoep JD, Tamsma JT, Huisman MV (2006).

Aspirin in the prevention and treatment of venous thromboembolism.

J Thromb Haemost; 4:1470

Howarth DM, Lan L, Thomas PA, Allen LW (1999).

99mTc technegas ventilation and perfusion lung scintigraphy for the diagnosis of pulmonary embolus.

J Nucl Med;40:579–584.

Hsiao SH, Chang SM, Lee CY, Yang SH et al. (2006).

Usefulness of tissue Doppler parameters for identifying pulmonary embolism in patients with signs of pulmonary hypertension.

Am J Cardiol;98:685–690.

Itti E, Nguyen S, Robin F et al. (2002).

Distribution of ventilation/perfusion ratios in pulmonary embolism: an adjunct to the interpretation of ventilation/perfusion lung scans.

J Nucl Med.;43:1596–1602

Jack H and John H (1996).

Management of Deep Vein Thrombosis and Pulmonary Embolism.

Circulation.;93:2212-2245.

Jardin F, Dubourg O, Bourdarias JP (1997).

Echocardiographic pattern of acute cor pulmonale .

Chest ;111:209–217.

Jay RL and Wellington KH (2005).

Prevention of Venous Thromboembolic Disease After Total Hip and Knee Arthroplasty.

J Bone Joint Surg Am.;87:2097-2112. doi:10.2106/JBJS.D.03033

Jeffrey IW (1997).

Low molecular weight heparin.

The New England Journal of Medicine; September 4 ,Volume 337, Number 10.

John AH (2008).

The Epidemiology of Venous Thromboembolism in the Community.

Arterioscler Thromb Vasc Biol;28;370-372.

Jun ZJ, Ping T, Lei Y, Li L, Ming SY, Jing W (2006).

Prevalence of factor V Leiden and prothrombin G20210A mutations in Chinese patients with deep venous thrombosis and pulmonary embolism.

Clin Lab Haematol;28:111–116. [PubMed: 16630215].

Kaboli P , Henderson MC, White RH (2003).

DVT prophylaxis and anticoagulation in the surgical patient.

Med Clin N Am; 87:77-110.

Kahn SR, Panju A, Geerts W, Pineo GF, Turpie AG et al. (2007).

CURVE study investigators. Multicenter evaluation of the use of venous thromboembolism prophylaxis in acutely ill medical patients in Canada.

Thromb Res; 119:145–55

Kalla K, Christ G, Karnik R, Malzer R, Norman G, Prachar H et al. (2006).

Implementation of guidelines improves the standard of care: the Viennese registry on reperfusion strategies in ST-elevation myocardial infarction (Vienna STEMI registry).

Circulation;113:2398–2405.

Karcz M and James ES (2009).

Perioperative Venous Thromboembolism Prevention and Treatment.

The New York State Society of Anesthesiologists;; 61 (2):48-51

Kaufman JA, Kinney TB, Streiff MB et al. (2006).

Guidelines for the use of retrievable and convertible vena cava filters: report from the Society of Interventional Radiology multidisciplinary consensus conference.

J Vasc Interv Radiol; 17: 449–59.

Kawada T, Kitagawa H, Hoson M et al. (2000).

Clinical application of argatroban as an alternative anticoagulant for extracorporeal circulation.

Hematol Oncol Clin No Amer.;14:445-457.

Kearon C (2003).

Natural history of venous thromboembolism.

Circulation;107 (Suppl 1): I22-I30.

Kearon C, Ginsberg JS, Douketis J, Turpie AG, Bates SM et al. (2006).

An evaluation of D-dimer in the diagnosis of pulmonary embolism: a randomized trial.

Ann Intern Med;144:812–821.

Kearon C, Ginsberg JS, Julian J et al. (2006).

Comparison of fixed dose weight adjusted unfractionated heparin and low molecular weight heparin for acute treatment of venous thromboembolism.

JAMA; 296:935–942.

King CS, Holley AB, Jackson JL et al. (2007).

Twice vs three times daily heparin dosing for thromboembolism prophylaxis in the general medical population.

Chest; 131:507.

Kinney TB (2003).

Update on inferior vena cava filters.

J Vasc Interv Radiol;14:425-440.

Kline JA, Hernandez-Nino J, Jones AE (2007).

Tenecteplase to treat pulmonary embolism in the emergency department.

J Thromb Thrombolysis;23:101–105.

Koh L, Ng H, Lee L (2003).

Successful control of postsurgical bleeding by recombinant factor VIIa in a renal failure patient given low molecular weight heparin and aspirin.

Ann Hematol; 82:257–258.

Kruip MJ, Leclercq MG, van der HC, Prins MH, Buller HR (2003).

Diagnostic strategies for excluding pulmonary embolism in clinical outcome studies.

A systematic review.

Ann Intern Med;138:941–951.

Kucher N, Printzen G, Doernhoefer T et al. (2003).

Low pro-brain natriuretic peptide levels predict benign clinical outcome in acute pulmonary embolism.

Circulation. ;107:1576–1578.

Kucher N, Rossi E, De Rosa M, Goldhaber SZ (2006).

Massive pulmonary embolism.

Circulation. Jan 31;113(4):577-82.

Kucher N, Windecker S, Banz Y, Schmitz-Rode T, Mettler D et al. (2005).

Percutaneous catheter thrombectomy device for acute pulmonary embolism: in vitro and in vivo testing.

Radiology;236:852–858.

Layish DT and Tapson VF (1997).

Pharmacologic hemodynamic support in massive pulmonary embolism.

Chest;111:218–224.

Leacche M, Unic D, Goldhaber SZ, Aranki SF, Couper GS et al. (2005).

Modern surgical treatment of massive pulmonary embolism: results in 47 consecutive patients after rapid diagnosis and aggressive surgical approach.

J Thorac Cardiovasc Surg;129:1018–1023.

Le Gal G, Righini M, Roy PM, Sanchez O et al. (2006).

Prediction of pulmonary embolism in the emergency department: the revised Geneva score.

Ann Intern Med;144:165–171.

Le Gal G, Righini M, Sanchez O, Baba-Ahmed M, Perrier A et al. (2006).

A positive compression ultrasonography of the lower limb veins is highly predictive of pulmonary embolism on computed tomography in suspected patients.

Thromb Haemost;95:963–966.

Lutsey PL, Cushman M, Steffen LM, Green D, Barr RG et al. (2006).

Plasma hemostatic factors and endothelial markers in four racial/ethnic groups: the MESA study.

J Thromb Haemost;4:2629–2635.

Macina, Orest T, Schardein, James L (2007).

"Warfarin". Human Developmental Toxicants. Boca Raton:

CRC Taylor & Francis. pp. 193–4.

Macintyre N, Crapo RO, Viegi G et al. (2005).

Standardisation of the single-breath determination of carbon monoxide uptake in the lung.

Eur. Respir. J. 26 (4): 720–35

Markel A, Manzo RA, Strandness DE (1992).

The potential role of thrombolytic therapy in venous thrombosis.

Arch Intern Med;152:1265-7

Masotti L, Righini M, Vuilleumier N, Antonelli F et al. (2009).

Prognostic stratification of acute pulmonary embolism.

Vascular Health and Risk Management :5 567–575.

Mayer E, HoeperMM, Simonneau G,Rubin L (2006).

Chronic thromboembolic pulmonary hypertension.

Circulation;113:2011–2020.

Mellor A and Soni N (2001).

Fat embolism.

Anaesthesia ;56:145–154.

Mirski MA, Lele AV, Fitzsimmons L, Toung TJ (2007).

Diagnosis and treatment of vascular air embolism.

Anesthesiology;106:164–177.

Munin MC, Rudy TE, Glynn NW, Crossett LS, Rubash HE (1998).

Early inpatient rehabilitation after elective hip and knee arthroplasty.

JAMA; 279:847-52.

Musset D ,Parent F, Meyer G et al. (2002).

diagnostic strategy for patients suspected with pulmonary embolism aprospective multicentre outcome study.

lancet 2002 Dec 14;360(9349):1914-20.

O'Shea A and Eappen S (2007).

Amniotic fluid embolism.

Int Anesthesiol Clin ;45:17–28.

Pengo V, Lensing AW, Prins MH et al. (2004).

Incidence of chronic thromboembolic pulmonary hypertension after pulmonary embolism.

N Engl J Med; 350: 2257-2264.

Perrier A, Roy PM, Sanchez O, Le Gal G, Meyer G et al. (2005).

Multidetector-row computed tomography in suspected pulmonary embolism.
N Engl J Med;352:1760–1768.

Pierre-Justin G and Pierard LA (2005).

Management of mobile right heart thrombi: a prospective series.
Int J Cardiol;99:381–388.

Piovella F, Wang CJ, Lu H, Lee K, Lee LH, Lee WC et al. (2005).

Deep-vein thrombosis rates after major orthopedic surgery in Asia. An epidemiological study based on postoperative screening with centrally adjudicated bilateral venography.
J Thromb Haemost;3:2664–2670. [PubMed: 16359505]

Prandoni P, Carnovali M, Marchiori A et al. (2004).

Subcutaneous adjusted-dose unfractionated heparin vs fixed dose low molecular weight heparin in the initial treatment of venous thromboembolism.
Arch Intern Med; 164:1077–1083.

Pupka A and Szyber PP (2007).

The treatment of deep vein thrombosis in iliofemoral segment- The athletic injuries.
Medicina Sportiva, Vol. 11 (1): 17-19.

Quinlan D, McQuillan A, Eikelboom J (2004).

Low-molecular-weight heparin compared with intravenous unfractionated heparin for treatment of pulmonary embolism.
Ann Intern Med; 140:175–183

Reinartz P, Kaiser HJ, Wildberger JE, Gordji C et al. (2006).

SPECT imaging in the diagnosis of pulmonary embolism: automated detection of match and mismatch defects by means of image-processing techniques.
J Nucl Med;47:968–973.

Richard HW (2003).

The Epidemiology of Venous Thromboembolism.
Circulation;107;I-4-I-8.

Riedel M (2000).

Diagnosing pulmonary embolism.
Postgraduate Medical Journal, 80, 309–319.

Riedel M, Gibson GJ, Geddes DM, Costabel U et al. (2003).

Pulmonary emboli disease.
Respiratory medicine.London: Saunders,:1711–58.

Righini M, Le Gal G, Aujesky D, Roy PM, Sanchez O et al. (2008).

Diagnosis of pulmonary embolism by multidetector CT alone or combined with venous ultrasonography of the leg: a randomised non-inferiority trial.
Lancet;371:1343–1352.

Righini M, Le Gal G, De Lucia S, Roy PM, Meyer G et al. (2006).

Clinical usefulness of D-dimer testing in cancer patients with suspected pulmonary embolism.
Thromb Haemost;95:715–719.

Robert L. Levine, David M , Marcie JH (2006).

How Frequently Is Venous Thromboembolism in Heparin-Treated Patients Associated With Heparin-Induced Thrombocytopenia?
Chest;130;681-687.

Rodger LB (2002).

Oral antithrombotic therapy:warfarin and related compound. Page:351-352
Disorders of thrombosis and hemostasis: clinical and laboratory practice.
3 rd edition Published by Lippincott Williams& Wilkins.

Rodger M, Makropoulos D, Turek M, Quevillon J et al. (2000).

Diagnostic value of the electrocardiogram in suspected pulmonary embolism.

Am J Cardiol;86:807–9, A10.

Rodger MA, Maser E, Stiell I, Howley HE, Wells PS (2005).

The interobserver reliability of pretest probability assessment in patients with suspected pulmonary embolism.

Thromb Res;116:101–107.

Ros E,Ramos JL, Alós R, Espín E, Rico P et al. (2009).

Prevention of venous thromboembolic disease in general surgery

Cirugía españolaJun;85 Suppl 1:51-1.

Rosendaal FR (2005).

Venous thrombosis: the role of genes, environment, and behavior.

Hematology Am Soc Hematol Educ Program :1-12.

Rossum DV (2001)

Diagnostic strategies for suspected pulmonary embolism.

Eur Respir J ; 18: 589–597.

Roy PM, Colombet I, Durieux P, Chatellier G et al. (2005).

Systematic review and meta-analysis of strategies for the diagnosis of suspected pulmonary embolism. Br Med J;331:259.

Roy PM, Meyer G, Vielle B, Le Gall C, Verschuren F, Carpentier F et al. (2006).

Appropriateness of diagnostic management and outcomes of suspected pulmonary embolism.

Ann Intern Med;144:157–164.

Runyon MS, Webb WB, Jones AE, Kline JA (2005).

Comparison of the unstructured clinician estimate of pretest probability for pulmonary embolism to the Canadian score and the Charlotte rule.

Acad Emerg Med;12:587–593.

Schulman S, Lindmarker P, Holmström M, Larfars S et al. (2006).

Post-thrombotic syndrome, recurrence, and death 10 years after the first episode of venous thromboembolism treated with warfarin for 6 weeks or 6 months.

J Thromb Haemost.;4:732–742.

Sergysels R (1994) .

Pulmonary gas exchange abnormalities in pulmonary embolism. In: Morpurgo M, ed. Pulmonary Embolism.

New York: Marcel Dekker,. p89–96.

Sevransky JE, Levy MM, Marini JJ (2004).

Mechanical ventilation in sepsis-induced acute lung injury/acute respiratory distress syndrome: an evidence-based review.

Crit Care Med;32(11 Suppl.):S548–S553.

Sivakumaran M, Coles C, Simpson K and Coleman B (1997).

Treatment of warfarin intolerance.

Postgrad Med J. September; 73(863): 607.

Smulders YM (2000).

Pathophysiology and treatment of haemodynamic instability in acute pulmonary embolism: the pivotal role of pulmonary vasoconstriction.

Cardiovasc Res ;48:23–33.

Snow V, Qaseem A, Barry P, Hornbake R, Rodnick JE et al. (2007).

Management of Venous Thromboembolism: A Clinical Practice Guideline from the American College of Physicians and the American Academy of Family Physicians

Ann Intern Med.;146:204-210.

Sorensen B and Ingerslev J (2006).

A direct thrombin inhibitor studied by dynamic whole blood clot formation: haemostatic response to ex-vivo addition of recombinant factor VIIa or activated prothrombin complex concentrate.

Thromb Haemost; 96:446–453.

Sostman HD, Stein PD, Gottschalk A, Matta F, Hull R, Goodman L (2008).

Acute pulmonary embolism: sensitivity and specificity of ventilation-perfusion scintigraphy in PIOPED II study.

Radiology;246:941–946.

Stein PD, Fowler SE, Goodman LR, Gottschalk A, Hales CA et al. (2006).

Multi-detector computed tomography for acute pulmonary embolism.

N Engl J Med;354:2317–2327.

Stein PD, Hull RD, Patel KC, Olson RE, GhaliWA, Brant R et al. (2004).

D-dimer for the exclusion of acute venous thrombosis and pulmonary embolism: a systematic review.

Ann Intern Med;140:589–602.

Stein PD, Kayali F, Olson RE, Milford CE (2004).

Pulmonary thromboembolism in Asians/Pacific Islanders in the United States: analysis of data from the National Hospital Discharge Survey and the United States Bureau of the Census.

Am J Med;116:435–442. [PubMed: 15047032]

Steiner S, Schwalen A, Heintzen MP, Strauer BE (2003).

Catheter embolization into the pulmonary artery. Interventional therapy of a rare form of pulmonary embolism.

Pneumologie ;57:19–21.

Strebel N, Prins M, Agnelli G, Buller HR (2002).

Preoperative or postoperative start of prophylaxis for venous thromboembolism with low-molecular-weight heparin in elective hip surgery?

Arch Intern Med;162:1451–6.

Streiff MB, Kinney TB, Kaufman JA, Sing RF, Proctor MC et al. (2007).

Guidelines for the Use of Retrievable and Convertible Vena Cava Filters.

World J Surg 31: 251–264.

Szold O, Khoury W, Biderman P, Klausner JM, Halpern P et al. (2006).

Inhaled nitric oxide improves pulmonary functions following massive pulmonary embolism: a report of four patients and review of the literature.

Lung;184:1–5.

Talmadge DB and Spyropoulos AC (2003).

Purple toes syndrome associated with warfarin therapy in a patient with antiphospholipid syndrome.

Pharmacotherapy 23 (5): 674–7

Tapson VF (2008).

Acute Pulmonary Embolism.

N Engl J Med ;358:1037-52.

Ten Wolde M, Hagen PJ, MacGillavry MR, Pollen IJ et al. (2004).

Non-invasive diagnostic work-up of patients with clinically suspected pulmonary embolism: results of a management study.

J Thromb Haemost;2:1110–1117.

Terese T, Horlocker and John A (1997).

low Molecular Weight Heparin: Biochemistry, Pharmacology, Perioperative Prophylaxis Regimens, and Guidelines for Regional Anesthetic Management.

Anesth analg;85:874&85.

The Matisse Investigators (2002).

Fondaparinux (Arixtra®) in comparison to low molecular weight heparin for the initial treatment of symptomatic deep venous thrombosis or pulmonary embolism.

The Matisse clinical outcome studies.

Blood.;100:83a.

Torbicki A, Chairperson ,Perrier A, Konstantinides S et al.(2008).

Guidelines on the diagnosis and management of acute pulmonary embolism

European Heart Journal doi:10.1093/eurheartj/ehn310 29, 2276–2315.

Torbicki A, Galie N, Covezzoli A, Rossi E, De Rosa M, Goldhaber SZ (2003).

Right heart thrombi in pulmonary embolism: results from the International Cooperative Pulmonary Embolism Registry.

J Am Coll Cardiol;41:2245–2251.

Toschi V, Lettino M, Gallo R et al. (1996).

Biochemistry and biology of hirudin.

Coron Artery Dis; 7:420–428.

Toung TJ, Rossberg MI, Hutchins GM (2001).

Volume of air in a lethal venous air embolism.

Anesthesiology;94:360–361.

Tran H, McRae S, Ginsberg J (2006).

Anticoagulant Treatment of Deep Vein Thrombosis and Pulmonary Embolism.

Clin Geriatr Med 22 113–134.

Turpie AG (2007).

Extended duration of thromboprophylaxis in acutely ill medical patients: optimizing therapy? J Thromb Haemost; 5:5–11.

Turpie AG (2008).

Selective factor Xa inhibition with fondaparinux: from concept to clinical benefit.
European Heart Journal Supplements (2008) 10 (Supplement C), C1–C7.

Turpie AG, Chin BS, Lip GY (2002).

Venous thromboembolism: pathophysiology, clinical features, and prevention.
BMJ; 325:887.

Uflacker R (2001).

Interventional therapy for pulmonary embolism.
J Vasc Interv Radiol;12:147–164.

Van Dongen C, van den Belt A, Prins M et al. (2004).

Fixed dose subcutaneous low molecular weight heparins versus adjusted dose unfractionated heparin for venous thromboembolism.
Cochrane Database Syst Rev; 18:DC01100.

Van Strijen MJ, de Monye W, Kieft GJ, Pattynama PM et al. (2005).

Accuracy of single-detector spiral CT in the diagnosis of pulmonary embolism.
J Thromb Haemost;3:17–25.

Vedantham S, Vesely TM, Parti N et al. (2002).

Lower extremity venous thrombolysis with adjunctive mechanical thrombectomy. J
Vasc Interv Radiol.;13:1001–1008.

Victor F. Tapson (2008).

Acute Pulmonary Embolism .
New England Journal of Medicine ;358:1037-52.

Wan S, Quinlan DJ, Agnelli G, Eikelboom JW (2004).

Thrombolysis compared with heparin for the initial treatment of pulmonary embolism:
a meta-analysis of the randomized controlled trials.
Circulation;110:744–749.

Weitz J (1997).

Low molecular weight heparins.

The New England Journal of Medicine, September 4, 337:688

Weitz J, Young E, Johnston M et al. (1999).

Vasoflux, a new anticoagulant with a novel mechanism of action.

Circulation; 99:682–689.

Weitz J I, Hirsh J and Samama MM (2004).

New Anticoagulant Drugs.

Chest;126;265S-286S.

Wells PS, Anderson DR, Rodger M, Ginsberg JS et al. (2000).

Derivation of a simple clinical model to categorize patients probability of pulmonary embolism: increasing the models utility with the SimpliRED D-dimer.

Thromb Haemost;83:416–420.

Wicki J, Perneger TV, Junod AF, Bounameaux H, Perrier A (2001).

Assessing clinical probability of pulmonary embolism in the emergency ward: a simple score.

Arch Intern Med;161:92–97.

Wildberger JE, Reinartz P, Schaefer W, Nowak B et al. (2004).

Tomographic imaging in the diagnosis of pulmonary embolism: a comparison between V/Q lung scintigraphy in SPECT technique and multislice spiral CT.

J Nucl Med;45:1501–1508.

Winer-Muram HT, Rydberg J, Johnson MS, Tarver RD et al. (2004).

Suspected acute pulmonary embolism: evaluation with multi-detector row CT versus digital subtraction pulmonary arteriography.

Radiology;233:806–815.

White RH (2003).

The epidemiology of venous thromboembolism.

Circulation, 107:14-18.

White RH, Gettner S, Newman JM, Trauner KB, Romano PS (2000).

Predictor of rehospitalization for symptomatic venous thromboembolism after total hip arthroplasty.

N Engl J Med; 343:1758-64.

White RH, Romano PS, Zhou H (2001).

A population-based comparison of the 3 month incidence of thromboembolism after major elective/urgent surgery.

J Thromb Haemost; 86:2255. [PubMed]

White RH, Zhou H, Kim J et al.(2000).

A population-based study of the effectiveness of inferior vena cava filter use among patients with venous thromboembolism.

Arch Intern Med; 160:2033-2041

Wolde M, Tulevski II, Mulder JW et al. (2003).

Brain natriuretic peptide as a predictor of adverse outcome in patients with pulmonary embolism.

Circulation.;107:2082–2084.

Wood K E (2002).

Major Pulmonary Embolism . Review of a pathophysiologic approach to the golden hour of hemodynamically significant pulmonary embolism.

Chest.121.3.877;121;877-905.

Yalamanchili K, Fleisher AG, Lehrman SG, Lafaro RJ et al. (2004).

Open pulmonary embolectomy for treatment of major pulmonary embolism.

Ann Thorac Surg;77:819–823.