

6-REFERENCES

- Abderrahman, W. A., Boder, T. A., Kahn, A. U. and Ajward, M. H. (1991).** Weather modification impact on reference evapotranspiration, soil salinity and desertification in arid regions. A case study (1991). *Journal of Arid Environments*, 1991, 20: 3, 277-286.
- Abdel-Salam , A. Z. E., El-seaudy, A. A., Auad, H. Y. and Gibely, R.H. (2005).** Evaluation of genetic diversity in some cotton genotypes through yield components, biochemical and molecular analysis. *Egypt J. Genet. Cytol.* 34: 219-236
- Abdel-Tawab, F. M., Rashed, M. A., Hewedy, A. M. Abdel Aziz S. H. and Aziz M.H.(2003)** Molecular fingerprinting of ten tomato cultivars(*Lycopersicon esculentum* Mill.). *Egypt J. Genet. Cytol.* 32: 101-117.
- AL-Jibouri, A. A. M. and Adham, K. M. (1990).** Biochemical classification date palm cultivars *J. Hort. Sci.* 65: 725- 729.
- A.O. A. C. (1975).** Official methods of analysis 12thed. Association of official analytical chemists, Washington, D. C., U. S. A.

- Bai, C.S., Lu, G.Y.(1991).** Effects of water stress on tomato (*L. esculentum* Mill.) Acta-Horticulture Sinica.1991,18: 4, 340-344.
- Bates, L. S. , Waldren and Teare I. D. (1973).** Rapid determination of free proline for water-stress studies. Plant and Soil 39: 205-207.
- Ben-Asher, J., Phene, C. J. and Kinarti, A. (1992).** Canopy temperature to assess daily evapotranspiration and management of high frequency drip irrigation systems. Agri.-Water-Management. 1992, 22: (4) 379-390.
- Bhatt, R. P., Biswas, V. R., Pandey, H, K., Verma, G. S. and Kumar, N. (1998).** Heteosis for V.C in tomato *L. esculentum*. Indian Journal of Agricultural Sciences 68: (3), 176-178.
- Bonnema, . B., Castillo, C., Reiter, N., Cunningham, M., Adams, H. P. and Oconnel, M (1995).** Molecular and ultra structural analysis of a nonchromosomal variegated mutant leaf development plant physiology. Rock, MD: American Society of Plant Physiologists, 1926- Oct 1995. v. 109 (2) P. 35-392.
- Boyer, J. S. and Mcpherson, H. G. (1975).** Physiology of water deficits in cereal crops. Adv. Agron. 27: 1-23.

- Bray, E. A., Coken, A., Plant, A. J., Moses, M. S. Jmai, R. and Griffiths, A. (1997).** Regulation of gene expression by endogenous ABA in tomato plants. In Molecular biology of plants under environmental stress. Acta Physiologiae Plantarum, 19: (4), 405-418.
- Briggs, N. F. and Knowles, P.F. (1977).** Introduction to plant breeding Reinhold Publishing Corporation. U.S.A..
- Chakrabarti, A. K. and Chattopadhyay, N. C. (1995).** Identification of some Indian tomato cultivars by polyacrylamide gel electrophoresis of seed proteins. Seed- Research 1992, Publ. 1993, 20: (1), 10-13.
- Chen, R. D. and Tabaeizadeh, Z. (1992a).** Alteration of gene expression in tomato plants (*L. esculentum*) by drought and salt stress. Genome. Ottawa. National Research Council of Canada. June 1992 35: (3), 385-391.
- Chen, R. D., and Tabaeizadeh, Z. (1992b).** Expression and molecular cloning of drought induced genes in the wild tomato *L. chilense*. Biochemistry and Cell Biology 70:(3-4) 199-206.
- Chen, R. D., Yu, L. X., Greer, A. F., Cheriti, H. and Tabaeizadeh, Z. (1994).** Isolation of an osmotic

stress and abscisic acid induced gene encoding an acidic endochitinase from *Lycopersicon chilense* MGG, Mol. Gent. Berlin, Germany v. 245. P 195-202.

Cherry, J. H. (1977). Differences in protein from normal and non-ripening tomatoes by acrylamidegel electrophoresis. Hortscience 12: 413.

Chloupek, O. and Rod, J. (1992). The root system as a selection criterion. Plant- Breeding -Abs. 1992, 62: (12) 1337-1341

Cohen, D. (1982). Water relation of cultivated tomato (*L. esculentum*) and two relative species (*solanum pennellii* corr., *Lycopersicon cheesmanii* Riley). Msc Thesis Faculty of Agriculture. H. U. J. Rehovot. (Hebrew with English summary).

Culter, J. M., Rains, D. W. and Loomis, R. S. (1977a). The importance of cell size in the water relations of plants. Physiol. Plant 40: 255-260.

Culter, J. M., Rains, D. W. and Loomis, R. S. (1977b). Role of changes in solute concentration in maintaining favorable water balance in field-grown cotton. Agron. J., 69, 773-779.

Donald, M. M., Douglas, P., Teri, W. and Donald, W. G. (1990). Moisture stress as it affects yield, soluble

solids and viscosity of tomato. Acta Horticulture 277, Processing tomatoes.

Fadeel, A. A. (1962). Location and properties of chloroplasts and pigments determination in roots. *Physiol. Plant*, 15: 130-147.

Fischer, R. A. and Turner, N. C. (1978). Plant productivity in the arid and semiarid zones. *Annu. Rev. plant Physiol.* 55: 455-459.

Foolad, M. R. and Chen, F. Q. (1997). RAPD markers associated with salt tolerance in an interspecific crosses of tomato (*Lycopersicon esculentum* X *L. pennellii*). *Plant cell Reports* 17: (4), 306-312.

Foolad, M. R. and Jones, R. A. (1993). Mapping salt-tolerance gene in tomato (*Lycoperscon esculentum*) using trait- based marker analysis. *Theer. Appl. Cenet.* 87: 187-192.

Foolad, M. R. and Lin, G. Y. (1997). Absence of a genetic relation ship between salt tolerance during seed germination and vegetative growth in tomato. *Plant Breeding* 116: (4), 363-367.

Gomez, K.A., and Gomez, A.A. (1984). Statistical procedures for agricultural research. John Wily & Sons. Inc. New York, U.S.A..

- Gurrier, G. (1998)**, Proline accumulation in salt treated tomato different proline precursors in *L. esculentum* and *L. pennellii*. Journal of Plant Nutrition, 21: (3), 505-513.
- Hashim, M. M., El-Beltagy, A. S. and Jones, R. A. (1988)**. Salt tolerance in *L. esculentum*1. The effect of salinity on growth. Egypt. J. Hort. 15, No. 1: 85-96.
- Hassan, A. A., Abdel- Fattah M. A. and Khalid. E. Ali. (1987)**. Inheritance of total soluble solids and Ascorbic acid content n tomato Egypt. J. Hort. 14: 2, 155 160.
- Hassan, A. A., Nassar, H. H., Barakat, M. A. and Tolba, M. S. (1999a)**. Tomato breeding for salinity tolerance . Genetic of tolerance. Egypt. J. Hort. 26: 3, 391-403.
- Hassan, M. M., El Masry, T. A. and Abou-Arab, A. A. (1999b)**. Effect of soil salinity growth, yield and elemental concentration in tomato. Egypt. J. Hort. 26: 2, 187- 198.
- Hewitt, J. D., Dinar, M. and Stevens, M. A. (1982)**. Sink strength of fruits of tomato genotypes differencing

in total fruit solids content. J. Amer. Soc. Hort. SCI., 107, 896-900.

Ho, L. G. (1988). The physiological basis for improving dry matter content and calcium status in tomato fruit. Applied agricultural research (USA), 3: (5), 257-261.

Iljin, W. C. (1957). Drought resistance in plants and physiological processes. Annu. Rev. Plant Physiol. 8: 257-274.

James, S. A. and Bell, D. T. (1995). Morphology and anatomy of leaves of *Eucalyptus camaldulensis* clones: variation between geographically separated locations. Australian J. of Botany, 43: 415-433.

John, M., John, C. O. and Jennifer, L. M (1998). Calibration of the Minolta SPDA- 502 leaf chlorophyll meter" photosynthesis Research 48: 467-472.

Kahn, L.; Feder, S. E.; Bray, E. A. and Oconnell, M. A. (1993). Characterization of expression of drought and abscisic acid regulated tomato genes in the drought-resistant species *Lycopersicon pennellii*. Plant physiology, 103: 2, 597-605.

- Karlen, D. L. and Robbins, M. L. (1983).** Management practices for fresh-market tomato production in the southeastern coastal plant. Hort. Sci. 18: 732-734.
- Kebede, H., Johnson, R. C., Carver, B. F. and Ferris, D. M. (1992).** Physiological and anatomical features of two *Triticum discocoides* wheat accessions differing in photosynthetic rate. Crop Sci. 32: 138-143.
- Kebede, H., Martin, B., James, N. and Gretchen, K. (1994).** Leaf anatomy of two *Lycopersicon* species with contrasting gas exchange properties. Crop Science 34: 108-113.
- Kibenko, T. Y., Zhuchenko, A. A. and Khariton, A. A. (1981).** Study of tomato proteins by mean of gel electrophoresis. Ekol. Genet. Rast, Zhivothykh. TZ. DOK. VSES. Konf., 2, 216-217.
- Kubis, J. and Krzywanski, Z. (1991).** Change in the content of polyamines in tomato leaves during increasing soil water deficit and after its removal. Roczniki. Akademi. Rolnizej W. Poznaniu-Ogrodniotwo. No. 19, 43: 50.
- Laemmli, U. K. (1970).** Cleavage of structural proteins during the assembly of the head of bacteriophage T2. Nature, 227: 680-685.

- Lagrimini, L. M., Vaughn, J., Finer, J., Klotz, K. and Rubaihayo, P. (1992).** Expression of a chimeric tobacco peroxidase gene in transgenic tomato. *J. Am. Soc. Hortic. Sci. Alexandria* 117: (6). 1012-1015.
- Lawrence, R. P. (1979).** Breeding for drought resistance what plant characteristics impart resistance. *Hort. Science*, 14: 5. October.
- Levitt, J. (1972).** Responses of plants to environmental stresses. New York: Academic Press.
- Liu, Y., Silverstone, A. L., Wu, Y. M. and Yang, S. F.(1995).** Formation of N-malonyl-L-tryptophan in water stressed tomato leaves. *Phytochemistry-oxford Elsevier science Hot. Oct.* 40: (3). 691-697.
- Lower, R. L. and Thompson, A. E. (1967).** Inheritance of acidity and solids content of small fruited tomato *American Society for Horticultural Science* 41: 486-471.
- Markovic, Z., Zdravrovic, j. and Damjanovic, M. (1997).** Correlation between the morphological characteristics and the biochemical of tomato fruit quality. *Acta Hort.* 462: 1545.
- Martin, B., Kebede, H. and Rilling, C. (1994).** Photosynthetic differences among *Lycopersicon*

species and *Triticum aestivum* are cultivars crop science. Society of American. Jan/Feb 1994. 34: 113-118.

Martin, B., Nienhuns, J., King, G. and Schaefer, A. (1989). Restriction fragment length polymorphisms associated with water use efficiency in tomato. Science 243: 48, 1725.

McCue, K. F. and Hanson, A. D. (1990). Drought and salt tolerance: towards understanding and application. Trends-Biotechnol.; (1990) 8: 12, 358-362.

Melchor, J., Rosa, M., Rivero, L. R. and Juan, M. R. (2004) Evaluation of some nutritional and biochemical indicators in selecting salt-resistant tomato cultivars. Environmental and Experimental Botany 54: 193-201.

Mitchell, J. P., Shennan, C. and Grattan, S. R. (1991). Developmental in changes in tomato fruit composition in response to water deficit and salinity. Physiologia plantum, 83: 1, 177-185.

Nasr, M. I., L El-shawaf, I. I. S., Nazeem, K. R. and Hassanein, S. H. (1982). Inheritance of some characters of tomato fruits and chlorophyll content in four *Lycopersicon* species. Annals of Agric. Sci., Moshtohor, 17: 1982.

- Nassar, H. H., Hassan, A. A., Barakat, M. A. and Tolba, M.S. (1999).** Tomato breeding for salinity tolerance. Assessment and Nature of tolerance Domestic and wild Accessions. Egypt J. Hort. 26: 3, 357-390.
- Natarajan, S. (1990a)** Genetic variability and association of physiological and biochemical traits in tomato under moisture stress. South- Indian- Horticulture, 38: 2, 66-69.
- Natarajan, S. (1990b).** Inheritance of certain growth parameters in tomato under moisture stress. South- Indian. Horticulture 38: 3, 133-136.
- Natarajan, S. (1991).** Genetic variability and heritability in F_2 generation of inter-varietal crosses of tomato under moisture stress. South Indian Horticulture 39: 1, 27-31.
- Natarajan, S. (1992a).** Association of traits in F_2 generation of tomato under moisture stress moisture. South- Indian- Horticulture, 40: 6, 355-356.
- Natarajan, S. (1992b).** Inheritance of yield and its components in tomato under moisture stress. Madras. Agricultural. Journal. 79: 12, 705-710.

- Natarajan, S. (1993).** Heterosis in tomato under moisture stress. *South. Indian. Horticulture*. 41: 5, 245-247.
- Nobel, P.S. (1991).** *Physicochemical and environmental plant physiology*. Academic Press, Inc. San Diego.
- Obreza, T. A., Pitts, D. J., McGovern, R. J. and Spreen, T. H. (1996).** Deficit irrigation of micro-irrigated tomato affects yield, fruit quality, and disease severity. *J. Prod. American Society of Agronomy*, 1987 Apr/June 1996. 9: (2). 270-275.
- Ong, H. T. (1979).** Gel electrophoresis patterns of proteins and peroxidase of excised tomato cotyledons subjected to mannitol induced water stress. *Biologia Plantarum* , 20: 5, 330-334.
- Oosterhuis, and Wullschleger, S. D. (1990).** Consideration of drought tolerance in irrigation management of vegetable crops. *Acta Horticulture*, 278.
- Orton, T. J. (1982).** New concepts in whole plant genetics. *Calif. Agric*, 36: 22-23.
- Patel, K. V., Talati, J. G. and Bhatanger, R. (2002).** Application of polyacrylamide gel electrophoresis technique for identification of varieties of chill, tomato, brinjal and bhindi (okra). *Journal of*

Moharashtra- Agricultural Universities, 26: (3),
266-268.

Perez-Alfocea, F. and Larher, F. (1995). Effects of phlorizin and p-chloromer curibenzenesulfonic acid on sucrose and praline accumulation in detached tomato leaves submitted to NaCl and osmotic stress. *Journal of Plant Physiology*. 145: 3, 367-373.

Plant, A. L., Cohen, A., Moses, M. S. and Bray, E. A. (1991). Nucleotide sequence and spatial expression pattern of a drought and abscisic acid induced gene of tomato. *Plant-Physiol.* Rockville, Md.: American Society of plant physiologists. Nov 1991, 97: 3, 900-906.

Rao, N.K. S. and Bhatt, M. RM. (1992). Responses of tomato to moisture stress. Plant water balance and yield. *Plant Physiology and Biochemistry-New-Delhi*, 19: 1, 36-41.

Rao, D. V. R. and Padma, S. S. V. (1991). Effect of induced moisture stress as different phonological stages on growth and yield tomato cultivars. *South, Indian- Horti*. 39: 2, 81-87.

Rendon-Poblete, E. (1980). Effect of soil water status on yield quality and root development of several

tomato genotypes. Ph. D. dissertation. University of California, Davis.

Richards, M. A., Phills, B. R. and Hill, W. (1979) In Chapter 1 Plant Improvement and Somatic Cell Genetics. (eds. K. Vasil, W. R. Scowcroft, and K. L. Frey), Academic Press (1982), NY, PP. 1-28.

Rick, C. M. (1969). Controlled introgression of chromosomes of *Solanum pennellii* into *Lycopersicon esculentum* segregation and recombination. Genetics 62: 753-768.

Rick, C. M. (1973). Potential genetic resource in tomato species: clues from observations in native habitats. In Genes, Enzymes and Population (eds A. Hollaender and A. M. Srp) Plenum Press, Newyork.

Rick, C. M. (1982). The potential of exotic germplasm for tomato improvement. In Plant Improvement and Somatic Cell Genetics (eds. K. Vasil, W. R. Scowcroft, and K. L. Frey), Academic Press (1982), NY, PP. 1-28.

Rick, C. M. and Lamm, R. (1955). Biosystematic studies on the status of *Lycopersicon chilense*. Am. J. Bot., 42, 663-675.

Rodriguez, M., Canales, E., Carlos, J. B., Elva C., Lopez, J., Pujol, M. and Borrás-Hidalgo, O.

(2005) Identification of genes induced upon water-deficit stress in a drought-tolerant rice cultivar. *Journal of Plant Physiology*. Accepted 5 July 2005 (In press).

Rohlf, J. (1993). Numerical Taxonomy and multivariate analysis system NTSys- PC. Version 1-80 Exeter Software, New York.

Rossi, M., Lijararetzhy, D., Bernacchi, D., Hopp, H. E. and Iusem, N. (1996). Asr genes belong to a gene family comprising at least three closely linked loci on chromosome 4 in tomato. *MGG- Mol. Gen. genet.* Berlin, Germany: Springer productions Gesellschaft sept 25, 252: (4). 489-492.

Rudich, J. and Luchinsky, V. (1994). Water economy. In the tomato crop A scientific basis for improvement. Edited by Artherton J. G. and Rudich, J. PP. 334-368.

Rudich, J., Kalmar, D., Geizenberg, C. and Harel, S. (1977). Low water tensions in defined growth of processing tomato plants and their effects on yield and quality. *J. of Hort. Sci.* 52: 391-399.

Rudich, J., Rendon- Poblete, E., Stevens, M.A. and Ambri, A. I. (1981). Use of leaf water potential to

determine water stress in field grown tomato plants.

J. Amer. Soc. Hort. SCI. 106, 732-736.

Sam, O., Jerez, E. and Varela, M. (1996). Characteristics anatomic as de hojas de papa (*Solanum tuberosum* L.) Y tomato (*Lycopersicon esculentum* Mill.) can differences grados de tolerancia a estres de humedad Y temperature. Cultivos Tropicales, 17: 2, 32-38. (Cited from Agricola DATA BASE INTERNET).

Smajstrla, A. G, and Locascio, S. J. (1996). Tensiometer controlled drip irrigation scheduling of tomato. Applied Engineering in Agriculture 1996 American Society of Agricultural Engineers. V. 12 (3): 315-319.

Smith, F., Gilles, M.A., Hamilton, J.K. and Godees P.A. (1956). Colorimetric method for determination of sugar and related substances. Anal. Chem. 28: 350-356.

Souza, D. MC., Head, M. K. and Ingle, M. (1991). Quality of processed field growth tomatoes as influenced by growth of plants under less than optimum soil moisture conditions. Journal of Sustainable Agriculture , 2: 1 41-48.

SPSS,(1998). Statistical Package for Social Science SPSS for Windows Computer Program.Ver.6.1 SPSS Inc. 444 North Michingan Avenue, Chicago, Illinois 60611, USA.

Srinivasa-Rao, N. K., Bhatt, R. M. and Rao, N. K. S. (1991). Stomatal frequency conductance and transpiration rate at different canopy positions of water stressed cultivars of tomato (*Lycopersicon esculentum*). Indian Journal of Agricultural Sciences. 61: 6, 434- 436.

Stanhill, G. (1986). Water use efficiency. A, dv. Agron. 39: 53-85.

Stegmann, H. A., Afify, M. R. and Hussein, K. R. E. (1985). Cultivars identification of dates (*Phoenix dactylifera*) by protein patterns. 2nd International symposium of Biochemical Approaches to Identification of cultivars. Braunschweig, West Germany, pp 44-54.

Studier, F. W. (1973). Analysis of bacteriophage T1 early RNA_s and proteins of slab gels. J. Mol. Biol., 79: 237-248.

- Tanksley, S. D. and Rick, C. M. (1980).** Genetic of esterases in species of *Lycopersicon*. Theor. Appl. Genet. 56: 209- 219.
- Tanksely, S. D., Medina-Filho, H. and Rick, C. M. (1982).** Use of naturally occurring enzyme variation to detect and map genes controlling quantitative traits in an inter specific backcross of tomato. Heredity, 49: 11-25.
- Taylor, S. A. (1952).** Use of mean soil moisture tension to evaluate the effect of soil moisture on crop yields. Soil Sci. 74: 217-226.
- Thakur, P. S. (1991).** Effect of water stress on proline and relative water content in tomato cultivars. Indian, Journal, of Horticulture. 48:1, 36-41
- Thakur, P,S., Anju-Thakur and Thakur. A. (1993).** Influence of triacontanol and mixtalol during plant moisture stress in *Lycopersicon esculentum* cultivars. Plant physiology and Biochemistry. Paris 31: 3, 433-439.
- Torrecillas, A., Guillaume, C., Alarcon, J. J. and Sanchez, M. C.R.(1995).** Water relations of two tomato species under water stress and recovery. Plant Science, Volume 105, Issue 2, February 169-179.

- Valpuesta, V., Berteli, F., Perez prat, E., Carrales, E., Narasimham, M., Botella, M. A., Brassan, R. A., Pliego, F. and Haseggawa, P. M. (1992).** Metabolic changes and gene expression in higher plants in response to salt stress. *Agriscientia* . 9: 1, 55- 63.
- Vogel, A. (1968).** A text book of quantitative inorganic analysis Longman Pub., New York, pp.1216.
- Wei, T. and Oconnel, M. A. (1996).** Structure and characterization of a putative drought inducible H1 Histone gene. *Plant- mol- boil.* Dordrecht: Kluwer Academic Publishers. Jan. 30: (2). 255-268.
- Weibel, E. R. (1979).** Stereological methods. Vol. 1 Practical methods for biological morphometry. Academic Press, New York.