

References

- 1- **Akabori, O. (2006):** The role of NK Tcells in the development of bronchial asthma: a transitional approach from animal models to human. *Allergy Journal* 61: 962-968.
- 2- **Arima, K.; Umeshita-Suyama, R.; Sakata, Y. et al.,(2002):** Upregulation of IL-13 concentration in vivo by the IL13 variant associated with bronchial asthma. *J Allergy Clin Immunol* ; 109:980-987.
- 3- **Bardley, BL.; Azzawi, M.; Jacobson, M.; et al., (1991):** Eosinophils, T-lymphocytes, mast cells, neutrophils, and macrophages in bronchial biopsy specimens from atopic subjects with asthma: comparison with biopsy specimens from atopic subjects without asthma and normal control subjects and relationship to bronchial hyper responsiveness. *J. Allergy Clin. Immunol*; 88:661-674.
- 4- **Bardora, JE. (1997):** Indoor air pollution and health. *Health& Fitness Book*.
- 5- **Barnes, PJ. (1989):** New concepts in the pathogenesis of bronchial hyperresponsiveness and asthma. *J Allergy Clin Immunol*; 83: 1013.
- 6- **Bautsch, W.; Hoymann, HG.; Zhang, Q. et al.,(2000):** Cutting edge: guinea pigs with a natural C3a-receptor defect exhibit decreased bronchoconstriction in allergic airway disease:

- evidence for an involvement of the C3a anaphylatoxin in the pathogenesis of asthma. *J Immunol*; (165): 5401-5405.
- 7- **Beeh, K-M.; Micke, P.; Ksoll, M. et al., (2001):** Cigarette smoking, but not sensitization to alternia, is associated with severe asthma in urban patients. *Journal of Asthma*; 38(1): 41-49.
- 8- **Blease, K.; Jakubzic, C.; Westwick, J.; et al., (2001):** Therapeutic effect of IL-13 immunoneutralization during chronic experimental fungal asthma. *J Immunol*; 166: 5219-5224.
- 9- **Booth, BW.; Adier, KB.; Bonner, JC.; et al., (2001):** Interleukin-13 induces proliferation of human airway epithelial cells in vitro via a mechanism mediated by transforming growth factor- α . *Am J Respir Cell Mol Biol*; 25: 739-743.
- 10- **Boulet, L-P and Sterk, PJ. (2007):** A new series on airway remodelling. *Eur Respir J*; 29: 231-232
- 11- **Busse, WW.; and Germ, JE. (1997):** Viruses in asthma. *J. Allergy Clin. Immunol*; (100): 147-150
- 12- **Busse, WW. and Lemanske, RF. (2001):** Advances in immunology: asthma. *N Engl J Med*; 344 (5): 350-362.
- 13- **Caldiera, RD.; Bettiol, H.; Barbier, MA. Et al., (2006):** Prevalence and risk factors for work related asthma in young adults. *Occup Environ Med*; publication number 16728501.
- 14- **Cameron, L.; Webster, RB.; Strempel, JM. et al., (2006):** Th2 cell-selective enhancement of human IL-13 transcription

- cell by IL13-1112C>T, a polymorphism associated with allergic inflammation. J Immunol; (12): 8633-8642.
- 15- **Chung, KF. and Barnes, PJ. (1999):** Cytokines in asthma. Thorax: 54: 825.
- 16- **Creticos, PS. (2002):** Immunotherapy in allergic asthma. National Asthma Campaign: Division of Allergy and Clinical immunology; Johns Hopkins University School Of Medicine Baltimore Maryland.
- 17- **Dabahay, H.; Atherton, H.; Jones, G.; et al., (2002):** Interleukin-13 induces a hypersecretory ion transport phenotype in human bronchial epithelial cells. Am J Physiol Lung Cell Mol Physiol; 282: 226-236.
- 18- **Dawson-Saunders, B.; and Trapp, R. (1994):** "Basic and Clinical Biostatistics" 2nd edition. Lange Medical Book, Prentice-Hall International Inc.
- 19- **Del Prete, G.; Maggi, E.; Parronchi, P.; et al., (1988):** IL-4 is an essential factor for the IgE synthesis induced in vitro by human T cell clones and their supernatants. J. Immunol; 140: 4193.
- 20- **Der VP.; Der, VJ.; Je, BL. (1998):** The role of IL-13 in IgE synthesis by allergic asthma patients. Clin. Exp. Immunol; 117-129.
- 21- **Derycks, S.; Malefyt, DW.; Gauchat, JF.; (1992):** Immunoregulatory functions of PAF, Inhibition of IL-4 induced human IgE synthesis. J. Immunol; 148: 1465.

- 22- **Dienger, KM. and Wills-Karp, M. (2002):** Interleukin-13 regulation of complement factor 3 production by human bronchial epithelial cells. *Am J Respir Crit Care Med*; A733.
- 23- **Doherty, TM.; Kastelein, R.; Menon, S.; et al., (1993):** Modulation of murine macrophage function by IL-13. *J Immunol*; 151: 7151-7160.
- 24- **Dudley, KL.; Chiaramonte, M.; Wills-Karp, M. et al., (2002) :** Upregulation of mucin related genes is associated with susceptibility to allergen-induced hyperresponsiveness. *Am J Respir Crit Care Med* ; A 693.
- 25- **Elias, JA.; Lee, CG.; Zheun, T. et al., (2003):** Interleukin 13 and leukotrienes. *American Journal of Respiratory cell and molecular Biology*; (28): 401-404.
- 26- **Elliott, KA; Osna, NA.; Scofield, MA. et al., (2001):** Regulation of IL-13 production by histamine in cloned murine T helper type 2 cells. *Int Immunopharmacol* :1923-1937.
- 27- **Empey, DW.; Laitinen, LA.; Jacobs, L. et al., (1979):** Mechanism of bronchial hyperreactivity in normal subjects following upper respiratory tract infection. *Am. Rev. Respir Dis*; 113: 131-139.
- 28- **Fishman, AP.; Elias, JA.; Fishman, JA. Et al., (1998):** Fishman's Pulmonary Diseases and disorders. Chapter 20: 757-776.
- 29- **Fraenkel, DJ. and Holgate, ST. (1996):** Etiology of asthma : Pathology and mediators. "Allergy, Asthma, and Immunology

from Infancy to Adulthood”^{3rd}ed. Saunders company, Philadelphia, p 443.

- 30- **Furkerson, PC.; Fischetti, CA.; Rothenberg, ME. (2006) :** Eosinophilia and CCR3 regulate interleukin-13transgene-induced pulmonary remodelling. *Am J Pathol*; (6): 2117-2126.
- 31- **Gauchat, JF.; Grunig, G.; Warnock, M. et al., (1998):** Requirement for IL-13 independently of IL-4 in experimental asthma. *Science*; 282: 2261-2262.
- 32- **Gavett, SH.; O'Hearn, DJ.; Karp, CL. et al., (1997):** Interleukin-4 receptor blockade prevents airway responses induced by antigen challenge in mice. *AM J Physiol Lung Cell Mol Physiol*; 272: 253-261
- 33- **Graber, P.; Gretener, D.; Herren, S. et al. (1998):** The distribution of IL-13 receptor alpha 1 expression on B cells, T cells, and monocytes and its regulation by IL-13 and IL-4. *Eur J Immunol*; 28: 4286-4298.
- 34- **Graves PE, Kabesch M, Halonen M, et al.(2000):** A cluster of seven tightly linked polymorphisms in the IL-13 gene is associated with total serum IgE levels in three populations of white children. *J Allergy Clin Immunol* ;105:506-513.
- 35- **Grunig, G.; Warnock, M.; Wakil, AE. et al.,(1998):** Requirement for IL-13 independently of IL-4 in expermental asthma. *Science*;(282):2258-2262.
- 36- **Grunstein, MM.; Hakonarson H.; Leiter J. et al., (2002):** IL-13-dependent autocrine signaling mediates altered responsiveness

- of IgE-sensitized airway smooth muscle. *Am J Physiol Lung Cell Mol Physiol*; 282: 520-528.
- 37- **Gueant-Rodriguez, R-M.; Romano, A.; Beri-Dexheimer, M. et al., (2006):** Gene-gene interactions of IL-13 and IL-4RA variants in immediate allergic reactions to beta lactam antibiotics. *Pharmacogenet Genomics*. 16 (10): 713-719.
- 38- **Guerin, B. and Watson, RD. (1988):** Skin tests. *Clin Rev. Allergy*; (6): 211-227.
- 39- **Hamid, A. and Minshall, M. (2000) :** PHA Molecular pathology of allergic disease *J. Allergy clin. Immunol*; 105: 20.
- 40- **Harterd, TV. and Peebles, RS. (2000) :** Epidemiology of asthma. *Curr Opin pulm Med*; (1) : 4-9.
- 41- **Hashimoto, T.; Kobayashi, N.; Nakjiyama, Y.; et al., (2006) :** IL-2 induced IL-13 production by allergic-specific human helper T cell clones. *Int Arch Allergy Immunol*; (140):51-54
- 42- **Hattori, T.; Hizawa, N.; Takahashi, D. et al., (2006):** Total serum IgE levels in eosinophilic lung diseases. *Areugi* 55 (6): 647.
- 43- **Heinzmann, A.; Mao, XQ.; Akaiwa, M. et al.,(2000):** Genetic variants of IL-13 signaling and human asthma and atopy. *Hum Mol Genet*; (9):549-559.
- 44- **Hirst, SJ.; Hallsworth, MP.; Peng, Ql. et al. (2002):** Selective induction of eotaxin release by interleukin-13 or interleukin-4 in human airway smooth muscle cells is synergistic with

- interleukin-1B and is mediated by the interleukin-4 receptor α -chain. *Am J Respir Crit Care Med*; 165:1161-1171.
- 45- **Hopkin, JM. (1989):** Genetics of atopy. *Clin Exp Allergy*; (19):263.
- 46- **Hosseini–Faraabadi, S.; Tavakkol–Afshari, J.; Rafatpanah, H. et al., (2007):** Association between the polymorphisms of IL-4 gene promoter (-590>T), IL-13 coding region (R 130Q) and IL-16 gene promoter (-295T>C) and allergic asthma. *Iran J Allergy Asthma Immunol*; 6(1): 9-14.
- 47- **Howard, TD.; Whittaker, PA.; Zaiman, AL. et al., (2001):** Identification and association of polymorphisms in the interleukin-13 gene with asthma and atopy in a Dutch population. *Am J Respir Cell Mol Biol* ; 25: 377-378.
- 48- **Humbles, AA.; Lu, B.; Nilsson, CA. et al.,(2000):** A role for the C3a anaphylatoxin receptor in the effector phases of asthma. *Nature*; (406): 998-1001.
- 49- **Hurst, SD.; Muchamuel, T.; Gorman, DM. et al., (2002):** New IL-17 family members promote Th1 or Th2 responses in the lung: in vivo function of the novel cytokine IL-25. *J Immunol*; 169: 443-453.
- 50- **Ishizaka, T. (1981) :** Analysis of triggering events in mast cell for immunoglobulin–E mediated histamine release. *J. Allergy Clin. Immunol*; 67 : 90.

- 51- **Izuhara, K.(2001):** Involvement of interleukin-4 and-13 in the pathogenesis of allergic diseases. *Rinsho Byori* Apr; 49(4):360-364.
- 52- **James, EF. and Stephen, PP. (1999) :** Airway remodelling and persistant airway obstruction in asthma. *J. Allergy Clin. Immunol*; 104:509
- 53- **Johnston, SL.; Pattemore, G.; Sanderson, S. et al.,(1995):** Community study of role of viral infections in exacerbations of asthma in 9-11 years old children. *B.M.J.*; (310): 1225-1228.
- 54- **Kavuru, MS.; Lang, DM.; Erzurum, SC. (2005):** Definition of bronchial asthma. *Pulmonary, Allergy, and Critical care, The Cleveland and Clinic Foundation.*
- 55- **Kim, HA.; Heo, YJ.; Lee, SH.; et al., (2006):** No potential role of genetic polymorphisms for IL-4, IL-13 and IL-4 receptor in respiratory allergy. *JUOEH*; 28 (4): 369-379
- 56- **Kimata, H.; Yoshida, A.; Ishioka, C. et al., (1992) :** Interleukin-8 (IL-8) selectively inhibits immunoglobulin E production induced by IL-4 in human cells.
- 57- **Kiniwa, M.; Gately, M.; Gubler, U. (1992) :** Recombinant interleukin-12 suppresses the synthesis of immunoglobulin E by interleukin-4 stimulated human lymphocytes. *J. Clin. Nvest*; 90: 262.
- 58- **Kioi, M.; Seetharam, A.; Puri, K. (2006) :** N linked glycosylation of IL-13R (alpha) 2 is essential for optimal IL-13 inhibitory activity. *FASEBJ*; publication number 17023392

- 59- **Kondo, M.; Tamaoki, J.; Takeyama, K. et al., (2006):** Elimination of IL-13 reverses established goblet cell metaplasia into ciliated epithelia in airway epithelial culture. *Allergol Int*; 55 (3):329-336.
- 60- **Koshino, T.; Arai, Y.; Miyamoto, Y. et al., (1996):** Airway basophil and mast cell density in patients with bronchial asthma: Relationship to bronchial hyperresponsiveness. *Journal of asthma*; 33(2) 89-95.
- 61- **Kotsimbos, TC.; Ernst, P.; Hamid, QA. (1996):** Interleukin-13 and interleukin-4 are coexpressed in atopic asthma. *Proc Assoc Am Physicians*; 108(5): 368-373.
- 62- **Kraan, TC.; Veen, A.; Boeije, LC. (1999):** An IL-13 promoter polymorphism associated with increased risk of allergic asthma. *Genes and Immunity*; (1):61-65.
- 63- **Kuperman, D.; Schofield, B.; Wills-Karp, M. et al.(1998):** Signal transducer and activator of transcription factors (Stat6)-deficient mice are protected from antigen-induced airway hyperresponsiveness and mucus production. *J Exp Med*; 187: 939-948.
- 64- **Kuperman, DA.; Huang, X.; Koth, LL. et al. (2002):** Direct effects of interleukin-13 on epithelial cells cause airway hyperreactivity and mucus overproduction in asthma. *Nat Med*; 8: 885-889.
- 65- **Laoukili, J.; Perret E.; Willems, T. et al. (2001):** IL-13 alters mucociliary differentiation and ciliary beating of human respiratory epithelial cells. *J Clin Invest*; 108: 1817-1824.

- 66- **Laporte, JC.; Moore, PE.; Baraldo, S. et al. (2001):** Direct effects of interleukin-13 on signaling pathways for physiologic responses in cultured human airway smooth muscle cells. *Am J Respir Crit Care Med*; 164: 141-148.
- 67- **Lee, GR and Flavell, RA. (2004) :** Transgenic mice which overproduce Th2 cytokines develop spontaneous atopic dermatitis and asthma. *International Immunology*; 16(8): 1155-1160.
- 68- **Lee, YC.; Lee, KH.; Lee, HB. et al., (2001):** Serum levels of interleukins IL-4, IL-5, IL-13 and interferon- γ in acute asthma. *Journal of Asthma*; 38(8): 665-671
- 69- **Leigh, R.; Ellias, R.; Wattie, J. et al., (2004):** Is interleukin-13 critical in maintaining airway hyperresponsiveness in allergen-challenged mice?. *American Journal of Respiratory and Critical Care Medicine*; (170): 851-856.
- 70- **Leo, J.; Ukili, J.; Perret, E. et al., (2001):** IL-13 alters mucociliary differentiation and ciliary beating of human respiratory epithelial cells. *J Clin Invest*; 108(12): 1817-1824.
- 71- **Leung DY (1994) :** Mechanisms of the human allergic response. *Ped. Clin. North. Am*; 41: 727.
- 72- **Li, L.; Xia, Y.; Nguyen, A. et al., (1999):** Effects of Th2 cytokines on chemokine expression in the lung: IL-13 potently induces eotaxin expression by airway epithelial cells. *J Immunol* ; 162:2477-2487.

- 73- **Lin,L.; Bin,W.; Hua-Jun, C. et al., (2004):** Relationship between IL-13 gene polymorphisms and asthma , the relationship between the level of plasma cytokine . Journal of Clinical Lung.
- 74- **Liu, MC.; Hubbard, WC.; Proud, D. (1991):** Immediate and late inflammatory responses to ragweed antigen challenge of the peripheral airways in allergic asthmatics.
Am.Rev.Respire.Dis.:(144):51.
- 75- **Liu,X.; Kohyama,T.; Wang, H. et al., (2002):** Th2 cytokine regulation of type I collagen gel contraction mediated by human lung mesenchymal cells. Am J Physiol Lung Cell Mol Physiol; 282: L 1049-1056.
- 76- **Mattes, J.; Yang, M., Mahalingani, S. et al., (2001)a:** Internisic defect in T cell production of interleukin IL-13 in the absence of both IL-5 and eotaxin precludes the development of eosinophilia and airways hyperreactivity in experimental asthma. J. Exp. Med; 195: 1433-1444.
- 77- **Mattes, J.; Yang, M.; Siqueira, A. et al., (2001)b:** IL-13 induces airways hyperreactivity independently of the IL-4R alpha chain in the allergic lung. J Immunol; 167: 1683-1692.
- 78- **Matzger, H. (1992) :** The receptor with high affinity for IgE. Immunol Rev; 125 : 37 .
- 79- **McKenzie, GJ.; Emson, CL.; Bell, SE. et al., (1998):** Impaired development ofTh2 cells in IL-13 deficient mice. Immunity; 9: 423-432.

- 80- **Menzies-Gowarrd, A. and Robinson, DS. (2001):** Eosinophil chemokines and chemokines receptors: their role in eosinophil; accumalation and activation in asthma and potential as therapeutic targets . Journal of Asthma; 38(8): 605 –613.
- 81- **Middleton, E. et al., (1993):** Allergy: principles and practice. 4th edition Mosby.
- 82- **Miyahara, N.; Takeda, K.; Kodama, T. et al., (2004):** Contribution of antigen primed CD⁸⁺ T cells to the development of airway hyperresponsiveness and inflammation is associated with IL-13 . The Journal of Immunology; 172: 2549 –2558.
- 83- **Moore, PE.; Church, TL.; Chism, DD.; et al., (2002):** IL-13 and IL-4 cause eotaxin release in human airway smooth muscle cells: a role for ERK. Am J Physiol Lung Cell Mol Physiol ; 282:847-853.
- 84- **Mohrs, M.; Poor, BCM.; Wang, ZE. et al. (2001):** Deletion of a coordinate regulation of type 2 cytokine expression in mice Nat Immunol ; 2: 842-847.
- 85- **Motojima, S.; Takeishi, K.; Koseki, T. et al., (1997):** Serum levels of eosinophilis cationic protien and IL-5 in patients with asthma without systemic corticosteroids. Int. Arch Allergy Immunol (114) 1: 55-59.
- 86- **Mould, A.; Matthaei, K.; Young, IG.; et al., (1997):** Relationship between IL-5 and eotaxin in regulating blood and tissue eosinophilia in mice . J Clin Invest; (99): 1064-1071.

- 87- **Nadel, JA.; and Busse, WW. (1998):** Asthma. Am.J.Respir. Crit. Care Med.; 157; (4): 130-138.
- 88- **Naghmi, R. (1992):** ABC of the manegement of acute asthma. Medicine Digest; 18: 3-8.
- 89- **Nakanishi, A.; Morita, S.; Iwashita, H. et al., (2001):** Role of gob-5 in mucus overproduction and airway hyperresponsiveness in asthma. Proc Natl Acad Sci USA; 98: 5175-5180.
- 90- **Nelson.; Richard, EB.; Robert, MK. et al., (2000) :** Chapter 145 Asthma Nelson Text book of pediatrics 16 th edition.
- 91- **Niitsuma, TA.; Nukaga, A.; Izawa, AM. et al.,(2004):** Thirty cases of bronchial asthma associated with exposure to pet hamsters. J Invest Allergol Clin Immunol vol 14(3):221-224.
- 92- **Noguchi, E.; Nishioy, N.; Jian, Z. et al., (2001) :** Haplotypes of the 5 region of the IL-4 gene and SNPs in integrene sequence between the IL-4 and IL-13 grenes arte associated with atopic asthma. Hum Immunol; (62): 1251-1257.
- 93- **O'Byrne, PM.; Dolovich, J.; Hargreave, FE. et al., (1987) :** Late asthmatic responses. Am. Rev. Respir. Dis; 136: 740.
- 94- **Oh, SY.; Zheng, T.; Bailey, ML.; et al., (2007) :** Src homology 2 domain containing inositol 5-phosphatase 1 deficiency leads to a spontaneous allergic inflammation in the murine lung . J Allergy Clin .Immunol; Puplication number 7208593.

- 95- **Ohshima Y; Yasutomi M; Omata N, et al. (2002):** Dysregulation of Il-13 production by cord blood CD4⁺ T cells is associated with the subsequent development of atopic diseases in infants. *Pediatr Res*, (51): 195-200.
- 96- **Palframan, R.; Collins, PD.; Williams, TJ. Et al., (1998):** Eotaxin induces a rapid release of eosinophils and their progenitors from the bone marrow. *Blood*; (91): 2240-2248.
- 97- **Payne, DNR.; and Balfour-Lynn, IM.(2001):** Children with difficult asthma. A practical approach. *Journal of Asthma*; 38(3): 189-203.
- 98- **Peat, JK. (1987):** Asthma predicts accelerated decline in FEV1. *Eur J Respir Dis*; (70):171-179.
- 99- **Pene, J.; Rousset, F.; Briere, F. (1988):** IgE production by normal human B cells is induced by interleukin 4 and suppressed by interferon γ and α and prostaglandin E2. *Proc. Nat. Acad. Sci. USA*; (85) : 6880.
- 100-**Phelan, PD.; Olinsky, A.; Robertson, C. (1994):** Respiratory illness in children. 4th editor Blackwell london: 113-183
- 101-**Pope, SM.; Brandt, EB.; Mishra, A. et al. (2001):** IL-13 induces eosinophil recruitment into the lung by an IL-5 and eotaxin-dependent mechanism. *J Allergy Clin Immunol*; 108:594-601.
- 102-**Punnonen, J.; Aversa, G.; Cocks, BG. (1993):** Interleukin-13 induces interleukin-4 independent IgG4 and IgE synthesis and

CD23 expression by human B cells. Proc. Nar. Acad. Sci. USA; 90: 3730.

- 103-**Punnonen, J.; Aversa, G.; Cocks, BG. (1994):** Role of interleukin-4 and interleukin-13 in synthesis of IgE and expression of CD23 by human B cells. J. Allergy; 49 : 576 – 86.
- 104-**Richter, A.; Puddicombe, SM.; Lordan, JL. et al., (2001):** The contribution of interleukin (IL)-4 and IL-13 to the epithelial-mesenchymal trophic unit in asthma. Am J Respir Cell Mol Biol; 25: 385-391.
- 105- **Samter, M. (1994):** Immunologic diseases 5th ed. Little Brow.
- 106-**Schmid-Grendelmeier, P.; Altnauer, F.; Fischer, B. et al., (2002):** Eosinophils express functional IL-13 in eosinophilic inflammatory diseases. J Immunol; 169: 1021-1027.
- 107-**Schon-Hegrad, MA.; Oliver, J.; McMenamin, PG. (1991) :** Studies on the density, distribution and surface phenotype of intra epithelial class II MHC antigen-bearing dendritic cells in the conductive airways. J. Exp. Med.; 173: 1345.
- 108-**Schwartz, LB. and Kepley, C. (1994) :** Development of markers for human basophils and mast cells. J. Allergy Clin. Immunol; 94: 123.
- 109-**Singer, M.; Lefort, J.; Vargaftig, BB. (2002):** Granulocyte depletion and dexamethasone differentially modulate airways hyperreactivity, inflammation, mucus accumulation, and secretion induced by rmIL-13 or antigen. Am J Respir Cell Mol Biol; 26: 74-84.

- 110-**Tag-Eldin, M.; Abdel-Rahman, G.; Nor-Eldin, A. et al., (1998):** Classification of bronchial asthma according to Egyptian society of chest diseases and tuberculosis. Egyptian Journal of Chest Diseases and Tuberculosis.46(1)
- 111-**Tamachi, T.; Ikeda, MK.; Iwamoto, I. Et al., (2006):** Interleukin 25 in allergic airway inflammation. J of Immunol; Publication number 16772729.
- 112-**Temann, DA.; Geba, GP.; Rankin, JA. et al., (1998):** Expression of interleukin 9 in the lungs of transgenic mice causes airway inflammation, mast cell hyperplasia, and bronchial hyperresponsiveness. J Exp Med; 188: 1307-1320.
- 113- **Temann, U-A.; Laouar, Y.; Elizabeth, E. et al., (2007):** IL-9 leads to airway inflammation by inducing IL-13 expression in airway epithelial cells. Internal Immunology;19: 1-10.
- 114- **Temann, UA.; Ray, P.; Flavell, RA. Et al., (2002):** Pulmonary overexpression of IL-9 induces Th2 cytokine expression, leading to immune pathology. J Clin Invest; 109: 29-39.
- 115- **Venkayya, R.; Lam, M.; Willkom, M. et al., (2002):** The TH2 lymphocyte products IL-4 and IL-13 rapidly induce airway hyperresponsiveness through direct effects on resident airway cells. Am J Respir Cell Mol Biol; 26: 202-208
- 116- **Vladich, FD.; Brazille, SM.; Stern, D. et al., (2005) :** IL-13 R 130Q, a common variant associated with allergy and asthma, enhances effector mechanisms essential for human allergic inflammation. J.Clin.Invest; (115): 747-754.

- 117-**Vonk, JM.; Koppelman, GH.; Boezen, HM. Et al., (2006):** Genetic variation in IL-4 receptor- α and IL-13 is associated with the outcome of asthma after 30 years. 3rd ed :149
- 118-**Walter, DM.; McIntire, JJ.; Berry, G. et al., (2001):** Critical role for IL-13 in the development of allergen-induced airway hyperreactivity. J Immunol; 167: 4668-4675.
- 119- **Walter, DM.; Jennifer, J.; Intire, M.; et al., (2002):** Critical role for IL-13 in the development of allergen-induced airway hyperreactivity. The Journal of Immunology; 167: 4668-4675.
- 120-**Wardlaw, A. (1999):** Molecular basis for selective eosinophil trafficking in asthma. J. Allergy Clin Immunol; (104): 917-926.
- 121-**Wen, FQ.; Kohyama, T.; Liu, X. et al., (2002):** Interleukin-4 and interleukin-13-enhanced transforming growth factor beta-2 production in cultured human bronchial epithelial cells is attenuated by interferon-gamma. Am J Respir Cell Mol Biol; 26: 484-490.
- 122-**Wenzel, S.(2005):** severe asthma in adults. American Journal of Respiratory and Critical Care Medicine (172): 149-160.
- 123- **Wills-Karp, M. and Chiaramonate, M. (2003):** Interleukin 13 in asthma. Pulmonary Medicine.
- 124-**Wills-Karp, M.; Luyimbazi, J.; Xu, X. et al.,(1998):** Interleukin-13: central mediator of allergic asthma. Science; 282: 2258-2261.

- 125- **Yamada, N.; Inoue, H. ; Takishima, T. et al., (1996):**Role of macrophages in pulmonary late-phase reaction in guinea pigs. *Journal of Asthma*; 33(6): 395-405.
- 126- **Yamauchi, K.; Tanifuzi, Y.; Pan, L-H. et al., (2001):**Effects of pranlukast, a leukotriene receptor antagonist on airway inflammation in mild asthmatics. *Journal of Asthma*; 38(1): 51–57.
- 127- **Yang, G.; Li, L.; Volk, A. et al., (2005) :** Therapeutic dosing with anti-interleukin-13 monoclonal antibody inhibit asthma progression. *Journal of Pharmacology and Experimental Therapeutics Fast Forward*; (313): 8-15.
- 128-**Yang, M.; Hogan, SP.; Henry, PJ. et al., (2001):** Interleukin-13 mediates airways hyperreactivity through the IL-4 receptor-alpha chain and STAT-6 independently of IL-5 and eotaxin. *Am J Respir Cell Mol Biol*; 25: 522-530.
- 129-**Zheng, T.; Zhu, Z.; Wang, Z. et al., (2000):** Inducible targeting of IL-13 to the adult lung causes matrix metalloproteinase- and cathepsin-dependent emphysema. *J Clin Invest*; 106: 1081-1093.
- 130-**Zhou, Y.; Dong, Q.; Louahed, J. et al., (2001):** Characterization of a calcium-activated chloride channel as a shared target of Th2 cytokine pathways and its potential involvement in asthma. *Am J Respir Cell Mol Biol*; 25: 486-491.
- 131-**Zhu, Z.; Homer, RJ.; Wang, Z. et al., (1999):** Pulmonary expression of interleukin-13 causes inflammation, mucus hypersecretion, subepithelial fibrosis, physiologic abnormalities, and eotaxin production. *J Clin Invest*; 102: 779-788.