

elicitor activates the gene for indole emission in maize.  
Proc. Nat. Acad. Sci. USA, 97:1481-14806.

- Gbolade A.A and Adebayo, T.A. (1993):** Fumigant effects of some volatile oils on fecundity and adult emergence of *Callosobruchus maculatus* F. Insect Science and its Application. 14 (5): 631-636.
- Geissman, T.A. (1961):** Chemistry of flavonoids Pergmon press Co. London, 126.
- Hammerschmidt, R. (1999):** phytoalexins: what have we learned after 60 years? ANNU. Rev. phytopathal, 37: 285-306.
- Hanson, J.R. (1972):** Chemistry of terpenes and terpenoids (Ed. A.A. Newman). Academic Press. New York. Pp. 155-206.
- Harborne, J.B. (1998):** Phytochemical methods. A. guide to modern texhniques of plant analysis. New York, USA pp 54-150.
- Hewady, M.A.A.; El-Sherif, L.S. and Omar, A.M. (1994):** Evaluation of four plant oils against newly hatched larvae of the cotton bollworms; *Pectinophora gossypiella* (Saund) and *Earias insulana* (Boisd.) (*Lepidoptera: Noctuidae*). Annals of Agricultural science, moshtohor, 32; 4: 2097-2104.
- Ibrahim, M. M. (1991):** Genetic studies on lemongrass. M. Thesis, Fac. Agric. Ain Shams Univ. Egypt.
- Inyang, Y.E. and Emosairue, S.O. (2005):** Laboratory assessment of the repellent and antifeedant properties of aqueous extracts of 13 plants against the banana weevil

- Cosmopolites sordidus* German (Coleoptera: Curculionidae). Tropical and Subtropical Agroecosystems, 5(1): 33-44.
- Khan, M.; Syamasundar, S.K.; Singh, K.V. and Nagvi, A.A. (2002):** Chemical composition of leaf and flower essential oil of *Lantana camara* from India. Flavour and Fragrance Journal, 17(1): 75-77.
- Khan, M.; Syamasundar, S.K.; Neetujain, S.K. and Yadav, A.K. (2003):** Chemical composition of fruit and stem essential oils of *Lantana camara* from northern India. Flavour and Fragrance Journal 18(5): 376-379.
- Kinsella, J.E. (1966):** Metabolic patterns of the fatty acids of *periplaneta americana* during the embryonic development. Can. J. Biochemistry 44: 247-258.
- Kulkarni, N.; Joshi, K.C. and Gupta, B.N. (1997):** Antifeedant property of *Lantana camara* var. *aculeata* and *Aloe vera* leaves against teak skeletonizer, *Eutectona machaeralis* Walk. (Lepidoptera: pyralidae). Entomon. 22 (1): 61-56.
- Kumari, D.A.; Kumar, S.T. and Reddy, V.S. (1998):** management of the groundnut bruchid, *Caryedon serratus* (Olivier) with some plant products, Pest-management-and-Economic-Zoology, 6:2,127-131.
- Lewis, H. and Smith, C. (1967):** Sugar alcohols in fungi and green plants. Methods of detection and estimation. New Phytol. ; 66, 185-204. Referred to Harbrone, J.B. (1998). Phytochemical Methods, a guide to modern techniques of plant analysis, p. 54, New York, USA.

- Lowry O.H.; Rosebrough, N. J.; Farr, A.L. and Randall, R.J. (1951):** Protein measurements with the palm phenol reagent J. Biol. Chem 193-265.
- Martin, J.S.; Martin, M. and Bernays, E. A. (1987):** Failure of tannic acid to inhibit digestion or reduce digestibility of plant protein ingut fluids of insect.
- Mehta P.K.; Sood, A.K.; Parmar, S. and Kashyap, N.P. (2002):** Antifeedant activity of some plants of North Western Himalayas against cabbage caterpillar, *Pieris brassicae* (L.). Journal-of-Entomological-Research, 26:1, 51-54.
- Mehta, P. K.; Vaidya, D.N. and Kashyap, N.P. (1995):** Antifeedant properties of some plant extracts against brinjal hadda beetle, *Henosepilachna vigintioctopunctata* F. Journal-of-Entomological-Research, 19:2, 147-150.
- Ming, L.C.; Figueiredo, R.O.; Marchado, S.R.; Andrade, R.M.C.; Craker, L.E. (ed.); Nolan, L. (ed.) and Shetty, K. (1996):** Yield of essential oil of and citral content in different parts of lemongrass leaves (*Cymbopogon citratus* (D.C.) Stapf.) Poaceae. International symposium on medicinal and aromatic plant, Amherst, Massachusetts, USA Acta-horticulture No. 426, 555-559.
- Mohd, A.; Indu, S. and Onkar, S. (2004):** Volatile constituents of *Cymbopogon citratus* Stapf. Leaves. Journal of Essential Oil Bearing Plants. 8(1): 56-59.
- Morsy T.A.; Mazyad, S.A. and El-Aharkawy, I.M.A. (1998):** The larvicidal activity of solvent extracts of three medicinal plants against third instar larvae of *Chrysomya*

*albiceps*. Journal-of-the Egyptian-Society-of-parasitology. 28:3, 699-709.

- Muhamed, S.; Nighat, A.; Anwar, M.A.; Ali, S.M.A.; Shahida, S. and Rahman, A. (2003):** Chemistry and biological significance of essential oils of *Cymbopogon citratus* from Pakistan-natural Product Research 17(3); 159-163.
- Osman, S.M. and Radwan, O.M. (2004):** Isolation and Identification of active components in some plant extracts and their effect on *Agrotis ipsilon* (Hufn). Egyptian Journal of Biological pest Control 14(1): 181-185.
- Pandey, N.D.; Singh, L.; Singh, Y. P. and Tripathi, R. A. (1987):** Effect of certain plant extracts against *Lipaphis erysimi* Kalt. Under laboratory conditions. Indian J. Entomol. 49 (2): 238-242.
- Pandey, N.D.; Mathur, K.K.; Pandey, S. and Tripathi, R. A. (1986):** Effects of some plant extracts against pulse beetle, *Callosobruchus chinensis* Linnaeus. India J. Entomol. 48 (1): 85-90.
- Pino, J.A.; R. Marbot; Rosado, A.; Romeu, C. and Marti, M.P. (2004):** Chemical composition of the essential oil of *Lantana camara* L. from Cuba. Journal of essential Oil Research 16(3): 216-218.
- Rajesh, K.V. and Verma, S. K. (2006):** Phytochemical and termiticidal study of *Lantana camara* var. *Aculeata* leaves. Fitoterapia 20: 1-3.

- Raman, K.V.; Booth, R.H. and Palacios, M. (1987):** Control of potato tuber moth *phthoremaea operculella* (zeller) in rustic potato stores. *Tropical. Sic.* 27 (3): 175-194.
- Randrianalyaona, J.A.; Ramanoebina, P.A.R.; Rasoarahona, J.R.E. and Gaydou, E.M. (2005):** Seasonal and chemotype influences on the chemical composition of *Lantana camara L.*: essential oils from Madagascar. *Analytica. Chimica Acta* 545(1): 46-52.
- Reddy, G.V.P.; Urs, K.C.D.; Grawal, O.P.A. and Shashi, D. (1991):** Antifeedant and repellent activity of some indigenous plant extracts against the furniture beetle, *Sinoxylon sudanicum*. Lesene (Coleopatera: Bostrychidae). Biodeterioration of cultural property. Proceeding of the International Conference India: 173-185.
- Rozzi, N.L.; Phippen, W.; Simon, J.E. and Sing, R.K. (2002):** Supercritical fluid extraction of essential oil components from lemon-scented botanicals. *Lebensmittel Wissenschaft and Technologie* 35(4): 319-324.
- Sargent S.R. and Lancas, F.M. (1997):** Supercritical fluid extraction of *Cymbopogon citratus*(D.C.) Stapf. *Chromatographia*, 46: 5-6, 285-290.
- SAS Institute (1988):** SAS user's guide statistics. SAS Institute, Cary N.C.
- Schaneberg, B.T. and Khan, I.A. (2002):** Comparison of extraction methods for marker compounds in the essential oil of lemon grass by GC. *J. of Agric. And Food Chemistry* 50(6):" 1345-1349.

- Sharaby A. (1988).** Anti-insect properties of the essential oil of lemongrass, *Cymbopogon-citratus* against the lesser cotton leaf worm *Spodoptera exigua* (Hbn). *Insect-science and-its-Applications*. 9:1,77-80.
- Shellard, E.J. (1957):** Practical plant chemistry. London Ritman Medical Publishing Co. Ltd.
- Shen, B.Z.; Zheng, Z. W. and Dooner, H. K. (2000):** A maize sesquiterpene cyclase gene induced by insect herbivory and volicitin: Characterization of wild- type and mutant alleles. *Proc. Nat. Acad. Sci. USA*, 97:14807-14812.
- Shenge, K. C.; Urah, I.I. and Adamu, R.S. (2003):** Effect of different lemongrass powders on *Sitophilus zeamais* moth infesting stored maize grain. *journal of sustainable agriculture and the environment* 4(1): 23-28.
- Stahle, E. (1969):** Thin-layer chromatography, Springer-Verlag. Berlin, Heidelberg, New York, P. 439- 441, 856 and 873.
- Stein, U. and Klingauf, F. (1990):** Insecticidal effect of plant extracts from tropical and subtropical species. Traditional methods are good as long as they are effective. *J. Appl. Entmol.* 110 (2): 160-166.
- Sundufu, A. J. and Shoushan, H. (2004):** Chemical composition of the essential oils of *Lantana camara* L. occurring in south China. *Flavor and fragrance Journal* , 19(3): 229-232.
- Tanzubil, P.B. and McCaffery, A.R. (1990):** Effect of azadirachtin and aqueous neem seed extract on survival