

CONTENTS

	Page
1. INTRODUCTION	1
2. REVIEW OF LITERATURE	3
2.1. Wheat flour	3
2.1.1. Chemical composition	3
2.1.2. Rheological properties of wheat flour dough	4
2.2. Barley	5
2.2.1. Chemical composition	5
2.2.2. Effect of barley flour addition to wheat flour on the rheological properties of dough	9
2.2.3. Effect of barley on serum cholesterol levels in experimental animals	15
2.3. Partially defatted soybean flour	25
2.3.1. Chemical composition	25
2.3.2. Effect of addition of soybean flour to wheat flour on the rheological properties and doughs	27
2.3.3. Effect of soybean addition diets on lowering cholesterol and triglycerides in experimental animals	35
3. MATERIALS AND METHODS	39
3.1. Materials	39
3.1.1. Wheat flour (82% extraction)	39
3.1.2. Hull- less barely flour (70% extraction)	39
3.1.3. Partially defatted soybean flour	39
3.1.4. Experimental animals	39
3.1.5. Diagnostic kits	39
3.2. Methods	40
3.2.1. Chemical composition	40
3.2.2. Rheological characteristics	43
3.2.2.1. Farinograph test	43
3.2.2.2. Extensograph test	43
3.3. Preparation of balady bread	43

	Page
3.4. Organoleptic evaluation of balady bread	44
3.5. Biological experiment	45
3.5.1. Induction of hyperlipaemia	45
3.5.2. Experiment design	45
3.5.3. The prophylactic effect of different treatments against hyperlipaemia	47
3.6. Blood sampling	47
3.7. Biochemical analysis	48
3.7.1. Serum triglycerides	48
3.7.2. Serum total cholesterol	48
3.7.3. Serum LDL-cholesterol	48
3.7.4. Serum HDL-cholesterol	48
3.7.5. Serum transaminases	48
3.7.6. Serum creatinine	48
3.7.7. Blood urea	49
3.7.8. Hemoglobin	49
3.8. Statistical analysis	49
4. RESULTS AND DISCUSSION	51
4.1. Chemical composition	51
4.1.1. Moisture content	51
4.1.2. Ash content	51
4.1.3. Crude protein content	53
4.1.4. Lipids content	53
4.1.5. Crude fiber content	53
4.1.6. Carbohydrate content	54
4.1.7. Sugar content	54
4.1.8. Gluten content (of total protein)	54
4.1.9. Pigments content	55
4.1.10. Phytate content	55
4.1.11. Falling number (Alpha-amylase activity)	56
4.2. Rheological characteristics of dough	56
4.2.1. Farinograph properties	56

	Page
4.2.1.1. Effect of mixing wheat flour (82% extraction) with hull-less barley flour (70% extraction) on farinograph properties	56
4.2.1.2. Effect of mixing wheat flour (82% extraction) with partially defatted soybean flour on farinograph properties	60
4.2.1.3. Effect of mixing wheat flour (82% extraction) with hull-less barley flour (70% extraction) and partially defatted soybean flour on farinograph properties	64
4.2.2. Extensograph properties	67
4.2.2.1. Effect of mixing wheat flour (82% extraction) with hull-less barley flour (70% extraction) on extensograph properties	67
4.2.2.2. Effect of mixing wheat flour (82% extraction) with partially defatted soybean flour on its extensograph properties	70
4.2.2.3. Effect of mixing wheat flour (82% extraction) with hull-less barley flour (70% extraction) and partially defatted soybean flour on extensograph properties	73
4.3. Organoleptic evaluation	76
4.3.1. Organoleptic evaluation of balady breads produced from wheat flour (82% extraction) mixed with hull-less barley flour (70% extraction)	76
4.3.2. Organoleptic evaluation of balady breads produced from wheat flour (82% extraction) mixed with partially defatted soybean flour	79
4.3.3. Organoleptic evaluation of balady breads produced from wheat flour (82% extraction) mixed with hull-less barley flour (70% extraction) + partially defatted soybean flour	82
4.4 Biological evaluation	85

	Page
4.4.1. Effect of hypercholesterolemic diet mixed with wheat flour (82% extraction), hull-less barley flour (70% extraction) and partially defatted soybean flour on body weight, food intake and feed conversion ratio in rats after the end of experimental period (8 week)	85
4.4.2. Effect of different experimental diets on serum triglycerides, total cholesterol, LDL-cholesterol, HDL-cholesterol and risk ratio in hypercholesterolemic rats after the end of experimental period (8 weeks)	88
4.4.3. Effect of different experimental diets on liver functions in hypercholesterolemic rats	94
4.4.4. Effect of replacing wheat flour (82% extraction) with hull-less barley flour (70% extraction) and partially defatted soybean flour at different levels on kidneys functions in hypercholesterolemic rats after 8 weeks	97
4.4.5. Effect of replacing wheat flour (82% extraction) with hull-less barley four (70% extraction) and partially defatted soybean flour at different levels on hemoglobin in hypercholesterolemic rats after 8 weeks	100
5. SUMMARY	103
6. REFERENCES	113
7. ARABIC SUMMARY	