

REFERENCES

- **Aguayo A and Patt YZ.** Non-surgical treatment of hepatocellular carcinoma. *Semin Oncol* 28: 503-513(**2001**).
- **Ahrar K, Newman RA, Pang J, et al.** Relative Thermosensitivity of cytotoxic drugs used in transcatheter arterial chemoembolization. *J Vasc Interv Radiol* 15: 901-905 (**2004**).
- **Allison DJ:**diagnostic radiology (3rd edn.)Churchill Livingstone, Edinburgh(**1997**).
- **Bedikian AY, Legha SS, Mavligit G et al.** Treatment of uveal melanoma metastatic to the liver: a review of the M. D. Anderson Cancer Center experience and prognostic factors. *Cancer* 76:1665–1670(**2005**).
- **Berber E, Pelley R, Siperstein A.** Predictors of survival after radiofrequency thermal ablation of colorectal cancer metastases to the liver: a prospective study. *J Clin Oncol* 23:1358–136 (**2005**) .
- **Bloomston M, Binitie O, Fraiji E, et al.** Transcatheter arterial chemoembolization with or without radiofrequency ablation in the management of patients with advanced hepatic malignancy. *Am Surg* 68 (9),827-831 (**2007**).
- **Brown DB, Cardella JF, Sacks D, et al.** Quality improvement guidelines for transhepatic arterial chemoembolization, embolization, and chemotherapeutic infusion for hepatic malignancy. *J Vasc Interv Radiol* 17: 225-232 (**2008**).
- **Brown DB, Geschwind JF, Soulen MC, et al.** Society of interventional radiology position statement on chemoembolization of hepatic malignancies. *J Vasc Interv Radiol* 17: 217-223 (**2006**).

- **Bruix J, Llovet JM, Castells A; et al.** Transarterial embolization versus symptomatic treatment in patients with advanced hepatocellular carcinoma: results of a randomized, controlled trial in a single institution. *Hepatology*. 27(6):1578-1583 (**2002**).
- **Bruix J and Sherman M.** Management of hepatocellular carcinoma. *Journal of Hepatology* 42: 1208-1236 (**2005**).
- **Bruix J, Sherman M, Llovet JM, et al.** Clinical Management of Hepatocellular carcinoma: conclusions of the Barcelona-2000 EASL Conference. *J Hepatol* 35: 421-430 (**2001**).
- **Cady B, Stone MD.** The role of surgical resection of liver metastases in colorectal carcinoma. *Semin Oncol* 18:399–406(**2002**).
- **Choi H, Loyer EM, DuBrow RA, et al.** Radiofrequency ablation of liver tumors: assessment of therapeutic response and complications. *RadioGraphics* 21: S41-S54 (**2001**).
- **Covey AM, Brody LA, Getrajdman GI, et al.** Incidence, patterns, and clinical relevance of variant portal vein anatomy. *AJR* 183: 1055-1064 (**2004**).
- **Covey AM, Brody LA, Maluccio MA, et al.** Variant hepatic arterial anatomy revisited: digital subtraction angiography performed in 600 patients. *Radiology* 224: 542-547 (**2002**).
- **Curley S, Izzo F, Delrio P et al.** Radiofrequency ablation of unresectable primary and metastatic hepatic malignancies: results in 123 patients. *Ann Surg*; 230:1–8 (**2003**).
- **Donna D'souza, Nagorney DM et al.** Tumor and liver drug uptake following hepatic artery and portal vein infusion. *J Clin Oncol* 5:1836–1840 (**2008**).

- Ensminger WD, Gyves JW. Clinical pharmacology of hepatic arterial chemotherapy. *Semin Oncol* 10:176–182 (1998).
- Florman S and Miller MC. Live donor liver transplantation. *Liver Transplantation* 12: 499-510 (2006).
- Gates J, Okamura J, Kuroda C. Transcatheter chemo-embolization effective for treating hepatocellular carcinoma: a radiological and histopathologic study. *Cancer*. 54(3):387-392 (2003).
- Gillams AR, Lees WR. Radiofrequency ablation of colorectal liver metastases. *Abdom Imaging* 30:419–426 (2005).
- Gilbert HA, Kagan AR. Metastases: incidence, detection, and evaluation without histologic confirmation. In: Weiss L, ed. *Fundamental Aspects of Metastasis*. Amsterdam: North-Holland 385–405(2000).
- Goldberg SN, Grassi CJ, Cardella JF, et al. Image guided tumor ablation: standardization of terminology and reporting criteria. *J Vasc Interv Radiol* 16: 765-778 (2005).
- Goldberg SN, Harrison LE, et al. Locoregional recurrences are frequent after radiofrequency ablation for hepatocellular carcinoma. *J Am Coll Surg* 197: 759-764(2000).
- Goldberg SN, Kamel IR, Kruskal JB, et al. Radiofrequency ablation of hepatic tumors: increased tumor destruction with adjuvant liposomal doxorubicin therapy. *AJR* 179: 93-101 (2002).
- Goldberg SN, Ono K, et al. Radiofrequency ablation of hepatocellular carcinoma: Correlation between local tumor progression after ablation and ablative margin. *Ann rad.* 24: 480-488 (1989).

- **Grieco A, Pompili M, Caminiti G, et al.** Prognostic factors for survival in patients with early-intermediate hepatocellular carcinoma undergoing non-surgical therapy: comparison of Okuda, CLIP, and BCLC staging systems in a single Italian centre. *Gut* 54: 411- 418 (2005).
- **Haaga JR, Lanzieri CF, Gilkeson RC, et al.** Computed Tomography and Magnetic Resonance Imaging of the Whole Body. Mosby-Year Book, Inc: St. Louis, Mo. 4th ed., 1318-1340 (2003).
- **Harnois DM, Ren ZG, et al.** Postoperative adjuvant arterial chemoembolization improves survival of hepatocellular carcinoma patients with risk factors for residual tumor: a retrospective control study. *World J Gastroenterol* 10: 2791-2794 (2003).
- **Hayashi PH and Di Bisceglie AM.** The progression of hepatitis B- and C-Infections to chronic liver disease and hepatocellular carcinoma: presentation, diagnosis, screening, prevention, and treatment of hepatocellular carcinoma. *Infect Dis Clin N Am* 20: 1-25 (2006).
- **Heslin MJ, Medina-Franco H, Parker M et al.** Colorectal hepatic metastases: resection, local ablation, and hepatic artery infusion pump are associated with prolonged survival. *Arch Surg* 2001;136:318–323. Erratum in: *Arch Surg* 136:809(2001).
- **Hussain SM, Zondervan PE, IJzermans JN, et al.** Benign versus malignant hepatic nodules: MR imaging findings with pathologic correlation. *RadioGraphics* 22: 1023-1039 (2002).

- **Hyo Soon Lim, Saitoh S, et al.** Effect of repeated transcatheter arterial embolization on the survival time in patients with hepatocellular carcinoma. An analysis by the Cox proportional hazard model. *Cancer* 68:2150-2154 (**2006**).
- **Irfan A and Dileep NL.** Malignant tumors of the liver. *Surgery* 25 (1): 34-41 (**2006**).
- **Jonathan Koea, John McCall, et al,** The role of radiofrequency ablation in the treatment of primary and metastatic tumours of the liver: initial lessons learned, *N Engl J Med* 324: 355–372 (**2007**).
- **Kim YS, Moretto P, et al.** Radiofrequency thermal ablation (RFA) after transarterial chemoembolization (TACE) as a combined therapy for unresectable non-early hepatocellular carcinoma (HCC). *Eur. Radiol* 16: 661-669 (**2006**).
- **Kobayahi M, Cheng BQ, et al.** Chemoembolization combined with radiofrequency ablation for patients with hepatocellular carcinoma larger than 3 cm: a randomized controlled trial. *JAMA* 299: 1669-1677(**2007**).
- **Kokudo N, Llovet JM, et al.** Systematic review of randomized trials for unresectable hepatocellular carcinoma: chemoembolization improves survival. *Hepatology*. 37(2):429-442 (**2000**).
- **Koops A, Wojciechowski B, Broering DC, et al.** Anatomic variations of the hepatic arteries in 604 selective celiac and superior mesenteric angiographies. *Surg Radiol Anat* 26: 239-244 (**2004**).
- **Koteish A and Thuluvath PJ.** Screening for hepatocellular carcinoma. *J Vasc Interv Radiol* 13: S185-S190 (**2002**).

- Lang EK, Brown CL Jr. Colorectal metastases to the liver: selective chemoembolization. *Radiology* 189:417–422(2000).
- Lau WY, Yu CH, Lai CH, et al. Transarterial chemoembolization for hepatocellular carcinoma. *J Am Coll Surg* 202: 155-168 (2006).
- Lencioni R, Pina CD and Bartolozzi C. Percutaneous image-guided radiofrequency ablation in the therapeutic management of hepatocellular carcinoma. *Abdom Imaging* 30: 401-408 (2005).
- Lencioni R, Kobayahi, et al. DEB-Enhanced RFA of HCC: A Pilot Study, *J Hepatol* 176: 34-63(2008).
- Lencioni R, Cioni D and Bartolozzi C. Focal Liver Lesions: Detection, Characterization, Ablation. Springer-Verlag, New York, 1st edn., 63-72 (2005).
- Lencioni R, Pina CD, Bartolozzi C, et al. Percutaneous image-guided radiofrequency ablation in the therapeutic management of hepatocellular carcinoma. *Abdom Imaging* 30:401-408 (2005).
- Lim HK, Di Stasi M, et al. Percutaneous radiofrequency interstitial thermal ablation in the treatment of small hepatocellular carcinoma. *Cancer J Sci Am* 18: 73-81 (2001).
- Limanond P, Zimmerman P, Raman S, et al. Interpretation of CT and MRI after radiofrequency ablation of hepatic malignancies. *AJR* 181: 1635-1640 (2003).
- Lippincott Williams & Wilkins Atlas of Anatomy, 8st Edn, 4:322 (2008).
- Liu LX, Zhang WH, Jiang HC. Current treatment for liver metastases from colorectal cancer. *World J Gastroenterol* 9:193–200 (2003).

- **Liu YM, Hsu CW, et al.** Randomised controlled trial comparing percutaneous radiofrequency thermal ablation, percutaneous ethanol injection, and transarterial chemoembolization to treat hepatocellular carcinoma of 3 cm or less. *Gut* 54: 1151-1156 (**2005**).
- **Livraghi T, Goldberg SN, Lazzaroni S, et al.** Hepatocellular carcinoma: radiofrequency ablation of medium and large lesions. *Radiology* 214: 761-768 (**2000**).
- **Louse ME, Lee RG, Duszlak EJ et al.** Peripheral hepatic artery embolization for primary and secondary hepatic neoplasms. *Radiology*;147: 407–411(2001).
- **Marrero JA, Fontana RJ, Barrat A, et al.** Prognosis of hepatocellular carcinoma: comparison of 7 staging systems in an American cohort. *Hepatology* 41: 707- 716 (**2005**).
- **Mavligit GM, Zukwiski AA, Ellis LM et al.** Gastrointestinal leiomyosarcoma metastatic to the liver. Durable tumor regression by hepatic chemoembolization infusion with cisplatin and vinblastine. *Cancer*;75: 2083–2088(**2002**).
- **McGahan JP and Goldberg BB.** Diagnostic ultrasound; a logical approach. Lippincott-Raven Publishers, Philadelphia, 599-691 (**1998**).
- **Moertel CG.** Karnofsky memorial lecture. An odyssey in the land of small tumors. *J Clin Oncol* 5:1502–1522(**1987**).
- **Mortelé KJ and Ros PR.** Anatomic variants of the biliary tree: MR cholangiographic findings and clinical applications. *AJR* 177: 389-394 (**2001**).

- Nakamura H, Hashimoto T, Oi H, et al. Treatment of hepatocellular carcinoma by segmental hepatic artery injection of Adriamycin-in-oil emulsion with overflow to segmental portal veins. *Acta Radiol* 31: 347-349(1990).
- Ni Y, Mulier S, Miao Y, et al. A review of the general aspects of radiofrequency ablation. *Abdom Imaging* 30: 381-400 (2005).
- Oshowo A, Gillams A, Harrison E et al. Comparison of resection and radiofrequency ablation for treatment of solitary colorectal liver metastases. *Br J Surg*;90: 1240–1243(2003).
- Perry LJ, Stuart K, Stokes KR et al. Hepatic arterial chemoembolization for metastatic neuroendocrine tumors. *Surgery* 116:1111–1116; discussion 1116–1117(2004).
- Prabhu VR and Burt AD. Pathology of liver tumors. *Surgery* 25 (1): 10-15 (2006).
- Reappraisal of hepatic arterial infusion in the treatment of nonresectable liver metastases from colorectal cancer. Meta-Analysis Group in Cancer. *J Natl Cancer Inst* 88:252–258(1996).
- Rhim H. Complications of radiofrequency ablation in hepatocellular carcinoma. *Abdom Imaging* 30: 409- 418 (2008).
- Rhim H, Dodd GD III. Radiofrequency thermal ablation of liver tumors. *J. Clin. Ultrasound* 27: 221-229 (1999).
- Rhim H, Goldberg SN, Dodd GD, et al. Essential techniques for successful radiofrequency thermal ablation of malignant hepatic tumors. *RadioGraphics* 21: S17-S39 (2001).

- Sahani D, D'souza R, Kadavigere R, et al. Evaluation of living liver transplant donors: method for precise anatomic definition by using a dedicated contrastenhanced MR imaging protocol. *RadioGraphics* 24: 957-958 **(2004)**.
- Sanz-Altamira PM, Spence LD, Huberman MS et al. Selective chemoembolization in the management of hepatic metastases in refractory colorectal carcinoma: a phase II trial. *Dis Colon Rectum* 40:770–775**(1999)**.
- Snell RS. *Clinical Anatomy*. Lippincott Williams, 7th edn, 183-274 **(2004)**.
- Solbiati L, Ierace T, Tonolini M et al. Radiofrequency thermal ablation of hepatic metastases. *Eur J Ultrasound* 13:149–158**(2001)**.
- Sugimori K, Nozawa A, Morimoto M, et al. Extension of radiofrequency ablation of the liver by transcatheter arterial embolization with iodized oil and gelatin sponge: results in a pig model. *J Vasc Interv Radiol* 16: 849-856 **(2005)**.
- Szklaruk J, Silverman PM, and Charnsangavej C. Imaging in the diagnosis, staging, treatment, and surveillance of hepatocellular carcinoma. *AJR* 180: 441-454 **(2003)**.
- Takayasu K, Adam R, Bismuth H, et al. Influence of preoperative transarterial lipiodol chemoembolization on resection and transplantation for hepatocellular carcinoma in patients with cirrhosis. *Ann Surg* 226: 688-703**(1998)**.

- Trevisani F, D'Intino PE, Morselli-Labate AM, et al. Serum alpha-fetoprotein for diagnosis of hepatocellular carcinoma in patients with chronic liver disease: influence of HBsAg and anti-HCV status. *J Hepatol* 34: 570-575 (2008).
- Yamaguchi A, Taniguchi H, Kunishima S et al. Correlation between angiographically assessed vascularity and blood flow in hepatic metastases in patients with colorectal carcinoma. *Cancer* 89:1236–1244 (2000).
- Yamakado K, Nakatsuka A, Akeboshi M, et al. Combination therapy with radiofrequency ablation and transcatheter chemoembolization for the treatment of hepatocellular carcinoma: Short-term recurrences and survival. *Oncology Reports* 11: 105-109 (2007).
- Yamakado K, Nakatsuka A, Ohmori S, et al. Radiofrequency ablation combined with chemoembolization in hepatocellular carcinoma: treatment response based on tumor size and morphology. *J Vasc Interv Radiol* 13: 1225-1232 (2004).