

References

Acharya UR, Yun WL, Yu W (2008): Imaging Systems of Human Eye: A Review. *J Med Syst*; 32: 301–315.

Acott TS and Kelley MJ (2008): Extracellular Matrix in the Trabecular Meshwork. *Exp Eye Res*; 86(4): 543–561.

Alencar LM, Bowd C, Weinreb RN et al (2008): Comparison of HRT-3 Glaucoma Probability Score and Subjective Stereophotograph Assessment for Prediction of Progression in Glaucoma. *Invest Ophthalmol Vis Sci*; 49: 1898–1906.

Antón A, Castany M, Pazos-Lopez M et al. (2009): Reproducibility of measurements and variability of the classification algorithm of Stratus OCT in normal, hypertensive, and glaucomatous patients. *Clinical Ophthalmology*; 3: 139–145.

Badala F, Nouri-Mahdavi K, Raoof DA et al. (2007): Optic disk and nerve fiber layer imaging to detect glaucoma. *Am J Ophthalmol*; 144: 724–732.

Baleriola J, García-Feijoo J, Martínez-de-la-Casa JM et al (2008): Apoptosis in the trabecular meshwork of glaucomatous patients. *Molecular Vision*; 14:1513-1516.

- Bertuzzi F, Hoffman DC, Fonseka AM (2009):** Concordance of Retinal Nerve Fiber Layer Defects between Fellow Eyes of Glaucoma Patients Measured by Optical Coherence Tomography. *Am J Ophthalmol*; 148: 148–154.
- Bizheva K, Pflug R, Hermann B, et al. (2006):** Optophysiology: depth-resolved probing of retinal physiology with functional ultrahighresolution optical coherence tomography. *Proc Natl Acad Sci USA*; 103: 5066-5071.
- Blumenthal EZ, Williams JM, Weinreb RN et al. (2000):** Reproducibility of nerve fiber layer thickness measurements by use of optical coherence tomography. *Ophthalmology*; 107(12): 2278–2282.
- Bohringer HJ, Lankenau E, Rohde V (2006):** Optical coherence tomography for experimental neuroendoscopy. *Minim Invasive Neurosurg*; 49: 269-275.
- Borque E, Ferreras A, Polo V et al. (2008):** Evaluation of four new discriminant functions for HRT II in glaucoma diagnosis. *Arch Soc Esp Oftalmol*; 83: 349-356.
- Bowd C, Chan K, Zangwill LM et al. (2002):** Comparing neural networks and linear discriminant functions for glaucoma detection using confocal scanning laser ophthalmoscopy of the optic disc. *Invest Ophthalmol Vis Sci*; 43(11): 3444–3454.
- Bower BA, Zhao M, Zawadzki RJ et al. (2007):** Real-time spectral domain Doppler optical coherence tomography and investigation of human retinal vessel autoregulation. *J Biomed Opt*; 12: 412-414.

- Budenz DL, Fredette MJ, Feuer WJ et al. (2008):** Reproducibility of peripapillary retinal nerve fiber thickness measurements with Stratus OCT in glaucomatous eyes. *Ophthalmology*; 115: 661-666.
- Bugueta A and Rumanet JP (1994):** 24-hour and sleep related variations of intraocular pressure in healthy white individuals. *Am. J. Ophthalmol*; 117: 342-347.
- Burgansky-Eliash Z, Wollstein G, Bilonick RA et al. (2007):** Glaucoma detection with the Heidelberg retina tomograph 3. *Ophthalmology*; 114: 466–471.
- Burk RO, Vihanninjoki K, Bartke T et al. (2000):** Development of the standard reference plane for the Heidelberg retina tomograph. *Graefes Arch Clin Exp Ophthalmol*; 238(5): 375–384.
- Carpineto P, Ciancaglini M, Zuppardi E et al. (2003):** Reliability of nerve fiber layer thickness measurements using optical coherence tomography in normal and glaucomatous eyes. *Ophthalmology*; 110 (1): 190–195.
- Chan A, Duker JS, Ishikawa H et al. (2006):** Quantification of photoreceptor layer thickness in normal eyes using optical coherence tomography. *Retina*; 26: 655-660.
- Chang RT and Budenz DL (2008):** Diagnosing glaucoma progression. *International Ophthalmology Clinics*; 28: 13-28.

- Chauhan BC, Blanchard JW, Hamilton DC et al. (2000):** Technique for detecting serial topographic changes in the optic disc and peripapillary retina using scanning laser tomography. *Invest Ophthalmol Vis Sci*; 41: 775–782.
- Chauhan BC and Marshall J (1999):** The Interpretation of Optical Coherence Tomography Images of the Retina. *Invest Ophthalmol Vis Sci*; 40: 2332–2342.
- Chauhan BC and McCormick TA. (1995):** Effect of the cardiac cycle on topographic measurements using confocal scanning laser tomography. *Graefes Arch Clin Exp Ophthalmol*; 233: 568–572.
- Chauhan BC, McCormick TA, Nicolela MT et al. (2001):** Optic disc and visual field changes in a prospective longitudinal study of patients with glaucoma: comparison of scanning laser tomography with conventional perimetry and optic disc photography. *Arch Ophthalmol*; 119(10): 1492–1499.
- Chauhan BC, Mikelberg FS, Balaszi AG et al. (2008):** Risk Factors for the Progression of Open-angle Glaucoma. *Arch Ophthalmol*; 126(8):1030-1036.
- Cheng E (2009):** For which glaucoma suspects is it appropriate to initiate treatment? *Ophthalmology*; 116(4): 710–717.
- Chisholm IA, Drance SM and To T (1989):** The glaucoma suspect: differentiation of the future glaucoma eye from the non-glaucomatous suspect eye. *Graefes Arch Clin Exp Ophthalmol*; 227: 110-113.

- Choi JA, Lee NY and Park CK (2009):** Interpretation of the Humphrey Matrix 24-2 Test in the Diagnosis of Pre-perimetric Glaucoma. *Jpn J Ophthalmol*; 53: 24–30.
- Choi SS, Zawadzki RJ, Greiner MA (2008):** Fourier-Domain Optical Coherence Tomography and Adaptive Optics Reveal Nerve Fiber Layer Loss and Photoreceptor Changes in a Patient With Optic Nerve Drusen. *J Neuroophthalmol*; 28(2): 120–125.
- Christopoulos V, Kagemann L, Wollstein G et al. (2007):** In vivo corneal high-speed, ultra high-resolution optical coherence tomography. *Arch Ophthalmol*; 125: 1027-1035.
- Coleman AL and Kodjebacheva G (2009):** Risk Factors for Glaucoma Needing More Attention. *The Open Ophthalmology Journal*; 3; 38-42.
- Coleman AL and Miglior S (2008):** Risk Factors for Glaucoma Onset and Progression. *Surv Ophthalmol*; 53: S3-S10.
- Coops A, Henson DB, Kwartz AJ et al (2006):** Automated Analysis of Heidelberg Retina Tomograph Optic Disc Images by Glaucoma Probability Score. *Invest Ophthalmol Vis Sci*; 47:5348–5355.
- Costa RA, Skaf M, Melo AS (2006):** Retinal assessment using optical coherence tomography. *Progress in Retinal and Eye Research* 25; 325–353.

- Cristina ML and Marvin JP (1993):** Intraocular pressure, Cardiovascular risk variables and Visual field defects. *Am. J. Epidemiol*; 118: 280-287.
- Daniel MS, Gadi W and Joel SS (2007):** Imaging in glaucoma. *Ophthalmol. Clin. North Am*; 17: 33-52.
- De Leon-Ortega JE, Sakata LM, Monheit BE et al. (2007):** Comparison of diagnostic accuracy of Heidelberg retina tomograph II and Heidelberg retina tomograph 3 to discriminate glaucomatous and nonglaucomatous eyes. *Am J Ophthalmol*; 144: 525–532.
- Devogelaere T and Dirven W (2007):** Drusen de la papille. *J Fr. Ophtalmol*; 30: 966-967.
- Dimitrios K, Gerasimos G, Antonios M et al. (2009):** New nonlinear multivariable model shows the relationship between central corneal thickness and HRTII topographic parameters in glaucoma patients. *Clinical Ophthalmology*; 3: 313–323.
- Ding Z, Ren H, Zhao Y et al. (2002):** High-resolution optical coherence tomography over a large depth range with an axicon lens. *Surv Ophthalmol*; 53: 93-110.
- Drexler W (2004):** Ultrahigh-resolution optical coherence tomography. *J Biomed Opt*; 9: 47-74.
- Drexler W and Fujimoto JG (2008):** State-of-the-art retinal optical coherence tomography. *Prog Ret Eye Res*; 27: 45-88.

- Drexler W, Morgner U, Schuman JS et al. (2001):** Ultrahigh-resolution ophthalmic optical coherence tomography. *Nat Med*; 7: 502-507.
- Drexler W, Sattmann H, Hermann B, et al. (2003):** Enhanced visualization of macular pathology with the use of ultrahigh-resolution optical coherence tomography. *Arch Ophthalmol*; 121: 695-706.
- Eduardo AA, Fernando CY and Mario RR (2001):** Evaluation and definition of physiologic macro cups with confocal optic nerve analysis (HRT). *International Ophthalmology* 23: 239–244.
- Eid TM, Spaeth GL, Katz LJ et al. (2000):** Quantitative estimation of retinal nerve fiber layer height in glaucoma and the relationship with optic nerve head topography and visual field. *J Glaucoma*; 6 (4): 221–230.
- Fayers T, Strouthidis NG, Garway-Heath DF. (2007):** Monitoring glaucomatous progression using a novel Heidelberg retina tomograph event analysis. *Ophthalmology*; 114: 1973–1980.
- Felipe AM, Linda MZ, Christopher B et al. (2006):** Influence of Disease Severity and Optic Disc Size on the Diagnostic Performance of Imaging Instruments in Glaucoma. *Invest. Ophthalmol. Vis. Sci*; 42:1063-1078.
- Fercher AF, Hitzenberger CK, Drexler W et al. (1993):** In vivo optical coherence tomography. *Am J Ophthalmol*; 116: 113-114.

- Fernandez EJ, Povazay B, Hermann B et al. (2005):** Three-dimensional adaptive optics ultrahigh-resolution optical coherence tomography using a liquid crystal spatial light modulator. *Vision Res*; 45: 3432-3444.
- Ferreras A, Pablo LE, Larrosa JM et al. (2007):** Discriminating between normal and glaucoma-damaged eyes with the Heidelberg retina tomograph 3. *Ophthalmology*; 15 September [Epub ahead of print].
- Ferreras A, Pablo LE, Pajari'n AB et al. (2008):** Logistic Regression Analysis for Early Glaucoma Diagnosis Using Optical Coherence Tomography. *Arch Ophthalmol*;126(4):465-470.
- Ford BA, Artes PH, McCormick TA et al. (2003):** Comparison of data analysis tools for detection of glaucoma with the Heidelberg Retina Tomograph. *Ophthalmology*; 110(6): 1145–1150.
- Fujimoto JG, Pitris C, Boppart SA et al. (2000):** Optical coherence tomography: an emerging technology for biomedical imaging and optical biopsy. *Neoplasia*; 2: 9-25.
- Gabriele ML, Ishikawa H, Wollstein G et al. (2008):** Optical Coherence Tomography Scan Circle Location and Mean Retinal Nerve Fiber Layer Measurement Variability. *Invest Ophthalmol Vis Sci* ; 49(6): 2315–2321.

- Gabriele ML, Wollstein G, Bilonick RA et al. (2007):** Comparison of parameters from Heidelberg retina tomographs 2 and 3. *Ophthalmology*; 24 August [Epub ahead of print].
- Galeana S, Weinreb N, Lam SC et al. (2001):** Longitudinal Variability of Optic Disc and Retinal Nerve Fiber Layer Measurements. (*Invest Ophthalmol Vis Sci*; 49: 4886–4892.
- Garway-Heath DF, Poinoosawmy D, Wollstein G et al. (1999):** Inter- and intraobserver variation in the analysis of optic disc images: comparison of the Heidelberg retina tomograph and computer assisted planimetry. *Br J Ophthalmol*; 83(6): 664–669.
- Giacomo S, Espana EM, Acosta AC et al. (2008):** Agreement between optical coherence tomography and digital stereophotography in vertical cup-to-disc ratio measurement. *Graefes Arch Clin Exp Ophthalmol*; 247: 377–383.
- Girkin CA, McGwin G Jr, Long C et al. (2004):** Subjective and objective optic nerve assessment in African Americans and whites. *Invest Ophthalmol Vis Sci*; 45: 2272–2278.
- Goldberg I (2009):** Stepping Up Glaucoma Management: When and How? *The Open Ophthalmology Journal*; 3: 67-69.
- Gonzalez C, Bueso S, Valle D (2006):** optic nerve drusen and deep visual field defects. *ARCH SOC ESP OFTALMOL*; 81: 269-274.

- Govindarajan B, Junk A, Algeciras M et al. (2009):** Increased isolevuglandin-modified proteins in glaucomatous astrocytes. *Molecular Vision*; 15:1079-1091.
- Greenfield DS, Bagga H and Knighton RW. (2003):** Macular thickness changes in glaucomatous optic neuropathy detected using optical coherence tomography. *Arch Ophthalmol*; 121: 41–6.
- Guedes V, Schuman JS, Hertzmark E et al. (2003):** Optical coherence tomography measurement of macular and nerve fiber layer thickness in normal and glaucomatous human eyes. *Ophthalmology*; 110 (1): 177–189.
- Gupta N (2009):** Playing tag with biomarkers in glaucoma. *Can J Ophthalmol*; 44: 372-377.
- Gupta N and Yucel YH (2007):** Glaucoma as a neurodegenerative disease. *Curr Opin Ophthalmol*; 18: 110-114.
- Gurses-Ozden R, Ishikawa H, Hoh ST et al. (1999):** Increasing sampling density improves reproducibility of optical coherence tomography measurements. *J Glaucoma*; 8(4): 238–241.
- Gyatsho J, Kaushik S, Gupta A et al. (2008):** Retinal Nerve Fiber Layer Thickness in Normal, Ocular Hypertensive, and Glaucomatous Indian Eyes An Optical Coherence Tomography Study. *J Glaucoma*; 17: 122–127.

- Harizman N, Zelefsky JR, Ilitchev E et al. (2006):** Detection of glaucoma using operator-dependent versus operator-independent classification in the Heidelberg retinal tomograph-III. *Br J Ophthalmol*; 90: 1390–1392.
- Harsha BL, Sekhar GC, Ganesh JB et al. (2009):** Clinical measurement and categorization of optic disc in glaucoma patients. *Indian J Ophthalmol*; 57:361-364.
- Harwerth RS, Carter-Dawson L, Shen F et al. (1999):** Ganglion cell losses underlying visual field defects from experimental glaucoma. *Invest Ophthalmol Vis Sci*; 40: 2242-2250.
- Hassenstein A and Meyer CH (2009):** Clinical use and research applications of Heidelberg retinal angiography and spectral domain optical coherence tomography – a review. *Clinical and Experimental Ophthalmology*; 37: 130–143.
- Hatch WV, Flanagan JG, Etchells EE et al. (1997):** Laser scanning tomography of the optic nerve head in ocular hypertension and glaucoma. *Br J Ophthalmol*; 81(10): 871–876.
- He Y, Leung KW, Zhuo YH et al. (2009):** Pro370Leu mutant myocilin impairs mitochondrial functions in human trabecular meshwork cells. *Molecular Vision*; 15:815-825.
- He Y, Leung KW, Zhang YH et al. (2008):** Mitochondrial Complex I Defect Induces ROS Release and Degeneration in Trabecular Meshwork Cells of POAG Patients: Protection by Antioxidants. *Invest Ophthalmol Vis Sci*; 49:1447–1458.

- Henry E, Newby DE, Webb DJ et al (2006):** Altered Endothelin-1 Vasoreactivity in Patients with Untreated Normal-Pressure Glaucoma. *Invest Ophthalmol Vis Sci*; 47:2528–2532.
- Hewick SA, Fairhead AC, Culy JC. Et al. (2004):** A comparison of 10 MHz and 20 MHz ultrasound probes in imaging the eye and orbit. *Br J Ophthalmol*; 88: 551-555.
- Hoffmann EM, Zangwill LM, Crowston JG et al. (2007):** Optic Disk Size and Glaucoma. *Surv Ophthalmol*; 52: 32—49.
- Hong S, Kang SY, Kyoung TM et al. (2008):** Latent Asymmetric Intraocular Pressure as a Predictor of Visual Field Defects. *Arch Ophthalmol*; 126(9):1211-1215.
- Huang D, Swanson EA, Lin CP, et al. (1991):** Optical coherence tomography. *Science*; 254: 1178-1181.
- Huang ML, Chen HY and Hung PT (2005):** Analysis of glaucoma diagnosis with automated classifiers using Stratus optical coherence tomography. *Optical and Quantum Electronics*; 37: 1239–1249.
- Huang P, Zhang SS and Zhang C (2009):** The two sides of cytokine signaling and glaucomatous optic neuropathy. *j ocul biol dis inform*; 2:78–83.
- Hudson CJ, Kim LS, Hancock SA et al. (2007):** Some dissociating factors in the analysis of structural and functional progressive damage in open-angle glaucoma. *Br J Ophthalmol*; 91: 624–628.

Iester M, De Ferrari R and Zanini M. (2001): Topographic analysis to discriminate glaucomatous from normal optic nerve heads with a confocal scanning laser: new optic disk analysis without any observer input. *Surv Ophthalmol*; 44: 433–439.

Ishikawa H, Stein DM, Wollstein G et al. (2005): Macular segmentation with optical coherence tomography. *Invest Ophthalmol Vis Sci*; 46: 2012-2017.

Janknecht P and Funk J. (1994): Optic nerve head analyser and Heidelberg retina tomograph: accuracy and reproducibility of topographic measurements in a model eye and in volunteers. *Br J Ophthalmol*; 78(10): 760–768.

Jansonius NM (2007): Progression detection in glaucoma can be made more efficient by using a variable interval between successive visual field tests. *Graefe's Arch Clin Exp Ophthalmol*; 245:1647–1651.

Johnson C (2002): Structure and function evaluation: criteria for glaucomatous visual field loss using standard automated perimetry and short wavelength automated perimetry. *Am. J. Ophthalmol*; 134: 177-185.4

Johnson DE, El-Defrawy SR, Almeida DR et al. (2009): Comparison of retinal nerve fibre layer measurements from time domain and spectral domain optical coherence tomography systems. *Can J Ophthalmol*; 44: 562–566.

- Jonas JB, Mardin CY and Grundler AE. (1998):** Comparison of measurements of neuroretinal rim area between confocal laser scanning tomography and planimetry of photographs. *Br J Ophthalmol*; 82(4): 362–366.
- Jorgensen TM, Thomadsen J, Christensen U et al. (2007):** Enhancing the signal-to-noise ratio in ophthalmic optical coherence tomography by image registration method and clinical examples. *J Biomed Opt*; 12: 412-428.
- Julio ED, Lisandro MS, Bobby K et al. (2007):** Effect of Glaucomatous Damage on Repeatability of Confocal Scanning Laser Ophthalmoscope, Scanning Laser Polarimetry, and Optical Coherence Tomography. . *Invest. Ophthalmol. Vis. Sci*; 48:1156-1163.
- Jun L and Cynthia JR (2005):** Influence of corneal biomechanical properties on intraocular pressure measurement. *Journal of cataract and refractive surgery*; 31: 146-155.
- Kagemann L, Wollstein G, Wojtkowski M et al. (2007):** Spectral oximetry assessed with high-speed ultra-high-resolution optical coherence tomography. *J Biomed Opt*; 12: 412-422.
- Kalaboukhova L, Fridhammar V and Lindblom B (2006):** Glaucoma follow-up by the Heidelberg retina tomograph--new graphical analysis of optic disc topography changes. *Graefes Arch Clin Exp Ophthalmol*; 244(6):654-662.

- Kamal DS, Viswanathan AC, Garway-Heath DF et al. (1999):** Detection of optic disc change with the Heidelberg retina tomograph before confirmed visual field change in ocular hypertensives converting to early glaucoma. *Br J Ophthalmol*; 83(3): 290–294.
- Kamio M, Meguro A, Ota M et al. (2009):** Investigation of the association between the GLC3A locus and normal tension glaucoma in Japanese patients by microsatellite analysis. *Clinical Ophthalmology*; 3: 183–188.
- Kanamori A, Nagai-Kusuhara A, Escaño FT et al. (2006):** Comparison of confocal scanning laser ophthalmoscopy, scanning laser polarimetry and optical coherence tomography to discriminate ocular hypertension and glaucoma at an early stage. *Graefe's Arch Clin Exp Ophthalmol*; 244: 58–68.
- Kim SH, Kim JY, Kim DM et al. (2006):** Investigations on the association between normal tension glaucoma and single nucleotide polymorphisms of the endothelin-1 and endothelin receptor genes. *Molecular Vision*; 12:1016-21.
- King JW, Bolton N, Aspinall P et al. (2000):** Measurement of Peripapillary Retinal Nerve Fiber Layer Volume in Glaucoma. *Am J Ophthalmol*; 129 :599–607.
- Ko TH, Witkin AJ, Fujimoto JG, et al. (2006):** Ultrahigh-resolution optical coherence tomography of surgically closed macular holes. *Arch Ophthalmol*; 124: 827-836.

- Kourkoutas D, Buys YM, Flanagan JG et al. (2007):** Comparison of glaucoma progression evaluated with Heidelberg retina tomograph II versus optic nerve head stereophotographs. *Can J Ophthalmol*; 42: 82–88.
- Kwon YH, Adix M, Zimmerman MB et al. (2009):** Variance Owing to Observer, Repeat Imaging, and Fundus Camera Type on Cup-to-disc Ratio Estimates by Stereo Planimetry. *J Glaucoma*;18: 305–310.
- Lalezary M, Medeiros FA, Robert N et al. (2006):** Baseline Optical Coherence Tomography Predicts the Development of Glaucomatous Change in Glaucoma Suspects. *Am J Ophthalmol*; 142: 576–582.
- Leung CK, Mohamed S, Leung KS, et al. (2006):** Retinal nerve fiber layer measurements in myopia: an optical coherence tomography study. *Invest Ophthalmol Vis Sci*; 47: 5171-5176.
- Linda MZ, Sonia J, Lyne R et al. (2007):** The Effect of Disc Size and Severity of Disease on the Diagnostic Accuracy of the Heidelberg Retina Tomograph Glaucoma Probability Score. *Invest Ophthalmol Vis Sci*.; 48: 2653–2660.
- Liu B and Brezinski ME (2007):** Theoretical and practical considerations on detection performance of time domain, Fourier domain, and swept source optical coherence tomography. *J Biomed Opt*; 12: 440-447.

- Lopez P, Ferreras A, Polo V et al. (2007):** Relationship between standard automated perimetry and HRT, OCT and GDx in normal, ocular hypertensive and glaucomatous subjects. *Arch. Soc. Esp. Oftalmol*; 82: 197-208.
- Mabuchi F, Sakurada Y, Kashiwagi K et al. (2009):** Lack of association between p53 gene polymorphisms and primary open angle glaucoma in the Japanese population. *Molecular Vision*; 15:1045-1049.
- Markovic V, Kontic D, Hentova-Sencanic P et al. (2009):** Contribution and significance of Heidelberg Retinal Tomography II in diagnostics of ocular hypertension and its conversion into primary open-angle glaucoma. *Vojnosanit Pregl*; 66(4):283-9.
- Manuel MH, David FG, Christian PJ et al. (2005):** Inter-observer variability in confocal optic nerve analysis (HRT). *Int Ophthalmol*; 26:143–149.
- Medeiros FA and Weinreb RN (2009):** Estimating the Risk of Developing Glaucoma. *The Open Ophthalmology Journal*; 3: 50-53.
- Medeiros FA, Zangwill LM, Bowd C et al. (2005):** Use of Progressive Glaucomatous Optic Disk Change as the Reference Standard for Evaluation of Diagnostic Tests in Glaucoma. *Am J Ophthalmol*; 139:1010–1018.

- Medeiros FA, Zangwill LM, Bowd C et al. (2007):** Agreement between stereophotographic and confocal scanning laser ophthalmoscopy measurements of cup/disc ratio: effect on a predictive model for glaucoma development. *J Glaucoma*; 16: 209–214.
- Medved N and Cvenkel B. (2007):** Diagnostic accuracy of the Moorfields regression analysis using the Heidelberg retina tomograph in glaucoma patients with visual field defects. *Eur J Ophthalmol*; 17: 216–222.
- Michael JD and Mohammed LZ (2000):** Human Corneal Thickness and Its Impact on Intraocular Pressure Measures. *Survey of ophthalmology*; 44: 367-408.
- Miglier S, Guareschi M, Romanazzi F et al. (2005):** The Impact of Definition of Primary Open angle Glaucoma on the Cross-sectional Assessment of Diagnostic Validity of Heidelberg Retinal Tomography. *Am J Ophthalmol* 2005; 139: 878–887.
- Miglier S, Guareschi M, Riva I et al. (2007):** Retinal Sensitivity and Retinal Nerve Fiber Layer Thickness Measured by Optical Coherence Tomography in Glaucoma. *Am J Ophthalmol*; 144: 733–740.
- Mikelberg FS, Parfitt CM and Swindale NV. (1996):** Ability of the Heidelberg Retina Tomograph to detect early glaucomatous visual field loss. *J Glaucoma*; 4: 242–247.

- Mohamed S, Lee GK, Rao SK, et al. (2007):** Repeatability and reproducibility of pachymetric mapping with visante anterior segment optical coherence tomography. *Invest Ophthalmol Vis Sci*; 48: 5499-5504.
- Moreno-Montan˜e's J, Anto'n A, Garc'ı'a N, et al. (2009):** Comparison of Retinal Nerve Fiber Layer Thickness Values Using Stratus Optical Coherence Tomography and Heidelberg Retina Tomograph-III. *J Glaucoma*; 18: 528–534.
- Mozaffarieh M, Grieshaber MC and Flammer J (2008):** Oxygen and blood flow: players in the pathogenesis of glaucoma. *Molecular Vision*; 14:224-33.
- Nayak J, Acharya UR, Bhat PS et al. (2009):** Automated Diagnosis of Glaucoma Using Digital Fundus Images. *J Med Syst*; 33: 337–346.
- Nicholas GS and David FG (2008):** New developments in Heidelberg retina tomograph for glaucoma. *Current Opinion in Ophthalmology*; 19:141–148.
- Nicolela MT, Soares AS, Carrillo MM et al. (2006):** Effect of moderate intraocular pressure changes on topographic measurements with confocal scanning laser tomography in patients with glaucoma. *Arch Ophthalmol*; 124: 633–640.
- Niemeijer M, Abràmoff MD and van Ginneken B (2009):** Fast detection of the optic disc and fovea in color fundus photographs. *Medical Image Analysis*; 13: 859–870.

- Nouri-Mahdavi K, Hoffman D, Gaasterland D (2004):** Prediction of Visual Field Progression in Glaucoma. *Invest Ophthalmol Vis Sci*; 45: 4346–4351.
- Nouri-Mahdavi K, Hoffman D, Tannenbaum DP et al. (2004):** Identifying Early Glaucoma with Optical Coherence Tomography. *Am J Ophthalmol*; 137: 228–235.
- Oddone F, Centofani M, Iester M et al. (2009):** Sector-based analysis with the Heidelberg Retinal Tomograph 3 across disc sizes and glaucoma stages: a multicenter study. *Ophthalmology*; 116(6):1106–1113.
- Ohkubo S, Takeda H, Higashide T et al. (2007):** A pilot study to detect glaucoma with confocal scanning laser ophthalmoscopy compared with nonmydriatic stereoscopic photography in a community health screening. *J Glaucoma*; 16: 531–538.
- Owen VM, Strouthidis NG, Garway-Heath DF et al. (2007):** Measurement variability in Heidelberg retina tomograph imaging of neuroretinal rim area. *Invest Ophthalmol Vis Sci*; 47: 5322–5330.
- Park KH and Caprioli J. (2002):** Development of a novel reference plane for the Heidelberg retina tomograph with optical coherence tomography measurements. *J Glaucoma*; 11(5): 385–391.

- Patterson AJ, Garway-Heath DF and Crabb DP. (2006):** Improving the repeatability of topographic height measurements in confocal scanning laser imaging using maximum-likelihood deconvolution. *Invest Ophthalmol Vis Sci*; 47: 4415– 4421.
- Paul JM and George AC (2007):** Measuring structure and function in patients with glaucoma. *Invest. Ophthalmol. Vis. Sci*; 47:1008-1015.
- Paul JM and Frederick SM (2009):** Evaluating Optic Nerve Damage: Pearls and Pitfalls. *The Open Ophthalmology Journal*; 3, 54-58.
- Prashant N, Ramanjit S, Parul S et al. (2007):** Evaluation of Optical Coherence Tomography and Heidelberg Retinal Tomography Parameters in Detecting Early and Moderate Glaucoma. . *Invest. Ophthalmol. Vis. Sci*; 48:3138-3145.
- Pueyo V, Polo V, Larrosa JM et al. (2006):** Diagnostic usefulness of OCT, HRT-II and GDx in operative glaucoma. *Arch. Soc. Esp. Oftalmol*; 81: 693-700
- Quigley HA, Katz J, Derick RJ et al. (1992):** An evaluation of optic disc and nerve fiber layer examinations in monitoring progression of early glaucoma damage. *Ophthalmology*; 99: 19-28.
- Racette L and Sample PA (2003):** Short-wavelength automated perimetry. *Ophthalmol. Clin. North Am*; 16: 227-236.
- Reus NJ, de Graaf M and Lemij HG. (2007):** Accuracy of GDx VCC, HRT I, and clinical assessment of stereoscopic optic nerve head photographs for diagnosing glaucoma. *Br J Ophthalmol*; 91: 313–318.

- Rohrschneider K, Burk RO, Kruse FE et al. (1994):** Reproducibility of the optic nerve head topography with a new laser tomographic scanning device. *Ophthalmology*; 101(6): 1044–1049.
- Roorda A, Metha AB, Lennie P et al. (2001):** Packing arrangement of the three cone classes in primate retina. *Vision Res*; 41: 1291-1306.
- Saglar E, Yucel D, Bozkurt B et al. (2009):** Association of polymorphisms in *APOE*, *p53*, and *p21* with primary open-angle glaucoma in Turkish patients. *Molecular Vision*; 15:1270-1276.
- Sanfilippo PG, Cardini A, Hewitt AW, et al. (2009):** Optic disc morphology - Rethinking shape. *Progress in Retinal and Eye Research*; 28: 227–248.
- Savini G, Espana EM, Acosta AC et al. (2009):** Agreement between optical coherence tomography and digital stereophotography in vertical cup-to-disc ratio measurement. *Graefes Arch Clin Exp Ophthalmol*; 247:377–383.
- Schuman JS (2008):** Spectral Domain Optical Coherence Tomography for Glaucoma (an AOS Thesis). *Trans Am Ophthalmol Soc*; 106: 426-458.
- Schuman JS, Hee MR, Arya AV et al. (1995):** Optical coherence tomography: a new tool for glaucoma diagnosis. *Curr Opin Ophthalmol*; 6: 89-95.

- Schuman JS, Pedut-Kloizman T, Hertzmark E et al. (1996):** Reproducibility of nerve fiber layer thickness measurements using optical coherence tomography. *Ophthalmology*; 103: 1889-1898.
- Scoles D, Gray DC, Hunter JJ et al. (2009):** In-vivo imaging of retinal nerve fiber layer vasculature: imaging – histology comparison. *BMC Ophthalmology*; 9: 9.
- Shaoqing HE, Prasanna G and Yorio T (2007):** Endothelin-1-Mediated Signaling in the Expression of Matrix Metalloproteinases and Tissue Inhibitors of Metalloproteinases in Astrocytes. *Invest Ophthalmol Vis Sc*; 48:3737–3745.
- Sharma P, Sample PA, Zangwill LM et al. (2008):** Diagnostic Tools for Glaucoma Detection and Management. *Surv Ophthalmol*; 53: S17--S32.
- Sherman J, Bass SJ, and Slotnick S (2004):** Glaucoma without cupping. *Optometry*; 75: 677-708.
- Siam GA, Gheith ME, De Barros DS et al. (2008):** Limitations of the Heidelberg Retina Tomograph. *Ophthalmic Surg Laser Imaging*; 39(3):262-4.
- Sigal IA, Flanagan JG, Tertinegg I et al. (2007):** Predicted extension, compression and shearing of optic nerve head tissues. *Exp Eye Res*; 85: 312-322.

- Sihota R, Sony P, Gupta V et al. (2006):** Diagnostic Capability of Optical Coherence Tomography in Evaluating the Degree of Glaucomatous Retinal Nerve Fiber Damage. *Invest Ophthalmol Vis Sci*; 47: 2006–2010.
- Singer AJ, Wang Z, McClain SA et al. (2007):** Optical coherence tomography: a noninvasive method to assess wound re-epithelialization. *Acad Emerg Med*; 14: 387-391.
- Soliman MA, Van Den Berg TJ, Ismaeil AA et al. (2002):** Retinal nerve fiber layer analysis: relationship between optical coherence tomography and red-free photography. *Am J Ophthalmol*; 133(2): 187–195.
- Stein DM, Wollstein G, Ishikawa H et al. (2006):** Longitudinal Variability of Optic Disc and Retinal Nerve Fiber Layer Measurements . *Ophthalmology*; 113: 985-991.
- Srinivasan VJ, Wojtkowski M, Fujimoto JG et al. (2006):** In vivo measurement of retinal physiology with high-speed ultrahighresolution optical coherence tomography. *Opt Lett*; 31: 2308-2310.
- Strouthidis NG, Scott A, Peter NM et al. (2006):** Optic disc and visual field progression in ocular hypertensive subjects: detection rates, specificity, and agreement. *Invest Ophthalmol Vis Sci*; 47: 2904–2910.
- Strouthidis NG, Scott A, Viswanathan AC et al. (2007):** Monitoring glaucomatous visual field progression: the effect of a novel spatial filter. *Invest Ophthalmol Vis Sci*; 48: 251–257.

Strouthidis NG, White ET, Owen VM et al. (2005): Factors affecting the test-retest variability of Heidelberg retina tomograph and Heidelberg retina tomograph II measurements. *Br J Ophthalmol*; 89: 1427–1432.

Sturmer J, Poinoosawmy D, Broadway DC (1992): Intra- and inter-observer variation of optic nerve head measurements in glaucoma suspects using disc-data. *International Ophthalmology*; 16: 227-233.

Swindale NV, Stjepanovic G, Chin A and Mikelberg FS. (2000): Automated analysis of normal and glaucomatous optic nerve head topography images. *Invest Ophthalmol Vis Sci*; 41: 1730–1742.

Tajunisah I, Reddy SC and Fathilah J (2007): Diurnal variation of intraocular pressure in suspected glaucoma patients and their outcome. *Graefes Arch Clin Exp Ophthalmol*; 245: 1851–1857.

Taliantzis S, Papaconstantinou D, Koutsandrea C et al. (2009): Comparative studies of RNFL thickness measured by OCT with global index of visual fields in patients with ocular hypertension and early open angle glaucoma. *Clinical Ophthalmology*;3: 373–379.

Tan JC, Garway-Heath DF, Fitzke FW et al. (2003): Reasons for rim area variability in scanning laser tomography. *Invest Ophthalmol Vis Sci*; 44(3): 1126–1131.

- Tanuj D, Ritu G, Aggarwal A et al. (2009):** Retinal Nerve Fiber Layer Thickness Measurement by Scanning Laser Polarimetry (GDxVCC) at Conventional and Modified Diameter Scans in Normals, Glaucoma Suspects, and Early Glaucoma Patients. *J Glaucoma*; 18: 448–452.
- Tanwar M, Dada T, Sihota R et al (2009):** Mutation spectrum of *CYP1B1* in North Indian congenital glaucoma patients. *Molecular Vision*; 15:1200-1209.
- Tzamalīs A, Kynigopoulos M, Schlote T et al. (2009):** Improved reproducibility of retinal nerve fiber layer thickness measurements with the repeat-scan protocol using the Stratus OCT in normal and glaucomatous eyes. *Graefes Arch Clin Exp Ophthalmol*; 247: 245–252.
- Varma R, Bazzaz S and Lai M. (2003):** Optical tomography-measured retinal nerve fiber layer thickness in normal Latinos. *Invest Ophthalmol Vis Sci*; 44(8): 3369–3373.
- Vessani RM, Moritz R, Batis L, et al. (2009):** Comparison of Quantitative Imaging Devices and Subjective Optic Nerve Head Assessment by General Ophthalmologists to Differentiate Normal From Glaucomatous Eyes. *J Glaucoma*; 18: 253–261.
- Ville S and Juhani PA (2008):** Heidelberg retina tomograph parameters of the optic disc in eyes with progressive retinal nerve fibre layer defects. *Acta Ophthalmol*; 86: 603–608.

- Vilupuru AS, Rangaswamy NV, Frishman LJ et al. (2007):** Adaptive optics scanning laser ophthalmoscopy for in vivo imaging of lamina cribrosa. *J Opt Soc Am A Opt Image Sci Vis*; 24: 1417-1425.
- Vizzeri G, Weinreb RN, Marteniz de la Casa et al (2009):** Clinicians agreement in establishing glaucomatous progression using the Heidelberg retina tomograph. *Ophthalmology*; 116(1):14-24.
- Wang Y, Bower BA, Izatt JA et al. (2007):** In vivo total retinal blood flow measurement by Fourier domain Doppler optical coherence tomography. *J Biomed Opt*; 12: 412-415.
- Weinreb RN, Lusky M, Bartsch DU et al. (1993):** Effect of repetitive imaging on topographic measurements of the optic nerve head. *Arch Ophthalmol*; 111(5): 636–638.
- Wilson MR and Singh K (2009):** Intraocular Pressure: Does it Measure Up? *The Open Ophthalmology Journal*; 3: 32-37.
- Wojtkowski M (2001):** Phase sensitive interferometry in optical coherence tomography. *Proc SPIE*; 4515.
- Wojtkowsky M, Leitgeb R, Kowalczyk A et al (2002):** In vivo human retinal imaging by Fourier domain optical coherence tomography. *J. Biomed Opt*; 7: 457-465.
- Wojtkowski M, Srinivasan V, Fujimoto JG et al. (2005):** Three-dimensional retinal imaging with high-speed ultrahigh-resolution optical coherence tomography. *Ophthalmology*; 112: 1734-1746.

Wollstein G, Garway-Heath DF and Hitchings RA. (1998): Identification of early glaucoma cases with the scanning laser ophthalmoscope. *Ophthalmology*; 105(8): 1557–1563.

Wollstein G, Ishikawa H, Wang J et al. (2005): Comparison of three optical coherence tomography scanning areas for detection of glaucomatous damage. *Am J Ophthalmol*; 139: 39-43.

Wollstein G, Schuman JS, Price LL et al. (2004): Optical Coherence Tomography (OCT) Macular and Peripapillary Retinal Nerve Fiber Layer Measurements and Automated Visual Fields. *Am J Ophthalmol*; 138: 218–225.

Wroblewski D, Francis BA, Chopra V et al. (2009): Glaucoma detection and evaluation through pattern recognition in standard automated perimetry data. *Graefes Arch Clin Exp Ophthalmol*; 247: 1517–1530.

Yang H, Downs JC, Girkin C et al. (2007): 3-D histomorphometry of the normal and early glaucomatous monkey optic nerve head: lamina cribrosa and peripapillary scleral position and thickness. *Invest Ophthalmol Vis Sci*; 48: 4597–4607.

Zalatimo NS (1999): Management of glaucoma suspect patients with congenital optic disc anomalies. *Clinical Eye and Vision Care*; 11: 63-74.

- Zangwill LM, Berry CC and Weinreb RN. (1999):** Optic disc topographic measurements after pupil dilation. *Ophthalmology*; 106(9): 1751–1755.
- Zangwill LM, Jain S, Racette L et al. (2007):** The effect of disc size and severity of disease on the diagnostic accuracy of the Heidelberg retina tomograph Glaucoma Probability Score. *Invest Ophthalmol Vis Sci*; 48: 2653– 2660.
- Zawadzki RJ, Choi SS, Fuller AR (2009):** Cellular resolution volumetric *in vivo* retinal imaging with adaptive optics–optical coherence tomography. *Opt Express* 2; 17(5): 4084–4094.
- Zeitz O, Galambos P, Matthiesen N et al (2008):** Effects of the systemic b-adrenoceptor antagonist nebivolol on ocular hemodynamics in glaucoma patients. *Med Sci Monit*; 14(5): 268-275.
- Zelevsky JR, Harizman N, Mora R et al. (2006):** Assessment of a race-specific normative HRT-III database to differentiate glaucomatous from normal eyes. *J Glaucoma*; 15: 548–551.
- Zvia BE, Gadi W, Richard AB et al. (2007):** Glaucoma Detection with the Heidelberg Retina Tomograph 3. *Ophthalmology*; 114: 466–471.