

6. REFERENCES

- A.A.C.C. (1990):** American Association of Cereal Chemists. Approved methods. 12th Ed. Pup. Univ. of Fam. St., Paul, Minn, USA.
- Abd El-Moniem, G.M. (1994):** Mathematical models for maximum improvement of *in vitro* protein digestibility of high dietary fibre cookies. *Nahrung*, 38(1): 32-37.
- Abd El-Razek, S.A.; El-Sahn, M.A.; Mesallam, A.A. and El-Tabey, S.A.M. (1997):** The chemical composition, the nutritive value and the functional properties of malt sprout and its components (Acrospires, Rootlets and Husks). *J. Sci. Food Agric.*, 75: 50-56.
- Abu El-Enean, H.M. (1998):** Chemical and technological studies on some food industry wastes. M.Sc. Thesis, in *Agric. Food Sci. and Technology*, Al-Azhar Univ., Egypt.
- Al-Jasser, M.S. and Mustafa, A.I. (1996):** Quality of hassawi rice bran. *Annals Agric. Sci.*, Ain Shams Univ., 41(2): 875-880.
- Alladi, S.; Gilbert, R. and Shanmugasundaram, K.R. (1989):** Lipids, lipoproteins and lipolytic activity in plasma with dietary protein changes. *Nutr. Rep. Inter.*, 40: 653.

- Amerine, M.A.; Panghorn, P.M. and Roessler, E.B. (1965):** Principles of sensory evaluation food. Academic Press, New York.
- Anderson, J.W. and Chen, W.J.L. (1986):** Cholesterol-lowering properties of oat products. In oats: Chemistry and Technology, Ed. Webster, F.H., American Association of Cereal Chemists, St Paul MN. PP. 309-333.
- Anon (1997):** El-Ahram Company for drinks statistical. Records for year (1991-1996) unpublished data (Arabic).
- A.O.A.C. (2000):** Association of Official Analytical Chemists, Official Methods of Analysis Washington, D.c. 17th Ed. U.S.A.
- Aoe, S.; Ohta, F. and Ayano, Y. (1989):** Effect of rice bran hemicellulose on the cholesterol metabolism in rats. J. Jap. Soc. Nutr. And Fd. Sci. 42: 55-61.
- Barber, S. and Benedito de Barber, C. (1980):** Rice bran: Chemistry and technology. In rice: Production and Utilization, Ed. By B.S. Luh. Westport, CT: AVI, PP. 790-862.
- Baurdon, I.; Yokoyama, W.; Davis, P.; Hudson, C.; Backus, R.; Richter, D.; Knuckles, B. and Schneeman, B.O. (1999):** Post prandial lipid, glucose, insulin and cholecystokinin responses in men fed barley pasta enriched with β -glucan. Am. J. Clin. Nutr., 69(1): 55-63.

- Bernfeld, P. (1955):** Methods in enzymology. Vol. 1: 149-154. (Acad Press, inc., New York). In S.P. Colowick and N.O. Koplan (Eds.).
- Beter, D.A. and Flucher, R.G. (2005):** The future of barley. Cereal Foods World, 50(5): 271-277.
- Bin, H.W.; Hsing, H.C.; Shene, P.H.; Ming, J.S. (1998):** Effect of different contents of rice bran on lipid metabolism in rats fed with exogenous cholesterol. Fd. Sci. Taiwan, 25(1): 1-13.
- Bjorck, I.; Ostergard, K. and Vainionpaa, G. (1990):** Effects of extrusion cooking on starch and dietary fiber in barley. Food Chem., 34(3): 215-227.
- Bhatty, R.S. (1992):** β -glucan content and viscosities of barley and their roller milled flour and bran products. Cereal Chem. 69(5): 469-471.
- Bhatty, R.S. (1993):** Physicochemical properties of roller-milled barley bran and flour. Cereal Chem., 70(4): 394-402.
- Buko, V.; Lukivskaya, O.; Niktin, V.; Tarasov, Y.; Zavodnik, L.; Borodassky, A.; Goren, S.B.; Janz, B. and Gundermann, K.J. (1996):** Hepatic and pancreatic effects of polyenoylphatidyl choline in rats with alloxan-induced diabetes. Cell Biochem. Funct., 14(2): 131-137.
- Campbell, J.A. (1961):** Methodology of protein evaluation. RAG Nutr. Document R. 101 add. June Meeting New York.

- Carroll, L. (1990):** Functional properties and applications of stabilized rice bran in bakery products. *Food Technol.*, 44(4): 74-76.
- Chaudhary, V. and Kandweber, F.E. (1990):** Barley bran flour evaluated as dietary fiber ingredient in wheat bread. *Cereal Foods World* 35(6): 560-563.
- Chen, F.; Yan, H. and Cai, T. (2003):** Study on making biscuit with rich fibre of rice bran. *Food Sci. and Technol.*, 4: 52-53.
- Chen, H.; Ruben Thaler, G.L. and Schanus, E.G. (1988):** Effect of apple fiber and cellulose on physical properties of wheat flour. *J. Food Sci.*, 53: 304.
- Daneil, E.; Verma, L.; Godber, S. and Wells, J. (1991):** Extraction processing of rice bran. *Amer. Soc. Of Agric. Eng. Paper No. 916028.*
- Dawoud, F.M.; El-Adawy, T.A. and Rabie, T.M. (2003):** Production and evaluation of high-fibre biscuit. *Minufiya J. Agric. Res.*, 28(3): 319-332.
- Doumas, B.T. (1975):** Standard methods of protein determination. *Clin. Chem.*, 7: 175-188.
- Doumas, B.T.; Watson, W.A. and Biggs, H.G. (1971):** Albumin standards and measurement of serum albumin with bromocresol green. *Clin. Chem. Acta*, 31: 87.
- Eastwood, M.A. (1992):** The physiological effects of dietary fiber: An update. *Ann. Rev. Nutr.*, 12: 19-35.

- Einely, M.K. (1978):** Enzymatic colorimetric determination of serum total cholesterol. *Clin. Chem.*, 24: 391.
- El-Sayed, H.M.A. (1998):** Biochemical studies on wheat milling by-products, M.Sc. Thesis, Fac. Of Agric. Moshtohor, Zagazig Univ., Egypt.
- El-Shirbeeny, A.A.; Alam, S.O. and Abd El-Salam, S.M. (2003):** Effect of malt and some barley by-products as hypolipidemic agents. Conference Comi-basal. Crops Technology Department, Food Technology Research Institute, A.R.C. Giza, Egypt, p. 965-982.
- El-Soukkary, F.A.H.M. and El-Sherif, S.A. (1998a):** Physical and sensory evaluation of high fibre biscuits. *Menofiya J. Agric. Res.*, 23(3): 651-667.
- Enochian, R.V.; Saunders, R.M.; Schultz, W.G.; Bealge, F.C. and Crowley, P.R. (1981):** Stabilization of rice bran with extrusion cooking and recovery of edible oil; a preliminary analysis of operations and financial feasibility. Marketing Res. Report 1120, ARS, USAD, Washington, D.C.
- Farinu, G.D.; Longe, O.G.; Feluga, B.L. and Legel, S. (1989):** Chemical composition of different rice processing by-products in Nigeria. *Archives of Animal Nutr.*, 39: 367-374.
- Fathi, A.Taha (1997):** Effect of different levels of fiber on pattern and lipid metabolism in rats. *Egypt. J. Nutr.*, 11(4):

- Fisher, R.A. (1970):** Statistical method for research workers. Edinburgh Ed. 14. Oliver and Boyd, P 140.
- Foda, F.F.A. (2005):** Effect of addition of freeze dried from young green barley leaves on Hypercholesterolemic rats. *Annals of Agric, Sc., Moshtohor*, 43(4): 1649-1663.
- Fossati, P. and Precipe, L. (1982):** The determination of triglyceride using enzymatic methods. *Clin. Chem.*, 28: 2077.
- Garcia-Diez, F.; Garcia-Mediavilla, V.; bayon, J.E.. and Gonzalez-Gallego, J. (1996):** Pectin feeding influences fecal bile acid excretion, hepatic bile acid and cholesterol synthesis and serum cholesterol in rats. *J. Nutr.*, 126(7): 1766-1771.
- Gerhardt, A.L. and Gallo, N.B. (1998):** Full fat rice bran and oat bran similarly reduce hypercholesterolemia in humans. *J. of Nutr.*, 128(5): 865-869.
- Haisman, P. and Muller, B.R. (1977):** Glossary of clinical chemistry term. P. 126. Butterworth, London.
- Hanaa, Sidky, M.A.; Afaf, A. Nassef; Ashgan, E.A. and Mahmoud, A.M. (1996):** Chemical and biological studies on producing high fiber biscuits from Irradiated potato peels. *Egypt. J. of Nutr.*, XI(1): 155-174.
- Hecker, K.D.; Meier, M.L.; Newman, R.K. and Newman, C.W. (1998):** Barley β -glucan is effective as

ahypocholesterolemic ingredient in foods. J. Sci. Food Agric., 77: 179-183.

- Hegsted, D.M.; Mills, R.C.; Elvhjem, G.A. and Hart, E.B. (1941):** Choline in the nutrition of chicks. J. Biol. Chem., 138: 495.
- Hegsted, M. and Kousik, E.S. (1994):** Rice bran and rice bran oil may lower heart disease risk by decreasing cholesterol synthesis in the body. Louisiana Agriculture, 37(2): 16-17.
- Hegsted, M. and Windhauser, M.M. (1993):** Reducing human heart disease risk with rice bran. Louisiana Agriculture, 36: 22-24.
- Hegsted, M.; Windhauser, M.M. and Lester, S.B. (1990):** Stabilized rice bran and oat bran lower cholesterol in humans. FASEB J., 4: A368 (Abs. 590).
- Hemavathy, J. and Prabhakar, J.V. (1987):** Lipid composition of rice (*Oryza sativa* L.) bran. J. AOCS, 64(7): 1016-1019.
- Henery, R.J.; Canon, D.C. and Winkelman, J.W. (1974):** Clinical Chemistry: Principles and techniques, 2nd Ed., New York, Harper and Row; PP. 422-424.
- Hicken, J.W. (1996):** Processing, types and uses of barley malt extracts and syrups. Cereal Foods World, 41(10): 788-790.
- Hudson, C.A.; Chiu, M.M. and Knuckles, B.E. (1992):** Development and characteristics of high fibre muffins

with oat bran, rice bran, or barley fibre fractions. *Cereal Foods World*, 37(5): 373-376, 378.

Hundemer, J.K.; Nabar, S.P.; Shriver, B.J. and Forman, L.P. (1991): Dietary fiber sources lower blood cholesterol in C57BL/6 mice. *J. Nutr.*, 121: 1360.

Ink, S.L. and Hurt, H.D. (1987): Nutritional implications of gums. *Food Technol.*, 41(1): 77-82.

James, C. and Ferguson, K. (1986): The use of rice or barley bran in corn tortilla chips. *Ark. Farm Res.*, 35(1): 8.

James, C.; Courtney, D. and Lorenz, K. (1989): Rice bran-soy blends as protein supplements in cookies. *International J. Food Sci. and Technol.*, 24(5): 495-502.

Johnson, I.T.; Gee, J.M. and Brown, J.C. (1989): Comparison of rice bran, wheat bran and cellulose as sources of dietary fiber in the rat. *Food Sci. Nutr.*, 42F: 153.

Jonnalagadda, S.S.; Thye, F.W. and Robertson, J.I. (1993): Plasma total and lipoprotein cholesterol, liver cholesterol and fecal cholesterol excretion in hamsters fed fiber diets. *J. Nutr.*, 123(8): 1377-1782.

Kahlon, T.S. (2003): Cholesterol lowering with grain fractions. *Getreide Mehl Und Brot.*, 57: 279-291.

Kahlon, T.S. and Chow, F.I. (1997): Hypocholesterolemic effects of oat, rice and barley dietary fibres and fractions. *Cereal Foods World*, 42(2): 86-92.

Kahlon, T.S. and Chow, F.I. (2000): Lipidemic response of hamsters to rice bran, uncooked or processed white

and brown rice and processed corn starch. *Cereal Chem.*, 77(5): 673-678.

- Kahlon, T.S.; Chow, F.I.; Knuckles, B.E. and Chiu, M.M. (1993):** Cholesterol lowering effects in hamsters of β -glucan enriched barley fraction, dehulled whole barley, rice bran, oat bran and their combinations. *Cereal Chem.*, 70(4): 435-440.
- Kahlon, T.S.; Chow, F.I.; Sayre, R.N. and Betschart, A.A. (1992a):** Cholesterol lowering in hamsters fed rice bran at various levels, defatted rice bran and rice bran oil. *J. of Nutr.*, 122(3): 513-519.
- Kahlon, T.S.; Chow, F.I. and Wood, D.F. (1999):** Cholesterol response and foam cell formation in hamsters fed rice bran, oat bran and cellulose, and soy protein diets with or without added vitamin E. *Cereal Chem.*, 76(5): 772-776.
- Kahlon, T.S.; Edwards, R.H. and Chow, F.I. (1998):** Effect of extrusion on hypocholesterolemic properties of rice, oat, corn, and wheat bran diets in hamsters. *Cereal Chem.*, 75(6): 897-903.
- Kahlon, T.S.; Saunders, R.M.; Chow, F.I.; Chiu, M.M. and Betschart, A.A. (1990):** Influence of rice bran, oat bran and wheat bran on cholesterol and triglycerides in hamsters. *Cereal Chem.*, 67(5): 439-443.
- Kahlon, T.S. and Smith, G.E. (2004):** Rice bran: a health-promoting ingredient. *Cereal Foods World*, 49(4): 188-194.

- Kahlon, T.S. and Woodruff, C.L. (2003):** *In vitro* binding of bile acids by rice bran, oat bran, barley and β -glucan enriched barley. *Cereal Chemistry*, 80: 260-263.
- Kalra, S. and Jood, S. (2000):** Effect of dietary barley β -glucan on cholesterol and lipoprotein fraction in rats. *J. Cereal Sci.*, 31(2): 141-145.
- Kennedy, L.L.; Chung, H.Y. and Hegsted, M. (1996):** New food products developed from rice bran fractions. Conference Proceedings. Dept. of Food Sci., Louisiana State Univ., Baton Rouge, LA 70803, USA.
- Kestin, M.; Moss, R.; Clifton, P.M. and Nestel, P.J. (1990):** Comparative effects of three cereal brans on plasma lipids, blood pressure and glucose metabolism in mildly hypercholesterolemic men. *Am. J. Clin. Nutr.*, 52: 661.
- Kim, Y.S.; Ha, T.Y.; Lee, S.H. and Lee, H.Y. (1997a):** Effect of rice bran dietary fiber on flour rheology and quality of wet noodles. *Korean J. Food Sci. and Technol.*, 29(1): 90-95.
- Kim, Y.S.; Ha, T.Y.; Lee, S.H. and Lee, H.Y. (1997b):** Effect of rice bran dietary fiber on gelatinization and retrogradation of wheat flour. *Korean J. Food Sci. and Technol.*, 29(3): 464-469.
- Kim, Y.S.; Ha, T.Y.; Lee, S.H. and Lee, H.Y. (1997c):** Properties of dietary fiber extract from rice bran and application in bread-making. *Korean J. Food Sci. and Technol.*, 29(3): 502-510.

- Kiriyama, S.; Okazak, Y. and Yoshida, A. (1968):** Hypocholesterolemic effect of polysaccharides and polysaccharide rich food stuffs in cholesterol fed rats. *J. Nutr.*, 97: 382.
- Kosugi, Y.; Kuneda, T. and Azuma, N. (1994):** Continual conversion of free fatty acid in rice bran oil to triacylglycerol by immobilized lipase. *J. AOCS*, 71(4): 445-448.
- Lane-Peter, W. and Pearson, A.E. (1971):** Dietary requirement. In "The laboratory animal principles and practice" P. 142, Academic Press, London and New York.
- Ledesma, L.; Fuertes, S.; Barrios, A.; Zumbado, H.; Hernandez, G. and Castillo, D. (1990):** Nutritive value of protein concentrate from rice bran. *Ciencia-Y-Tecnica-en-la-Agricultura Arroz*, 13(1-2): 7-14. (c.f. Rice Abstracts, N528666; 1992).
- Lima, I.; Guraya, H. and Champagne, E. (2002):** The functional effectiveness of reprocessed rice bran as an ingredient in bakery products. *Nahrung*, 46(2): 112-117.
- Lopez-virella, M.F.; Stone, S.; Ellis, S. and Collwel, J.A. (1977):** Cholesterol determination in high density lipoproteins separated by three deifferent methods. *Clin. Chem.*, 23(5): 882.
- Macrae, R.; Robinson, R.K. and Sadler, M.J. (1994):** Dietetic foods. In *Encyclopedia of Food Sciences and food*

Technology and Nutrition, Vol. 2. New York: Academic press.

- Maillard, M.N.; Soum, M.H.; Boivin, P. and Berset, C. (1996):** Antioxidative activity of barley and malt: relationship with phenolic content. *Lebensm. Wiss-Technol.*, 29: 238-244.
- Mansour, M.A. (1991):** Waste product treatment from beer industry in Giza governorate. M.Sc. Thesis, Department of Biology and Natural Sciences. Institute of Environmental Studies and Research Ain Shams University.
- Martin, D.J.; Verma, L.R.; Godber, S.J. and Wells, J.H. (1993):** Extrusion processing of rice bran. *Amer. Soc. Agric. Engineers*, Paper No. 916028.
- McDougall, G.J.; Morrison, I.M.; Stewart, D. and Hillman, J.R. (1996):** Plant cell wall as dietary fiber: Range, structure, processing and function. *J. Sci. Food Agric.*, 70: 133-150.
- Miller, G.L. (1959):** Use of dinitrosalicylic acid reagent for determination of reducing sugars. *Anal. Chem.*, 31: 426-428.
- Mongeau, R.; Brassard, R.; Malcolm, S. and Shah, B.G. (1991):** Effect of dietary cereal brans on body weight and blood lipids in a long-term rat experiment. *Cereal Chem.*, 68(5): 448-453.

- Newman, R.K.; Lewis, S.E.; Newman, C.W.; Boik, R.J. and Ramage, R.T. (1989):** Hypocholesterolemic effect of barley foods on healthy men. *Nutr. Rep. Int.* 39: 749.
- Newman, R.K. and Newman, C.W. (1991):** Barley as a food grain. *Cereal Foods World*, 36: 801-805.
- Newman, R.K.; Newman, C.W.; El-Negoumy, A.M. and Aastrup, S. (1984):** Nutritional quality of proanthocyanidin free barley. *Nutrition Reports International*, 30(4): 809. Montana State Univ., Bozeman, Montana 59717, USA.
- Nicolosi, R.J. Ausman, L.M. and Hegsted, D.M. (1991):** Rice bran oil lowers serum total and low density lipoprotein cholesterol and apo-B levels in nonhuman primates. *Atherosclerosis*, 88: 133-142.
- Oda, T.; Aoe, S.; Sanada, H. and Ayano, Y. (1993):** Effects of soluble and insoluble fiber preparations isolated from oat, barley and wheat on liver cholesterol accumulation in cholesterol-fed rats. *J. Nutr. Sci. Vitaminol-Tokyo*, 39(1): 73-79.
- Orthoefer, F.T. (1996):** Rice bran oil: Healthy lipid source. *Food Technol.*, Dec., 62-64.
- Peterson, D.M. (1994):** Barley Tocols: effects of milling, malting and mashing. *Cereal Chem.*, 71(1): 42-44.
- Peterson, D.M. and Qureshi, A.A. (1993):** Genotype and environment effects on totals of barley and oats. *Cereal Chem.*, 70: 157.

- Pomeranz, Y.; Ke, H. and Ward, A.B. (1971):** Composition and utilization of milled barley products. 1. Gross composition of roller-milled and air-separated fractions. *Cereal Chem.*, 48: 47-57.
- Prabhakar, J.V. and Venkatesh, V.L. (1986):** A simple chemical method for stabilization of rice bran. *J. Amer. Oil. Chem. Soc.*, 63(5): 644-646.
- Prakash, J. (1996):** Rice bran proteins: properties and food uses. *Critical Rev. in Food Sci. and Nutr.*, 36(6): 537-552.
- Prakash, J. and Ramantham, G. (1995):** Proximate composition and protein quality of stabilized rice bran. *J. Food Sci and Technol., India*, 32(5): 416-419.
- Prentice, N. and D'apponia, B.L. (1977):** High-fiber bread containing brewer's "spent grain". *Cereal Chem.*, 54(5): 1084-1095.
- Prosky, L.; Asp, N.G.; Furda, I.; Devries, J.W.; Schweizer, T.F. and Harland, B.F. (1984):** Determination of total dietary fibre in foods, laboratory study. *J.A.O.A.C.*, 67: 1044.
- Raghuram, T.C.; Rao, U.B. and Rukmini, C. (1989):** Studies on hypolipidemic effects of dietary rice bran oil in human subjects. *Nutr. Rep. Inter.*, 39(5): 889-895.
- Retiman, S. and Frankel, S. (1957):** A colorimetric method for the determination of serum glutamic oxaloacetic and glutamic pyruvic transaminases. *Am. J. Clin. Path.*, 28: 56.

- Riser, S. (1984):** Metabolic aspects of nonstarch polysaccharides. *Food Technol.*, 38(1): 107-113.
- Riser, S. (1987):** Metabolic effects of dietary pectins related to human health.. *Food Technol.*, 41(2): 91-99.
- Rukmini, C. (1988):** Chemical, nutritional and toxicological studies of rice bran oil. *Food Chem.*, 30(4): 257-268.
- Salama, A.A.; El-Sahn, M.A.; Mesallam, A.S. and Shehata, A.M.E. (1997):** Evaluation of the quality of bread, biscuit and butcher's sausage supplemented with rootlets of malt sprouts. *Nahrung*, 41(4): 228-231.
- Sangronis, E.; Cafiero, J. and Mosqued, M. (1997):** Cooking quality of pastas supplemented with rice bran. *Arch Latinoam Nutr.*, 47(2): 146-151.
- Sangronis, E. and Sancio, M. (1990):** Development and characterization of rice bran cookies. *Acta Venez*, 41(3): 199-202.
- Saunders, R.M. (1986):** Rice bran composition and potential foods uses. *Food Reviews International*, 1(3): 465-483.
- Saunders, R.M. (1990):** The properties of rice bran as a food stuff. *Cereal Fd. World*, 35: 632-636.
- Sayre, R.N.; Earl, L.; Kratzer, F.H. and Saunders, R.M. (1988):** Effect of diets containing raw and extrusion-cooked rice bran on growth and efficiency of food utilization of broilers. *Brit. Poultry Sci.*, 29(4): 815-823.

- Scavariello, S.E. and Daniel, B.A. (1998):** γ -oryzanol: an important component in rice bran oil. *Arch Latinoam Nutr.*, 48(1): 7-12.
- Seetharamaiah, G. and Chandrasekhara, N. (1989):** Studies on hypocholesterolemic activity of rice bran oil. *Atherosclerosis*, 78: 219-223.
- Seetharamaiah, G. and Prabhakar, J.V. (1986):** Oryzanol content of Indian rice bran oil and its extraction from soap stock. *J. Food Sci. Technol.*, 23: 270.
- Schnell, M.; Pacheco-de-Delahaye, E. and Mezones, Y. (2005):** Metabolic responses to Venezuelan corn meal and rice bran supplemented arepas (breds). *Cereal Chemistry*, 82(1): 77-80.
- Sharma, H.R. and Chauhan, G.S. (2002):** Effects of stabilized rice bran fenugreek blends on the quality of breads and cookies. *J. of Food Sci. and Technol.*, 39(3): 225-233.
- Sharma, R.D. and Rukmini, C. (1987):** Hypocholesterolemic activity of unsaponifiable matter of rice bran oil. *Indian J. Med. Res.*, 85: 278.
- Sharp, C.Q. and Kitchens, K.J. (1990):** Using rice bran in yeast bread in a home baker. *Cereal Foods World*, 35(10): 1021-1024.
- Shin, T.S.; Godber, S.; Martin, D. and Wells, J. (1997):** Hydrolytic stability and changes in E vitamers and oryzanol of extruded rice bran during storage. *J. Food Sci.*, 62(4): 704-708.

- Skurray, G.R.; Wooldridge, D.A. and Nguyen, M. (1986):** Rice bran as a source of dietary fiber in bread. *J. Food Technol.*, 21: 727.
- Soliman, A.E. (1997):** Raising the nutritive value of some cereals and its baking products. M.Sc. Thesis, Fac. Of Agric. Moshtohor, Egypt.
- Stauffer, C.E. (1993):** Dietary fiber: analysis, physiology and calorie reduction. *Advances in baking technology 1st* Black Academic. Professional London. Glasgow. New York. Tokyo. Melbourne. Madras.
- Steinberg, D. (1981):** Metabolism of lipoprotein at the cellular level in relation to atherogenesis. In *lipoproteins, atherosclerosis and coronary heart disease*. 1, 2, 31- Elsevier-North Holland.
- Sugano, M. and Tsuji, E. (1997):** Rice bran oil and cholesterol metabolism. *J. Nutr.*, 127: 521S-524S.
- Swanson, R.B. and Penfield, M.P. (1988):** Barley flour level and salt level selection for a whole-grain bread formula. *J. of Food Sci.*, 53(3): 896-901.
- Tabacco, A.; Searcy, R.L. and Reardon, J.E. (1979):** Standard Methods of Clinical Chemistry, Edited Seligson, D., *Clin. Chem.*, 25: 336.
- Tae, Y.H.; Sung, H.L. and Hyun, Y.L. (1997):** The effect of various rice bran extracts on the lipid contents of serum and liver in rat fed with high fat diet. *Korean J. Food Sci. and Technol.*, 29(1): 178-182.

- Tashiro, M. and Maki, Z. (1990):** Effects of dietary rice bran trypsin inhibitor on the growth and pancreatic weights of rats. *Nutr. Res.*, 10(11): 1325-1330. (c.f. *Rice Abst.*, 061-03785; 7V).
- Tietz, N.W. (1986):** "Textbook of Clinical Chemistry". P. 796. Saunders, W.B. Co., London-Philadelphia.
- Topping, D.L.; Illman, R.J.; Roach, P.D.; Trimble, R.P.; Kambouris, A. and Nestel, P.J. (1990):** Modulation of the hypolipidemic effect of fish oils by dietary fiber in rats: studies with rice and wheat bran. *J. Nutr.*, 120: 325-330.
- Topping, D.L.; Oakenfull, D.; Trimble, R.P. and Illman, R.J. (1988):** A viscous fiber (methyl cellulose) lowers blood glucose and plasma triacylglycerols and increases liver glycogen independently of volatile fatty acid production in the rat. *Br. J. Nutr.*, 59: 21-30.
- Tsai, C.E.; Ting, H.; Wang, L.J. and Lin, T. (1992):** Effect of rice bran on blood lipids in man. *J. Chinese Agric. Chem. Soc.*, 30(4): 484-495.
- Urooj, A.; Vinutha, S.R.; Shashikala, P.; Leelavathy, K. and Haridasa, R.P. (1998):** Effect of barley incorporation in bread on its quality and glycemic responses in diabetics. *International J. of Food Sci. and Nutr.*, 49: 265-270.
- Vahouny, G.V. and Kritchevsky, D. (1986):** Dietary fiber: Basic and clinical aspects, plenum. Press, New York.

- Wade, P. (1988):** Biscuits, cookies and crackers. Recipe of biscuit used during investigation. Vol. 1. Applied Science. Publisher, Ltd, London, UK.
- Waller, W.M. and Duncan, D.B. (1969):** A boys role for symmetric multiple composition problem. Am. State Assoc. J. 65: 1485.
- Wilson, T.A.; Idreis, H.M.; Taylor, C.M. and Nicolosi, R.J. (2002):** Whole fat rice bran reduces the development of early aortic atherosclerosis in hypercholesterolemic hamsters compared with wheat bran. Nutr. Res., 22(11): 1319-1332.
- Wilson, T.A.; Nicolosi, R.J.; Delaney, B.; Chadwell, K.; Moolchandani, V.; Kotyla, T.; Ponduru, S.; Zeng, G. Hess, R.; Knutson, N.; Curry, L.; Kolberg, L.; Goulson, M. and Ostergren, K. (2004):** Reduced and high molecular weight barley β -glucans decrease plasma total and non-HDL-cholesterol in hypercholesterolemic Syrian golden. J. Nutr., Vol. 134: 2617-2622.
- Windhauser, M.M. and Hegsted, M. (1993):** Examining the effects of rice bran components on lipids. Louisiana Agric., 36(3): 19.
- Yoshino, G.; Kazumi, T.; Amano, M.; Tateiwa, M.; Yamasaki, T.; Takashima, S.; Iwai, M.; Hatanaka, H. and Baba, S. (1989):** Effects of γ -oryzanol and probucol on hyperlipidemia. Current Ther. Res., 45: 975.

- Young, K.B.; Cramer, G.L. and Walles, E.J. (1991):**
Prospects for increasing returns from rice bran and other rice milling by-products. Arkansas Agricultural Experiment Station Fayetteville, Arkansas 72701 pp: 11-12.
- Zumbado, H.; Ledesma, L.; Fuertes, F. and Ventura, J. (1997):** Manufacture of a bakery product with incorporation of high levels of precooked rice bran. *Alimentaria*, 280: 21-23.