

References

Bastos I, Gomes D, Gouveia H, and Freitas D. (1998). Colonic metastasis of a lung carcinoma with gastro-colic fistula. J Clin Gastroenterol . 26:348.

Baxter AD, Fishwick NG, and Jeyapalan K. (1998). Case report; ileobronchial fistula. Clin Radiol. 53:231-233

Berna JD , Sanchez J, and Albarracin A. (2001). Electrical nerve stimulation in the management of the enterocutaneous low-output fistulae : a report of two cases .Am J Gastroenterol . Mar;96(3):900-1

Berry SM and Fischer JE .(1996) . Classification and pathophysiology of enterocutaneous fistulae . Surg Clin North Am.76:1009-18 .

Binnekade JM, Tepaske R, and Bruynzeel P. (2005). Daily enteral feeding practice in the ICU; attainment of goals and interfering factors. Critical Care. 9: 218-225.

Bloom A.(2002) . Abscess and enterocutaneous fistula. Radiology Ucsf Edu J .68; 241-245.

Branagau G & Asenapati Ash. (2008). Colonic (diverticulosis) surgery 142: 357:360.

Buffington C . (1998). Multi-center study: food intake of hospitalized patients. Clin nutri. 17; 14-16.

Burnand K and Young E. (2004) . Small Bowel Fistulae management. Gut . 49 : 811-816.

Carol E.H. (2002). Small bowel resection and anastomosis . Chassin's Operative Surgery in General Surgery – An expositive atlas, 3^{ed} edition . Springer. (37):331-337 .

Castel H, Tienqou LE, and Besancon I. (2005): What is the risk of nocturnal supine enteral nutrition? Clin Nutr. 24: 1014-1018.

Chamberlain RS , Kaufman HL and Danforth DN.(1998). Enterocutaneous fistula in cancer patients; etiology, management, outcome and impact on further treatment . Am Surg J .Dec; 64(12): 1204-

Chander J,Lal P, and Ramtek Vk.(2004) . Rectus abdominis ms flap for high-output fistula ; novel technique.World J Surg.28:179-182.

Chang P, Chun J, and Bell J.(2000) . Complex enterocutaneous fistula: Closure with rectus abdominus flap (electronic version). Southern Medical Journal. 93 (6): 599-604.

Chao G, Jian A, Wang X, and Shou J .(2004).Refeeding syndrome . nutrition . 20 ;346-350

Cherr GS, Hall C, Pineau BC, and Waters GS. (2001). Rectourethral fistula and massive rectal bleeding from iodine-125 prostate brachytherapy: a case report. Am Surg. 67:131-134.

Chintamani ,Rohini Badran, Daniel RK , Vinay Singhl and Dearlove JL.(2003) . Post irradiation intestinal fistula . Surg Clin north Am.83:1095-110 .

Chintapatla and Scott N. (2003).intestinal fistula ; common abd condition . Surg .137;46-50.

Chitale VR. (1993). enterocutaneous fistula treated successfully by a turnover flap from surrounding tissue reinforced by musclocutaneous flaps and other local flaps. Plast Reconster Surg. 96;1171-1172.

Clarence E , Daniel ,and Lefor T .(1996).General management of GI fistula . Surg clin north AM . 76; 1019-1033.

Class R.E. (1999). The management of fistulae in Crohn's disease.Br J Surg. 86;72-93 .

Clouse R.E . (1998). Nutritional therapy. Duranagan W.C. and Rindner M,L (Eds.) Manual of Medial therapeutics (16th Ed). Little, Brown and Company, Boston chap. 2pp. 35-51.

Cohen RD, Larson LR ,andRoth JM .(2000).The cost of hospitalization in crohn's Disease .Am J Gastroentrol . 95:524-530.

Corson D and Williamson R (2001) Abnormal compartmentalization , Extracellular water deficit .west . Afr . J surg . 20 ;98-109.

Cro C , George K J , Donnelly J , Irwin S T, and Gardiner.(2002). Vacuum assisted closure system in the management of enterocutaneous fistulae (case report). Postgraduate Medical Journal.78: 364-365.

David J, Karen J, and Kristine S. (2002) . Porcine small intestinal submucosa as a treatment for enterocutaneous fistulae. Journal of the American College of Surgeons. April; 194(4):32-37

David E , Radha V, and Asya S .(2008). Intestinal fistula. Emedicine gastro-ent .6;1-15

Dearlove J L.(1996). Skin care management of gastrointestinal fistulas. Surg Clin North Am.;Vol.76,N(5):1095-1109.

Debatin JF, Luboldt W, and Bauerfeind P. (1999). virtual colonoscopy, Computed Tomography or Magnetic Resonance Imaging?. Endoscopy. 31(2):174-179 .

Disa JJ ,Goldberg NH,and Carlton JM .(1998). Restoring abdominal wall integrity in contaminated tissue-deficient wound using antogous fascia grafts .Plast Reconstr Surg. 101:979-986.

Dioniqi G, dioniqi R ,and rovera F. (2008) . Treatment of high output enterocutaneous fistula (international J of surgery .16:51-65 .

Douglas Wilmore.(2003). Indication for specific therapy in rehabilitation of patient with the short bowel syndrome .Clin Gastroenterology (17) (6) 895:906

Dudrich SJ, Maharaj AR, and McKelvey AA. (1999). Artificial nutritional support in patients with gastrointestinal Fistulas. World Journal of Surgery .Jun; 23 (6): 201-210.

Dykes CM. (2000). Virtual colonoscopy; a new approach for colorectal cancer screening. Gastroenterol Nurs; 24 (1): 5-11,.

Eisenberg. J. (1993). Parenteral Enteral Nutrition: Calculation of metabolic expenditure and substrate utilization from gas exchange measurements. Nutrition .17: 201-210.

El-Bahar T.(1988). Intestinal fistula . Recent advances in Surgery,Vol. 13, Ch. 7 :103-124.

Evenson AR , Fischer JE. (2006). Current management of entero cutaneous fistula .J Gastro Intest Surg . Mar ; 10; 455-64.

(F)

Fadiran OA, Dare FO, Jeje EA, Nwosu SO, Oyero TO. (1993). Amoebic recto-vaginal fistula: a case report and review of the literature. Cent Afr J Med.39:172-175.

Falconi M and Pederzoli P.(2001) . Gastrointestinal fistulae, the relevance of gastrointestinal fistulae in clinical practice: a review. Gut;48:2-10 .

Familiari ,P,Marci A, and Consolo P .(2003). Endoscopic clipping of colocutaneous fistula following necrotizing pancreatitis case report Gastro enterologica Italiana .25 907-910.

Fazio VW,Coutsoftides T,and Steiger E.(1983). factors influencing the outcome of treatment of small bowel cutaneous fistula .World J Surg. 7:481-488.

Fenlon HM, Clarke PD, and Ferrucci JT.(1998). Virtual colonoscopy: Imaging features with colonoscopic correlation. AJR. 170(5): 1303-1309,.

Ferrucci JT. (2001). Colon cancer screening with virtual colonoscopy;Promise, Polyps, politics. AJR; 177(5): 975-988,

Fischer J E. (2001) . Gastrointestinal-Cutaneous Fistula . Mastery of surgery 4th edition . Library of cataloging-in-publication data .Small intestine .3(123) : 1435-1441.

Fischer JE. (2001) . Metabolism in surgical patients: Protein , Carbohydrate and Fat utilization by oral and parenteral routes , Sabiston textbook of surgery (*The biological basis of modern practice*) 16th edition . W.B. Saunders Company.(6):90-130 .

Fletcher JG, Johnson CD, MacCarty RL, Welch TJ, Hara AK, and Reed JE. (1998) . CT colonography overcoming the problems of collapse and colonic fluid. Radiology.16; 209-296.

Foster CE and Lefor AT. (2001).General management of gastrointestinal fistulas. Surg Clin North Am 81;1019-1034.

Fry DE and Osler T.(1996).Abdominal wall considerations and complications in preoperative surgery.Surg Clin North Am. 76:1-11.

Fustin P,Steinhart, and Cohen.(2003). Home total parental nutrition :an alternative to early surgery for complicated inflammatory bowel disease Gastrointestinal Surg. 138:562-566.

Garcia-de-Lorenzo A, Zara Zaga A, and Garcia W. (2003). Clinical evidence for enteral nutritional support with Glutamine; a systematic review. Nutrition. 19: 809-811.

Garcia GD, Freeman IH , Zagorski SM and Chung MH.(2004) . A laparoscopic approach to the surgical management of enterocutaneous fistula in a wound healing by secondary intention. Surg Endosc .Mar;18(3);554-6 .

Ghahremani GG. (1993). Radiologic evaluation of suspected gastrointestinal perforations. Radiol Clin North Am. 31:1219-1234.

Gillen P , Peel A, and Reeder A .(1985). Value of HIDA scanning in intestinal fistula . Annals of Royal College Of Surgery. 67;330-331.

Glass RE,Richic JK,Lennard-jones JE.(1999).internal fistula in CD .Dis.colon Rectum. 28(8):58.

Gonzalez I Pinto and Moreno E Gonzalez.(2002). Optimizing the treatment of upper Gastrointestinal fistulae. Gut. 49; 21- 28.

Goldman SM, Fishman EK, Gatewood OMB, Jones B and Siegelman SS. (1995). CT in the diagnosis of enterovesical fistulae. AJR Am J Roentgenol. 139:1229-1233.

Gollub MJ, Ginsberg MS, Cooper C and Thaler HT. (2002). Quality of virtual colonoscopy in patients who have undergone radiation therapy or surgery how successful we are?. AJR. 178 (5): 1109-1116,

Gollub MJ and Bains MS. (1997).Barium sulfate; a new (old) contrast agent for diagnosis of G.I.T leaks. Radiology; 202:360-362.

Grindrod K.A. Patel P, and Martin J.E. (2006). What interventions should pharmacists Employ to impact Health practitioners' Prescribing Practices? Ann. Pharmacother. September ; 40 (9); 1546 – 1557.

Guillem P , Binto D, and Dupuy J . (2001). Duodeno caval fistula . j vasc surg . 33 ; 643-645.

(H)

Hardcastle MR.(1998). Vacuum assisted closure therapy ; the application of negative pressure in the wound healing .Canad .plas. J .15;102-109 .

Ham M , Horton K , and Kaunitz J. (2007). Fistuloclysis . Nutr Clin Pract . 22; 553-7.

Hill L. (2002). choosing an enteral feed for the critically ill .S Afr Crit Care . 18:17-22.

Hollington P, Mawdsley J, Lim W , Gabe SM , Forbes A, and Windsor AJ. (2004) . An 11-years experience of enterocutaneous fistula. British. Journal of Surgery. Dec; 91 (12): 1646-51.

Hwang TL and Chen MF. (1999) .Short note randomized trial of fibrin tissue glue for low output enterocutaneous fistula. Br J Surg .86:112-113.

Iglesias S.B.d.O., Leite H.P., Meneses J.F.S.e, and De Carvalho W.B. (2007). Enteral nutrition in critically ill children; are prescription and delivery. Nutr Clin Pract. April 1, 22 (2): 233-239.

Inal M, Oguz M, Aksungur E, Soyupak S, Boruban S, and Akgul E.(1999). Biliary-enteric fistulas: report of five cases and review of the literature. Eur Radiol. 9 ; 1145-1151.

Jarrett TW and Vaughan ED. (1999). Accuracy of computerized tomography in the diagnosis of colovesical fistula secondary to diverticular disease. J Urol. 153:44-46.

Jacobs M, Verdiqa JC, and Golstein HS.(1991). minimally invasive colon resection . Surg Laparoscope . 1;144-150.

John M ,Drau Jr , Sara A and Niall J.(2006).Enterocutaneous fistula (are treatment improving) surgery (140) (4); 570-78.

Johnson CD and Dachman AH. (2000).CT Colonography; The next colon screening examination. Radiology. 216: 331-341.

Julie A. Margenthaler , Virginia M. Herrmann, and Michael S. Hickey : (2003). Surgical Metabolism and Nutrition , Current Surgical Diagnosis and Treatment 11th edition, Lange Medical Books/Mc Graw-Hill. 1(11) :156-190.

Kapadia BJ, Agarwal M, and Silva RD.(2000). Primary aortoduodenal fistulas in minimally aneurysmal aortas: imaging diagnosis. Abdom Imaging. 25:51-54.

Kaur N, Minocha VR, and Mundu M .(2004). Improving the outcome in patients of high output small bowel fistula. Trop Gastroenterology . Apr-Jun;25(2):92-5 .

Kay CL, Kulling D, Hawes RH, Young JR, and Cotton PB. (2000) . Virtual Endoscopy- Comparison with colonoscopy in the detection of space-occupying lesions of the colon. Endoscopy. 32(3): 226-232.

Kearny R ,Payne W, and Rosemurgy A.(1997). External abdominal closure of enterocutaneous fistula .Am Surg. 63:406-409.
Gastroenterology . Apr-Jun;25(2):92-5 .

Kess P, Mawdsley J, Agarwal M, and Silva RD, and Windsor AJ. (2000)
Vesico-colic fistula. British. Journal of Surgery. Dec; 88 (12): 196-51.

Kirk R.M .(2006).General surgical operations.5th edition .Churchill livingstone.Stomach and duodenum. M.C Winslet.(13): 173-174.

Kuvishinoff BW, Brodish RJ, and McFadden DW.(1993). Serum transferrin as a prognostic indicator of spontaneous closure and mortality in gastrointestinal cutaneous fistulas. Ann Surg. 217:615-623.

Larsen A, Johansen TEB, Solheim BM and Urnes T. (1999). Diagnosis and treatment of enterovesical fistula. Eur Urol. 31:318-321.

Lawrence W. Way and Doherty M. (2003). Small intestinal fistula ,Current Surgical Diagnosis& Treatment, 11th edition , Lange Medical Books/Mc Graw-Hill . 1(30):696-700.

Lee JT, Sarovan RM, Belzberg G, Pianim NA, and Bunqard FS. (2001).

Primary aortoenteric fistula: computed tomographic diagnosis of an atypical presentation. Ann Vasc Surg. 15; 251-254.

Lee YC , Na HG , Suh JH , Park IS , Chung KY, and Kim NK .(2001).

Three cases of fistulae arising from gastrointestinal tract treated with endoscopic injection of Histoacryl . Endoscopy. Feb;33(2); 184-6 .

Leone S, Barabino M, Marsetti M, Rampinelli L, Arnoldi D, Opocher E.(2003). Gastrojejunocolic fistula as rare complication of recurrent peptic ulcer. Chir Ital. Nov-Dec;55(6):923-8

Levin L, Brooks D, Smith RA, and Stone A.(2003). Emerging technologies in screening for colorectal cancer: CT colonography, immunochemical fecal occult blood tests, and stool screening using molecular markers. Cancer J Clin. 53:44-55.

Levine MS, Kelly MR, Laufer I, Rubesin SE, and Herlinger H. (1999).

Gastrocolic fistulas : the increasing role of aspirin. Radiology. 192; 359-361.

Lloyd DA , Gabsem SM and Windsor AC .(2006).

Nutritional management of entero cutaneous fistula. Br j surg. 93 ; 1045-55.

Lo Casto A, Salemo S, Grisanti M, and Mastrandrea G. (1997).

Hydatid cyst of the liver communicating with the left colon. Br J Radiol. 70; 650-651.

Loby I, Gomes D, Smith RA, and Freitas D. (2002).

Malignant gastrocolic fistula. Gastroenterology . Apr-Jun;25(2):92-5 .

Lui RC, Friedman R, and Fleischer A. (1989). Management of post-irradiation recurrent enterocutaneous fistula by muscle flaps .Am Surg. 55:403-407.

Maconi G, Parente F, and Bollani S. (1999) . Abdominal ultrasound in the assessment of extent and activity of Crohn's disease ; clinical significance and implication of bowel wall thickening . Am J Gastroenterology . 91:1604-1609 .

Malanqoni MA, shumate CR, and Thomas HA . (1990) . factors influencing the treatment of intrabdominal abscess. Am J Surg. 159:167-171.

Martinez Ordaz JL, Luque de leon E, and Suarez Moreno RM . (2003). Post-operative enterocutaneous fistula .Gaceca Medica De Mexico. Mar-Apr ; 139 (2) ; 1440-51.

Matie, Dordea , Arcotc ,Sue E, Jas S and Vavna .(2008).(delayed rectal anastamotic dehiscence presenting as colocutaneous fistula in popliteal fistula. Can J Surgery .51 (3);65- 66.

Maureen A and Harold J. (1996) . Surgical management and treatment of gastric and duodenal fistulas . Surg Clin North Am . Vol. 76.no.(5):1137-1145.

McClave SA. and Heyland D.K. (2006). Parenteral Enteral Nutr; . JPEN J. November 1; 30 (6): 537 – 538.

Meissner K. (1999). Late radiogenic small bowel damage . Dig Surg. 26 (3); 169-74.

Memon AS and Siddiqui EG . (2004). Causes and management of postoperative enterocutaneous fistulas . J Coll Physicians Surg Pak . Jan;14(1); 25-8 .

Merine D, Fishman EK, Kuhlman JE, Jones B, Bayless TM, and Siegelman S. (2001). Bladder involvement in Crohn disease: detection and evaluation by C.T. J Comput Assist Tomogr . 25 ; 90-93.

Metcalfe C. (1999). Enterocutaneous fistulae; A review of the types of enterocutaneous fistula and their management (electronic version). Journal of Wound Care. 10 (2); 157-161.

Michael H. Humphreys .(2003) . Fluid and Electrolyte management . Current Surgical Diagnosis and Treatment 11th edition , Lange Medical Books/Mc Graw-Hill. 1(10) :142-155 .

Michael M. Mequid, and Antonio C.L. Campos .(1996) . nutritional management of patients with gastrointestinal fistulas . Euro Nutri. .76(5);1035- 1046.

Morris J, Spencer JA, and Ambrose NS. (2000). MR imaging classification of pelvic fistulas and its implications for patient management. RadioGraphics. 20:623-635.

Mucha P. (1996) . Small intestinal obstruction .Surg.Clin. north Am . 67:597-620.

Mughal MM ., Bancewicz J., and Irving MH. (1996). "Laparostomy", a technique for the management of intractable intra-abdominal sepsis . British Journal of Surgery . 73 :253 -259 .

Ohanaka CE, Momoh IM, and Osime U. (2001) . Management of enterocutaneous fistulae in Benin City Nigeria . Trop Doct. Review . Apr;31(2); 104-6.

Ono Y, Sekihara T, Kurokawa K, and Saruki K. (1995). Renocolic fistula caused by xanthogranulomatous pyelonephritis. Int J Urol. 2:277-278.

Orton DF, LeVeen RF and Saigh JA. (2000). Aortic prosthetic graft infections; radiologic manifestations and implications for management. RadioGraphics . 20:977-993.

Paauw JD, Mc Camish MA, Dean RE, and Ouellette RT. (1984). Assessment of caloric need in stressed patients. J. Am Coll Nutr. 3;182-194.

Parvey HR, Cochran ST, Payan J, Goldman S, and Sandler CM .(1997). Renocolic fistulas ; complementary roles of computed tomography and direct pyelography. Abdom Imaging. 22:96-99.

Pickhardt PJ, Gouma D, and Bhalla S. (2000) .Spontaneous pneumopericardium secondary to penetrating benign gastric ulcer. Clin Radiol .55;798-800.

Poulin EC,Schlachta CM, and Mamazza J. (2000). Should enteric fistula from CD or diverticulitis be treated laparoscopically or by open surgery ? A matched cohort study .Dis Colon Rectum . 43:621-736.

Radin DR, Ray MJ and Halls JM. (1986). Strangulated diaphragmatic hernia with pneumothorax due to colopleural fistula. AJR Am J Roentgenol; 146:321-322.

Regan JP and Salky BA. (2004) . laparoscopic treatment of enteric fistulas.Surg Endosc.18 ; 252-254.

Reignier J, Thenoz-Jost N, and Fiancette M. (2004). Early enteral nutrition in mechanically ventilated patients in the prone position. Crit Care Med . 32: 94-99.

Rex DK. (2000). Virtual Colonoscopy: time for some tough question for Radiologists and Gastroenterologists. Endoscopy. 32(3): 260-263,.

Rinsema W, Gouma DJ and Meyenfeldt MF.(1990). primary conservative management of external small bowel fistulas.Acta Chir Scand. 156:457-462.

Rolando Rolandelli and Joel J . (1996). Surgical management and treatment of the sepsis associated with gastrointestinal fistulas.Clin North Am . vol.76,No(5):1111-1121.

Rubesin SE, Scotiniotis I, Birnbaum BA and Ginsberg GG. (2001). Radiologic and endoscopic diagnosis of Crohn's disease. Surg Clin North Am. 81:39-70.

Ruben G J, Visscher, and Steven V M . (2008). olde damink treatment strategies in 135 consecutive patients with entero cutaneous fistulas . world surgery. 32:445-453.

Russell R.C.G , Williams N.S, and Bulstrode C.J.K. (2000). Enterocutaneous faecal fistula . Br J Surg . (87) ;650-657 .

Sabharwal T, Hatzidakis A, Papageorgiou G . (2004). Trans-oral embolization of a post-operative duodenocutaneous fistula. Clinical Radiology. 59;102-104.

Sakr M . (1998). Principles of surgery. Small intestine . intestinal fistula .1st edition . Dar Elkatab . (4) (3) ;263-293.

Sarfeh IJ and Jakowatz JG .(1992) . Surgical treatment of enteric "bud" fistula in contaminated wounds; a risk less extraperitoneal method using split-thickness skin grafts. Arch Surg . 127:1027-1031.

Sanchez J, Albarracin A, loby H . (2003). Electrical nerve stimulation in the management of the enterocutaneous low-output fistulae. Canad J Surg. 49 : 185 – 198.

Schein M and Decker GA .(1991). postoperative external alimentary tract fistulas. Am. J Surg . 161:435-138.

Schneider . (2006). Nutrition Support Teams: An evidence-based practice.. Nutr Clin Pract. 21: 62-67.

Schraut WG, Champman C, and Abraham VS . (2000). operative treatment of Crohn's ileocolitis complicated by ileosigmoid and ileovesical fistulae. Ann Surg . 207:48-49.

Sciarretta G , Furno A, and Mazzoni M . (1993). scintigraphy in C.D . Gut .34 ; 1364-1369.

Semelka RC, Hricak H, and Kim B .(1997) . Pelvic fistulas; appearances on MR images. Abdom Imaging. 22:91-95

Sheldon GF. And Gardiner BN .(2002). Management of gastrointestinal fistulae. Canad J Surg. 45 : 385 – 398.

Shou SE, Scotiniotis I, and Ginsberg GG . (2001). Management of perianal fistulae. Surg Clin North Am. 81:345-49.

Sims G J, Furno A, and Steven V M . (2004). Colo -vaginal fistulae. world surgery. 28:485-490.

Singer G .(1998) . Fluid and electrolyte management . The Washington Manual of Medical Therapeutics 29th edition . Lippincott-Raven;(3):39-60 .

Slade D and Scott N (2005). Intestinal fistula management . recent advance in surgery 28; 87-98.

Sloan S, Bartlett BSN, and Wei RN .(2002). Management of the patient with an enterocutaneous fistula . Br J Surg . Dec;(89);183-196.

Sou S , Matsui T , and Yao T .(2006).Iodocyanine green as marker in entero cutaneous fistula. Gastro Enterol Hepatol. Dec 21 ; 1850-3.

Soeter P B ., Ebeid AM ., & Fisher JE .(1999). Review of 404 Patients with gastrointestinal fistula . Impact of Parenteral nutrition .Annals of Surgery ,190 :189 – 202.

Stein TP. (1985). Why measure the respiratory quotient of patients on total parenteral nutrition? Am Coll Nutr 4: 501-502.

Stephen A. Mc Clave, MD, Wei-Kuo chang, MD, Rupinder Dhaliwal, RD, and Daren K. (2006) . Nutrition support in acute pancreatitis. Journal of parental and enteral nutrition. Vol 30, No 2: 143-156.

Stollman NH, Raskin JB .(1999). Diverticular disease of the colon. J Clin Gastroenterol. 29:241-252.

Stroud M, Dun Can H,and Night tingale J. (2003). Guidelines for enteral feeding in adult hospital patients. Gut; 52 ; 56-63.

Schner U, Katz DP, and Furst P . (2001) . Effects of intravenous fat emulsions on lung function & liver in patients with acute respiratory distress syndrome or sepsis. Crit Care Med . 29: 1569-1574.

Swanimar DL, Phunq PT, and Jones RL. (1987). Twenty-four hour energy expenditure in critically ill patients. Crit Care Med .15: 637-638.

Tachi M and Hirabayashi S. (2002). Enterocutaneous fistula treated with a fasciocutaneous turnover flap. Ann Plast Surg. 48:554-556.

Triantafillidis JK, Pomonis E, Emmanouilidis A, Merikas EG, Tsitsa C and Hanjiandreou E. (1999). Surgery for ulcerative colitis in Greece; clinicoepidemiological features and long-term outcome of 69 consecutive patients. Hepatogastroenterology. 22:961-968.

Tsiotos GG, Smith CD, and Sarr MG. (1995). Incidence and management of pancreatic and enteric fistulas after surgical management of severe necrotizing pancreatitis. Arch Surg . 130:48-52.

Tuney D, Furst P, and Cimsit C. Bouveret's syndrome.(2000). CT findings. Eur Radiol. 10:1711-1712.

Tweedle D . (1998) . Fluid and nutrient replacement . Journal of the American College of anath. (21):23-45 .

Vagholkar K. (2001). Small intestinal fistula (Case report): Bombay Hospital Journal Oct.61;231-248.

Valeri M and Christopher (2002). Gastrointestinal function. Europ physio J. 25;231-248.

Van Way C.W. (2006). Perioperative immunonutrition: good idea about

Parenter Enteral Nutrition. Canad Nutri. November 1, 30 (6): 539-540.

Vargish T and Baker J. (2001). Fluid and electrolytes disorders in acquired intestinal fistula . South Afr J Surg . 28 ; 238-249.

Vining DJ and Gelfand DW. (1994). Non invasive colonoscopy using helical CT scanning, 3D reconstruction, and virtual reality. Presented at the 23 rd Annual meeting and postgraduate course of the society of gastrointestinal radiologists, Mani, Hawaii,

Walker CW, Purnell GL, and Diner WC. (1999). Imaging of acquired G.I fistula. Radiology . 73:618-620.

Wang WY, Hernandez JE, Denny B, Gerard PS and Horovitz J. (1998). Postradiation colospinal fistula; a rare cause of spinal epidural abscess. AJNR Am J Neuroradiol . 19:1959-1960.

Wang Bo , Jian An, and Shou Li .(2002).changes in body composition in entero cutaneous fistula . Euro Nutri. 15 ; 1149-1152.

Weissenfluh G.M, Brundage S.I, and Spain D.A. (2006). Early enteral nutrition after abdominal trauma: effects on septic morbidity and practicality. Nutr Clin Pract, October 1; 21 (5): 522 – 528.

Wong Sk, Lam YH, and Lau JY.(2000). Diagnosis and therapeutic fistuloscopy. an adjuvant management in postoperative fistulas and abscesses after UGI surgery.Endoscopy. 32(4)311-3.

Yamashita H, Chijiwa K, and Ogawa Y.(1997). The internal biliary fistula: reappraisal of incidence, type, diagnosis, and management of 33 consecutive cases. HPB Surg .10:143-147.

Yee J, Kumar NN, Hung RK, Akerkar GA, Kumar PG, and Wall SD. (2003). Comparison of supine and prone scanning separately and in combination at CT colonography. Radiology. 226; 653-661.

Yee LF, Birnbaum EH, Read TE, Kodner IJ, Fleshman JW. (1999). Use of endoanal ultrasound in patients with rectovaginal fistulas. Dis Colon Rectum; 42:1057-1064.

Yeh TS , Jan YY and Jenq LB. (1999). Pancreaticojejunal anastomosis leak after pancreaticoduodenectomy-multivariate analysis of peri-operative risk factors. J Surg Res . 67:119-25 .

List of abbreviation

G.I	Gastrointestinal
BCM	Body cell mass
BEE	Basal energy expenditure
BIA	Bioelectrical impedance analysis
BLP	Blood pressure
BMI	Body mass index
C.T.C	Computed tomographic colonography
CD	Crohn`s disease
CT	Computed tomography
CVP	Central venus pressure
DM	Diabetes mellitus.
ECW	Extra cellular water
ECW	Extra cellular water
EN	Enteral Nutrition
ERCP	Endoscopic retrograde cholangio pacreatography
FFM	Fat free mass
GIT	Gastrointestinal tract.
HIDA	Hydroxy imino di-acetic acid
HOF	High output fistula
HPN	Home Parenteral nutrition
IBD	Inflammatory bowel disease
ICW	Intra cellular water
LN	Lymph node.
LOF	Low output fistula
MRCP	Magnetic resonance cholangio pancreatography
MRI	Magnetic resonance imaging

PN	Parenteral nutrition
PPN	Peripheral parenteral nutrition
SBS	Short Bowel Syndrome
TB	Tuberculosis
TBW	Total body water
TPN	Total parenteral nutrition
U/S	Ultrasonography
UOP	Urine out put
UTI	Urinary tract infection
VAC	Vacuum assisted closure
Na	Sodium
K	Potassium
CL	Chloride
HCO ₃	Bicarbonate
Ca	Calcium
Mg	Magnesium
KCl	Potassium chloride
SOWATS	Sepsis, optimizing nutritional state, wound care, anatomy of the fistula, timing of surgery, and surgical strategy
SNAP	Sepsis and skin care, Nutrition, Anatomical determination, and Performing procedure
H ₂ antagonists	Histamine receptors antagonists
H-K atpase inhibitors	Hydrogen-potassium atpase inhibitors
TSH	Thyroid stimulating hormone
Wt	Weight

A	Age
Ht	Height
rh,GH	Recumbnant human growth hormon
MP	Mercapto Purine
TNF	Tumour Necrosis Factor
PO	Per oral
ENS	Electrical nerve stimulation
CBD	Common bile duct
NSAIDS	Non steroidal anti inflammatory drugs
Ph	phosphrus
Zn	Zinc
D	Dimension

