

References

- Abel, E. and Lebwohl, M. (2005):** Psoriasis. In: Dale DC, Federman DD, eds. ACP Medicine; sec(2),ch(3): 401-410. New York.
- Abel, E.A.; DiCicco, L.M. and Orenberg, E.K. (2001):** Drugs in exacerbation of psoriasis. J Am Acad Dermatol; 15(5): 1007-22.
- Aizawa, S.; Nakano, H. and Ishida, T. (1999):** Tumor necrosis factor Receptor-Associated Factor (TRAF)-5 and TRAF-2 are involved in CD₃₀-mediated NF-kappa-B activation". J. Biol. Chem.; 272(4): 2042-5.
- Al-Khenaizan, S. and Al-Mubarak, L. (2008):** Ulcerative lichen planus of the sole: Excellent response to topical tacrolimus Inter. J. of Derm; 6: 626-628.
- Ansieau, S.; Scheffrahn, I. and Mosialos, G. et al. (2000):** Tumor necrosis factor Receptor-Associated Factor (TRAF)-1, TRAF-2, and TRAF-3 interact in vivo with the CD₃₀ cytoplasmic domain; TRAF-2 mediates CD₃₀-induced nuclear factor kappa B activation. Proc. Natl. Acad. Sci. USA; 93(24): 14053-8.
- Antonella Tosti; Bianca Maria Piraccini and Stefano Cambiaghi (2001):** Nail Lichen planus in children clinical features, response to treatment, and long-term follow-up Arch dermatol; 137: 1027-1032.
- Armitage, F. (1998):** Statistical methods in scientific researches, Blackwell, Scientific Publication, New York.
- Asarch, A.; Gottlieb, A. and Lee, J. (2008):** Lichen planus-like eruptions: An emerging side effect of tumor necrosis factor- α antagonists. JAAD; 61(1): 104-111.
-

- Bal, N.; Tuncer, I. and Baba, M. (2008):** Bcl-2 expression in dermal lymphocytes in lichen planus and psoriasis vulgaris. *J Eur Acad Dermatol Venereol*; 22(5): 640.
- Barranco, C.; Toll, A. and Pujol, R.M. (2002):** CD₃₀ antigen expression in cutaneous inflammatory infiltrates of scabies: A dynamic immunophenotypic pattern that should be distinguished from lymphomatoid papulosis. *J Cutan Pathol*; 29(6): 368-73.
- Bascones-Ilundain, C.; Gonzalez-Moles, M.A. and Esparza-Gmez, G. (2006):** Importance of apoptotic mechanisms in inflammatory infiltrate of lichen planus lesions. *Anticancer Res*; 26: 357-363.
- Battino, M.; Greabu, M. and Totan, A. (2008):** Oxidative stress markers in oral lichen planus; 33(4): 301-10.
- Bekkenk, M.W.; Geelen, F.A. and van Voorst Vader, P.C. (2000):** Primary and secondary cutaneous CD₃₀⁺ lymphoproliferative disorders. *Blood*; 95: 3653-3661.
- Beljaards, R.C.; MeiJer, C.J. and Van Der Putte, S.C. (2000):** Primary cutaneous T-cell lymphoma: Clinicopathological features and prognostic parameters of 35 cases other than mycosis fungoides and CD30-positive large cell lymphoma. *J pathol*; 72: 53-55.
- Benoist, C. and Mathis, D. (2002):** Mast cells in autoimmune disease. *Nature*; 420: 875-878.
- Bernard, A.; Boumsell, L. and Dausett, J. (2000):** Leucocyte Typing: Human Leucocyte Differentiation Antigens Detected by Monoclonal Antibodies. Berlin.
-

- Betina Werner; Cesare Massone and Helmut Kerl (2006):** Large CD₃₀-positive cells in benign, atypical lymphoid infiltrates of the skin. *Journal of Cutaneous Pathology*; 35(12): 1100-1107.
- Bezold, G.; Lange, M. and Schiener, R. (2001):** Hidden scabies: Diagnosis by polymerase chain reaction. *Br J Dermatol*; 144(3): 614-8.
- Biondi Oriente, C.; Scarpa, R. and Pucino, A. (1999):** Psoriasis and psoriatic arthritis. Dermatological and rheumatological co-operative clinical report. *Acta Derm Venereol Suppl (Stockh)*; 146: 69-71.
- Black, M.M. (2002):** Lichen planus and lichenoid disorders. In: *Textbook of Dermatology*. Champion RH, Burton JL and Ebling FJG (eds.), 5th ed. Blackwell Scientific Publications, Oxford; 3: 1670-1675.
- Borchmann, P.; Schnell, R. and Fuss, I. (2002):** Phase 1 trial of the novel bispecific molecule H22xKi-4 in patients with refractory Hodgkin lymphoma. *Blood*; 100(3): 3101-3107.
- Boucher, L.M.; Marengère, L.E. and Lu, Y. (2001):** Binding sites of cytoplasmic effectors TRAF-1, 2, and 3 on CD₃₀ and other members of the TNF receptor superfamily. *Biochem. Biophys. Res. Commun.*; 233(3): 592-600.
- Bowcock, A.M. and Cookson, W.O. (2004):** The genetics of psoriasis, psoriatic arthritis and atopic dermatitis. *Human Mol Genet*; 13(1): 43-55.
- Bruce, A.J. and Rogers, R.S. (2003):** Oral psoriasis. *Dermatol Clin*; 21(1): 99-104.
-

- Burkhart, C.G.; Burkhart, C.N. and Burkhart, K.M. (2005):** An epidemiologic and therapeutic reassessment of scabies. *Cutis*; 65(4): 233-40.
- Cavagni, G.; Caffarelli, C. and Facchetti, F. (2000):** *Int Arch Allergy Immunol*; 121: 224-241.
- Cavalli, F.; Isaacson, P.G. and Gascoyne, R.D. (2001):** MALT lymphomas. *Hematology Am Soc Hematol Educ Program*; 120: 241-58.
- Cendras, J. and Bonnetblanc, J.M. (2008):** Erosive oral lichen planus. *Ann Dermatol Venereol*; 136(5): 458-70.
- Cepeda, L.T.; Pieretti, M. and Chapman, S.F. (2003):** CD₃₀-positive atypical lymphoid cells in common non-neoplastic cutaneous infiltrates rich in neutrophils and eosinophils; 27(7): 912-8.
- Chiarle, R.; Antonello Podda and Gabriel Prolla (2002):** CD₃₀ in normal and neoplastic cells. *Clinical Immunology*; 90(157): 164-189.
- Chosidow, O. (2006):** Clinical practices. Scabies. *N Engl J Med*; 354(16): 1718-27.
- Chouela, E.; Abeldano, A. and Pellerano, G. (2002):** Diagnosis and treatment of scabies: A practical guide. *Am J Clin*; 3(1): 9-18.
- Conrotto, D.; Carbone, M. and Carrozzo, M. (2006):** Ciclosporin versus clobetasol in the topical management of atrophic and erosive oral lichen planus: A double-blind, randomized controlled trial. *Br J Dermatol*; 154(1): 139-45.
- De Rie; Menno and Amber Goedkoop (2004):** Overview of psoriasis. *Dermatologic Therapy*; 17: 341-359.
-

- DeBruin, P.C.; Gruss, H.J. and van der Valk, P. (2000):** CD₃₀ expression in normal and neoplastic lymphoid tissue: Biological aspects and clinical implications. *Leukemia*; 9(6): 1620-1636.
- Del Prete, G.; Maggi, E. and Pizzolo, G. (2000):** CD₃₀, Th₂ cytokines and HIV infection: A complex and fascinating link. *Immunol Today*; 16: 76-81.
- Delves, P.J. and Roitt, I.M. (2005):** The immune system. Second of two parts. *N Engl J Med*; 343(2): 108-17.
- Diamantis, S.A.; Morrell, D.s. and Burkhart, C.N. (2007):** Pediatric infestations. *Pediatric Ann*; 38(6): 326-32.
- Diaz-Cascajo, C. (2001):** Strong immunoexpression of the monoclonal antibody CD₃₀ in lymphocytic infiltrates of the skin not by itself evidence for diagnosing malignant lymphoma. *Am J Dermatopathol*; 23(5): 179-190.
- Duckett, C.S. and Thompson, C.B. (2001):** CD₃₀-dependent degradation of TRAF₂: Implications for negative regulation of TRAF signaling and the control of cell survival. *Genes Dev*; 11: 2810-2821.
- Duckett, C.S.; Gedrich, R.W. and Gilfillan, M.C. (1997):** Induction of nuclear factor kappa B by the CD₃₀ receptor is mediated by TRAF₁ and TRAF₂. *Mol Cell Biol*; 17: 1535-1542.
- Dummer, W.; Rose, C. and Brocker, E. (2002):** Expression of CD₃₀ on T-helper cells in the inflammatory infiltrate of acute atopic dermatitis but not of allergic contact dermatitis. *Arch Dermatol Res*; 290: 598-608.

- Fan, Y.; Zhan, Z. and Peng, T. (2004):** The expression of apoptosis-associated proteins Bcl-2, Bax in oral leukoplakia and lichen planus. *Shanghai Kou Qiang Yi Xue*; 13(6): 497-506.
- Femiano, F. and Scully, C. (2006):** Oral lichen planus: Clinical and histological evaluation in an open trial using a low molecular weight heparinoid (sulodexide). *Int J Dermatol*; 45(5): 986-989.
- Fernandez, N.; torres, A. and Ackerman, A.B. (1997):** Pathologic findings in human scabies. *Arch Dermatolo*; 113(6): 320-335.
- Flávia, C.; Passos, D.O. and Garcia, V.S. (2004):** Biochemical characterization and biosynthesis of the Ki-1 antigen in Hodgkin-derived and virus-transformed human B and T lymphoid cell lines. *J. Biol. Chem.*; 279(4): 11444-11455.
- Fox, G. (2005):** Diagnosis of scabies by dermoscopy: Case reports. *BMJ*; 331(6): 972-972.
- Gedrich, R.W.; Gilfillan, M.C. and Duckett, C.S. (2001):** CD₃₀ contains two binding sites with different specificities for members of the tumor necrosis factor receptor-associated factor family of signal transducing proteins. *J. Biol. Chem.*; 271(22): 12852-8.
- Giardina, E.; Sinibaldi, C. and Novelli, G. (2004):** The psoriasis genetics as a model of complex disease. *Curr Drug Targets Inflamm Allergy*; 3(6): 129-36.
- Gonzalez-Moles, M.A.; Scully, C. and Gil-Montoya, J.A. (2008):** Oral lichen planus: Controversies surrounding malignant transformation. *Oral Dis*; 14(3): 229-43.
- Graham Johnston (2005):** Scabies: Diagnosis and treatment. *BMJ*; 331(4): 619-622.
-

- Granados, S. and Hwang, S.T. (2004):** Roles for CD₃₀ in the biology and treatment of CD₃₀ lymphoproliferative diseases. *J invest Dermatol*; 122(6): 1345-7.
- Guðjónsson, J.E. (2002):** HLA-Cw6-positive and HLA-Cw6-negative patients with psoriasis vulgaris have distinct clinical features. *Journal of Investigative Dermatology*; 118(2): 362-365.
- Habib, F.; Stoeber, P.E. and Picot, E. (2005):** Narrow band UVB phototherapy in the treatment of widespread lichen planus. *Ann Dermatol Venereol*; 132(1): 17-24.
- Haneke, E. (2007):** Noninfectious inflammatory disorders of the nail apparatus. *J Dtsch Dermatol Ges*; 349(2):15-23.
- Heath Christian Kurts, W.R.; Irina Caminschi and Carbone, F.R.** CD₃₀ prevents T-cell responses to non-lymphoid tissues. *Immunological. Rev.*; 169(3): 23-29.
- Heath Christian Kurts, W.R.; Irina Caminschi and Carbone, F.R.** CD₃₀ prevents T-cell responses to non-lymphoid tissues. *Immunological. Rev.*; 169(4): 23-29.
- Henseler, T. and Christophers, E. (2000):** Psoriasis of early and late onset: characterization of two types of psoriasis vulgaris. *J Am Acad Dermatol*; 13: 450-60.
- Hicks, M.I. and Elston, D.M. (2008):** Scabies. *Dermatol Ther*; 22(4): 279-92.
- Holsinger, L. (2004):** A battle with my skin. *Journal of the American Academy of Dermatology*; 51(1): S41-42.
- Horie, R. and Watanabe, T. (1998):** CD₃₀ expression and function in health and disease. *Semin Immunol*; 10(1): 457-470.
-

- Horie, R.; Higashihara, M. and Watanabe, T. (2003):** Hodgkin's lymphoma and CD₃₀ signal transduction.. *Int. J. Hematol.*; 77(1): 37-47.
- Huang, B.B.; Bonish, B.K. and Chaturvedi, V. (2001):** Keratinocyte CDw60 expression is modulated by both a Th-1 type cytokine IFN-gamma and Th-2 cytokines IL-4 and IL-13: Relevance to psoriasis. *J Invest Dermatol*; 116(6): 305-12.
- Inaloz (2006):** Lichen planus; 12(5): 463-8.
- Ingafou, M.; Leao, J.C. and Porter, S.R. (2006):** Oral lichen planus: A retrospective study of 690 British patients. *Oral Dis*; 12(5): 463-8.
- Ismail, S.B.; Kumar, S.K.S. and Zain, R.B. (2007):** Oral lichen planus and lichenoid reactions: Etiopathogenesis, diagnosis, management and malignant transformation. *Journal of Oral Science*; 49: 89-106.
- Jang, K.A.; Kim, S.H. and Choi, J.H. (2000):** Lichenoid keratosis: A clinicopathologic study of 17 patients. *J Am Acad Dermatol*; 43(3): 511-16.
- Jariwala, S.P. (2007):** The role of dendritic cells in the immunopathogenesis of psoriasis. *Arch Dermatol Res*; 299(8): 359-366.
- Jiaravuthisan, M.M.; Sasseville, D. and Vender, R.B. (2007):** Psoriasis of the nail: anatomy, pathology, clinical presentation, and a review of the literature on therapy. *J Am Acad Dermatol*; 57(1): 1-27.
- Jobling, R. (2007):** A patient's journey: Psoriasis. *Br Med J*; 334: 953-962.
- Johnson, R.M. and Brown, E.J. (2000):** Cell-mediated immunity in host defense against infectious diseases. In: Mandell GL, Bennett JE,

- Dolin R, eds. Principles and Practice of Infectious Disease. 5th ed. Philadelphia, Pa: Churchill Livingstone; 55:131-134.
- Johnston, G. and Sladden, M. (2005):** Scabies: Diagnosis and treatment. *BMJ*; 331(7517): 619-22.
- Josimovic, O.; Durkop, H. and Schwarting, R. (1999):** Ki-1 (Cd₃₀) antigen is released by Ki-1 positive tumor cell in vitro and in vivo: partial characterization of soluble Ki-1 antigen and detection of the antigen in cell culture supernatants and in serum by an enzyme-linked immunosorbent assay. *Eur. J. Immunol.*; 19(8): 157-162.
- Kadin, M.E.; Levi, E. and Kempf, W. (2001):** Progression of lymphomatoid papulosis to systemic lymphoma is associated with escape from growth inhibition by transforming growth factor-beta and CD₃₀ ligand. *Ann N Y Acad Sci*; 20(7): 59-68.
- Kimball, A.; Gladman, D. and Gelfand, J. (2008):** National Psoriasis Foundation clinical consensus on psoriasis comorbidities and recommendations for screening. *Journal of the American Academy of Dermatology*; 58(6): 1031-1042.
- Krathen, R.A.; Ward, S. and Duvic, M. (2003):** Bexarotene is a new treatment option for lymphomatoid papulosis. *Dermatology*; 206: (5)142-147.
- Krueger, G., and Ellis, C.N. (2005):** Psoriasis: Recent advances in understanding its pathogenesis and treatment. *J. Am. Acad. Dermatol.*; 53(1): 94-100.
- Kumar, V.; Nangia, V. and Nangia, A. (2000):** Lichen planus: A clinico-histopathological correlation. *Indian Journal of Dermatology Venereology and Leprology*; 66(2): 193-195.
-

- Kurts, C.; Carbone, F.R. and Krummel, M.F. (2001):** Signalling through CD₃₀ protects against autoimmune diabetes mediated by CD₈ T-cells. *Nature*; 398(6725): 341-340.
- Laporte, M.; Galand, P. and Fokan, D. (2000):** Apoptosis in established and healing psoriasis. *Dermatology*; 200(4): 314-322.
- Lee, C.S. and Koo, J. (2007):** Papulosquamous disorders. In: Rakel RE, Bope ET, eds. *Conn's Current Therapy*. Philadelphia: Elsevier Saunders; 12(9):931-936.
- Lee, D. (2002):** Mast cells: A cellular link between autoantibodies and inflammatory arthritis. *Science*; 297(4): 1689-1692.
- Levi, E.; Pfeifer, W.M. and Kadin, M.E. (2006):** CD₃₀-activation-mediated growth inhibition of anaplastic large-cell lymphoma cell line: Apoptosis or cell-cycle arrest? *Blood*; 98(7): 1630-1632.
- Levi, E.; Wang, Z. and Petrogiannis-Haliotis, T. (2000):** Distinct effects of CD₃₀ and Fas signaling in cutaneous anaplastic lymphomas: A possible mechanism for disease progression. *J Invest Dermatol*; 115(6): 1034-1040.
- Lewis, F.M. (2002):** Vulval lichen planus. *Br J Dermatol*; 138(4): 569-75.
- Lin, A.N.; Reimer, R.J. and Carter, D.M. (2001):** Scabies. *J Am Acad Dermatol*; 18(9): 553-58.
- Liu, H.N.; Sheu, W.J. and Chu, T.L. (1999):** Scabietic nodules: A dermatopathologic and immunofluorescent study. *J Cutan Pathol*; 19: 124-130.
- Lodi, G.; Scully, C. and Carrozzo, M. (2005):** Current controversies in oral lichen planus: Report of an international consensus meeting.
-

- Part 1. Viral infections and etiopathogenesis. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*; 100(5): 40-51.
- Louden, B.A.; Pearce, D.J. and Lang, W. (2004):** A Simplified Psoriasis Area Severity Index (SPASI) for rating psoriasis severity in clinic patients. *Dermatol. Online J.*; 10(2): 7-15.
- Manolache, L.; Seceleanu-Petrescu, D. and Benea, V. (2008):** Lichen planus patients and stressful events. *J Eur Acad Dermatol Venereol*; 22(4): 437-41.
- Markell, E.K.; John, D.C. and Petri, W.H. (2006):** Markell and Voge's Medical Parasitology 9(6):212-220
- Marshall, J.S., King, C.A., and McCurdy, J.D. (2003):** Mast cell cytokine and chemokine responses to bacterial and viral infection. *Curr. Pharm. Des.*; 9(2): 11-24.
- Mary Tonsager and Crutchfield, C.E.III (2004):** Atrophic Lichen planus. *Dermatol Nurs*; 16(1): 73-74.
- Mathieu, M.E. and Wilson, B.B. (2005):** Scabies. In: Mandell GL, et al., eds. *Mandell, Douglas, and Bennett's Principles and Practices of Infectious Diseases*; 6(292): 3304-3307.
- McCalmont, T.H. and LeBoit, P.E. (2006):** Lymphomatoid papule but not lymphomatoid papulosis. *Am J Dermatopathol*; 22: 188-202.
- McCarthy, J.S.; Kemp, D.J. and Walton, S.F. (2004):** Scabies: More than just an irritation. *Postgrad Med J*; 80: 382-407.
- McMichael, A.J. (2000):** Leukocyte typing III: White cell differentiation antigens. *Oxford University Press*; 3:336-338.
- Mehlis, S.L. and Gordon, K.B. (2003):** The immunology of psoriasis and biologic immunotherapy. *J Am Acad Dermatol*; 49: 44-50.
-

- Michale, J.T.; Fisher, G.J. and Lindquist, P.B. (2007):** Overexpression of transforming growth factor alpha in psoriatic epidermis. *Science*; 243(4892): 811-814.
- Mir, S.S.; Richter, B.W. and Duckett, C.S. (2000):** Differential effects of CD₃₀ activation in anaplastic large cell lymphoma and Hodgkin disease cells. *Blood*; 96: 4307-4312.
- Mizushima, S.; Fujita, M. and Ishida, T. (2001):** Cloning and characterization of a cDNA encoding the human homolog of Tumor necrosis factor Receptor-Associated Factor-5 (TRAF5). *Gene*; 207(2): 135-40.
- Moretó, M. and Pérez-Bosque, A. (2008):** Dietary plasma proteins, the intestinal immune system and the barrier functions of the intestinal mucosa. *J Anim Sci*;106(2)128-142
- Mori, M.; Manuelli, C. and Pimpinelli, N. (1999):** CD₃₀-CD₃₀ ligand interaction in primary cutaneous CD₃₀⁺ T-cell lymphomas: A clue to the pathophysiology of clinical regression. *Blood*; 94: 3077-3083.
- Mumcuoglu, K.Y.; Friger, M. and Ioffe-Uspensky, I. (2001):** Louse comb versus direct visual examination for the diagnosis of head louse infestations. *Pediatr Dermatol*; 18(2): 9-12.
- Munk, M.E.; Kern, P. and Kaufmann, S.H.E. (1997):** Human CD₃₀⁺ cells are induced by mycobacterium tuberculosis and present in tuberculosis lesions . *Int Immunol*; 9: 713-735.
- Murphy, M.; Kerr, P. and Grant-Kels, J. (2007):** The histopathologic spectrum of psoriasis. *Clinics in Dermatology*; 25(6): 524-528.

- Nagao, Y.; Sata, M. and Kage, M. (2002):** Histopathological and immunohistochemical study of oral lichen planus-associated HCV infection. *European Journal of Internal Medicine*; 6(11): 277-282.
- Naldi, L. (2004):** Epidemiology of psoriasis. *Curr Drug Targets Inflamm Allergy*; 3: 121-128.
- Namazi, M.R. (2004):** Paradoxical exacerbation of psoriasis in AIDS: proposed explanations including the potential roles of substance P and gram-negative bacteria. *Autoimmunity*; 37: 67-71.
- Nathan, C. (2002):** Points of control in inflammation. *Nature*; 420(5): 846-852.
- Nathan, D. and Belsto, D. (1998):** Carbamazepine induced pseudo lymphoma with CD₃₀ positive cells. *Jam Acad Dermatol*; 38(7): 806-826.
- Pabst, R. (2001):** Lymphocyte migration to the gut: Oversimplifications and controversial aspects. *Immunol Res*; 10(3): 279-81.
- Pavlotsky, F.; Nathansohn, N. and Kriger, G. (2008):** Ultraviolet-B treatment for cutaneous lichen planus: our experience with 50 patients. *Photodermatol Photoimmunol Photomed*; 24(2): 83-6.
- Picardi, A. and Abeni, D. (2001):** Stressful life events and skin diseases: Disentangling evidence from myth. *Psychother Psychosom*; 70(2): 118-136.
- Polysangam, T.; Breneman, D.L. and Mutasim, D.F. (1998):** Cutaneous pseudolymphomas. *J AM Acad Dermatol*; 38(4): 880-903.

- Roberts, A.B. and Wakefield, L.M. (2003):** The two faces of transforming growth factor beta in carcinogenesis. *Proc Natl Acad Sci USA*; 100(4): 8621-8623.
- Roberts, L.J.; Huffam, S.E. and Walton, S.F. (2005):** Crusted scabies: Clinical and immunological findings in seventy-eight patients and a review of the literature. *J Infect* ;50(4): 375-81.
- Rodriguez-Diaz, E.; Roman, C. and Yuste, M. (1998):** Cutaneous lymphadenoma. An adnexal neoplasm with intralobular activated lymphocytes. *Am J Dermatopathol*; 20(5): 74-92.
- Romagnani, S. (2001):** Th₁ and Th₂ in human diseases. *Clin Immunol Immunopathol*; 80(5): 225-228.
- Romero, M.A.; Seoane, J. and Varela-Centelles, P. (2002):** Clinical and pathological characteristics of oral lichen planus in hepatitis C +ve and –ve patients. *Clin Otolaryngol*; 27(1): 22-26.
- Rose, C.; Starostik, P. and Brocker, E. (1999):** Infection with parapoxvirus induces CD₃₀ positive cutaneous infiltrates in human. *J Cutan Pathol*; 26(4): 520-525.
- Saaeed, S.A.; Bazza, M. and Zaman, M. (2000):** Cefuroxime induced lymphomatoid hypersensitivity reaction. *Postgrad Med J*; 76(7): 577-581.
- Sabat, R.; Philipp, S. and Höflich, C. (2007):** Immunopathogenesis of psoriasis. *Exp Dermatol*; 16(10): 779-98.
- Sallusto, F.; Lenig, D. and Forster, R. (2000):** Two subsets of memory T-lymphocytes with distinct homing potentials and effector functions. *Nature*; 401(4): 708-715.
-

- Scarpa, R.; Oriente, P. and Pucino, A. (2005):** Psoriatic arthritis in psoriatic patients. *Br J Rheumatol*; 23(11): 246-50.
- Scheinfeld, N.S. (2004):** Controlling scabies in institutional settings: A review of medications, treatment models, and implementation. *Amer J Clin Dermatol*; 5(1): 31-37.
- Schleicher, S.M. and Stewart, P. (2001):** Scabies: The mite that roars. *Emerg Med*; 6: 54-58.
- Schneider, C. and Hübinger, G. (2003):** Pleiotropic signal transduction mediated by human CD₃₀: A member of the Tumor Necrosis Factor Receptor (TNFR) family. *Leuk. Lymphoma*; 43(7): 1355-66.
- Schon, M. and Henning Boehncke, W. (2005):** Psoriasis. *The New England Journal of Medicine*; 352: 1899-1912.
- Schwarting, R.; Gerdes, J. and Durkop, H. (1999):** Ber-H₂: A new anti-Ki-1 (CD₃₀) monoclonal antibody directed at a formal resistant epitope. *Blood*; 74(7): 1678-1682.
- Scully, C. (2002):** Lichen planus: Review and update on pathogenesis. *J Oral Pathol*; 14(6): 431-438.
- Shulman, T.S. (2006):** Skin diseases. *Pediatr Ann*; 38(6): 299-308.
- Silverman, S.Jr. (2000):** Oral lichen planus: A potentially premalignant lesion. *J Oral Maxillofac Surg*; 58(11): 1286-1299.
- Sladden, M.J. and Johnson, G.A. (2004):** Common skin infections in children. *BMJ*; 329: 95-99.
- Smith, C. and Barker, J.N.W.N. (2006):** Psoriasis and its management. *British Medical Journal*; 333(10): 380-389.
- Smith, K.J.; Barrett, T.L. and Neafie, R. (1998):** Is CD₃₀ (Ki-1) immunostaining in cutaneous eruption useful as marker of Th1 to
-

- Th2 cytokine swittching and/or as a marker of adavnced HIV-1 disease? *BR J Dermatol*; 138(9): 774-789.
- Smoller, B.R.; longarce, T.A. and Warnke, R.A. (1999):** Ki₁ (CD₃₀) expression in differentiation of lymphomatoid papulosis from insect bite reactions. *Mod Pathol*; 5: 492-503.
- Stebbing, J.; Woolfson, A. and Tom, B.D. (2005):** Conservation of unique cell-surface CD antigen mosaics in HIV-1-infected individuals. *Blood*; 106: 1003-1007.
- Steer, A.C.; Jenney, A.W. and Kado, J. (2006):** High burden of impetigo and scabies in a tropical country. *PLoS Negl Trop Dis*; 3(6): 467-470.
- Stein, H; Mason, D. and Gerdes, J. (1998):** Expression of Hodgkin's disease-associated antigen Ki-1 in reactive and neoplastic lymphoid tissue: Reed-Sternberg cells and histocytic malignancies are derives from activated lymphoid cells. *Blood*; 66: 848-853.
- Steven Kossard; Carol Thompson and Duncan, G.M. (2004):** Pathway to neoplasia. *Arch Dermatol*; 140: 1262-1267.
- Sugerman, P.; Savage, N. and Walsh, L. (2002):** Pathogenesis of oral lichen planus. *Crit Rev Oral Biol Med*; 13: 350-365.
- Tag-El-Din Anbar; Manal Barakat and Ghannam, S.F. (2003):** Clinical and epidemiological study of lichen planus among Egyptians of Al-Minya province. *Dermatology Online Journal*; 11(2): 4-6.
- Tan, E.; Malik, R. and Quirk, C.J. (2000):** Hypertrophic lichen planus mimicking squamous cell carcinoma. *The Australasian Journal of Dermatology*; 39(1):245-253.
-

- Taneja, A. and Taylor, C.R. (2002):** Narrow-band UVB for lichen planus treatment. *Int J Dermatol*; 41(5): 282-283.
- Tarkowski, M. (2004):** Expression and a role of CD₃₀ in regulation of T-cell activity. *Curr. Opin. Hematol.*; 10(4): 267-271.
- Tesner, B.; Williams, N.O. and Brodell, R.T. (2007):** The pathophysiologic basis of scabietic nodules. *J Am Acad Dermatol*; 57(2): 56-70.
- Thompson, J.; Cochrane, T. and Cochran, R. et al. (1999):** Histology simulating reticulosis in persistent nodular scabies. *Br J Dermatol*; 90: 421.
- Ulmer, A.; Schanz, S. and Röcken, M. (2007):** A papulovesicular rash in a farmer and his wife. *Clin Infect Dis*; 45(3): 395-396.
- Van der Putte, S.C.; Toonstra, J. and van Wichen, D.F. (2005):** The expression of the Hodgkin's disease-associated antigen Ki-1 in cutaneous infiltrates. *Acta Derm Venereol*; 68(3): 202-208.
- Villarroel-Dorrego, M.; Correnti, M. and Delgado, R. (2002):** Lichen planus: Immunohistology of mucosal lesions. *J Oral Pathol Med*; 31(9) 410-414.
- Weed B.R and Folpe, A.L. (2008):** Cutaneous CD₃₀-positive epithelioid angiosarcoma following breast-conserving therapy and irradiation. *The American Journal of Dermatopathology*; 30: 370-372.
- Weedon, D. and Strutton, J. (2002):** The psoriasiform reaction pattern. In: *David Weedon Skin Pathology*; 42(2): 75-83.
- Willemze, R. and Beljaards, R.C. (1999):** Spectrum of primary cutaneous CD₃₀ (Ki-1)-positive lymphoproliferative disorders: A

proposal for classification and guidelines for management and treatment. J Am Acad Dermatol; 28(4): 973-978.

Woolfson, A.; Ellmark, P. and Chrisp, J.S. (2006): The application of CD antigen proteomics to pharmacogenomics. Pharmacogenomics; 7: 759-771.

Yosipovitch, G.; Goon, A. and Wee, J. (2000): The prevalence and clinical characteristics of pruritus among patients with extensive psoriasis. Br J Dermatol; 143(7): 969-973.

Zola, H.; Swart, B. and Nicholson, I. (2005): CD molecules: Human cell differentiation molecules. Blood; 106(9): 3123-6.