

RESULTS

This study comprised 420 volleyball players. They were 300 males (71.4 %) and 120 females (28.6 %).

A- Men's Tournament :

During the 1993-1994 season, 76 players reported a total of 84 injuries. During competitions, 303 hours of volleyball were played. The injury rate was 0.28 / hour.

A breakdown of injuries by anatomical site showed that the ankle was the most frequently injured joint 24% (20 of 84), followed by the knee 19 % (16 of 84), hands and wrists 17 % (14 of 84), low back 14 % (12 of 84) and shoulder 11 % (9 of 84). There was no significant difference ($p > 0.05$) between male and female volleyball players as regards to the anatomical site of injury (Table 10 and Fig. 1).

A breakdown of the injuries by type showed that strain accounted for 36% (30 of 84) of injuries, sprains for 27% (23 of 84) and inflammation and overuse (together) for 26% (22 of 84) of the reported injuries. There was no significant difference ($p > 0.05$) between male and female volleyball players as regards to the type of injury (Table 11 and Fig. 2).

Position on the court and attempted play at the time of injury affected the rate of injury. Position 3 (middle hitter / middle blocker) had the highest rate of injury, followed by position (4) and position (2). There was no significant difference ($P > 0.05$) between male and female

volleyball players according to injuries related to position (Table 12 and Fig. 3a & 3b).

Eighty five percent (75 /88) of the injuries were related to a particular type of volleyball maneuvers. Attempt to block accounted for 55% (41/75) of these injuries. Attempt to hit accounted for 25% (19/75). Attempt to defense accounted for 15 % (11/ 75). There was no significant difference ($p > 0.05$) between male and female volleyball players

according to injuries related to maneuvers (Table 13 and Fig. 4).

B- Female Tournament :

During the 1993-1994 season, 11 female players reported a total of 13 injuries. During competitions, 36 hours of volleyball were played. The injury rate was 0.36 / hour.

A breakdown of the injuries by anatomical site showed that the ankle was the most frequently injured area 31% (4 of 13), followed by the knee 23% (3 of 13), hands and wrists 15% (2 of 13), thigh 15% (2 of 13), low back 8% (1 of 13) and shoulder 8% (1 of 13) (Table 10 and Fig. 1).

A breakdown of the injuries by type showed that strains accounted for 46% (6 of 13) of injuries, sprains for 39% (5 of 13) and inflammation and overuse together for 15% (2 of 13) of the reported injuries (Table 11 and Fig. 2).

Position on the court and attempted play at the time of injury

had the highest rate of injury, followed by position (4) and position (1) (Table 12 and Fig. 3).

All injuries were related to a particular type of volleyball maneuver. Attempting to block accounted for 54% (7/13) of these injuries. Attempting to hit accounted for 23% (3/13) and defense for 15% (2/13) (Table 13 and Fig. 4).

Table (1) : Mean (\bar{X}) and Standard deviation (\pm S.D.) of age among the studied Egyptian volleyball players (EVBPs) according to sex.

| Age (y) | Sex | Males | Females |
|-------------|-----|--------|---------|
| | | | |
| (\bar{X}) | | 23.91 | 20.76 |
| \pm S.D. | | 1.72 | 2.16 |
| t | | 5.91 | |
| p | | < 0.01 | |

Table (2) : Mean \pm S.D. of age among the studied EVBPs according to type of injury.

| Age Type of injury | (\bar{X}) | \pm S.D. | Range | |
|-----------------------|------------------|------------|---------|---------|
| | | | Minimum | maximum |
| Strain | 23.16 | 2.23 | 17.00 | 27.00 |
| Sprain | 23.57 | 1.98 | 19.00 | 28.00 |
| Inflam.and overuse | 23.95 | 2.15 | 18.00 | 28.00 |
| Contusion | 23.00 | 1.41 | 21.00 | 24.00 |
| Fracture | 22.50 | 0.70 | 22.00 | 23.00 |
| Dislocation | 24.33 | 1.52 | 23.00 | 26.00 |
| F | 0.646 | | | |
| P | > 0.05 | | | |

Table (3) : Mean \pm S.D. of age among the studied EVBPs according to anatomical site of injury.

| Age (y) Anatomical site | \bar{X} | \pm S.D. | Range | |
|----------------------------|-----------|------------|---------|---------|
| | | | Minimum | Maximum |
| Ankle | 23.3 | ± 2.1 | 19.0 | 28.0 |
| Knee | 23.6 | ± 2.4 | 18.0 | 28.0 |
| Hand, wrist | 23.5 | ± 1.9 | 20.0 | 27.0 |
| Back | 23.1 | ± 2.0 | 19.0 | 26.0 |
| Shoulder | 24.3 | ± 1.9 | 20.0 | 26.0 |
| Thigh | 22.7 | ± 2.5 | 17.0 | 25.0 |
| Abd.ms. | 26.0 | ± 0.0 | 26.0 | 26.0 |
| Foot | 23.5 | ± 1.3 | 22.0 | 25.0 |
| Elbow | 23.5 | ± 2.1 | 22.0 | 25.0 |
| F | 0.551 | | | |
| P | > 0.05 | | | |

Table (4):Mean and standard deviation (\pm S.D.) of duration of participation among the studied EVBPs according to sex.

| Duration of participation (y) | Sex | |
|----------------------------------|--------|---------|
| | Males | Females |
| (\bar{X}) | 4.97 | 4.00 |
| \pm S.D. | 1.49 | 1.29 |
| t | 2.22 | |
| p | < 0.05 | |

Table (5) : Mean \pm S.D of duration of participation among the studied EVBPs according to type of injury.

| Type of Injury | Duration of participation (y) (\bar{X}) | \pm S.D. | Range | |
|---------------------|--|------------|---------|---------|
| | | | Minimum | maximum |
| Strain | 4.94 | 1.49 | 2.00 | 10.00 |
| Sprain | 5.00 | 1.65 | 3.00 | 10.00 |
| Inflam. and overuse | 4.62 | 1.46 | 2.00 | 9.00 |
| Contusion | 4.75 | 0.95 | 4.00 | 6.00 |
| Fracture | 4.50 | 0.70 | 4.00 | 5.00 |
| Dislocation | 4.33 | 2.08 | 2.00 | 6.00 |
| F | 0.277 | | | |
| P | > 0.05 | | | |

Table (6) : Mean \pm S.D of duration of participation among the studied EVBPs according to anatomical site of injury.

| Anatomical site of injury | Duration of participation | (\bar{X}) | \pm S.D. | Range | |
|---------------------------|---------------------------|-------------|------------|---------|---------|
| | | | | Minimum | maximum |
| Ankle | (n = 24) | 4.9 | ± 1.6 | 3.0 | 10.0 |
| Knee | (n = 19) | 5.0 | ± 2.0 | 2.0 | 10.0 |
| Hand, wrist | (n = 16) | 5.1 | ± 1.3 | 3.0 | 8.0 |
| Back | (n = 13) | 5.1 | ± 1.3 | 3.0 | 8.0 |
| Shoulder | (n = 10) | 4.6 | ± 1.2 | 3.0 | 6.0 |
| Thigh | (n = 8) | 4.1 | ± 1.4 | 2.0 | 6.0 |
| Foot | (n = 4) | 4.2 | ± 1.0 | 3.0 | 5.0 |
| Elbow | (n = 2) | 4.0 | ± 0.0 | 4.0 | 4.0 |
| Abd. ms. | (n = 1) | 6.0 | ± 0.0 | 6.0 | 6.0 |
| F | | 0.586 | | | |
| P | | > 0.05 | | | |

Table (7) : Side of injury among the studied EVBPs according to sex.

| Side of injury \ Sex | Males | | Females | | Total | |
|----------------------|-------|-------|---------|-------|-------|-------|
| | No. | % | No. | % | No. | % |
| Rt | 44 | 61.97 | 8 | 66.67 | 52 | 62.65 |
| Lt | 27 | 38.03 | 4 | 33.33 | 31 | 37.35 |
| Total | 71 | 100.0 | 12 | 100.0 | 83 | 100.0 |

$$X^2 = 0.097$$

$$P > 0.05$$

Table (8) : Side of injury among the studied EVBPs according to type of injury.

| Side Type of injury | Rt | | LT | | Total | |
|------------------------|-----|-------|-----|-------|-------|-------|
| | No. | % | No. | % | No. | % |
| Strain | 14 | 26.92 | 9 | 29.03 | 23 | 27.71 |
| Sprain | 14 | 26.92 | 13 | 41.94 | 27 | 32.53 |
| Inflam. and overuse | 18 | 34.62 | 6 | 19.34 | 24 | 28.92 |
| Contusion | 3 | 5.77 | 1 | 3.23 | 4 | 4.82 |
| Fracture | 2 | 3.85 | 0 | 0 | 2 | 2.41 |
| Dislocation | 1 | 1.92 | 2 | 6.46 | 3 | 3.61 |
| Total | 52 | 62.65 | 31 | 37.35 | 83 | 100.0 |

Table (9) : Side of injury among the studied EVBPs according to anatomical site of injury.

| Side anatomical site | Rt | | Lt | | Total | |
|-------------------------|-----|-------|-----|-------|-------|-------|
| | No. | % | No. | % | No. | % |
| Ankle | 13 | 25.00 | 11 | 35.48 | 24 | 28.92 |
| knee | 8 | 15.38 | 11 | 35.48 | 19 | 22.89 |
| Hand, wrist | 10 | 19.23 | 6 | 19.35 | 16 | 19.28 |
| Shoulder | 9 | 17.31 | 1 | 3.23 | 10 | 12.05 |
| Thigh | 7 | 13.46 | 1 | 3.23 | 8 | 9.64 |
| Foot | 3 | 5.77 | 1 | 3.23 | 4 | 4.82 |
| Elbow | 2 | 3.85 | 0 | 0 | 2 | 2.41 |
| Total | 52 | 62.65 | 31 | 37.35 | 83 | 100.0 |

Table (10) : Anatomical site of injury among the studied EVBPs
according to sex.

| Anatomical site | Sex | | Males | | Females | | Total | | Z | P |
|-----------------|-----|-------|-------|-------|---------|-------|-------|---|-------|--------|
| | No. | % | No. | % | No. | % | No. | % | | |
| Ankle | 20 | 23.81 | 4 | 30.77 | 24 | 24.74 | | | 0.541 | > 0.05 |
| Knee | 16 | 19.05 | 3 | 23.08 | 19 | 19.59 | | | 0.341 | > 0.05 |
| Hand, wrist | 14 | 16.67 | 2 | 15.38 | 16 | 16.49 | | | 0.116 | > 0.05 |
| Back | 12 | 14.29 | 1 | 7.69 | 13 | 13.40 | | | 0.649 | > 0.05 |
| Shoulder | 9 | 10.71 | 1 | 7.69 | 10 | 10.31 | | | 0.333 | > 0.05 |
| Thigh | 6 | 7.14 | 2 | 15.38 | 8 | 8.25 | | | 1.005 | > 0.05 |
| Foot | 4 | 4.76 | 0 | 0 | 4 | 4.12 | | | 0.804 | > 0.05 |
| Elbow | 2 | 2.38 | 0 | 0 | 2 | 2.06 | | | 0.562 | > 0.05 |
| Abd, ms. | 1 | 1.19 | 0 | 0 | 1 | 1.03 | | | 0.393 | > 0.05 |
| Total | 84 | 100.0 | 13 | 100.0 | 97 | 100.0 | | | -- | -- |




Fig.(1): Anatomical site of injury among the studied EVBPs according to sex

Table (11) : Type of injury among the studied EVBPs according to sex.

| Type of injury | Males | | Females | | Total | | Z | P |
|------------------|-------|-------|---------|-------|-------|-------|-------|-------|
| | No. | % | No. | % | No. | % | | |
| Strain | 30 | 35.72 | 6 | 46.16 | 36 | 37.12 | 0.725 | >0.05 |
| Sprain | 23 | 27.38 | 5 | 38.46 | 28 | 28.87 | 0.820 | >0.05 |
| Inflam., overuse | 22 | 26.19 | 2 | 15.38 | 24 | 24.74 | 0.840 | >0.05 |
| Contusion | 4 | 4.76 | 0 | 0 | 4 | 4.12 | 0.804 | >0.05 |
| Dislocation | 3 | 3.57 | 0 | 0 | 3 | 3.09 | 0.692 | >0.05 |
| Fracture | 2 | 2.38 | 0 | 0 | 2 | 2.06 | 0.562 | >0.05 |
| Total | 84 | 100.0 | 13 | 100.0 | 97 | 100.0 | -- | -- |




Fig.(2): Type of injury among the studied EVBPs according to sex.

Table (12) : Positions at the time of injury among the studied EVBPs according to sex.

| Sex Position | Males | | Females | | Total | | Z | P |
|-----------------|-------|-------|---------|-------|-------|-------|-------|--------|
| | No. | % | No. | % | No. | % | | |
| 1 | 4 | 5.63 | 2 | 15.38 | 6 | 7.14 | 1.255 | > 0.05 |
| 2 | 11 | 15.49 | 1 | 7.69 | 12 | 14.29 | 0.739 | > 0.05 |
| 3 | 31 | 43.66 | 6 | 46.15 | 37 | 44.05 | 0.166 | > 0.05 |
| 4 | 17 | 23.94 | 3 | 23.08 | 20 | 23.81 | 0.067 | > 0.05 |
| 5 | 3 | 4.23 | 0 | 0 | 3 | 3.57 | 0.755 | > 0.05 |
| 6 | 5 | 7.04 | 1 | 7.69 | 6 | 7.14 | 0.084 | > 0.05 |
| Total | 71 | 100.0 | 13 | 100.0 | 84 | 100.0 | -- | -- |




Fig.(3): Positions at the time of injury among the studied EVBPs according to sex.




Fig. (3b): Positions at the time of injury among the studied EVBPs according to sex.

Table (13) : Maneuvers at the time of injury among the studied EVBPs according to sex.

| Maneuver \ Sex | Males | | Females | | Total | | Z | P |
|----------------|-------|-------|---------|-------|-------|--------|-------|--------|
| | No. | % | No. | % | No. | % | | |
| Block | 41 | 54.67 | 7 | 53.85 | 48 | 54.55 | 0.055 | > 0.05 |
| Hit | 19 | 25.33 | 3 | 23.08 | 22 | 25.00 | 0.173 | > 0.05 |
| Defense | 11 | 14.67 | 2 | 15.38 | 13 | 14.77 | 0.067 | > 0.05 |
| Serve | 4 | 5.33 | 1 | 7.69 | 5 | 5.68 | 0.339 | > 0.05 |
| Total | 75 | 85.23 | 13 | 14.77 | 88 | 100.00 | -- | -- |




Fig.(4):Maneuvers at the time of injury among the studied EVBPs according to sex

Table (14): Type of injury among the studied EVBPs according to anatomical site of injury.

| Site Type | Ankle | | Knee | | Hand wrist | | Back | | Shoulder | | Thigh | | Abd. ms. | | Foot | | Elbow | | Total | |
|--------------|-------|-------|------|-------|------------|-------|------|-------|----------|-------|-------|-------|----------|--------|------|-------|-------|--------|-------|--------|
| | No | % | No | % | No | % | No | % | No | % | No | % | No | % | No | % | No | % | No | % |
| Strain | 4 | 16.67 | 4 | 21.05 | 8 | 50.00 | 13 | 100.0 | 0 | 0 | 7 | 87.50 | 0 | 0 | 0 | 0 | 0 | 0 | 36 | 37.11 |
| Sprain | 20 | 83.33 | 2 | 10.53 | 4 | 25.00 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 100.00 | 1 | 25.00 | 0 | 0 | 28 | 28.87 |
| Inflam. | 0 | 0 | 11 | 57.89 | 0 | 0 | 0 | 0 | 9 | 90.00 | 0 | 0 | 0 | 0 | 2 | 50.00 | 2 | 100.00 | 24 | 24.74 |
| Overuse | | | | | | | | | | | | | | | | | | | | |
| Contusion | 0 | 0 | 1 | 5.26 | 2 | 12.50 | 0 | 0 | 0 | 0 | 1 | 12.50 | 0 | 0 | 0 | 0 | 0 | 0 | 4 | 4.12 |
| Fracture | 0 | 0 | 0 | 0 | 1 | 6.25 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 25.00 | 0 | 0 | 2 | 2.06 |
| Dislocation | 0 | 0 | 1 | 5.26 | 1 | 6.25 | 0 | 0 | 1 | 10.00 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 3 | 3.09 |
| Total | 24 | 24.74 | 19 | 19.59 | 16 | 16.49 | 13 | 13.40 | 10 | 10.31 | 8 | 8.25 | 1 | 1.03 | 4 | 4.12 | 2 | 2.06 | 97 | 100.00 |

Table (15): Type of injury among the studied EVBPs according to positions.

| Position Type of injury | 1 | | 2 | | 3 | | 4 | | 5 | | 6 | | Total | |
|----------------------------|----|-------|----|-------|----|-------|----|-------|----|-------|----|-------|-------|-------|
| | No | % | No | % | No | % | No | % | No | % | No | % | No | % |
| Strain | 1 | 16.67 | 2 | 16.67 | 13 | 35.14 | 6 | 30.00 | 2 | 66.67 | 1 | 16.67 | 25 | 29.76 |
| Sprain | 2 | 33.33 | 4 | 33.33 | 13 | 35.14 | 8 | 40.00 | 0 | 0 | 1 | 16.67 | 28 | 33.33 |
| Inflam | | | | | | | | | | | | | | |
| Overuse | 3 | 56.00 | 6 | 50.00 | 7 | 18.92 | 5 | 25.00 | 0 | 0 | 1 | 16.67 | 22 | 26.19 |
| Contusion | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 5.00 | 1 | 33.33 | 2 | 33.33 | 4 | 4.76 |
| Fracture | 0 | 0 | 0 | 0 | 2 | 5.41 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 2.38 |
| Dislocation | 0 | 0 | 0 | 0 | 2 | 5.41 | 0 | 0 | 0 | 0 | 1 | 16.67 | 3 | 3.57 |
| Total | 6 | 7.14 | 12 | 14.29 | 37 | 44.05 | 20 | 23.81 | 3 | 3.57 | 6 | 7.14 | 84 | 100 |

Table (16): Type of injury among the studied EVBPs according to maneuvers.

| Maneuver Type of injury | Block | | Hit | | Defense | | Serve | | Total | |
|----------------------------|-------|-------|-----|-------|---------|-------|-------|-------|-------|--------|
| | No | % | No | % | No | % | No | % | No | % |
| Strain | 17 | 35.42 | 5 | 22.73 | 6 | 46.15 | 0 | 0 | 28 | 31.82 |
| Sprain | 19 | 39.58 | 6 | 27.27 | 1 | 7.69 | 2 | 40.00 | 28 | 31.82 |
| Inflam., overuse | 9 | 18.75 | 9 | 40.91 | 2 | 15.38 | 3 | 60.00 | 23 | 26.14 |
| Contusion | 1 | 2.08 | 0 | 0 | 3 | 23.08 | 0 | 0 | 4 | 4.55 |
| Fracture | 1 | 2.08 | 1 | 4.55 | 0 | 0 | 0 | 0 | 2 | 2.27 |
| Dislocation | 1 | 2.08 | 1 | 4.55 | 1 | 7.69 | 0 | 0 | 3 | 3.41 |
| Total | 48 | 54.55 | 22 | 25.00 | 13 | 14.77 | 5 | 5.68 | 88 | 100.00 |

Table (18) : Anatomical site of injury among the studied EVBPs according to maneuvers.

| Anatomical site | Block | | Hit | | Defense | | Serve | | Total | |
|-----------------|-------|-------|-----|-------|---------|-------|-------|------|-------|-------|
| | No. | % | No. | % | No. | % | No. | % | No. | % |
| Ankle | 16 | 33.33 | 5 | 22.73 | 2 | 15.38 | 0 | 0 | 23 | 26.14 |
| Knee | 9 | 18.75 | 3 | 13.64 | 5 | 38.46 | 1 | 20.0 | 18 | 20.45 |
| Hand, wrist | 9 | 18.75 | 2 | 9.09 | 3 | 23.08 | 0 | 0 | 14 | 15.91 |
| Back | 8 | 16.67 | 2 | 9.09 | 1 | 7.69 | 0 | 0 | 11 | 12.50 |
| Shoulder | 1 | 2.08 | 7 | 31.82 | 0 | 0 | 1 | 20.0 | 9 | 10.23 |
| Thigh | 3 | 6.25 | 1 | 4.55 | 2 | 15.38 | 0 | 0 | 6 | 6.82 |
| Foot | 1 | 2.08 | 2 | 9.09 | 0 | 0 | 1 | 20.0 | 4 | 4.55 |
| Elbow | 1 | 2.08 | 0 | 0 | 0 | 0 | 1 | 20.0 | 2 | 2.27 |
| Abd.ms | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 20.0 | 1 | 1.14 |
| Total | 48 | 54.55 | 22 | 25.00 | 13 | 14.77 | 5 | 5.68 | 88 | 100.0 |

Table (17) : Anatomical site of injury among the studied EVBPs according to positions.

| Position Site of injury | 1 | | 2 | | 3 | | 4 | | 5 | | 6 | | Total | |
|----------------------------|----|-------|----|-------|----|-------|----|-------|----|-------|----|-------|-------|-------|
| | No | % | No | % | No | % | No | % | No | % | No | % | No | % |
| Ankle | 0 | 0 | 4 | 33.33 | 10 | 27.03 | 7 | 35.00 | 0 | 0 | 1 | 16.67 | 22 | 26.19 |
| Knee | 1 | 16.67 | 4 | 33.33 | 6 | 16.22 | 2 | 10.0 | 2 | 66.67 | 2 | 33.33 | 17 | 20.24 |
| Hand, wrist | 0 | 0 | 1 | 8.33 | 7 | 18.92 | 3 | 15.0 | 1 | 33.33 | 2 | 33.33 | 14 | 16.67 |
| Back | 1 | 16.67 | 1 | 8.33 | 4 | 10.81 | 4 | 20.0 | 0 | 0 | 0 | 0 | 10 | 11.90 |
| Shoulder | 1 | 16.67 | 2 | 16.67 | 4 | 10.81 | 2 | 10.0 | 0 | 0 | 0 | 0 | 9 | 10.71 |
| Thigh | 0 | 0 | 0 | 0 | 3 | 8.11 | 1 | 5.0 | 0 | 0 | 1 | 16.67 | 5 | 5.95 |
| Abd. ms. | 1 | 16.67 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1.19 |
| Foot | 1 | 16.67 | 0 | 0 | 2 | 5.41 | 1 | 5.0 | 0 | 0 | 0 | 0 | 4 | 4.76 |
| Elbow | 1 | 16.67 | 0 | 0 | 1 | 2.70 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 2.38 |
| Total | 6 | 7.14 | 12 | 14.29 | 37 | 44.05 | 20 | 23.81 | 3 | 3.57 | 6 | 7.14 | 84 | 100.0 |

Table (19): Maneuver at the time of injury among the studied EVBPs according to positions.

| Position | 1 | | 2 | | 3 | | 4 | | 5 | | 6 | | Total | |
|-----------|-----|-------|-----|-------|-----|-------|-----|-------|-----|-------|-----|-------|-------|-------|
| | No. | % | No. | % | No. | % | No. | % | No. | % | No. | % | No. | % |
| Maneuvers | | | | | | | | | | | | | | |
| Block | 0 | 0 | 8 | 66.67 | 24 | 64.86 | 15 | 75.0 | 0 | 0 | 0 | 0 | 47 | 55.95 |
| Hit | 0 | 0 | 4 | 33.33 | 13 | 35.14 | 5 | 75.0 | 0 | 0 | 0 | 0 | 22 | 26.19 |
| Defense | 1 | 16.67 | 0 | 0 | 0 | 0 | 0 | 0 | 3 | 100.0 | 6 | 100.0 | 10 | 11.90 |
| Serve | 5 | 83.33 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 5 | 5.95 |
| Total | 6 | 7.14 | 12 | 14.29 | 37 | 44.05 | 20 | 23.81 | 3 | 3.57 | 6 | 7.14 | 84 | 100.0 |