

REFERENCES

- Abdel Hadi M, Derballa S, El Halafawy A, Abdel Hamid E. Sonographic assessemnt of cesarean section scar; evaluation of new approach. J Egy, Soc. Obstet Gynecol,1985; 67:314.
- Acher, DB, Hoss S, O'Brien E, Donabye CL, Prorfell MM, Sacks BP. Cesarean birth rate; small geographic area analysis. Am J Obstet Gynecol;1999, 159:386-388
- ACOG: Induction of Labor for Vaginal Birth After Cesarean Delivery. Committee Opinion No. 342: Obstet Gynecol 2006; 108:465.
- ACOG: Dystoci, bulletin 137, Obestet Gynacol, DC, ACOG, 1990;105:452.
- ACOG : Vaginal birth after previous cesarean. Practice Bulletin #54: Obstet Gynecol 2004; 104:203.
- ACOG: Management of Postterm Pregnancy. Practice Bulletin #55. Obstet Gynecol 2004; 104:639.
- ACOG : Vaginal birth after previous cesaern birth . Practice patterns No. 1; Obestet Gynacol, DC:ACOG, 1995;142:352.
- ACOG. Induction of labor with misoprostol. ACOG Committee Opinion #228, Obestet Gynacol, 1999 ;104:235.
- ACOG. Vaginal birth after previous Cesarean delivery. ACOG Practice Bulletin No. 5. Obestet Gynacol: ACOG; 1999;102:322.

- .Agency for Healthcare Research and Quality .AHRQ .Publication No. 03-E018. March 2003, Rockville,MD.
- Al Sakka, M, Dauleh, W, Al Hassani, S. Case series of uterine rupture and subsequent pregnancy outcome. *Int J Fertil Womens Med* 1999; 44:297.
- Ananth, CV, Smulian, JC, Vintzileos, AM. The association of placenta previa with history of cesarean delivery and abortion: a metaanalysis. *Am J Obstet Gynecol* 1997; 177:1071.
- Asukura H, Myers SA. More than one previous cesarean delivery :A 5-year experience with 435 patients. *Obstet Gynecol*;1995,85: 924.
- Asakura, H, Nakai, A, Ishikawa, G, . Prediction of uterine dehiscence by measuring lower uterine segment thickness prior to the onset of labor: evaluation by transvaginal ultrasonography. *J Nippon Med Sch* 2000; 67:352.
- Bedoya C, Bartha JL, Rodriguez I, Fontan I, Bedoya JM, Sanchez – Ramos J. Atrial of labour after cesarean section in patients with or without a previous vaginal delivery. *Int. J Obstet Gynecol*;1992, 39:285-289.
- Bond, A.L.; Edersheim, T.G.; Curry, L.; Druzin, M.L. and Huston, J.M.: Expectant management of abruption placentae before 35 weeks gestation am. *J. Perinatol.*;1989, 6:121.
- Brown L, Karrison T, Cibils LA. Mode of delivery and perinatal results in breech presentation.A m J Obestet Gynaecol;1995, 171 : 28-34.

- Biswass A. Management of previous Cesarean section. *Curr Opin Obstet Gynecol* 2003;15:123–9.
- Bujold E, Gauthier R. Should we allow a trial of labor after a previous Cesarean for dystocia in the second stage of labor? *Obstet Gynecol* 2001;98:652–5.
- Bujold, E, Blackwell, SC, Gauthier, RJ. Cervical ripening with transcervical foley catheter and the risk of uterine rupture. *Obstet Gynecol* 2004; 103:18.
- Bujold, E, Bujold, C, Hamilton, EF, et al. The impact of a single-layer or double-layer closure on uterine rupture. *Am J Obstet Gynecol* 2002; 186:1326.
- Bujold, E, Hammoud, AO, Hendler, I, .Trial of labor in patients with a previous cesarean section: Does maternal age influence the outcome?. *Am J Obstet Gynecol* 2004; 190:1113.
- Bujold, E, Mehta, SH, Bujold, C, Gauthier, RJ. Interdelivery interval and uterine rupture. *Am J Obstet Gynecol* 2002; 187:1199.
- Cahill, AG, Stamilio, DM, Odibo, AO, et al. Does a maximum dose of oxytocin affect risk for uterine rupture in candidates for vaginal birth after cesarean delivery?. *Am J Obstet Gynecol* 2007; 197:495.
- Cahill, AG, Stamilio, DM, Odibo, AO, et al. Is vaginal birth after cesarean (VBAC) or elective repeat cesarean safer in women with a prior vaginal delivery?. *Am J Obstet Gynecol* 2006; 195:1143.

- Cahill, AG, Waterman, BM, Stamilio, DM, et al. Higher maximum doses of oxytocin are associated with an unacceptably high risk for uterine rupture in patients attempting vaginal birth after cesarean delivery. *Am J Obstet Gynecol* 2008; 199:32.
- Callen P.W. and Filly, R.A.: The placental-subplacental complex: A specific indicator of placental position on ultrasound. *J. Clin. Ultrasound*;1980, 8:21.
- Carroll CS, Sr, Magann, EF, Chauhan, SP, et al. Vaginal birth after cesarean section versus elective repeat cesarean delivery: Weight-based outcomes. *Am J Obstet Gynecol* 2003; 188:1516.
- Caughey, AB, Shipp, TD, Repke, JT, et al. Rate of uterine rupture during a trial of labor in women with one or two prior cesarean deliveries. *Am J Obstet Gynecol* 1999; 181:872.
- Chamberlain G., Steer P., : Cesarean Section ,in Turnbolls Obstetrics third edition .Harcourt Publishers,p;2001,601 -618 .
- Chauhan, SP, Martin, JN Jr, Henrichs, CE, et al. Maternal and perinatal complications with uterine rupture in 142,075 patients who attempted vaginal birth after cesarean delivery: A review of the literature. *Am J Obstet Gynecol* 2003; 189:408.
- Chauhan SP, Martin Jr Jn, Henrichs CE, Morrison JC, Magann EF. Maternal and perinatal complications with uterine rupture in 142, 075 patients who attempted vaginal birth after cesarean delivery : a review of the literature. *Am J obstet Gynecol*;2007, 189 : 408 - 417.
- Chaung VY: Sonographic measurements of the lower uterine

segment thickness in women with previous caesarean section. *J. Obstet Gynaecol Can.* Jul; 2005,27 (7): 674-81.

- Clark SL., Koonings PP, Phelan JP. Placenta previa accreta and prior cesarean section. *Obstet Gynecol*; 1995,66: '89 -92.
- Coassolo, KM, Stamilio, DM, Pare, E, et al. Safety and Efficacy of Vaginal Birth After Cesarean Attempts at or Beyond 40 Weeks of Gestation. *Obstet Gynecol* 2005; 106:700.
- Committee opinion. Induction of labor for vaginal birth after cesarean delivery. *Obstet Gynecol* 2002; 99:679.
- Constantinescu OC, and Ahluwalia BS : Sonographic evaluation of the lower uterine segment in patient with previous cesarean delivery. *J. Ultrasound . Med.* Nov;2004,23 (11):1441-7.
- Cunningham FG, Larry CG, Norman FG' Kenneth JL. Cesarean delivery and Postpartum, hysterectomy. In *Williams Obstetrics* :21th ed . Appleton and Lange , California;2001, page 538-545.
- De Costa, C. Vaginal birth after classical Caesarean section. *Aust N Z J Obstet Gynaecol* 2005; 45:182.
- De Mcndonca, L.K : Sonographic diagnosis of placenta accrete, presentation of six cases. *J.Ultrasound Med*;1988 , 7-211.
- Dodd, J, Crowther, C. Induction of labour for women with a previous Cesarean birth: a systematic review of the literature. *Aust N Z J Obstet Gynaecol* 2004; 44:392.
- Duff P. Antibiotic prophylaxis for abdominal cesarean section. A review of the American literature *sex, Med. Today*;1985, 9: 18 .

- Dunn EA, O'Herlihy C. Comparison of maternal satisfaction following vaginal delivery after caesarean section and caesarean section after previous vaginal delivery. *Eur J Obstet Gynecol Reprod Biol* 2005;121:56–60.
- Durnwald, C, Mercer, B. Uterine rupture, perioperative and perinatal morbidity after single-layer and double-layer closure at cesarean delivery. *Am J Obstet Gynecol* 2003; 189:925.
- Durnwald, CP, Rouse, DJ, Leveno, KJ, et al. The Maternal-Fetal Medicine Units Cesarean Registry: safety and efficacy of a trial of labor in preterm pregnancy after a prior cesarean delivery. *Am J Obstet Gynecol* 2006; 195:1119.
- Ekpo, EE. Uterine rupture as seen in the University of Calabar Teaching Hospital, Nigeria: a five-year review. *J Obstet Gynaecol* 2000; 20:154.
- Elkousy, MA, Sammel, M, Stevens, E, Peipert, JF. The effect of birth weight on vaginal birth after cesarean delivery success rates. *Am J Obstet Gynecol* 2003; 188:824.
- El-Sayed, YY, Watkins, MM, Fix, M, et al. Perinatal outcomes after successful and failed trials of labor after cesarean delivery. *Am J Obstet Gynecol* 2007; 196:583.
- Esposito, MA, Menihan, CA, Malee, MP. Association of interpregnancy interval with uterine scar failure in labor: a case-control study. *Am J Obstet Gynecol* 2000; 183:1180.

- Farmer, RM, Kirschbaum, T, Potter, D, et al. Uterine rupture during trial of labor after previous cesarean section. *Am J Obstet Gynecol* 1991; 165:996.
- Flamm, BL, Goings, JR, Fuelberth, NJ, et al. Oxytocin during labor after previous cesarean section: results of a multicenter study. *Obstet Gynecol* 1987; 70:709.
- Flamm BL, Newman LA, Thomas SJ. Vaginal birth after cesarean delivery: Results of a 5 – years multi center Collaborative study. *Obstet Gynecol*;1990,76:750.
- Frank F: Suprasymphysial delivery and its relation to other operation in the presence of contracted pelvis. *Arch Gynaecol*;1907,81:46.
- Fukuda M, Shimizu TY, NatsuymamaE, Mochizuki M, Hara. Ultrasound examination of cesarean section scar during pregnancy. *Asch. Gynecol Obstet*;1991, 248 129.
- Geerts LT,Brand EJ,Theron Gb. Routine obstetric ultrasound examinations in pregnancy .A Prospective randomized controlled study .*Br J Obstet Gynecol*;1996,103:501-507.
- Gillbert L, Sounders N, Sharp F: The management of multiplel Pregnancy in women with a lower segment cesarean scar . Is repeated section really the safe option *Br. J Obstet Gynaecol*;1988, 95: 1312-1316.
- Gilliam M, Rosenberg D, Davis F. The likelihood of placenta previa with greater number of cesarean deliveries and higher parity. *Obstet Gynecol* 2002;99:976–80.

- Goetzl, L, Shipp, TD, Cohen, A, et al. Oxytocin dose and the risk of uterine rupture in trial of labor after cesarean. *Obstet Gynecol* 2001; 97:381.
- Gotoh, H, Masuzaki, H, Yoshida, A, et al. Predicting incomplete uterine rupture with vaginal sonography during the late second trimester in women with prior cesarean. *Obstet Gynecol* 2000; 95:596.
- Greene, MF. Vaginal birth after cesarean section. *N Engl J Med* 2004; 351:2647.
- Greene R, Gardeil F, Turner Mj: Long term implications of cesarean delivery. Letter of the Editors. *Am J : Gynecol*; 1997, 176 : 254.
- Greenhill J.P. and Friedman: Effects of pregnancy on the maternal organisms. *Biological principles and modern practice of Obstetrics*. 1st ed. Philadelphia: Saunders;1974,105.
- Grobman, WA, Gilbert, S, Landon, MB, et al. Outcomes of induction of labor after one prior cesarean. *Obstet Gynecol* 2007; 109:262.
- Grobman, WA, Lai, Y, Landon, MB, et al. Development of a nomogram for prediction of vaginal birth after cesarean delivery. *Obstet Gynecol* 2007; 109:806.
- Grobman, WA, Lai, Y, Landon, MB, et al. Prediction of uterine rupture associated with attempted vaginal birth after cesarean delivery. *Am J Obstet Gynecol* 2008; 199:30.

- Guise, JM, McDonagh, MS, Osterweil, P, et al. Systematic review of the incidence and consequences of uterine rupture in women with previous caesarean section. *BMJ* 2004; 329:19.
- Halperin, ME, Moore, DC, Hannah, WJ. Classical versus low-segment transverse incision for preterm caesarean section: maternal complications and outcome of subsequent pregnancies. *Br J Obstet Gynaecol* 1988; 95:990.
- Harold Gee.' Trial of Scar: Yearbook ed, Pag. 263 - 2,2001. Harrington LC, Miller DA, McClain CJ, Paul RH. Vaginal birth after cesarean in a hospital-based birth center staffed by certified nurse midwives. *J Nurse Midwifery*;1997, 42: 304 – 307.
- Hamilton, EF, Bujold, E, McNamara, H, et al. Dystocia among women with symptomatic uterine rupture. *Am J Obstet Gynecol* 2001; 184:620.
- Hashima, JN, Eden, KB, Osterweil, P, et al. Predicting vaginal birth after cesarean delivery: A review of prognostic factors and screening tools. *Am J Obstet Gynecol* 2004; 190:547.
- Hendrix SL, Schimp V, Martin J, Singh A, Kruger, M, McNeeley SO: The legendary superior strength of the Pfannenstiel incision: A myth? *Am J Obstet Gynecol*;2000, 182: 1446.
- Hibbard JU, Ismail M, Wang Y, Te C, Karrison T, Ismail M. Failed vaginal birth after a cesarean section: How risky is it? I. Maternal morbidity. *Am J Obstet Gynecol* 2001;184:1365–73.

- Hoffman, MK, Sciscione, A, Srinivasana, M, et al. Uterine rupture in patients with a prior cesarean delivery: the impact of cervical ripening. *Am J Perinatol* 2004; 21:217.
- Hofmeyr, GJ, Say, L, Gulmezoglu, AM. WHO systematic review of maternal mortality and morbidity: the prevalence of uterine rupture. *Br J Obstet . Gynaecol.* 2005; 112:1221.
- Hofmeyr, GJ. Obstructed labor: using better technologies to reduce mortality. *Int J Gynaecol Obstet* 2004; 85 Suppl 1:S62.
- Hook B, Kiwi R, Amini SB, Fanaroff A, Hack M. Neonatal morbidity after elective repeat cesarean section and trial of labor. *Pediatrics* 1997;100:348–53.
- Hoskins, IA, Gomez, JL. Correlation between maximum cervical dilatation at cesarean delivery and subsequent vaginal birth after cesarean delivery. *Obstet Gynecol* 1997; 89:591.
- Juntunen, K, Makarainen, L, Kirkinen, P. Outcome after a high number (4-10) of repeated caesarean sections. *Br J Obstet. Gynaecol.* 2004; 111:561.
- Kayani, SI, Alfirevic, Z. Induction of labour with previous caesarean delivery: where do we stand?. *Curr Opin Obstet Gynecol* 2006; 18:636.
- Kerr JMM: The technic of cesarean section with special reference to the lower uterine segment incision. *Am J Obstet Gynecol*;1926, 2:729.
- Krishnamurthy S, Fairlie F, Cameron AD, Walker JJ, Mackenzie

- JR. The role of post natal x-ray pelvimetry after cesarean section in the management of subsequent delivery. *Br J. Obstet Gynecol*;1991, 98:716-718.
- Kwee A, Bots ML, Visser GHA, Bruinse WH. Emergency peripartum hysterectomy: a prospective study in the Netherlands. *Eur J Obstet Gynecol Reprod Biol* 2006;124:187–92.
 - Laing, F.C.: Ultrasound evaluation of the Obstetric problems relating to the lower uterine segment and cervix . The principles and practice of Ultrasoundography in Obstetric and Gynecology. 4th ed. Appleton and Lang; U.S.A., (1991);36:155-158.
 - Landon MB, Hauth JC, Leveno KJ, Spong CY, Leindecker S, Varner MW, *et al.* Maternal and perinatal outcomes associated with a trial of labor after prior cesarean delivery. *N Engl J Med* 2004;351:2581–9.
 - Landon, MB, Hauth, JC, Leveno, KJ, *et al.* Maternal and perinatal outcomes associated with a trial of labor after prior cesarean delivery. *N Engl J Med* 2004; 351:2581.
 - Landon, MB, Leindecker, S, Spong, CY, *et al.* The MFMU Cesarean Registry: factors affecting the success of trial of labor after previous cesarean delivery. *Am J Obstet Gynecol* 2005; 193:1016.
 - Landon, MB, Spong, CY, Thom, E, *et al.* Risk of uterine rupture with a trial of labor in women with multiple and single prior cesarean delivery. *Obstet Gynecol* 2006; 108:12.
 - Latzko W: Ueber den extraperitoealen Kaiserschnitt. *Zentralbl*

Gynaecol;1909,33:275.

- Lavin JP, Stephan RJ, Miodpvik M, Barden TP. Vaginal delivery in patients with a prior cesarean section. *Obstet Gynecol*; 1982 ,59(2): 135.
- Leung, AS, Leung, EK, Paul, RH. Uterine rupture after previous cesarean delivery: maternal and fetal consequences. *Am J Obstet Gynecol* 1993; 169:945.
- Leveno KJ: Controversies in OB-Gyn: Should we rethink the cirteria for VBAC? *Contemporary OB-Gyn*; January, 1999.
- Lieberman, E, Ernst, EK, Rooks, JP, et al. Results of the national study of vaginal birth after cesarean in birth centers. *Obstet Gynecol* 2004; 104:933.
- Lydon-Rochelle, M, Holt, VL, Easterling, TR, Martin, DP. Risk of uterine rupture during labor among women with a prior cesarean delivery. *N Engl J Med* 2001; 345:3.
- Macones GA, Hausman N, Edelstein R, Stamilio DM, Marder SJ. Predicting outcomes of trials of labor in women attempting vaginal birth after cesarean delivery: a comparison of multivariate methods with neural networks. *Am J Obstet Gynecol* 2001;184:409–13. doi:10.1067/mob.2001.109386
- Macones, GA, Cahill, A, Pare, E, et al. Obstetric outcomes in women with two prior cesarean deliveries: is vaginal birth after cesarean delivery a viable option?. *Am J Obstet Gynecol* 2005; 192:1223.

- Macones, GA, Cahill, AG, Stamilio, DM, et al. Can uterine rupture in patients attempting vaginal birth after cesarean delivery be predicted?. *Am J Obstet Gynecol* 2006; 195:1148.
- Macones, GA, Peipert, J, Nelson, DB, et al. Maternal complications with vaginal birth after cesarean delivery: a multicenter study. *Am J Obstet Gynecol* 2005; 193:1656.
- Makoha, FW, Felimban, HM, Fathuddien, MA, et al. Multiple cesarean section morbidity. *Int J Gynaecol Obstet* 2004; 87:227.
- Mankuta, DD, Leshno, MM, Menasche, MM, Brezis, MM. Vaginal birth after cesarean section: trial of labor or repeat cesarean section? A decision analysis. *Am J Obstet Gynecol* 2003; 189:714.
- Martin JN , Harris BA, Huddleston JF, Morisson JC, Prospst MG, Davidson JT. Vaginal delivery following pervios cesarean birth. *Am J Obstet Gynecol*;1996, 146(3): 255.
- Martin JN, Perry KG, Roberts WE, Meydrech EF. The care for trail of labour in the paients with a prior low- segment vertical cesarean incision . *Am J Obstet Gynecol*;1997, 177:144.
- McCarth S.Magnetic resonance imaging in obstetric and gynecology.*Magnetic Resonance Imaging* 1996;4 :59.
- McDonagh, MS, Osterweil, P, Guise, JM. The benefits and risks of inducing labour in patients with prior caesarean delivery: a systematic review. *Br J Obestet. Gynaecol.* 2005; 112:1007.

- McMahon, MJ, Luther, ER, Bowes, WA Jr, Olshan, AF. Comparison of a trial of labor with an elective second cesarean section. *N Engl J Med* 1996; 335:689.
- Mercer, BM, Gilbert, S, Landon, MB, et al. Labor Outcomes With Increasing Number of Prior Vaginal Births After Cesarean Delivery. *Obstet Gynecol* 2008; 111:285.
- Michaels WH, Thompson HO, Boutt A, Schreiber FR, Michaels SLR, Karo J. Ultrasound diagnosis of defects in the scarred lower-uterine segment during pregnancy. *Obstet Gynecol*:1988, 71;112.
- Miller DA, Chollet JA, Goodwin TM. Clinical risk factors for placenta previa-placenta accreta. *Am J Obstet Gynecol* 1997;177:210–4.
- Miller, DA, Goodwin, TM, Gherman, RB, Paul, RH. Intrapartum rupture of the unscarred uterus. *Obstet Gynecol* 1997; 89:671.
- Montgomery, AA, Emmett, CL, Fahey, T, et al. Two decision aids for mode of delivery among women with previous cesarean section: randomised controlled trial. *BMJ* 2007; 334:1305.
- Morales, KJ, Gordon, MC, Bates, GW Jr. Postcesarean delivery adhesions associated with delayed delivery of infant. *Am J Obstet Gynecol* 2007; 196:461.
- Mozurkewich, EL, Hutton, EK. Elective repeat cesarean delivery versus trial of labor: a meta-analysis of the literature from 1989 to 1999. *Am J Obstet Gynecol* 2000; 183:1187.

- Nielsen, TF, Hagberg, H, Ljungblad, U. Placenta previa and antepartum hemorrhage after previous cesarean section. *Gynecol Obstet Invest* 1989; 27:88.
- Nielsen TF, Holkegard KA. Post operative cesarean section morbidity . A postoperative. *Am J Obstet Gynecol*;1994,146:911.
- Nisenblat, V, Barak, S, Griness, OB, et al. Maternal complications associated with multiple cesarean deliveries. *Obstet Gynecol* 2006; 108:21.
- O'Connor, RA, Gaughan, B. Pregnancy following simple repair of the ruptured gravid uterus. *Br J Obstet Gynaecol* 1989; 96:942.
- Oyelese, Y, Ikomi, A, Manyonda, IT. Third consecutive complete posterior uterine rupture. A case report. *J Reprod Med* 2001; 46:694.
- Pare E, Quinones JN, Macones GA. Vaginal birth after caesarean section *versus* elective repeat caesarean section: assessment of maternal downstream health outcomes. *Br J Obstet Gynaecol* 2006;113:75–85.
- Pasto,M.E.; Kurtz,A.B.and Rilkin,M.D.: Ultrasonographic findings in placenta increta. *J.Ultrasound Med*;1983, 2:155.
- Peaceman, AM, Gersnoviez, R, Landon, MB, et al. Cesarean Registry: impact of fetal size on trial of labor success for patients with previous cesarean for dystocia. *Am J Obstet Gynecol* 2006; 195:1127.

- Petrikovsky BM. Endoscopic assessment of the integrity of the post cesarean uterine wall before a trial of labour .Transcervical Endocopy Registry. J Reprod Med; 1994 ,39:464-466.
- Phelan JP, Enlinton GS, Horenstein JM. Previous cesarean birth: Trial of labor in women with macrosomic infants. J Reprod Med; 1984,29:36-40.
- Phelan JP. Uterine rupture. Clin. Obstet Gynecol; 1990,33: 432.
- Pierrat, V, Haouari, N, Liska, A, et al. Prevalence, causes, and outcome at 2 years of age of newborn encephalopathy: population based study. Arch Dis Child Fetal Neonatal Ed 2005; 90:F257.
- Poma P.A, : Effect of departmental policies on cesarean section rates. Obstet Gynecol 1998, 91: 1013-18.
- Pruett K M, Kinshan B, Cotton D. Unknown uterine scar and trial of labor. Am J Obstet Gynecol; 1988,159:807.
- Quilligan EJ. Vaginal birth after Cesarean section: 270 degrees. J Obstet Gynecol Res 2001;27:169–73.
- Rageth, JC, Juzi, C, Grossenbacher, H. Delivery after previous cesarean: a risk evaluation. Swiss Working Group of Obstetric and Gynecologic Institutions. Obstet Gynecol 1999; 93:332.
- Rashid, M, Rashid, RS. Higher order repeat caesarean sections: how safe are five or more?. *Br J Obstetric Gynaecol* 2004; 111:1090.
- Rezenberg P, Goffinet F, Philippe HJ; N isand I. Ultrasonographic measurement of lower uterine segment to assess

risk of defects of scarred uterus . Lancet;1969, 347:281- 284.

- Ridgeway, JJ, Weyrich, DL, Benedetti, TJ. Fetal heart rate changes associated with uterine rupture. Obstet Gynecol 2004; 103:506.
- Ritchie, EH. Pregnancy following simple repair of the ruptured gravid uterus. Br J Obstet Gynaecol 1989; 96:1361.
- Roberts LJ, Beardsworth SA, Trew G. Labour following Caesarean section: current practice in the United Kingdom. Br J Obstet Gynaecol 1994;101:153–5.
- Rodriguez MH, Masaki DI, Phelan JP, Diaz FG. Uterine rupture: Are intrauterine pressure catheters usefal in diagnosis? Am J Obstet Gynecol;1994, 161: 666-669.
- Rosemond, R.L. and Kepple, D.M.: Transvaginal color dopplcr sonography in the prenatal diagnosis of placenta previa. Obstet. Gynecol.;1992, 80:508.
- Rosen MG, Dickinson JC, Westhoff CL. Vaginal birth after cesarean : ameta-analysis of morbidity and mortality. Obstet Gynecol;1991,77:465.
- Rossi, AC, D'Addario, V. Maternal morbidity following a trial of labor after cesarean section vs elective repeat cesarean delivery: a systematic review with metaanalysis. Am J Obstet Gynecol 2008; 199:224.
- Rowzberg P, Goffinet, F, Phillippe HJ, Nisand I : Ultrasonographic measurements of lower uterine segment to assess risk of defects of scarred uterus. Lancet Feb 3;2004, 347

(8997):281- 4.

- Royal College of Obstetricians and Gynaecologists Induction of labour. Evidence Based Clinical Guideline No. 9, Clinical Effectiveness Support Unit, RCOG, London, Royal College of Obstetricians and Gynaecologists 2001.
- Rozenberg, P, Goffinet, F, Phillippe, HJ, Nisand, I. Ultrasonographic measurement of lower uterine segment to assess risk of defects of scarred uterus. Lancet 1996; 347:281.
- Sachs BP , Kobelin C, Castro MA. Frigoletto F: The risks of lowering the cesarean – delivery rate . N E ngl J Med;1999, 340: 54, .
- Sambaziotis H, Conway C, Figueroa R, Elimian A Carry D,): Second – trimester songographic comparision of the lower uterine segment in pregnant women with and without a previous cesarean delivery. J. Ultrasound med;2007, 23 (7): 907-11 ; 913-4.
- Sajjad, Y, Sharma, SD, Thomas, K. Three consecutive uterine ruptures in the same woman, each with fetal survival. Br J Obstetric Gynaecol ;2005; 112:1005.
- Sanchez-Ramos, L, Gaudier, FL, Kaunitz, AM. Cervical ripening and labor induction after previous cesarean delivery. Clin Obstet Gynecol 2000; 43:513.
- Schwartz H, Gajewski C. Vaginal delivery following cesarean section. Am J Obstet Gynecol;1983, 61: 1232.
- Sen S, Malik S, Salhan S. Ultrasonographic evaluation of lower

uterine segment thickness in patients of previous cesarean

section. *Int J Gynaecol Obstet*;2004 ,87(3):215-9

- Sewell JE; Cesarean section—a brief history. Washington, DC. American College of Obstetricians and Gynecologists. 1993.
- Shipp, TD, Zelop, C, Cohen, A, et al. Post-cesarean delivery fever and uterine rupture in a subsequent trial of labor. *Obstet Gynecol* 2003; 101:136.
- Shipp TD, Zelop CM, Repke JT, Cohen A, Caughey AB, Lieberman E: Intrapartum uterine rupture and dehiscence in patients with prior lower uterine segment vertical and transverse incisions. *Obstet Gynecol*;1999 ,94: 735.
- Shipp, TD, Zelop, C, Repke, JT, et al. The association of maternal age and symptomatic uterine rupture during a trial of labor after prior cesarean delivery. *Obstet Gynecol* 2002; 99:585.
- Shipp, TD, Zelop, CM, Repke, JT, Cohen, A. Interdelivery interval and risk of symptomatic uterine rupture. *Obstet Gynecol* 2001; 97:175.
- Silver, RM, Landon, MB, Rouse, DJ, et al. Maternal Morbidity Associated With Multiple Repeat Cesarean Deliveries. *Obstet Gynecol* 2006; 107:1226.
- Sims, EJ, Newman, RB, Hulsey, TC. Vaginal birth after cesarean: to induce or not to induce. *Am J Obstet Gynecol* 2001; 184:1122.

- Smith GC, Pell, JP, Cameron, AD, Dobbie, R. Risk of perinatal death associated with labor after previous cesarean delivery in uncomplicated term pregnancies. JAMA 2002; 287:2684.
- Smith, GC, Pell, JP, Pasupathy, D, Dobbie, R. Factors predisposing to perinatal death related to uterine rupture during attempted vaginal birth after caesarean section: retrospective cohort study. BMJ 2004; 329:375.
- Socol ML, Peaceman AM. Vaginal birth after Cesarean: an appraisal of fetal risk. Obstet Gynecol 1999;93:674–9.
- SOGC clinical practice guidelines. Guidelines for vaginal birth after previous caesarean birth. Number 155 (Replaces guideline Number 147), February 2005. Int J Gynaecol Obstet 2005; 89:319.
- Soltan, MH, Khashoggi, T, Adelusi, B. Pregnancy following rupture of the pregnant uterus. Int J Gynaecol Obstet 1996; 52:37.
- Speert H: Edward Porro and cesarean hysterectomy Surg. Gynecol. Obstet. 1; 1958 : 106-245.
- Stark DD, M c Carrthy SM, Filly RA, Parer J nT, Hricak H, Callen PW. Plevimetry by magnetic resonance imaging. Am. J.Radiol, ;1995 ,144,947.
- Stark DD, MC Carthy SM, Parer JT, Gallon PW. Delivery after previous cesarean section. Evaluation and management Am - J Obstet Gynecol;1991 , 69: 696 – 700.

- Stotland, NE, Lipschitz, LS, Caughey, AB. Delivery strategies for women with a previous classic cesarean delivery: a decision analysis. *Am J Obstet Gynecol* 2002; 187:1203.
- Stutchfield P, Whitaker R, Russell I. Antenatal Steroids for Term Elective Caesarean Section Research Team Antenatal betamethasone and incidence of neonatal respiratory distress after elective caesarean section: pragmatic randomised trial. *BMJ* 2005;331:662.
- Stutchfield, P, Whitaker, R, Russell, I. Antenatal betamethasone and incidence of neonatal respiratory distress after elective caesarean section: pragmatic randomised trial. *BMJ* 2005; 331:662.
- Taffel SM, Placek PJ, Moien M, Kosary CL, 19089 US cesarean section rate steadies. VBAC rises to nearly one in five birth;1991 , 18:73.
- Thubisi, M, Ebrahim, A, Moodley, J, Shweni, PM. Vaginal delivery after previous cesarean section: is X-ray pelvimetry necessary?. *Br J Obstet Gynaecol* 1993; 100:421.
- Tita, AT, Landon, MB, Spong, CY, et al. Timing of elective repeat cesarean delivery at term and neonatal outcomes. *N Engl J Med* 2009; 360:111.
- Traci L., Hugh r., William G.Michael L.: Vaginal birth after cesarean among women with gestational diabetes. *Am J Obstet Gynecol*;2002, 184; 1104-7.
- Tucker JM, Hauth JC, Hodgkins P, Owen J, Winkler CL.Trial of labor after one-or two layer closure, of a low transverse uterine

incision. Am J Obstet Gynecol;1993, 168: 545 -546 .

- Usta, IM, Hamdi, MA, Musa, AA, Nassar, AH. Pregnancy outcome in patients with previous uterine rupture. Acta Obstet Gynecol Scand 2007; 86:172.
- Uygur D, Gun O, Kelekci S, Ozturk A, Ugur M, Mungan T. Multiple repeat caesarean section: is it safe? *Eur J Obstet Gynecol Reprod Biol* 2005;119:171–5.
- Velasco VR, Beltron ER , Bejuranom OT. Vaginal delivery after cesarean section . J Obstet Gynecol , Biol. Reprod (Paris);1990, 2:6473.
- Walsh, CA, Baxi, LV. Rupture of the primigravid uterus: a review of the literature. Obstet Gynecol Surv 2007; 62:327.
- Washington DC: Repeat cesarean birth . U. Governement Printing Office;1998, 351 -374.
- Wen, SW, Rusen, ID, Walker, M, et al. Comparison of maternal mortality and morbidity between trial of labor and elective cesarean section among women with previous cesarean delivery. Am J Obstet Gynecol 2004; 191:1263.
- Wing MD, Deborah A, Paul MD and Richard H. Vaginal birth after cesarean section: Selection and management. Clinical. Obstet Gynecol.;1999, Volume 42 (4): 846 – 848.
- Zelop, CM, Shipp, TD, Repke, JT, et al. Effect of previous vaginal delivery on the risk of uterine rupture during a subsequent trial of labor. Am J Obstet Gynecol 2000; 183:1184.

- Zelop, CM, Shipp, TD; Repke, JT, et al. Uterine rupture during induced or augmented labor in gravid women with one prior cesarean delivery. Am J Obstet Gynecol 1999; 181:882.
- Zemlyn,S.: The length of the uterine cervix and its significance. J. Clin, Ultrasound;1981, 9:267.