

REFERENCES

1. **Agrons GA, Wagner BJ, Lonergan GJ, Dickey GE, Kaufman MS. (1997):** Genitourinary rhabdomyosarcoma in children: radiologic-pathologic correlation. *RadioGraphics*;17:919–937.
2. **Ahmed El-Assmy Mohamed E. Abou-El-Ghar Ahmed Mosbah (2009):**Bladder tumour staging: comparison of diffusion- and T2-weighted MR imaging *European Society of Radiology*.
3. **Amendola MA, Glaser GM, Grossman HB, et al.(1985):** Staging of **bladder** carcinoma: MRI-CT-Surgical correlation. *AJR* 1986; 146: 1179-1183. 2. **Arger PH.** Computed tomography of the lower urinary tract. *Urologic clinics of North America*; 12: 677-686.
4. **Asbury WL Jr, Hatcher PA, Gould HR, Reeves WA, Wilson DD. (1996):**Bladder pheochromocytoma with ring calcification. *Abdom Imaging*;21: 275–277.
5. **Babaian RJ, Johnson DE, Liamas L et al.(1991):** Metastases from transitional cell carcinoma of the urinary bladder. *Urology* 1990; 26: 142-150. 4. **Barbaric ZL.** Bladder "principles of genito-urinary radiology". Thieme medical publishers inc., New York; 358-386.
6. **Bammer R.(2003):** Basic principles of diffusion-weighted imaging. *Eur J Radiol*; 45:169–184.
7. **Bardales RH, Pitman MB, Stanley MW, Korourian S, Suhrland MJ. (1998):** Urine cytology of primary and secondary urinary bladder adenocarcinoma. *Cancer*;84:335–343.

8. **Barentsz J.O., Engelbrecht, M.R. & Jager,G.J. (1999):** Fast dynamic Gadolinium-enhanced MR imaging of urinary bladder and prostate cancer. *J Magnetic Resonance imaging*, 10, 3, 295-304.
9. **Barentsz JO, Berger-Hartog O, Witjes JA et al.(1998):** Evaluation of chemotherapy in advanced urinary bladder cancer with fast dynamic contrast-enhanced MR imaging. *Radiology*; 207: 791-797.
- 10.**Barentsz JO, Debruyne FMJ, Ruijs SHJ et al. (1990):**MR imaging of carcinoma of the urinary bladder; Kluiver Academic publishers, Dordrecht.
- 11.**Barentsz JO, Jager GJ, Mugler JP et al.(1994):** Staging urinary bladder cancer: Value of T1-weighted three-dimensional magnetization prepared rapid gradient-echo and two dimensional spin-echo sequence. *AJR*; 164: 109-115.
- 12.**Barentsz JO, Jager GJ, Vierzen PBJ et al. (1996a):** Staging urinary bladder cancer after transurethral biopsy: Value of fast dynamic contrastenhanced MR imaging. *Radiology*; 201: 185-193.
- 13.**Barentsz JO, Jager GJ, Vierzen PBJ et al.(1996a):** Staging urinary bladder cancer after transurethral biopsy: value of fast dynamic contrast-enhanced MR imaging. *Radiology*; 201:185-193.
- 14.**Barentsz JO, Jager GJ, Witjes JA et al. (1996b):** Primary staging of urinary bladder carcinoma: The role of MRI and a comparison with CT. *European Radiology* 6: 129-133.
- 15.**Barentsz JO, Jager GJ, Witjes JA et al.(2000):** MR imaging of the urinary bladder. *Magnetic Resonance imaging*; 8: 853-867.

16. **Barentsz JO, Jager GJ, Witjes JA, Ruijs JHJ (1996):** Primary staging of urinary bladder carcinoma: the role of MR imaging and a comparison with CT. *Eur Radiol* 6:134–139.
17. **Barentsz JO, Ruijs SHJ, Strijk SP.(1993):** The role of MR imaging in carcinoma of the urinary bladder. *AJR*; 160: 937- 947.
18. **Barentsz JO, Van Erning LJ, Ruijs SHJ.(1992):** Dynamic turbo-FLASH subtraction MR imaging. *Radiology*; 185: 340-345.
19. **Barentsz JO, Witjes JA, Ruijs SHJ et al.(1997):** What is new in bladder cancer imaging. *Urologic Clinics of North America*; 221: 583-602.
20. **Bates AW, Baithun SI.(2000):** Secondary neoplasms of the bladder are histological mimics of nontransitional cell primary tumours: clinicopathological and histological features of 282 cases. *Histopathology*;36:32–40.
21. **Bates AW, Norton AJ, Baithun SI.(2000):** Malignant lymphoma of the urinary bladder: a clinicopathological study of 11 cases. *J Clin Pathol*;53:458– 461.
22. **Beahrs OH, Hensen DE, Hutter RVP et al.(1990):** Manual for staging of cancer; 193-195.
23. **Bellin MF, Roy C, Kinkel K et al.(1998):** Lymph node metastases: safety and effectiveness of MR imaging with ultra-small super-paramagnetic iron oxide particles- initial clinical experience. *Radiology*; 207: 799808.
24. **Bennett JK, Wheatly JK and Walton KN. (1984):** 10-year experience with adenocarcinoma of the bladder. *Urology*; 131: 262-275.

- 25.Binsaleh S, Corcos J, Elhilali MM, Carrier S.(2004):** Bladder leiomyoma: report of two cases and literature review. *Can J Urol*;11:2411–2413.
- 26. Bloch F, Hansen WW, Packard M.(1946):** The nuclear induction experiment. *Phys Rev* ;70:474-485.
- 27.Blute MI, Engen DE and Traris WD.(1988):** Primary signet ring adenocarcinoma of the bladder. *Urology* 1989; 141: 17-20. 20.Brick SH, Frideman AC and Pollacic HM. Urachal carcinoma. *Radiology*; 169: 377-382.
- 28.Bo W and Krueger W.(1979):** Anatomical concepts of the urogenital system: Gross and cross sectional anatomy, *Ultrasound in urology*.
- 29.Borden, L.S., Clark, P.E., Hall. C.M. (2003):** Bladder cancer, Current opinion.*Oncology*, 15, 227-233.
- 30.Brick SH, Friedman AC, Pollack HM, et al.(1988):** Urachal carcinoma: CT findings. *Radiology*; 169:377–381.
- 31. Bryan PJ, Butler HE, Resnick MI et al.(1987):** CT and MR imaging in staging bladder neoplasms. *J Comp Assist Tomography*; 11: 96101.
- 32.Buckley BT, Wainwright A, Meagher T, Briley D.(2003):** Audit of a policy of magnetic resonance imaging with diffusion-weighted imaging as first-line neuroimaging for inpatients with clinically suspected acute stroke. *Clin Radiol*;58(3):234-237.
- 33.Bullock NE, Sibley G and Whitaker R.(1990):** Urothelial tumors. *Essential Urology*; 16: 214-226.

- 34.Buy JN, Moss AA, Guinet C et al.(1988):** MR staging of bladder carcinoma: Correlation with pathological findings. *Radiology*; 169: 695-700.
- 35.Cancer facts and figures 2005 (online database) (2005):** American Cancer Society Web site.
[http://www.cancer.org/docroot/STT/content/STT_1x_Cancer_Facts_Figures .asp](http://www.cancer.org/docroot/STT/content/STT_1x_Cancer_Facts_Figures.asp). Published 2005.
- 36.Chang SD and Hricak H.(2001):** Radiological evaluation of the urinary bladder, prostate and urethra. Grenger and Allison (ED). *Diagnostic radiology; A text book of medical imaging* (4th edition). Churchill livingstone, Harcourt publisher limited; 1615-1652.
- 37. Chen M, Lipson SA, Hricak H.(1997):** MR imaging evaluation of benign mesenchymal tumors of the urinary bladder. *AJR Am J Roentgenol*;168: 399–403.
- 38. Cheng L, Nascimento AG, Neumann RM, et al.(1999):** Hemangioma of the urinary bladder. *Cancer*;86:498–504.
- 39. Cheng L, Pan CX, Yang XJ, et al.(2004):** Small cell carcinoma of the urinary bladder: a clinicopathologic analysis of 64 patients. *Cancer*;101:957.
- 40. Chernoff, Hricak H(1997):** The male pelvis prostate and seminal vesicles(In) Higgins CB, Hricak H, Helms CA: *Magnetic resonance imaging of the body*, rd edition. Lippincott-Roven press, New work: 875-900.
- 41.Choi H, Lamb S and Pintar K.(1984):** Primary signet-ring cell carcinoma of urinary bladder. *Cancer*; 53: 1989-1990.

- 42.Clark SS, Marlet MM, Prudenico RF et al.(1989):** Neurofibromatosis of the bladder in children, case report. *Urology*; 118: 654.
- 43.Connor JP, Olsson CA and Kressel HY.(1989):** Long term follow-up in patients treated with chemotherapy for transitional cell carcinoma of the urinary bladder. *Urology*; 34: 353-363.
- 44.Cornella JL, Larson TR, Lee RA, Magrina JF, Kammerer- Doak D. (1997):** Leiomyoma of the female urethra and bladder: report of twenty-three patients and review of the literature. *Am J Obstet Gynecol*;176:1278–1285.
- 45. Corrington MB, HricakH (1991):**MRI of the pelvis.(In) Hricak& Corrington: *MRI of the pelvis A text atlas*, Applenton & large Nor walk, Connection scan mateo, California:45-64.
- 46.Cousins J. P. Clinical MR Spectroscopy (1995):** Fundamentals, Current Applications, and Future Potential*AJR*;164:1337-1347.
- 47.Cova M, Squillaci E, Stacul F, et al.(2004):** Diffusion-weighted MRI in the evaluation of renal lesions: preliminary results. *Br J Radiol*;77(922):851-857.
- 48.Crawford ED.(1996):** Diagnosis and treatment of suspected bladder cancer: An update seminar in urologic oncology; 14: 1-9.
- 49.Crecelius SA, Bellah R.(1995):** Pheochromocytoma of the bladder in an adolescent: sonographic and MR imaging findings. *AJR Am J Roentgenol*;165: 101–103.
- 50. Cummings KB, Barone JG and Word WS. (1992):**Diagnosis and staging of bladder cancer. *Urologic Clinics of North America*; 455-465.

- 51.Dahm P, Gschwend JE.(2003):** Malignant non-urothelial neoplasms of the urinary bladder: a review. *Eur Urol*;44:672–681.
- 52. Deklerk DP, Catalona WJ and Cohen EK.(1988):** Malignant pheochromocytoma of the bladder. *Urology*; 113: 864-870.
- 53.Deserno WM, Harisinghani MG, Taupitz M, et al.(2004):** Urinary bladder cancer: preoperative nodal staging with ferumoxtran-10-enhanced MR imaging. *Radiology* ;233:449–456.
- 54. Dondalski M, White EM, Ghahremani GG, Patel SK. (1993):** Carcinoma arising in urinary bladder diverticula: imaging findings in six patients. *AJR Am J Roentgenol* ;161:817–820.
- 55. Doms GC, Hricak H, Crooks LE, et al (1984):**Magnetic resonance imaging of lymph nodes:Comparison with CT.*radiology*;153:719-728.
- 56.Doms GC, Hricak H, Tscholakoff D (1986):** Adnexial structure: MR imaging.*Radiology*;158:639-646.
- 57.Doran F, Varinli S, Bayazit Y, Bal N, Ozdemir S.(2002):** Pheochromocytoma of the urinary bladder. *APMIS*;110:733–736.
- 58.Doringer E, Joss H and Scher HI.(1991):** MRI of the bladder carcinoma: Tumor staging and gadolinium contrast behavior. *Clinical radiology*; 154: 357-363.
- 59. Eble JN, Sauter G, Epstein JI, Sesterhenn IA, eds.(2004):** Pathology and genetics of tumours of the urinary system and male genital organs. World Health Organization Classification of Tumours. Lyon, France: IARC Press.

- 60. El-Assmy A, Abou-El-Ghar ME, Refaie HF, El-Diasty (2008):** Diffusion-weighted MR imaging in diagnosis of superficial and invasive urinary bladder carcinoma: apreliminary prospective study. Scientific World J Urol 8:364–370.
- 61.EL-Bolkainy MN, Ghoneim M and Mochtar NM.(1981):** Pathology of bladder cancer associated with schistosomiasis. Medical J. Cairo Univ.; 49: 361-374.
- 62. EL-Sebaei I.(1983):** Cancer of bilharzial bladder, Bladder cancer volume II. CRC press inc., Florida, 16-18.
- 63.Elwi AM.(1976):** Pathology of Shistosomiasis, Compliled review on Shistosomiasis. The national information and documentation centre, Dokki, Cairo. Egypt; 4: 47-80.
- 64.Evans CP and Swanson DA. (1996):** What to do if lymph nodes are positive. Seminars in urologic oncology; 14: 96-102.
- 65.Fisher MR, Hricak H and Tanagho EA.(1985):** Urinary bladder imaging II. Neoplasms. Radiology; 157: 471-477.
- 66.Fisher MR, Hricak H, Crooks LE. (1985):** Urinary bladder MR imaging. I. Normal and benign conditions. Radiology 157: 467-470.
- 67.Flamm J, Bucher A, Read G et al.(1990):** Recurrent superficial transitional cell carcinoma of the bladder. Urology; 144:260-268.
- 68.Fletcher BD, Kaste SC and Johnson RJ.(1992):** Magnetic resonance imaging for diagnosis and follow-up of genitourinary, pelvic and perineal rhabdosarcoma. Radiology; 14:263-272.

- 69.Friedell GH, Soloway MS and Hilgar AG.(1986):** Summary of workshop on carcinoma in situ of the bladder. *Urology*; 136: 1047-1050.
- 70.Friedman AC, Radecki PD, Toaff ASL et al.(1990):** Clinical pelvic imaging: CT, US and MRI.; 346- 380.
- 71.Gordon NS, Sinclair RA and Snow RM.(1990):** Pelvic lipomatosis with cystitis cystica, cystitis glandularis and adenocarcinoma of the bladder. *J. Urology*; 60: 229-236.
- 72. Gray H.(2000):** the urogenital apparatus, anatomy of urinary bladder. Reedited by warren HL, *Anatomy of the human body* (20th edition). Bartleby comp, New York, 956-974.
- 73. Grignon DJ, Ro JY, Ayala AG, Johnson DE, Ordonez NG. (1991):** Primary adenocarcinoma of the urinary bladder: a clinicopathologic analysis of 72 cases. *Cancer*;67:2165–2172.
- 74.Grignon DJ, Ro JY, Ayala AG, Johnson DE.(1991):** Primary signet-ring cell carcinoma of the urinary bladder. *Am J Clin Pathol*;95:13–20.
- 75.Grossfeld GD, Carroll PR. (1998):** Evaluation of asymptomatic microscopic hematuria. *Urol Clin North Am*. 25:661–676.
- 76.Grossfeld GD, Litwin MS, Wolf JS Jr, et al.(2001):** Evaluation of asymptomatic microscopic hematuria in adults: the American Urological) Association best practice policy. II. Patient evaluation, cytology, voided markers, imaging, cystoscopy, nephrology evaluation, and follow-tip. *Urology*;57(4):604-610.

- 77. Guo Y, Cai YQ, Cai ZL, et al.(2002):** Differentiation of clinically benign and malignant breast lesions using diffusion- weighted imaging. *J Magn Reson Imaging*; 16:172–178.
- 78.Hahn D.(1990):** Neoplasms of the urinary bladder. *Clinical urography*: 1353-1380.
- 79.Hall TB and Mac Vicar AD.(2001):** Imaging of bladder cancer. *Imaging*; 13: 1-10.
- 80. Hardeman SW, Perry A and Soloway MS.(1990):** Transitional cell carcinoma of the prostate following intravesical therapy for transitional cell carcinoma of the bladder. *J. Urology*; 140: 289-295.
- 81.Harisinghani MG, Barentsz JO, Hahn PF et al.(2002):** MR lymphangiography for detection of minimal nodal disease in patients with prostate cancer. *Academic Radiology*; 9 (supplement 2): 312-313.
- 82. Heiken JP and Lee JKT. (1988):** MR imaging of the pelvis. *Radiology*; 166: 11-16.
- 83. Heiken JP, Forman HP and Broum JJ.(1994).** Neoplasms of the bladder, prostate and testis. *Radiologic clinics of North America*; 32: 81-98.
- 84.Helzlsouer KJ, Comstock GW and Morris JS.(1989):** Selenium, lycopene, alpha-tocopherol, beta-carotene and subsequent bladder cancer recurrences. *J. Urology*; 140: 434-444.
- 85.Hemal AK, Singh I, Pawar R, Kumar M, Taneja P.(2000):** Primary malignant bladder carcinoid: a diagnostic and management dilemma. *Urology*; 55:949.

- 86.Hendry WF and Bloom HJG.(1985):** Urothelial neoplasia, Textbook of genito-urinary surgery. Churchill Livingstone: 970-1000.
- 87.Hricak H, Alpers C, Crooks LE, et al (1983):** Magnetic resonance imaging of the female pelvis: initial experience AJR; 141:1119-1128.
- 88. Hricak H. Alpers C, Crooks LE, et al (1990):** The bladder and female urethra. MRI of the pelvis; 417-461.
- 89.Huang WL, Ro JY and Grignon DJ.(1990):** Post-operative spindle nodule of the prostate and bladder. J. Urology; 143: 824-840.
- 90.Hughes MJ, Fisher C, Sohaib SA.(2004):** Imaging features of primary nonurachal adenocarcinoma of the bladder. AJR Am J Roentgenol;183: 1397–1401.
- 91. Husband JE, Ollif JFC, Williams MP et al.(1989):** Bladder cancer: Staging with CT and MR imaging. Radiology; 173: 435-440.
- 92. Husband JE. Ollif JFC, Williams MP et al (1995):** Staging urinary bladder cancer. Clinical Radiology; 46: 153-159.
- 93.Hwang JJ, Uchio EM, Patel SV, Linehan WM, Walther MM, Pacak K.(2003):** Diagnostic localization of malignant bladder pheochromocytoma using 6-18F fluorodopamine positron emission tomography. J Urol;169:274–275.
- 94.Ishikawa K, Saitoh M, Chida S.(2003):** Detection of bladder hemangioma in a child by blood-pool scintigraphy. Pediatr Radiol;33:433–435.

- 95. Issa B.(2002):** In vivo measurement of the apparent diffusion coefficient in normal and malignant prostatic tissues using echo-planar imaging. *J Magn Reson Imaging*;16(2):196-200.
- 96.Jager GJ, Barentsz JO, Oosterhof G et al.(1996):** 3D MR imaging in nodal staging of urinary bladder and prostate carcinoma. *AJR* 6; 167: 1503-1507.
- 97.Jalil ND, Pattou FN, Combemale F, et al.(1998):** Effectiveness and limits of preoperative imaging studies for the localisation of pheochromocytomas and paragangliomas: a review of 282 cases. French Association of Surgery (AFC), and The French Association of Endocrine Surgeons (AFCE). *Eur J Surg*;164:23–28.
- 98. Jocham D.(1994):** Malignant tumors of the bladder. Ed. Jocham D and Miller K. Stuttgart publishers,: 49-115.
- 99.Julie A, Trevor B, Carrington BM et al.(1995):** The role of MRI in tumor staging. *Journal of Pathology*; 175: 357-363.
- 100. Kabalin JN, Freiha FS and Niebel JD.(1990):** Leiomyoma of the urinary bladder. *Urology*; 35: 210-217.
- 101. Kakioze T, Matsumoto K, Audoh M et al.(1983):** Adenocarcinoma of the urachus. *Urology*; 21: 360-365.
- 102. Kalble T, Tricher AR and Friedel P.(1990):** Risk of carcinoma and etiologic factors for carcinogenesis. *J. Urology*; 144: 1110-1120.
- 103. Kantor AF, Hartge P, Hoover RN et al.(1988):** Epidemiological characteristics of squamous cell carcinoma of the bladder. *Cancer*; 48: 3853-3866.

104. **Kay R, Lattanzi C and Nerlon BG.(1985):** Nephrogenic adenoma in children. *J. Urology*; 133: 99-119.
105. **Keane P, Waxman J and Sikora K.(1995):** Urological system, Treatment in Cancer (3rd edition). Chapman and Hall medical press; 557-586.
106. **Keep JC, Piehl M, Miller A et al.(1989):** Invasive carcinomas of the urinary bladder. *J. Urology*; 141: 33-50.
107. **Kent DL, Shacher R, Sox HC et al.(1990):** Efficient scheduling of cystoscopies in monitoring for recurrent bladder cancer. *Med. Decision Making*; 9: 26-36.
108. **Kim B, Semelka RC, Ascher SM et al.(1994):** Bladder tumor staging: Comparison of contrast-enhanced CT, T1-, T2-weighted, dynamic Gadolinium-enhanced images and late Gadolinium-enhanced images. *Radiology*; 193: 239-245.
109. **Kim JC, Kim KH, Jung S.(2003):** Small cell carcinoma of the urinary bladder: CT and MR imaging findings. *Korean J Radiol*;4:130–135.
110. **Kim JC, Kim KH, Jung S.(2004):** Small cell carcinoma of the urinary bladder: CT and MR imaging findings. *Korean J Radiol* 2003;4:130–135.
111. **Kim JC. CT(2004):** features of bladder small cell carcinoma. *Clin Imaging*;28:201–205.
112. **Kim JK, Ahn JH, Park T, Ahn HJ, Kim CS, Cho KS.(2002):** Virtual cystoscopy of the contrast material-filled bladder in patients with gross hematuria. *AJR Am J Roentgenol*;179(3):763-768.

113. **Kim JK, Park SY, Ahn HJ, Kim CS, Cho KS. Bladder (2004):** Cancer: analysis of multi-detector row helical CT enhancement pattern and accuracy in tumor detection and perivesical staging. *Radiology*;231:725–731.
114. **Kim SH, Cha KB, Choi YD, Cho NH.(2004):** Solitary fibrous tumor of the urinary bladder. *Yonsei Med J*;45:573–576.
115. **Kim, B., Semelka, R.C., Ascher, S.M., Chalpin, D.B., Carroll, P.R., Hrica, H. (1994):** Bladder Tumor Staging: Comparison of Contrast-enhanced CT, T1 and T2-weighted MR Imaging, Dynamic Gadolinium-enhanced Imaging, and Late Gadolinium-enhanced Imaging. *Radiology*, 193, 239-245.
116. **Knoll LD, Segura JW, Scheithauer BW.(1986):** Leiomyoma of the bladder. *J Urol*;136:906–908.
117. **Koelbel G, Schmeidel U, Griebel J et al.(1988):** MR imaging of urinary bladder neoplasms. *J Comput Assist Tomogr*; 12: 98-103.
118. **Koh D.M.& Collins D.J. (2007):** Diffusion-Weighted MRI in the Body: Applications and Challenges in Oncology *AJR*;188,. June.
119. **Koh DM, Scurr E, Collins DJ, et al.(2006):** Colorectal hepatic metastases: quantitative measurements using single-shot echo-planar diffusion-weighted MR imaging. *Eur Radiol*; 16:1898–1905
120. **Kosuda S, Kison PV, Greenough R, Grossman HB, Wahl RL. (1997):** Preliminary assessment of fluorine-18 fluorodeoxyglucose positron emission tomography in patients with bladder cancer. *Eur J Nucl Med* ;24:615–620.

121. **Krestin GP, Steinbach W and Friedman G.(1990):** Recurrent rectal cancer: Diagnosis with MR imaging versus CT. *Radiology*; 168: 307-311.
122. **Kuroda M, Saiki S, Kinouchi T, et al. (1988):** Treatment of transitional cell carcinoma of the urinary bladder. *Jpn J Urol* 79: 507-512.
123. **Lämmle M, Beer A, Settles M, Hannig C, Schwaibold li, Drews C.(2002):** Reliability of MR imaging-based virtual cystoscopy in the diagnosis of cancer of the urinary bladder. *AJR Am J Roentgenol*;178(6):1483-1488.
124. **Last JR.(1984):** Anatomy: Regional and applied (7th edition). Churchill livingstone, London;.
125. **Le Bihan D, Breton E, Lallemand D, Aubin ML, Vignaud J, Laval-Jeantet M.(1988):** Separation of diffusion and perfusion in intravoxel incoherent motion MR imaging. *Radiology*; 168:497–505.
126. **Leiskovsky G, Ahlering T and Skinner DG.(1988):** Diagnosis and staging of bladder cancer, Diagnosis and management of Genito-urinary cancer. WB Saunders CO, Philadelphia; P.264.
127. **Leite KR, Srougi M, Miotto A, Camara-Lopes LH.(2004):** Solitary fibrous tumor in bladder wall. *Int Braz J Urol*;30:406–409.
128. **Lerner SP, Skinner DG, Lieskovsky G et al.(1993):** The rationale for en bloc pelvic lymph node dissection for bladder cancer patients with nodal metastases. *J. Urology*; 149: 758-765.

129. **Levy AD, Patel N, Dow N, Abbott RM, Miettinen M, Sobin LH. (2005):** Abdominal neoplasms in patients with neurofibromatosis type 1: radiologic-pathologic correlation. *RadioGraphics*;25:455– 480.
130. **Lipson SA& Hricak H(1997):**The urinary bladder and female urethra(In) Higgins CB, Hricak H, Helms CA: *Magnetic resonance imaging of the body*, 3 red, Lippincott Romapress, New work:815-900.
131. **Maeda H, Kinukawa T, Hattori R et al.(1995):** Detection of muscle layer invasion with submillimeter pixel MR imaging: Staging of bladder carcinoma. *Magnetic Resonance imaging*; 13: 9-19.
132. **Mallampati GK, Siegelman ES. MR(2004):** imaging of the bladder. *Magn Reson Imaging Clin N Am*;12:545–555.
133. **Marcus PM, Hayes RB, Vineis P, et al.(2000):** Cigarette smoking, N-acetyltransferase 2 acetylation status, and bladder cancer risk: a case-series meta-analysis of a gene-environment interaction. *Cancer Epidemiol Biomarkers Prev*;9:461–467.
134. **Mariani AJ.(1998):** The evaluation of adult hematuria: a clinical update. In: *AUA update sed ries. Vol XVII. Lesson 24.* Houston, Tex: AUA Office of Education, 185-192.
135. **Martignoni G, Eble JN.(2003):** Carcinoid tumors of the urinary bladder: immunohistochemical study of 2 cases and review of the literature. *Arch Pathol Lab Med*;127:e22–e24.

136. **Martin SA, Sears DL, Sebo TJ, Lohse CM, Cheville JC.(2002):** Smooth muscle neoplasms of the urinary bladder: a clinicopathologic comparison of leiomyoma and leiomyosarcoma. *Am J Surg Pathol*;26:292–300.
137. **Matsuki M, Inada Y, Tatsugami F, Tanikake M, Narabayashi I, Katsuoka Y.(2007):** Diffusion-weighted MR imaging for urinary bladder carcinoma: initial results. *Eur Radio*; 17(1):201-204.
138. **Mc Carthy S(1992):**Magnetic resonance imaging of the normal female pelvis .*Radiol. clin. North Am*; 30:769-773.
139. **Millan-Rodriguez F, Chechile-Toniolo G, Salvador- Bayarri J, Huguet-Perez J, Vicente-Rodriguez J.(2000):** Upper urinary tract tumors after primary superficial bladder tumors: prognostic factors and risk groups. *J Urol*;164:1183 1187.
140. **Montei JE.(1990):** High-stage bladder cancer: Bladder preservation or reconstruction. *Clinical Jour. Med.*; 57: 280-290.
141. **Montironi R, Lopez-Beltran A.(2005):** The 2004 WHO classification of bladder tumors: a summary and commentary. *Int J Surg Pathol*;13:143–153.
142. **Moon WK, Kim SH, Cho JM, Han MC.(1992):** Calcified bladder tumors: CT features. *Acta Radiol*; 33:440–443.
143. **Muller-Lisse UG, Heuck AF and Barentsz JO.(2001):** Bladder "Medical Radiology, Diagnostic imaging, Abdominal and pelvic MRI. Heuck A and Reiser M (Ed.). Springer Verlag Berlin, Heidelberg, New York; 16: 209-226.

144. **Murphy WM, Grignon DJ, Perlman EJ.(2004):** Tumors of the kidney, bladder, and related urinary structures. Washington, DC: American Registry of Pathology; 394.
145. **Nanumi Y, Kadota T, Inoue E, et al (1993):** Bladder tumors: staging with gadolinium-enhanced oblique MR imaging. *Radiology* 187:145-150.
146. **Narumi Y, Kadota T and Inoue E.(1993):** Bladder wall morphology: In vitro MR imaging with histopathological correlation. *Radiology*; 187: 151-155.
147. **Narumi Y, Kadota T, Inoue E, et al (1995):** Bladder wall morphology: In vitro MR imaging histological correlation. *Radiology*; 187:151-155.
148. **Narumi Y, Sato T, Hori S, et al.(1989):** Squamous cell carcinoma of the uroepithelium: CT evaluation. *Radiology*;173:853–856.
149. **Nasu K, Kuroki Y, Kuroki S, Murakami K, Nawano S, Moriyama N.(2004):** Diffusion-weighted single shot planar imaging of colorectal cancer using a sensitivity-encoding technique. *Jpn J Clin Oncol*;34(10):620-626.
150. **Netter FH. Urinary system. Netter FH (Ed)(2002):** Interactive atlas of human anatomy (4th edition). The graphic witness Inc, south Florida; 430475.
151. **Neuerburg JM, Bohondrof K and Sohn M.(1992):** Urinary bladder neoplasms: evaluation with contrast-enhanced MRI. *Radiology*; 172: 739-743.

152. **Neurerburg JM, Bohndorf K, Sohn M, et al (1989):** Urinary bladder neoplasms: evaluation with contrast enhanced MR imaging. *Radiology*; 172:739-743.
153. **Nicolas V, Spielmann R and Mass R.(1991):** The diagnostic value of MR tomography following Gd-DTPA compared to CT in bladder tumors. *Radiology*; 164; 357-363.
154. **Nigro KG, MacLennan GT.(2005):** Rhabdomyosarcoma of the bladder and prostate. *J Urol*;173:1365.
155. **Nikken JJ, Krestin GP.(2007):** MRI of the kidney state of the art. *Eur Radiol*;17(11): 2780-2793.
156. **Oldbring J, Glifberg I, Mikulowski P et al.(1990):** Carcinoma of the renal pelvis and ureter following bladder carcinoma. *J. Urology*; 141: 1311-1321.
157. **Olliff JFC, Husband JES, Williams MP et al.(1989):** Bladder cancer: staging with CT and MR imaging. *Radiology*; 173: 435-440.
158. **Outwater EC(1992):**Magnetic resonance image of pelvis Haaya JR, Lanzier CF, Sartoris DJ,et al:Computed tomography and magnetic resonance imaging of the whole body,vol.2 Mosby, St Louis;1350-1404.
159. **Park SW, Lee JH, Ehara S, et al.(2004):** Single shot fast spin echo diffusion-weighted MR imaging of the spine: is it useful in differentiating malignant metastatic tumor infiltration from benign fracture edema? *Clin Imaging*; 28:102–108.
160. **Pashos CL, Botteman MF, Laskin BL, Redaelli A.Bladder (2002):** cancer epidemiology, diagnosis, andmanagement. *Cancer Pract*;10:311–322.

161. **Patel MD, Hricak H.(1995):** Current role of magnetic resonance imaging in urology. *Curr Opin Urology*; 5: 67-74.
162. **Persad R, Kabala J, Gillat D et al.,(1993):** Magnetic resonance imaging in the staging of urinary bladder cancer. *British J. Urology*; 71:556-573.
163. **Planz B, Synek C, Robben J, Bocking A, Marberger M.(1993):** Diagnostic accuracy of DNA image cytometry and urinary cytology with cells from voided urine³⁰. Stein JP, Skinner EC, Boyd SD, Skinner DG. Squamous cell carcinoma of the bladder associated with cyclophosphamide therapy for Wegener's granulomatosis: a report of 2 cases. *J Urol*;149:588–589.
164. **Prout GR, Griffm PP, Shipley WV et al.(1988):** Bladder carcinoma as a systemic disease. *Cancer*; 48: 2532-2540.
165. **Prout GR.(1997):** Bladder carcinoma and a TNM system for classification. *J. Urology*; 117:583-590.
166. **Purcell EM, Torrey HC, Pound RV.(1946):** Resonance absorption by nuclear magnetic moments in a solid. *Phys Rev*;69:37-38.
167. **Putman CE, Ravin CE and Lawrie CA.(1994):** Textbook of diagnostic imaging.: 2011-2029.
168. **Rafal RB, Markisz JA.(1991):** Urachal carcinoma: the role of magnetic resonance imaging. *Urol Radiol*; 12:184–187.
169. **Raghavan D, Shipley WV, Garnick MB et al.(1990):** Biology in management of bladder cancer. *North Engl. J.Medicine*; 322: 1129-1140.

170. **Reuter VE. Bladder: (1999):** risk and prognostic factors- a pathologist's perspective. *Urol Clin North Am* ;26:481–492.
171. **Riccardi VM.(1992):**Type 1 neurofibromatosis and the pediatric patient. *Curr Probl Pediatr*;22:66–106.
172. **Richard S, Jones S, Sherwood I et al(1993):** The bladder and prostate, *Textbook of radiology and medical imaging* (5th edition). Sutton D and Sherwood I (Ed.), Churchill Livingstone; Edinburgh, London: 1139-1158.
173. **Richie JP, Shipley WU, Yagoda A (1992):** Cancer of the bladder. In: Walsh PC, Retik AB, Stamey TA, Vaughan ED, eds. *Campbell's urology*. 6th ed. Vol 2. Philadelphia, Pa: Saunders, 1(X) 8-1022.
174. **Robbins SL and Cotran RS.(1979):** *Pathological basis of disease*. WS Saunders Co, Philadelphia, London, Toronto: 1197-1204.
175. **Rosi P, Sell] C, Carini M et al.(1983):** Myxoid liposarcoma of the bladder. *J. Urology*.; 130: 560-577.
176. **Rosser CJ, Slaton JW, Izawa JI, Levy LB, Dinney CP.(2003):** Clinical presentation and outcome of highgrade urinary bladder leiomyosarcoma in adults. *Urology*;61:1151–1155.
177. **Rotman M, Porrazzo M and Aziz H.(1990):** Treatment of advanced transitional cell carcinoma of the bladder with irradiation. *Magnetic resonance imaging*; 18: 1131-1141.
178. **Sanz LE.(1988):** *Anatomy of the retro peritoneum, Gynecologic surgery*. Medical Economics Books, Oradell; New Jersey: 3-10.

179. **Sato C, Naganawa S, Nakamura T, et al.(2005):** is Differentiation of non-cancerous tissue and cancer lesions by apparent diffusion coefficient values in transition and peripheral zones of the prostate. *J Magn Reson Imaging*;21(3):258-262.
180. **Scattoni V, Da-Pozzo LF, Colombo R et al.(1996):** Dynamic Gadolinium enhanced magnetic resonance imaging in staging of superficial bladder cancer.*J Urology*; 155:1594-1599.
181. **Schalow EL, Broecker BH.(2003):** Role of surgery in children with rhabdomyosarcoma. *Med Pediatr Oncol*;41:1–6.
182. **Schnall MD, Imai Y, Tomaszewski J, et al (1991):** Prostate cancer: local staging with endorectal surface coils MR imaging. *Radiology*; 178:797-801.
183. **Schoder H, Larson SM.(2004):** Positron emission tomography for prostate, bladder, and renal cancer. *Semin Nucl Med*;34:274–292.
184. **See WA and Fuller JR.(1992):** Staging of advanced bladder cancer: Current concepts and pitfalls. *Urologic clinics of North America*; 19: 663-683.
185. **Serretta V, Pomara G, Piazza F, Gange E.(2000):** Pure squamous cell carcinoma of the bladder in western countries: report on 19 consecutive cases. *Eur Urol* ;37:85–89.
186. **Shatov AV.(1996):** Use of low field MR imaging in the comprehensive diagnosis of disease of the pelvis. *Vestn-Rentgenol-Radiology*; 3:57-60.

187. **Sheldon CA, Clayman RV, Gonzalez R, Williams RD, Fraley EE. (1984):** Malignant urachal lesions. J Urol;131:1–8.
188. **Shokeir AA. (2004):** Squamous cell carcinoma of the bladder: pathology, diagnosis and treatment. BJU Int 2004;93:216–220.
189. **Shonnard KM, Jelinek JS, Benedikt RA, Kransdorf MJ. CT and MR (1992):** of neurofibromatosis of the bladder. J Comput Assist Tomogr;16:433–438.
190. **Smith FW.(1988):** Magnetic resonance imaging of the pelvis. Radiology, 8: 367-378.
191. **Smith H, Weaver G, Barenbruch et al.(1990):** Routine excretory urography in follow-up of superficial transitional cell carcinoma OF bladder. J. urology; 34: 193-203.
192. **Smith RC, Reinhold C, McCauley TR et al.(1992):** Multi-coil high resolution fast spin echo MR imaging of the pelvis. Radiology; 184: 665-669.
193. **Sohn M, Neuerburg J, Teufl F et al.(1990):** Gadolinium-enhanced MRI in the staging of urinary bladder neoplasms. Urology Int; 45: 142-147. 109.
194. **Song JH, Francis IR, Platt JF, et al.(2001):** Bladder tumor detection at virtual cystoscopy. Radiology; 218:95–100.
195. **Sparenberg A, Hamm B, Scher HI et al.(1991):** The diagnosis of bladder carcinoma by MR imaging. Fortschr Rontgenstr; 155: 117-122.
196. **Stein JP, Skinner EC, Boyd SD, Skinner DG. (1993):** Squamous cell carcinoma of the bladder associated with cyclophosphamide therapy for Wegener's granulomatosis: a report of 2 cases. J Urol 1993;149:588–589.

197. **Sudakoff GS, Dunn DP, Guralnick ML, Hellman RS, Eastwood D, See WA.(2008):** Multidetector computerized tomography urography as the primary imaging modality for detecting urinary tract neoplasms in patients with asymptomatic hematuria. *J Urol*; 179(3):862-867.
198. **Sundaram CP, Rawal A, Saltzman B.(1998):** Characteristics of bladder leiomyoma as noted on magnetic resonance imaging. *Urology*;52:1142–1143.
199. **Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Incidence-SEER 9 Regs Public-Use. Bethesda, Md: (2004):** National Cancer Institute, Division of Cancer Control and Population Sciences, Surveillance Research Program, Cancer Statistics Branch.
200. **Sved P, Gomez P, Manoharan M, Civantos F, Soloway MS.(2004):** Small cell carcinoma of the bladder. *BJU Int*;94:12–17.
201. **Swanson DA, Liles A and Zagar GK.(1990):** Pre-operative irradiation and radical cystectomy of squamous cell carcinoma of the bladder. *J. Urology*; 143: 37-42.
202. **Tachibana M, Baba S, Deguchi E et al.(1991):** Efficacy of gadolinium DTPA- enhanced magnetic resonance imaging for differentiation between superficial and muscle invasive tumors of the bladder. *J Urology*; 145: 1169-1173.
203. **Takahara T, Imai Y, Yamashita T, Yasuda S, Nasu S, Van Cauteren M.(2004):** Diffusion weighted whole body imaging with background body signal suppression (DWIBS): technical improvement using free breathing, STIR and high resolution 3D display. *Radiat Med*; 22:275–282.

204. **Takkouche B, Etminan M, Montes-Martinez A.(2005):** Personal use of hair dyes and risk of cancer: a meta-analysis. *JAMA*;293:2516–2525.
205. **Tanagho EA.(1992):** Anatomy of the lower urinary tract. *Campbell's Urology*; 40-47.
206. **Tanimoto A, Yuasa Y, Imai Y, et al. (1992):** Bladder Tumor Staging: Comparison of Conventional and Gadolinium-enhanced Dynamic MR Imaging and CT. *Radiology* 185: 741-747.
207. **Taouli B, Vilgrain V, Dumont E, Daire JL, Fan B, Menu Y. (2003):**Evaluation of liver diffusion isotrophy and characterization of focal hepatic lesions with two single-shot echo-planar MR imaging sequences: prospective study in 66 patients. *Radiology*;226(1): 71-78.
208. **Teeger S, Sica GT and Peter HA. (1996):** MR imaging of Bladder diseases. *MRI Clinics of North America*; 4: 565-581.
209. **Tekes A, Kamel I, Imam K, et al.(2005):** Dynamic MRI of bladder cancer: evaluation of staging accuracy. *AJR Am J Roentgenol*;184:121–127.
210. **Tekes A, Kamel IR, Imam K, Chan TY, Schoenberg MP, Bluemke DA.(2001):** MR imaging features of transitional cell carcinoma of the urinary bladder. *AJR*Huguet-Perez J, Palou J, Millan-Rodriguez F, Salvador- Bayarri J, Villavicencio-Mavrich H, Vicente- Rodriguez J. Upper tract transitional cell carcinoma following cystectomy for bladder cancer. *Eur Urol*;40:318–323.

211. **Thali-Schwab CM, Woodward PJ, Wagner BJ.(2005):** Computed tomographic appearance of urachal adenocarcinomas: review of 25 cases. *Eur Radiol*;15:79–84.
212. **Thoeny HC, De Keyzer F, Oyen RH, Peeters RR.(2005):** Diffusion-weighted MR imaging of kidneys in healthy volunteers and patients with parenchymal diseases: initial experience. *Radiology*;235(3):911-917.
213. **Thoeny HC, De Keyzer F, Vandecaveye V, et al.(2005):** Effect of vascular targeting agent in rat tumor model: dynamic contrast-enhanced versus diffusion- weighted MR imaging. *Radiology*; 237:492–499.
214. **Tsarnopoulos C, Tsili AC, Giannakis D, Alamanos Y, Sofikitis N, Efremidis SC.(2008):** MDCT cystoscopy in the evaluation of neoplasms of the urinary bladder. *AJR Am J Roentgenol*;190(3):729 -735.
215. **Tsukamoto T, Lieber MM and Tricker AR.(1990):** Staging systems of urinary bladder tumors. *Cancer*; 50: 44-50.
216. **Vandecaveye V, de Keyzer F, Vander Poorten V, et al.(2006):** Evaluation of the larynx for tumour recurrence by diffusion-weighted MRI after radiotherapy: initial experience in four cases. *Br J Radiol*; 79:681–687.
217. **Vinnicombe SJ, Norman A, Nicolson V et al.(1995):** Normal pelvic lymph nodes: Evaluation with CT after Bipodal lymphangiography. *Radiology*; 194: 349-355.
218. **Walsh JW.(1994):** Computed tomography of the pelvis, Computed tomography and magnetic resonance imaging of the whole body. Haggag JR, Lanzieri CF, Sartoris DJ and Zerhoun EA (Ed.). Mosby, St. Louis, Baltimore, Berlin, Boston 3rd edition;: 1327-1349.

219. **Webb JA.(1997):** Imaging in hematuria. Clin Radiol;52(3):167-171.
220. **Wegener OH.(1992):** The urinary bladder, Whole Body computed tomography. Blackwell scientific pub., Boston, Oxford, London 2nd edition: 415-424.
221. **Weiner DP, Koss LG, Sablay B et al.(1979):** The prevalence and significance of Brunn's nests, cystitis cystica and squamous metaplasia in normal bladder. J. Urology; 122:317-320.
222. **Weinstein RS, Miller AW, Paule BV et al.(1980):** Carcinoma in situ. Cancer; 43: 661-670.
223. **Wilkinson LM, Manson D, Smith CR.(2004):** Plexiform neurofibroma of the bladder. RadioGraphics;24(Spec Issue):S237–S242.
224. **Wolfe JS, Cher M and Era M.(1995):** The use and accuracy of cross sectional imaging and fine needle aspiration cytology for detection of pelvic lymph node metastases. J. Urology; 153: 993-999.
225. **Wong JT, Wasserman NF, Padurean AM.(2004):** Bladder squamous cell carcinoma. RadioGraphics;24:855–860.
226. **Yip Sk, Peh WC, Tam PC et al. (1999):** Role of ultrasonography in screening for urological malignancies in patients presenting with painless Hematuria. Ann Academic Med Singapore; 28: 174-177.
227. **Yoshida S, Masuda H, Ishii C, Saito K, Kawakami S, Kihara K (2008):** Initial experience of functional imaging of upper urinary tract neoplasm by diffusion-weighted magnetic resonance imaging. Int J Urol 15:140–143.

228. **Young RH.(1987):** Carcinosarcoma of the urinary bladder. *Cancer*; 9: 1333-1350.
229. **Zeegers MP, Kellen E, Buntinx F, van den Brandt PA.(2004):** The association between smoking, beverage consumption, diet and bladder cancer: a systematic literature review. *World J Urol*;21:392– 401.
230. **Zuk RJ, Baithum SI, Martin JE et al.(1989):** The demonstration of basement membrane deposition in transitional cell carcinoma of the bladder. *J. Urology*; 129: 505-518.