



REFERENCES



REFERENCES

1. Badgley, C.E.:
Etiology of congenital dislocation of the hip.
J. Bone Joint Surg., 31 (Am.) : 341-356 (1949).
2. Barret, W.P.; Staheli, L.T.; and Chow, D.E.:
The effectiveness of the Salter innominate osteotomy in the
treatment of congenital dislocation of the hip.
J. Bone Joint Surg., 68 (Am.) : 79-81 (1986).
3. Barry, K.; McManus, F.; and O'Brien, T.:
Leg lengthening by the transiliac method.
J. Bone Joint Surg., 74 (Br.): 275-278 (1992).
4. Bos, C.F.; and Bloem, J.L.:
Treatment of dislocation of the hip, detected in early
childhood based on magnetic resonance imaging.
J. Bone Joint Surg., 71 (Am.) : 1523-1529 (1989).
5. Bost, F.C.; Hagey, H.; Schottstaedt, E.R.; and Larsen, J.J.:
The results of treatment of congenital dislocation of the hip
in infancy.
J. Bone Joint Surg., 30 (Am.) : 454-468 (1948).
6. Broughton, W.S.; Cole, W.G.; and Menelaus, M.B.:
Reliability of radiological measurements in the assessment
of the child's hip.
J. Bone Joint Surg., 71 (Br.): 6-8 (1989).

7. Browning, W.H.; Rosenkrantz, H.; and Tarquinio, T.:
Computed tomography in congenital hip dislocation: The
role of acetabular anteversion.
J. Bone Joint Surg., 64 (Am.) : 27-31 (1982).
8. Buchanan, J.R.; Greer, R.B.III; and Colter, J.M.:
Management strategy for prevention of avascular necrosis
during treatment of congenital dislocation of the hip.
J. Bone Joint Surg., 63 (Am.) : 140-146 (1981).
9. Bucholz, R.W.; and Ogden, J.A.:
Patterns of ischaemic necrosis of the proximal femur in
nonoperatively treated congenital hip disease. In : The Hip.
Proceedings of the 6th Meeting of the Hip Society, St.
Louis, The C.V. Mosby Co., p. 43-63 (1978).
10. Buckley, S.L.; Sponseller, P.D.; and Magid, D.:
The acetabulum in congenital and neuromuscular hip
instability.
J. Pediatr. Orthop., 77 : 498-501 (1991).
11. Cage, J.R.; and Winter, R.B.:
Avascular necrosis of the capital femoral epiphysis as a
complication of closed reduction of the hip.
J. Bone Joint Surg., 54 (Am.) : 373-388 (1972).
12. Caplin, R.D.; Roach, J.W.; Wenger, D.R.; Herring, J.A.;
and Birch, J.G.:
One-stage treatment of congenital dislocation of the hip in
older children, including femoral shortening.
J. Bone Joint Surg., 71 (Am.) : 734-741 (1989).

13. Carlloz, H.; and Filipe, G.:
The natural history of the limbus in congenital dislocation of the hip: An arthrographic study. In : Tachdjian, M.O. (Ed.): Congenital Dislocation of the Hip. New York, Churchill Livingstone, p. 247-262 (1982).
14. Chapchal, G.:
Indications for the various types of pelvic osteotomy. Clin. Orthop., 98 : 111-115 (1974).
15. Chapchal, G.:
The intertrochanteric osteotomy in the treatment of congenital dysplasia of the hip. Clin. Orthop., 119 : 54-59 (1976).
16. Coleman, S.S.:
The incomplete pericapsular (Pemberton) and innominate (Salter) osteotomies. Clin. Orthop., 98 : 116-123 (1974).
17. Cooperman, D.R.; Wallenstein, R.; and Stulberg, S.D.:
Post-reduction avascular necrosis in congenital dislocation of the hip. J. Bone Joint Surg., 61 (Am.) : 247-258 (1980).
18. Core, D.R.:
Iatrogenic avascular necrosis of the hip in young children. J. Bone Joint Surg., 56(A) : 493-502 (1974).
19. Crego, C.H.; and Schwartzmann, J.R.:
Follow-up study of the early treatment of congenital dislocation of the hip. J. Bone Joint Surg., 30(A) : 428-442 (1948).

20. Crellin, R.Q.:
Innominate osteotomy for congenital dislocation and
subluxation.
Clin. Orthop., 98 : 171-177 (1974).
21. Denham, R.A.:
Hip mechanics.
J. Bone Joint Surg., 41 (Br.) : 550-557 (1959).
22. Denton, J.R.; and Ryder, C.T.:
Radiographic follow-up of Salter innominate osteotomy
for congenital dysplasia of the hip.
Clin. Orthop., 98 : 210-213 (1974).
23. De Rosa, G.P.; and Feller, N.:
Treatment of congenital dislocation of the hip: Management
before the walking age.
Clin. Orthop., 225 : 77-85 (1987).
24. Edelson, J.G.; Hirsch, M.; Weinberg, H.; Attar, D.; and
Barmeir, E.:
Congenital dislocation of the hip and computerized axial
tomography.
J. Bone Joint Surg., 66 (Br.) : 472-478 (1984).
25. Esteve, R.:
Congenital dislocation of the hip: A review and assessment
of results of treatment with special reference to frame
reduction as compared with manipulative reduction.
J. Bone Joint Surg., 42 (Br.) : 253-263 (1960).

26. Ferré, R.L.; and Schachter, S.:
Congenital dislocation of the hip: Innominate osteotomy.
Clin. Orthop., 98 : 183-194 (1974).
27. Fisher, R.L.; and Cary, J.M.:
Avascular necrosis complicating congenital dislocation of
the hip: Course, prognosis and orthopaedic management.
Int. Orthop., 2 : 229-240 (1978).
28. Fixen, J.A.:
Anterior and posterior displacement of the hip after
innominate osteotomy.
J. Bone Joint Surg., 69 (Br.) : 361-364 (1987).
29. Gallien, R.; Bertin, D.; and Lirette, R.:
Salter procedure in congenital dislocation of the hip.
J. Ped. Orthop., 4 : 427-430 (1984).
30. Ganz, R.; Klaue, K.; Vinh, T.S.; and Mast, J.W.:
A new periacetabular osteotomy for the treatment of hip
dysplasias.
Clin. Orthop., 232 : 26-36 (1988).
31. Garray, G.G.; Baeza, E.D.; and Hierro, A.S.:
Acetabular dysplasia in the rat induced by injury to the
triradiate growth cartilage.
Acta Orthop. Scand., 59 (5) : 516-519 (1988).
32. Gibson, P.; and Benson, M.:
Congenital dislocation of the hip.
J. Bone Joint Surg., 64 (Br.) : 169-175 (1982).

33. Graf, R.:
The diagnosis of congenital hip joint dislocation by
ultrasonic compound treatment.
Arch. Orthop. Trauma Surg., 97 : 117-133 (1980).
34. Graf, R.:
New possibilities for the diagnosis of congenital hip joint
dislocation by ultrasonography.
J. Pediatr. Orthop., 3 : 354-359 (1983).
35. Graf, R.:
Fundamentals of sonographic diagnosis of infant hip
dysplasia.
J. Pediatr. Orthop., 4 : 735-740 (1984).
36. Graf, R.:
Guide to sonography of the infant hip. Stuttgart, Federal
Republic of Germany, Thieme (1987).
37. Grill, F.:
Personal communication (1992).
38. Hansson, G.; Althoff, B.; Bylund, P.; Jacobsson, B.;
Lofberg, A.; and Lonnerholm, D.:
The Swedish experience with Salter's innominate osteotomy
in the treatment of congenital subluxation and dislocation
of the hip.
J. Pediatr. Orthop., 10 : 159-162 (1990).
39. Hansson, L.I.; Olsson, T.H.; Selvik, G.; and Sundén, G.:
A roentgen-stereo-photogrammetric investigation of
innominate osteotomy (Salter).
Acta Orthop. Scand., 49 : 68-72 (1978).

40. Harris, W.H.:
Acetabular growth potential in congenital dislocation of the hip and some factors upon which it may depend.
Clin. Orthop., 119 : 99-106 (1976).
- 41.. Harris, N.H.; Lloyd-Roberts, G.C.; and Gallien, R.:
Acetabular development in congenital dislocation of the hip with special reference to the indications for acetabuloplasty and pelvic or femoral realignment osteotomy.
J. Bone Joint Surg., 57 (Br.) : 46-52 (1975).
42. Herold, H.Z.:
Unilateral congenital hip dislocation with contralateral avascular necrosis.
Clin. Orthop., 148 : 196-202 (1980).
43. Hesinger, R.N.:
Congenital dislocation of the hip: Treatment in infancy to walking age.
Orthop. Clin. N. Amer., 18 : 597-616 (1987).
44. Howorth, B.:
Development of present knowledge of congenital displacement of the hip.
Clin. Orthop., 125 : 68-87 (1977).
45. Hughes, J.R.:
Intrinsic obstructive factors in congenital dislocation of the hip: The role of arthrography. In : Tachdjian, M.O. (Ed.): Congenital Dislocation of the Hip. New York, Churchill Livingstone, p. 227-245 (1982).

46. Izadpanch, M.:
Beckenosteotomie nach Salter zur Behandlung der
Kongenitalen Huftluxation und Subluxation.
Z. Orthop., 113 : 295-305 (1975).
47. Jacquemier, M.; Louve, J.L.; Bellini, G.; Panuel, M.; and
Migliani, R.:
Acetabular anteversion in children.
J. Pediatr. Orthop., 12 (3) : 373-375 (1992).
48. Kalamchi, A.:
Modified Salter osteotomy.
J. Bone Joint Surg., 64 (Am.) : 183-187 (1982).
49. Kalamchi, A.; and MacEwen, G.D.:
Avascular necrosis following treatment of congenital
dislocation of the hip.
J. Bone Joint Surg., 62 (Am.) : 876-888 (1980).
50. Kasset, M.R.; Whiteside, L.A.; Schoenecker, P.L.; and
Simmons, D.J.:
Salter innominate osteotomy: The effect of blood supply to
the roof of the acetabulum.
Clin. Orthop., 183 : 262-266 (1983).
51. Kim, Y.H.:
Acetabular dysplasia and osteoarthritis developed by an
eversion of the acetabular labrum.
Clin. Orthop., 215 : 289-295 (1987).

52. Klapsch, W.; Tschauner, C.; and Graf, R.:
Behandlungsergebnisse d₂zentrierter Huftgelenke seit
Einführung der Huftsonographie 1980.
Orthop. Praxis, 6 : 353-354 (1991).

53. Kollmann, K.:
Frühergebnisse Fehler and Gefahren der Beckenosteotomie.
Z. Orthop., 102 : 262-267 (1966).

54. Krismer, M.; and Hingsammer, R.:
Langzeitergebnisse der Beckenosteotomie nach Salter.
Orthop. Praxis, 6 : 355-360 (1991).

55. Lindholm, T.S.; Laurent, L.E.; Osterman, K.; and Snellmann,
O.:
Perthes' disease of a severe type developing after satisfactory
closed reduction of congenital dislocation of the hip: A
report of three cases.
J. Bone Joint Surg., 60 (Br.) : 15-17 (1978).

56. Lindstrom, J.R.; Ponesti, I.V.; and Wagner, S.R.:
Acetabular development after reduction in congenital
dislocation of the hip.
J. Bone Joint Surg., 61 (Am.) : 112-118 (1979).

57. Lloyd-Roberts, G.C.; and Swan, M.:
Pitfalls in the management of congenital dislocation of the
hip.
J. Bone Joint Surg., 48 (Br.) : 666-681 (1966).

58. Mackenzie, I.G.; Seddon, H.J.; and Trevor, D.:
Congenital dislocation of the hip.
J. Bone Joint Surg., 42 (Br.) : 689-705 (1960).

59. Marchetti, P.S.:
Open reduction of congenital dislocation of the hip. In :
Tachdjian, M.O. (Ed.): Congenital Dislocation of the Hip.
New York, Churchill Livingstone, p. 401-415 (1982).
60. Massie, W.K.; and Howorth, M.B.:
Congenital dislocation of the hip. III. Results of open
reduction as seen in early adult period.
J. Bone Joint Surg., 33 (Am.) : 171-198 (1951).
61. McCarrol, H.R.; and Crego, C.H.Jr.:
Primary anterior congenital dislocation of the hip.
J. Bone Joint Surg., 21 : 648-664 (1939).
62. McKay, W.D.:
A comparison of the innominate and pericapsular osteotomy
in the treatment of congenital dislocation of the hip.
Clin. Orthop., 98 : 124-132 (1974).
63. McKibbin, B.:
Anatomical factors in the stability of the hip joint in the
newborn.
J. Bone Joint Surg., 52 (Br.) : 148-159 (1970).
64. Melzer, C.; and Wulker, W.:
Potential mistakes in hip joint sonography.
Arch. Orthop. Trauma Surg., 109 : 126-130 (1990).
65. Melzer, C.; Sturz, H.; and Sniezynski, R.:
Langzeitergebnisse nach Offener Einstellung der
Sogenannten Angeborenen Huftluxation.
Orthop. Praxis, 6 : 361-365 (1991).

66. Metaizeau, J.P.; and Prevot, J.:
Le retentissement des ostéotomies innominées sur le cotyle
contralatéral.
Rev. Chir. Orthop., 66 : 453 (1980).
67. Millis, M.B.; and Murphy, S.B.:
Use of computed tomographic reconstruction in planning
osteotomies of the hip.
Clin. Orthop., 274 : 154-159 (1992).
67. Millis, M.B.; and Share, J.C.:
Use of ultrasonography in dysplasia of the immature hip.
Clin. Orthop., 274 : 160-171 (1992).
68. Monticelli, G.M.:
Intertrochanteric femoral osteotomy with concentric
reduction of the femoral head in treatment of residual
acetabular dysplasia.
Clin. Orthop., 119 : 48-59 (1976).
69. Moore, F.H.:
Examining infant's hips - Can it do harm ?
J. Bone Joint Surg., 71 (Br.) : 4-5 (1989).
70. Morel, G.:
The treatment of congenital dislocation and subluxation of
the hip in the older child.
Acta Orthop. Scand., 46 : 364-399 (1975).
71. Morrissey, R.T.; and Cowie, G.H.:
Congenital dislocation of the hip: Early detection and
prevention of late complications.
Clin. Orthop., 222 : 79-84 (1987).

72. Morscher, E.:
Our experience with Salter's innominate osteotomy in the treatment of hip dysplasia. In : Weil, U.H. (Ed.): Progress in Orthopaedic Surgery. Berlin, Springer, vol. 2, p. 107-119 (1978).
73. Mose, K.:
Methods of measuring in Legg-Calve Perthes disease with special regard to prognosis.
Clin. Orthop., 150 : 103-109 (1980).
74. Moulin, P.; and Morscher, E.:
Langzeitresultate der Becken-Osteotomie nach Salter.
Orthopade, 17 : 479-484 (1988).
75. Moulton, A.; and Upadhyay, S.S.:
A direct method of measuring femoral anteversion using ultrasound.
J. Bone Joint Surg., 64 (Br.) : 469-472 (1982).
76. Muller, G.M.; and Seddon, H.J.:
Late results of treatment of congenital dislocation of the hip.
J. Bone Joint Surg., 35 (Br.) : 242-262 (1953).
77. Murphy, S.B.; Krjewski, P.K.; Millis, M.B.; and Harless, A.:
Acetabular dysplasia in the adolescent and young adult.
Clin. Orthop., 261 : 214-223 (1990).

78. Ogden, J.A.:
Changing patterns of proximal femoral vascularity.
J. Bone Joint Surg., 56 (Am.) : 941-950 (1974).
79. O'Hara, J.N.:
Congenital dislocation of the hip: Acetabular deficiency
(absence of the lateral acetabular epiphysis) after
limbectomy in infancy.
J. Pediatr. Orthop., 9 : 640-648 (1989).
80. O'Hara, J.N.; Bernard, A.A.; and Dwyer, N.:
Congenital dislocation of the hip: Open reduction by the
medial approach in infancy.
J. Bone Joint Surg., 68 (Br.) : 668 (1986).
81. O'Hara, J.N.; Bernard, A.A.; and Dwyer, N.:
Early results of medial approach, open reduction of
congenital dislocation of the hip: Use before walking.
J. Pediatr. Orthop., 8 : 288-294 (1988).
82. Paterson, D.C.:
Innominate osteotomy: Its role in the treatment of congenital
dislocation and subluxation of the hip joint.
Clin. Orthop., 98 : 198-209 (1974).
83. Paterson, D.C.:
The early diagnosis and treatment of congenital dislocation
of the hip.
Clin. Orthop., 119 : 28-38 (1976).
84. Pemberton, P.A.:
Pericapsular osteotomy of the ilium for treatment of
congenital subluxation and dislocation of the hip.
J. Bone Joint Surg., 47 (Am.), 65-86 (1965).

85. Peterson, H.A.; Klassen, R.A.; McLeod, R.A.; and Hoffman, A.D.:
The use of computerized tomography in dislocation of the hip and femoral neck anteversion in children.
J. Bone Joint Surg., 63 (Br.) : 198-208 (1981).

86. Ponseti, I.V.:
Morphology of the acetabulum in congenital dislocation of the hip: Gross, histological and roentgenographic studies.
J. Bone Joint Surg., 60 (Am.) : 586-599 (1978).

87. Rab, G.T.:
Biomechanical aspects of Salter osteotomy. In: Proceedings of the Fourth Open Scientific Meeting of the Hip Society. The C.V. Mosby Co., St. Louis, p. 67-74 (1976).

88. Rab, G.T.:
Biomechanical aspects of Salter osteotomy.
Clin. Orthop., 132 : 82-88 (1978).

89. Rab, G.T.:
Preoperative roentgenographic evaluation for osteotomies about the hip in children.
J. Bone Joint Surg., 63 (Am.) : 306-309 (1981).

90. Radin, E.L.; and Paul, I.L.:
The biomechanics of congenital dislocated hips and their treatment.
Clin. Orthop., 98 : 32-38 (1974).

91. Reynolds, D.A.:
Chiari innominate osteotomy in adults: Technique indications and contraindications.
J. Bone Joint Surg., 43 (Br.) : 45-54 (1986).

92. Roth, A.; Gibson, D.A.; and Hall, J.E.:
The experience of five orthopaedic surgeons with innominate osteotomy in the treatment of congenital dislocation and subluxation of the hip.
Clin. Orthop., 98 : 178-182 (1974).
93. Rungee, J.L.; and Reinker, K.A.:
Ossific nucleus eccentricity in congenital dislocation of the hip.
J. Pediatr. Orthop., 12 : 61-67 (1992).
94. Salter, R.B.:
Innominate osteotomy in the treatment of congenital dislocation of the hip.
J. Bone Joint Surg., 43 (Br.) : 518-539 (1961).
95. Salter, R.B.:
Role of innominate osteotomy in the treatment of congenital dislocation and subluxation of the hip in the older child.
J. Bone Joint Surg., 48 (Am.) : 1413-1438 (1966).
96. Salter, R.B.:
Specific guidelines in the application of the principle of innominate osteotomy.
Orthop. Clin. N. Amer., 3 : 149-156 (1972).
97. Salter, R.B.; and Dubos, J.P.:
The first fifteen years personal experience with innominate osteotomy in the treatment of congenital dislocation and subluxation of the hip.
Clin. Orthop., 98 : 72-103 (1974).

98. Salter, R.B.; Hansson, G.; and Thompson, G.H.:
Innominate osteotomy in the management of residual
congenital subluxation of the hip in young adults.
Clin. Orthop., 182 : 53-68 (1984).
99. Salter, R.B.; Kostuik, J.; and Dallas, S.:
Avascular necrosis of the femoral head as a complication of
treatment for congenital dislocation of the hip in young
children: A clinical and experimental investigation.
Can. J. Surg., 12 : 44-60 (1969).
100. Schuler, P.; Feltes, E.; Kienapfel, H.; and Griss, P.:
Ultrasound examination for the early determination of dysplasia
and congenital dislocation of neonatal hips.
Clin. Orthop., 258 : 18-25 (1990).
101. Schwartz, D.B.:
Acetabular development after reduction of congenital
dislocation of the hip.
J. Bone Joint Surg., 47 (Am.) : 705-714 (1965).
102. Serafimov, L.:
Biomechanical influence of the innominate osteotomy on
the growth of the upper part of the femur.
Clin. Orthop., 98 : 39-40 (1974).
103. Severin, E.:
Contribution to the knowledge of congenital dislocation of
the hip joint.
Acta Orthop. Scand., 84 (Suppl. 63) : 1-141 (1941).

104. Smith, W.S.; Badgley, C.E.; Orwig, J.B.; and Harper, J.M.:
Correlation of postreduction roentgenograms and 31-year follow-up in congenital dislocation of the hip.
J. Bone Joint Surg., 50 (Am.) : 1081-1097 (1968).
105. Staheli, L.T.:
Surgical management of acetabular dysplasia.
Clin. Orthop., 264 : 111-121 (1991).
106. Steel, H.H.:
Triple osteotomy of the innominate bone.
J. Bone Joint Surg., 55 (Am.) : 343-350 (1973).
107. Suzuki, S.; Kashara, Y.; Futami, T.; Ushikubo, S.; and Tsuchiya, T.:
Ultrasonography in congenital dislocation of the hip.
J. Bone Joint Surg., 73 (Br.) : 879-883 (1991).
108. Tachdjian, M.O.:
Salter's innominate osteotomy to derotate the maldirected acetabulum. In : Tachdjian, M.O. (Ed.): Congenital Dislocation of the Hip. New York, Churchill Livingstone, p. 525-541 (1982).
109. Tachdjian, M.O.:
Treatment of hip dysplasia in the older child and adolescent: Factors in decision making. In : Tachdjian, M.O. (Ed.): Congenital Dislocation of the Hip. New York, Churchill Livingstone, p. 625-46 (1982).

110. Teuffer, A.P.; and Noguera, J.G.:
Experience with innominate osteotomy (Salter) and medial displacement osteotomy (Chiari) in the treatment of acetabular dysplasia.
Clin. Orthop., 98 : 133-136 (1974).
111. Thomas, I.H.; Dunnin, A.J.; Cole, S.; and Menelaus, M.B.:
Avascular necrosis after open reduction for congenital dislocation of the hip.
J. Pediatr. Orthop., 9 : 525-531 (1989).
112. Tonnis, D.:
Normal values of the hip joint or the evaluation of X-rays in children and adults.
Clin. Orthop., 119 : 39-47 (1976).
113. Tonnis, D.:
An evaluation of conservative and operative methods in the treatment of congenital hip dislocation.
Clin. Orthop., 119 : 76-88 (1976).
114. Tonnis, D.:
Die angeborene Huft dysplasie und Huftluxation im Kindes und Erwachsenenalter. Springer, Berlin, Heidelberg, New York, Tokyo (1984).
115. Tonnis, D.:
Surgical treatment of congenital dislocation of the hip.
Clin. Orthop., 268 : 33-39 (1990).

116. Tonnis, D.; Behrens, K.; and Tascharani, F.:
Eine neue Technik der Dreifachosteotomie Schwenkung
dysplastischer Huftpfnen bei Jugendlichen und
Erwachsenen.
Z. Orthop., 119 : 253-263 (1981).

117. Trevor, D.:
Treatment of congenital hip dislocation in older children.
Proc. Roy. Soc. Med., 54 : 481-490 (1960).

118. Utterback, T.D.; and MacEwen, G.D.:
Comparison of pelvic osteotomies for the surgical correction
of the congenital hip.
Clin. Orthop., 98 : 104-110 (1974).

120. Viser, J.D.; Jonkers, A.; and Hillen, B.:
Hip joint measurements with computerized tomography.
J. Pediatr. Orthop., 2 (2) : 143-146 (1982).

121. Waters, P.; Kurica, K.; Hall, J.; and Micheli, L.J.:
Salter innominate osteotomies in congenital dislocation of
the hip.
J. Pediatr. Orthop., 8 : 650-655 (1988).

122. Wedge, J.H.; and Salter, R.B.:
Innominate osteotomy: Its role in the arrest of secondary
degenerative arthritis of the hip in the adults.
Clin. Orthop., 98 : 214-224 (1974).

123. Weiner, D.S.; Hoyt, W.A.Jr.; and O'Dell, H.W.:
Congenital dislocation of the hip: The relationship of
premanipulation traction and age to avascular necrosis of
the femoral head.
J. Bone Joint Surg., 59 (Am.): 306-311 (1977).

124. Weinstein, S.L.:
Natural history of congenital hip dislocation and hip dysplasia..
Clin. Orthop., 225 : 62-86 (1987).
125. Weintroub, S.; Green, I.; Terdiman, R.; and Weissmann, S.L.:
Growth and development of congenitally dislocated hips reduced in early infancy.
J. Bone Joint Surg., 61 (Am.) : 125-130 (1979).
126. Westin, G.W.; Ilfeld, F.W.; and Provost, J.:
Total avascular necrosis of the capital femoral epiphysis in congenitally dislocated hips.
Clin. Orthop., 119 : 93-98 (1976).
127. Wilkinson, J.A.:
Prime factors in the aetiology of congenital dislocation of the hip.
J. Bone Joint Surg., 45 (Br.) : 268-283 (1963).
128. Wong-Chung, J.; Ryan, M.; and O'Brien, T.M.:
Movement of the femoral head after Salter osteotomy for acetabular dysplasia.
J. Bone Joint Surg., 72 (Br.) : 563-567 (1990).
129. Yamamuro, T.:
Personal communication (1992).

130. Yamamuro, T.; and Ishida, K.:
Recent advances in the prevention, early diagnosis and
treatment of congenital dislocation of the hip in Japan.
Clin. Orthop., 184 : 34-40 (1984).

131. Zions, L.E.; and MacEwen, G.D.:
Treatment of congenital dislocation of the hip in children
between the ages of one and three years.
J. Bone Joint Surg., 68 (Am.) : 829-846 (1986).